American Samoa Model Checks

2022-08-29

This is a summary report for the VALO base model run.

Model Output

Input Data





Convergence Check

```
  Converged     MaxGrad
1      TRUE 1.39802e-05
```

```
[1] "1 NOTE:  Max data length bin: 51  < max pop len bins: 57; so will accumulate larger pop
[2] "N warnings: 1"
```

Fit to Model

CPUE

```
Residual Runs Test (/w plot) stats by Index:
```

```
RMSE stats by Index:
```

Length Comp


Index
0.0
0.1
0.2
0.3
0.4
0.5
0.6
2016
2018
2020
Year


FISHERY
Index residuals
−0.6
−0.2
0.2
0.4
0.6
2016
2018
2020
Year


RMSE = 67.3%
Index residuals
−0.4
−0.2
0.0
0.2
0.4
FISHERY
Loess
2017
2018
2019
2020
Year

#Factor	Fleet	New_Var_adj	Type	Name
4	1	0.103658	len	FISHERY

```
Residual Runs Test (/w plot) stats by Mean length:

    Index runs.p   test  sigma3.lo sigma3.hi type
1 FISHERY    0.5 Passed -0.2509701 0.2509701  len

RMSE stats by Index:

# A tibble: 2 x 3
  Fleet    RMSE.perc  Nobs
  <chr>        <dbl> <int>
1 FISHERY        7.9     6
2 Combined       7.9     6
```



Retrospective and Hindcasting


FISHERY (whole catch)
Mean length
35
30
25
2011
2012
2013
2014
2015
2016
Year


FISHERY
Sum of N input=368.8
Sum of N adj.=260.5
Proportion
0.20
0.15
0.10
0.05
0.00
20
30
40
50
Length (cm)



Retrospective

`Mohn's Rho stats, including one step ahead forecasts:`



```
Mohn's Rho stats, including one step ahead forecasts:

  type     peel         Rho ForecastRho
1    F     2020 -0.02117726 -0.02091191
2    F     2019 -0.07413450 -0.07393367
3    F     2018  0.00000000  0.00000000
4    F Combined -0.03177059 -0.03161519
```

Hindcasting

```
Plotting Hindcast Cross-Validation (one-step-ahead)

 Computing MASE with only 2 of 3  prediction residuals for Index FISHERY

 Warning:  Unequal spacing of naive predictions residuals may influence the interpretation o
```




```
MASE stats by Index:
Plotting Hindcast Cross-Validation (one-step-ahead)

No observations in evaluation years to compute prediction residuals for Index FISHERY
```

MASE stats by Index:

Recruitment Deviations

Likelihood Profile

```
[1] "SR_LN"
                     frac_change include                          label
TOTAL                     1.0000    TRUE                          Total
Catch                     0.0636    TRUE                          Catch
Equil_catch               0.0000   FALSE                Equilibrium catch
Survey                    0.0318    TRUE                       Index data
Length_comp               0.9678    TRUE                      Length data
Recruitment               0.0000   FALSE                      Recruitment
InitEQ_Regime             0.0000   FALSE Initital equilibrium recruitment
Forecast_Recruitment      0.0000   FALSE             Forecast recruitment
Parm_priors               0.0052   FALSE                           Priors
Parm_softbounds           0.0001   FALSE                      Soft bounds
Parm_devs                 0.0000   FALSE             Parameter deviations
Crash_Pen                 0.0000   FALSE                    Crash penalty
```





Management Quantities

```
starter.sso with Bratio: SSB/SSBMSY and F: _abs_F
```


0%
0%
0%
100%
F/F_MSY
SSB/SSB_MSY
0.0
0.5
1.0
1.5
2.0
0
1
2
3
4
5


MVLN
SSB/SSB_MSY
Year
0
1
2
3
4
1970
1980
1990
2000
2010
2020


MVLN
F/F_MSY
0.0
0.5
1.0
1.5
2.0
1970
1980
1990
2000
2010
2020
Year


MVLN
SSB (t)
0
2
4
6
8
12
1970
1980
1990
2000
2010
2020
Year




```
null device
          1
```

Jitter


Likelihood
138.375
138.350
138.325
138.300
1.0
1.5
2.0
2.5
3.0
Run


Spawning biomass (t)
10
8
6
4
2
0
1970
1980
1990
2000
2010
2020
Year
model 1
model 2
model 3


abs F; with F=sum(full Fs)
0.0
0.1
0.2
0.3
0.4
0.5
1970
1980
1990
2000
2010
2020
Year
model 1
model 2
model 3


Age−0 recruits (1,000s)
0
5
10
15
20
25
30
1970
1980
1990
2000
2010
2020
Year
model 1
model 2
model 3