Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Sat Jul 23 09:15:50 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
0
10
20
30
40
0
1
2
3
4
5
6
Age (yr)


Age (yr)
0
1
2
3
4
5
6
CV of lengths
0
0.02
0.06
Length (cm, beginning of the year)
40
30
20
10
0
0.0
0.5
1.0
1.5
2.0
SD of lengths
CV of lengths
0.08
0.04
0
2.0
1.0
0.0
Age (yr)


Age (yr)
0
1
2
3
4
5
6
Length (cm, beginning of the year)
0
10
20
30
40
Mean weight
0
0.5
1
1.5
Frac. mature
0.0
0.5
1.0
Mean weight
0
0.5
1
1.5
Age (yr)
0
1
2
3
4
5
6
Frac. mature
0.0
0.5
1.0


Length
0
10
20
30
40
0
1
2
3
4
5
6
Age


Length
0
10
20
30
40
0
1
2
3
4
5
6
Age


Mean weight (kg) in last year
0.0
0.5
1.0
1.5
0
10
20
30
40
Length (cm)


1.0
0.8
0.6
0.4
0.2
0.0
Maturity
0
10
20
30
40
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0.0
0.5
1.0
1.5
Female weight (kg)


Fecundity
0.0
0.5
1.0
1.5
0.0
0.5
1.0
1.5
Female weight (kg)


2.0
1.5
1.0
0.5
0.0
Fecundity
0
10
20
30
40
Female length (cm)


Spawning output
0.0
0.5
1.0
1.5
0
10
20
30
40
Length (cm)


Spawning output
0.00
0.05
0.10
0.15
0.20
0.25
0
1
2
3
4
5
6
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
Length (cm)
FISHERY
SURVEY2


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
1
2
3
4
5
6
Age (yr)
FISHERY
SURVEY2


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
Length (cm)


Length (cm)
0
10
20
30
40
1
0.2
0.4
0.6
0
1
2
3
4
5
6
Age (yr)


Length (cm)
0
10
20
30
40
1
0.2
0.4
0.6
0
1
2
3
4
5
6
Age (yr)


Spawning biomass (mt)
0
2
4
6
8
10
12
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
0.5
1.0
1.5
2.0
2.5
1970
1980
1990
2000
2010
2020
Year


Total biomass (mt)
0
5
10
15
20
25
1970
1980
1990
2000
2010
2020
Year


Summary biomass (mt)
0
5
10
15
20
25
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
50
150
250
350
1970
1980
1990
2000
2010
2020
Year


Summary Fishing Mortality
0
1
2
3
4
5
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
2
4
6
8
10
12
14
1970
1980
1990
2000
2010
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
0.4
0.2
0.0
−0.2
−0.6
1970
1980
1990
2000
2010
2020
Year


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
0
50
100
200
300
0
1
2
3
4
5
6
Spawning biomass (mt)


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
0
50
100
200
300
0
1
2
3
4
5
6
Spawning biomass (mt)
2021
1987
2015


Log recruitment deviation
0.6
0.4
0.2
0.0
−0.2
−0.4
−0.6
0.0
0.2
0.4
0.6
0.8
1.0
Spawning output (relative to B_0)
2021
1967
2015


Landings (mt)
0
2
4
6
8
10
1970
1980
1990
2000
2010
2020
Year


Landings (mt)
0
2
4
6
8
10
1970
1980
1990
2000
2010
2020
Year


FISHERY
FISHERY obs.
Observed and expected Landings (mt)
0
2
4
6
8
10
12
1970
1980
1990
2000
2010
2020
Year


Continuous F
0
1
2
3
4
5
1970
1980
1990
2000
2010
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0.0
0.5
1.0
1.5
2.0
2.5
Relative spawning output: B/B_MSY
2000
1967
2021


Index
0
10
20
30
40
50
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
10
20
30
40
50
1990
1995
2000
2005
2010
2015
2020
Year


Expected index
0
2
4
6
8
0
5
10
15
20
25
30
Observed index


Log index
0
1
2
3
4
1990
1995
2000
2005
2010
2015
2020
Year


Log index
0
1
2
3
4
1990
1995
2000
2005
2010
2015
2020
Year


Log expected index
2.2
2.0
1.8
1.6
1.4
1.2
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
Log observed index


Residual
10
5
0
−5
−10
1990
1995
2000
2005
2010
2015
2020
Year


Deviation
2
1
0
−1
−2
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
10
20
30
40
50
1990
1995
2000
2005
Year


Index
0
10
20
30
40
50
1990
1995
2000
2005
Year


Expected index
0
2
4
6
8
0
5
10
15
20
25
Observed index


Log index
4
2
0
−2
−4
1990
1995
2000
2005
Year


Log index
4
2
0
−2
−4
1990
1995
2000
2005
Year


Log expected index
1.2
1.4
1.6
1.8
2.0
−1
0
1
2
3
Log observed index


Residual
6
4
2
0
−2
−4
−6
1990
1995
2000
2005
Year


Deviation
3
2
1
0
−1
−2
−3
1990
1995
2000
2005
Year


FISHERY
SURVEY2
Standardized index
Year
0
1
2
3
4
1990
1995
2000
2005
2010
2015
2020


50
150
250
350
Age
0
1
2
3
4
5
6
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


50
150
250
Age
0
1
2
3
4
5
6
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


50
150
250
350
Length
0
10
20
30
40
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


20
60
100
40
30
20
10
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


Numbers at age at equilibrium
0
50
100
150
200
250
0
1
2
3
4
5
6
Age


FISHERY
Sum of N adj.=4510.8
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
5
10
15
20
25
30
35
40
Length (cm)


FISHERY
0.01
0.2
0.4
Length (cm)
40
35
30
25
20
15
10
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


2004
N adj.=247.8
2010
N adj.=154.6
2016
N adj.=352.8
2005
N adj.=157.9
2011
N adj.=269.6
2017
N adj.=169.7
2006
N adj.=125.2
2012
N adj.=56.3
2018
N adj.=163.8
2007
N adj.=275.5
2013
N adj.=172.2
2019
N adj.=220.9
2008
N adj.=297.4
2014
N adj.=552.7
2020
N adj.=167.2
2009
N adj.=472.1
2015
N adj.=563.6
2021
N adj.=91.6
Proportion
0.00
0.10
0.20
0.30
5
10
20
30
40
Length (cm)


0.01
0.2
0.4
Length (cm)
40
35
30
25
20
15
10
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


FISHERY (whole catch)
23.0
22.5
22.0
21.5
21.0
20.5
Mean length
2005
2010
2015
2020
Year


FISHERY
Sum of N adj.=4510.8
Sum of N eff.=2060.1
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
5
10
15
20
25
30
35
40
Length (cm)


FISHERY
−500
−300
100
300
500
40
35
30
25
20
15
10
Length (cm)
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


2004
N adj.=247.8
N eff.=165
2010
N adj.=154.6
N eff.=38.1
2016
N adj.=352.8
N eff.=101.2
2005
N adj.=157.9
N eff.=92.9
2011
N adj.=269.6
N eff.=231.3
2017
N adj.=169.7
N eff.=35.1
2006
N adj.=125.2
N eff.=75.8
2012
N adj.=56.3
N eff.=34.8
2018
N adj.=163.8
N eff.=55.3
2007
N adj.=275.5
N eff.=173.5
2013
N adj.=172.2
N eff.=215.5
2019
N adj.=220.9
N eff.=94.4
2008
N adj.=297.4
N eff.=105
2014
N adj.=552.7
N eff.=204.4
2020
N adj.=167.2
N eff.=28.8
2009
N adj.=472.1
N eff.=88.8
2015
N adj.=563.6
N eff.=304.6
2021
N adj.=91.6
N eff.=15.6
Proportion
0.00
0.10
0.20
0.30
5
10
20
30
40
Length (cm)


−500
−300
100
300
500
40
35
30
25
20
15
10
Length (cm)
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


Effective sample size
Observed sample size
0
50
150
250
350
0
100
200
300
400
500
600
2015
2014
2011
2013
2007
2004
2008
2016
2009
2005
2019
2006
2018
2012
2021
2020
harmonic mean
arithmetic mean


FISHERY (whole catch)
Mean length
23.0
22.5
22.0
21.5
21.0
20.5
2005
2010
2015
2020
Year


Equilibrium yield (mt)
0
2
4
6
8
10
14
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0
2
4
6
8
10
14
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
8
6
4
2
0
−4
0
5
10
15
20
25
Total biomass (mt)


Yield per recruit (kg)
0.000
0.010
0.020
0.030
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
SURVEY2
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


Catches
FISHERY
Abundance indices
FISHERY
SURVEY2
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


SR_LN(R0)
prior
max. likelihood
5
10
15
20
Size_inflection_FISHERY(1)
18
20
22
24
26
28
InitF_seas_1_flt_1FISHERY
0
2
4
6
8
10
Size_95%width_FISHERY(1)
1
2
3
4
5
6
LnQ_base_FISHERY(1)
−10
−5
0
5
10
LnQ_base_SURVEY2(2)
−10
−5
0
5
10
Density
Parameter value