Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Tue Aug 02 09:54:10 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
0
20
40
60
0
2
4
6
8
10
12
Age (yr)


Age (yr)
0
2
4
6
8
10
12
Length (cm, beginning of the year)
0
20
40
60
CV of lengths
0
0.02
0.04
0.06
SD of lengths
0
1
2
3
4
CV of lengths
0
0.02
0.04
0.06
SD of lengths
0
2
4
Age (yr)
0
2
4
6
8
10
12


Age (yr)
0
2
4
6
8
10
12
Length (cm, beginning of the year)
0
20
40
60
Mean weight
0
2
4
6
8
0.0
0.5
1.0
Frac. mature
Mean weight
0
4
8
Frac. mature
0.0
0.5
1.0
Age (yr)


80
60
40
20
0
Length
0
2
4
6
8
10
12
Age


80
60
40
20
0
Length
0
2
4
6
8
10
12
Age


Mean weight (kg) in last year
0
2
4
6
8
0
10
20
30
40
50
60
70
Length (cm)


1.0
0.8
0.6
0.4
0.2
0.0
Maturity
0
10
20
30
40
50
60
70
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0
2
4
6
8
Female weight (kg)


Fecundity
0
2
4
6
8
0
2
4
6
8
Female weight (kg)


Fecundity
0
2
4
6
8
10
0
10
20
30
40
50
60
70
Female length (cm)


Spawning output
0
2
4
6
8
0
10
20
30
40
50
60
70
Length (cm)


Spawning output
0
2
4
6
8
0
2
4
6
8
10
12
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
50
60
70
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
2
4
6
8
10
12
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
50
60
70
Length (cm)


Length (cm)
0
10
20
30
40
50
60
70
0.8
0.6
0.4
0.2
0
2
4
6
8
10
12
Age (yr)


Spawning biomass (mt)
0.0
0.5
1.0
1.5
1990
2000
2010
2020
Year


Spawning biomass (mt)
0.0
0.5
1.0
1.5
2.0
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
0.5
1.0
1.5
2.0
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
1.0
2.0
3.0
1990
2000
2010
2020
Year


Total biomass (mt)
0
1
2
3
4
5
1990
2000
2010
2020
Year


Summary biomass (mt)
0
1
2
3
4
5
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0.0
0.5
1.0
1.5
2.0
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
1
2
3
4
5
1990
2000
2010
2020
Year


Summary Fishing Mortality
0
1
2
3
4
5
1985
1990
1995
2000
2005
2010
2015
2020
Year


Spawning biomass (mt)
0.0
0.5
1.0
1.5
2.0
2.5
1985
1990
1995
2000
2005
2010
2015
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
−0.4
−0.2
0.0
0.2
0.4
1980
1990
2000
2010
2020
Year


Log recruitment deviation
1.0
0.5
0.0
−0.5
−1.0
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.1
0.2
0.3
0.4
0.5
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
1.0
0.5
0.0
$1 - SE(\hat{r}_y)^2/\sigma_R^2$
1980
1990
2000
2010
2020
Year


Recruitment (1,000s)
0.0
0.5
1.0
1.5
2.0
0.0
0.1
0.2
0.3
0.4
0.5
0.6
Spawning biomass (mt)
Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium


Recruitment (1,000s)
0.0
0.5
1.0
1.5
2.0
2006
1986
2021
Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
0.0
0.1
0.2
0.3
0.4
0.5
0.6
Spawning biomass (mt)


2006
1986
2021
0.4
0.2
0.0
−0.2
−0.4
Log recruitment deviation
0.0
0.1
0.2
0.3
0.4
0.5
0.6
Spawning output (relative to B_0)


Landings (mt)
0.0
0.5
1.0
1.5
1985
1990
1995
2000
2005
2010
2015
2020
Year


Landings (mt)
0.0
0.5
1.0
1.5
1985
1990
1995
2000
2005
2010
2015
2020
Year


FISHERY
FISHERY obs.
Observed and expected Landings (mt)
0.0
0.5
1.0
1.5
2.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


Continuous F
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0.0
0.5
1.0
1.5
2.0
Relative spawning output: B/B_MSY
1986
1990
2000
2010
2020
2021


Index
0
5
10
15
20
25
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
5
10
15
20
25
1990
1995
2000
2005
2010
2015
2020
Year


Expected index
Observed index
0
1
2
3
4
5
6
7
0
2
4
6
8
10


Log index
3
2
1
0
−1
1990
1995
2000
2005
2010
2015
2020
Year


Log index
3
2
1
0
−1
1990
1995
2000
2005
2010
2015
2020
Year


Log expected index
0.0
0.5
1.0
1.5
Log observed index
0.0
0.5
1.0
1.5
2.0


Effective catchability
4.989500
4.989490
4.989480
0.2
0.4
0.6
0.8
1.0
1.2
1.4
Vulnerable biomass


Residual
2
1
0
−1
−2
1990
1995
2000
2005
2010
2015
2020
Year


Deviation
1.0
0.5
0.0
−0.5
−1.0
1990
1995
2000
2005
2010
2015
2020
Year


FISHERY
Standardized index
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1990
1995
2000
2005
2010
2015
2020
Year


0.1
2
Age
12
11
10
9
8
7
6
5
4
3
2
1
0
1985
1989
1993
1997
2001
2005
2009
2013
2017
2021
Year


0.1
2
Age
12
11
10
9
8
7
6
5
4
3
2
1
0
1985
1989
1993
1997
2001
2005
2009
2013
2017
2021
Year


80
60
40
20
0
Length
0.1
2
1986
1990
1994
1998
2002
2006
2010
2014
2018
2022
Year


100
300
500
700
Length
0
20
40
60
80
1986
1990
1994
1998
2002
2006
2010
2014
2018
2022
Year


Numbers at age at equilibrium
0.0
0.5
1.0
1.5
0
2
4
6
8
10
12
Age


FISHERY
Sum of N adj.=457.8
Proportion
0.30
0.25
0.20
0.15
0.10
0.05
0.00
10
20
30
40
50
60
70
Length (cm)


FISHERY
0.01
0.4
Length (cm)
65
60
55
50
45
40
35
30
25
20
15
2008
2011
2013
2015
2017
2019
Year


2007
N adj.=54.6
2014
N adj.=37.1
2008
N adj.=21.2
2015
N adj.=59.6
2009
N adj.=50.9
2016
N adj.=29.4
2011
N adj.=31.9
2017
N adj.=33.6
2012
N adj.=72.2
2019
N adj.=31.1
2013
N adj.=36.1
Proportion
Length (cm)


0.01
0.4
65
60
55
50
45
40
35
30
25
20
15
Length (cm)
2008
2011
2013
2015
2017
2019
Year


FISHERY (whole catch)
Mean length
50
45
40
35
2008
2010
2012
2014
2016
2018
Year


FISHERY
Sum of N adj.=457.8
Sum of N eff.=403.4
Proportion
0.30
0.25
0.20
0.15
0.10
0.05
0.00
10
20
30
40
50
60
70
Length (cm)


FISHERY
−14
−10
−6
2
6
10
14
Length (cm)
65
60
55
50
45
40
35
30
25
20
15
2008
2011
2013
2015
2017
2019
Year


2007
N adj.=54.6
N eff.=58.4
2014
N adj.=37.1
N eff.=35.1
2008
N adj.=21.2
N eff.=31.1
2015
N adj.=59.6
N eff.=77.4
2009
N adj.=50.9
N eff.=34.4
2016
N adj.=29.4
N eff.=11.7
2011
N adj.=31.9
N eff.=27.7
2017
N adj.=33.6
N eff.=54.5
2012
N adj.=72.2
N eff.=32.5
2019
N adj.=31.1
N eff.=19.2
2013
N adj.=36.1
N eff.=21.4
Proportion
0.0
0.1
0.2
0.3
0.4
10
20
30
40
50
60
70
Length (cm)


−14
−10
−6
2
6
10
14
65
60
55
50
45
40
35
30
25
20
15
Length (cm)
2008
2011
2013
2015
2017
2019
Year


Effective sample size
Observed sample size
0
20
40
60
80
harmonic mean
arithmetic mean
2015
2007
2017
2014
2009
2012
2008
2011
2013
2019
2016


FISHERY (whole catch)
Mean length
50
45
40
35
2008
2010
2012
2014
2016
2018
Year


Equilibrium yield (mt)
0.0
0.2
0.4
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0.0
0.2
0.4
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current

Surplus production (mt)
0.0
0.4
0.8
1.2
0
1
2
3
4
Total biomass (mt)


Yield per recruit (kg)
0.0
0.1
0.2
0.3
0.4
0.5
1985
1990
1995
2000
2005
2010
2015
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1985
1990
1995
2000
2005
2010
2015
2020
Year


Catches
Abundance indices
Length compositions
FISHERY
FISHERY
FISHERY
1985
1990
1995
2000
2005
2010
2015
2020
Year


SR_LN(R0)
prior
max. likelihood
2
4
6
8
10
12
14
Size_inflection_FISHERY(1)
30
35
40
45
50
55
60
InitF_seas_1_flt_1FISHERY
0.0
0.1
0.2
0.3
0.4
0.5
Size_95%width_FISHERY(1)
5
10
15
20
25
LnQ_base_FISHERY(1)
−4
−2
0
2
4
Q_extraSD_FISHERY(1)
0.0
0.5
1.0
1.5
2.0
Density
Parameter value