Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Thu Jun 30 11:09:08 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
120
100
80
60
40
20
0
0
5
10
15
20
25
30
Age (yr)


Age (yr)
0
5
10
15
20
25
30
Length (cm, beginning of the year)
120
100
80
60
40
20
0
CV of lengths
0.12
0.08
0.04
0
SD of lengths
8
4
0
CV of lengths
0
0.04
0.08
0.12
SD of lengths
0
2
4
6
8
Length (cm, beginning of the year)


Age (yr)
Mean weight
Length (cm, beginning of the year)
Mean weight
Frac. mature
Age (yr)
Frac. mature
Length (cm, beginning of the year)


120
100
80
60
40
20
0
Length
0
5
10
15
20
25
30
Age


120
100
80
60
40
20
0
Length
0
5
10
15
20
25
30
Age


Mean weight (kg) in last year
0
5
10
15
20
25
0
20
40
60
80
100
Length (cm)


1.0
0.8
0.6
0.4
0.2
0.0
Maturity
0
20
40
60
80
100
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0
5
10
15
20
25
Female weight (kg)


Fecundity
0
5
10
15
20
25
0
5
10
15
20
25
Female weight (kg)


Fecundity
30
25
20
15
10
5
0
0
20
40
60
80
100
Female length (cm)


Spawning output
0
5
10
15
20
25
0
20
40
60
80
100
Length (cm)


Spawning output
0
2
4
6
8
10
12
0
5
10
15
20
25
30
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
100
Length (cm)
FISHERY
SURVEY2


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
5
10
15
20
25
30
Age (yr)
FISHERY
SURVEY2


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
100
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
100
Length (cm)


Length (cm)
0
20
40
60
80
100
1
0.8
0.6
0.4
0.2
0
5
10
15
20
25
30
Age (yr)


Length (cm)
0
20
40
60
80
100
1
0.8
0.6
0.4
0.2
0
5
10
15
20
25
30
Age (yr)


Spawning biomass (mt)
0
2
4
6
8
10
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
5
10
15
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
1.0
2.0
3.0
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Total biomass (mt)
0
5
10
15
20
25
1970
1980
1990
2000
2010
2020
Year


Summary biomass (mt)
0
5
10
15
20
25
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
1
2
3
4
5
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
5
10
15
1970
1980
1990
2000
2010
2020
Year


Summary Fishing Mortality
0.0
0.2
0.4
0.6
0.8
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
2
4
6
8
10
12
1970
1980
1990
2000
2010
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
0.6
0.4
0.2
−0.2
−0.6
1970
1980
1990
2000
2010
2020
Year


Log recruitment deviation
−1.0
0.0
0.5
1.0
1.5
1970
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.1
0.2
0.3
0.4
0.5
0.6
1970
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
1.0
0.5
0.0
−0.5
$1-SE(\hat{r}_y)^2/\sigma_R^2$
1970
1980
1990
2000
2010
2020
Year


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
0
1
2
3
4
5
Spawning biomass (mt)
0
1
2
3
4
5


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
1998
2021
1967
2010
Recruitment (1,000s)
Spawning biomass (mt)
0
1
2
3
4
5


Log recruitment deviation
0.6
0.4
0.2
0.0
−0.2
−0.4
−0.6
1998
2021
2010
0.0
0.2
0.4
0.6
0.8
1.0
Spawning output (relative to B_0)


Landings (mt)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Landings (mt)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


FISHERY
FISHERY obs.
Observed and expected Landings (mt)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Continuous F
0.00
0.05
0.10
0.15
0.20
1970
1980
1990
2000
2010
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.5
1.0
1.5
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0
1
2
3
Relative spawning output: B/B_MSY
1969
2021


Index
0
5
10
15
20
25
30
35
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
5
10
15
20
25
30
35
1990
1995
2000
2005
2010
2015
2020
Year


Expected index
0
1
2
3
4
0
2
4
6
8
10
Observed index


Log index
3
2
1
0
−1
−2
1990
1995
2000
2005
2010
2015
2020
Year


Log index
3
2
1
0
−1
−2
1990
1995
2000
2005
2010
2015
2020
Year


Log expected index
0.8
1.0
1.2
1.4
−0.5
0.0
0.5
1.0
1.5
2.0
Log observed index


Residual
2
0
−2
1990
1995
2000
2005
2010
2015
2020
Year


Deviation
1.5
1.0
0.5
−0.5
−1.5
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
5
10
15
20
1990
1995
2000
2005
Year


Index
0
5
10
15
20
1990
1995
2000
2005
Year


Expected index
Observed index
0
2
4
6
8
0
2
4
6
8
10
12


Log index
2
0
−2
−4
1990
1995
2000
2005
Year


Log index
2
0
−2
−4
1990
1995
2000
2005
Year


Log expected index
2.0
1.8
1.6
1.4
−3
−2
−1
0
1
2
Log observed index


Residual
2
0
−2
1990
1995
2000
2005
Year


Deviation
4
2
0
−2
−4
1990
1995
2000
2005
Year


FISHERY
SURVEY2
Standardized index
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1990
1995
2000
2005
2010
2015
2020
Year


0.1
4
30
25
20
15
10
5
0
Age
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
2
Age
30
25
20
15
10
5
0
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
4
120
100
80
60
40
20
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


0.1
2
120
100
80
60
40
20
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


Numbers at age at equilibrium
0.0
0.5
1.0
1.5
2.0
2.5
3.0
0
5
10
15
20
25
30
Age


FISHERY
Sum of N adj.=983.8
Proportion
0.00
0.05
0.10
0.15
0
20
40
60
80
100
Length (cm)


FISHERY
0.01
0.2
Length (cm)
95
85
75
65
55
45
35
25
15
5
2008
2010
2012
2014
2016
2018
Year


2007
N adj.=93.4
2013
N adj.=49.6
2019
N adj.=65.5
2008
N adj.=79.8
2014
N adj.=72.2
2009
N adj.=110
2015
N adj.=99.1
2010
N adj.=41.2
2016
N adj.=81.5
2011
N adj.=61.3
2017
N adj.=105
2012
N adj.=86.5
2018
N adj.=38.6
Proportion
0.00
0.10
0.20
0
20
40
60
80
100
Length (cm)


0.01
0.2
Length (cm)
95
85
75
65
55
45
35
25
15
5
2008
2010
2012
2014
2016
2018
Year


FISHERY (whole catch)
Mean length
60
55
50
45
2008
2010
2012
2014
2016
2018
Year


FISHERY
Sum of N adj.=983.8
Sum of N eff.=1324.4
Proportion
0.15
0.10
0.05
0.00
0
20
40
60
80
100
Length (cm)


FISHERY
−4
0.1
4
Length (cm)
95
85
75
65
55
45
35
25
15
5
2008
2010
2012
2014
2016
2018
Year


2007
N adj.=93.4
N eff.=65.9
2013
N adj.=49.6
N eff.=90.9
2019
N adj.=65.5
N eff.=27.4
2008
N adj.=79.8
N eff.=53.1
2014
N adj.=72.2
N eff.=23.5
2009
N adj.=110
N eff.=281.7
2015
N adj.=99.1
N eff.=470.6
2010
N adj.=41.2
N eff.=74.8
2016
N adj.=81.5
N eff.=34.1
2011
N adj.=61.3
N eff.=34.5
2017
N adj.=105
N eff.=80.2
2012
N adj.=86.5
N eff.=52.3
2018
N adj.=38.6
N eff.=35.6
Proportion
0.00
0.10
0.20
0.30
0
20
40
60
80
100
Length (cm)


−4
0.1
4
95
85
75
65
55
45
35
25
15
5
Length (cm)
2008
2010
2012
2014
2016
2018
Year


Effective sample size
0
100
300
500
0
20
40
60
80
100
120
Observed sample size
harmonic mean
arithmetic mean
2015
2009
2013
2010
2017
2007
2008
2012
2018
2016
2014


FISHERY (whole catch)
Mean length
60
55
50
45
2008
2010
2012
2014
2016
2018
Year


Equilibrium yield (mt)
0.0
0.4
0.8
1.2
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


MSY
B target
SPR target
Current
Equilibrium yield (mt)
0.0
0.4
0.8
1.2
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Surplus production (mt)
0.0
0.5
1.0
1.5
2.0
0
5
10
15
20
25
Total biomass (mt)


Yield per recruit (kg)
0.0
0.2
0.4
0.6
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
SURVEY2
Length compositions
FISHERY
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
SURVEY2
Length compositions
FISHERY
1970
1980
1990
2000
2010
2020
Year


SR_LN(R0)
prior
max. likelihood
5
10
15
20
25
30
Size_95%width_FISHERY(1)
0
5
10
15
20
LnQ_base_FISHERY(1)
−6
−4
−2
0
2
4
LnQ_base_SURVEY2(2)
−6
−4
−2
0
2
4
Size_inflection_FISHERY(1)
20
30
40
50
60
Density
Parameter value