Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Sun Feb 19 15:07:32 2023
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
0
5
10
15
20
25
30
0
2
4
6
8
Age (yr)


Age (yr)
0
2
4
6
8
CV of lengths
0
0.01
0.03
Length (cm, beginning of the year)
0
5
10
15
20
25
30
CV of lengths
0.00
0.02
0.04
SD of lengths
0.0
0.4
0.8
Age (yr)


Age (yr)
Mean weight
Length (cm, beginning of the year)
Mean weight
Frac. mature
Age (yr)
Frac. mature
Length (cm, beginning of the year)


35
30
25
20
15
10
5
0
Length
0
2
4
6
8
Age


35
30
25
20
15
10
5
0
Length
0
2
4
6
8
Age


Length
35
30
25
20
15
10
5
0
0
2
4
6
8
Age


35
30
25
20
15
10
5
0
Length
0
2
4
6
8
Age


35
30
25
20
15
10
5
0
Length
0
2
4
6
8
Age


35
30
25
20
15
10
5
0
Length
0
2
4
6
8
Age


Mean weight (kg) in last year
0.0
0.1
0.2
0.3
0.4
0.5
0.6
5
10
15
20
25
30
Length (cm)


1.0
0.8
0.6
0.4
0.2
0.0
Maturity
5
10
15
20
25
30
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0.0
0.1
0.2
0.3
0.4
0.5
0.6
Female weight (kg)


Fecundity
0.0
0.1
0.2
0.3
0.4
0.5
0.6
0.0
0.1
0.2
0.3
0.4
0.5
0.6
Female weight (kg)


Fecundity
0.0
0.1
0.2
0.3
0.4
0.5
0.6
0.7
5
10
15
20
25
30
Female length (cm)


Spawning output
0.0
0.1
0.2
0.3
0.4
0.5
0.6
5
10
15
20
25
30
Length (cm)


Spawning output
0.00
0.05
0.10
0.15
0.20
0.25
0.30
0
2
4
6
8
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
5
10
15
20
25
30
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
2
4
6
8
Age (yr)
FISHERY
FISHERY
FISHERY


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
5
10
15
20
25
30
Length (cm)


Length (cm)
0
5
10
15
20
25
30
0.2
0.4
0
2
4
6
8
Age (yr)


Spawning biomass (mt)
0
5
10
15
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
5
10
15
20
25
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
1
2
3
4
5
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
2
4
6
8
1970
1980
1990
2000
2010
2020
Year


Total biomass (mt)
0
10
20
30
40
50
60
70
1970
1980
1990
2000
2010
2020
Year


Summary biomass (mt)
0
10
20
30
40
50
60
70
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
200
600
1000
1400
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
1000
2000
3000
4000
1970
1980
1990
2000
2010
2020
Year


Summary Fishing Mortality
0.0
0.1
0.2
0.3
0.4
0.5
0.6
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
5
10
15
20
1970
1980
1990
2000
2010
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
−0.2
0.0
0.2
0.4
0.6
1970
1980
1990
2000
2010
2020
Year


Log recruitment deviation
2.0
1.5
1.0
0.5
0.0
−1.0
1970
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.2
0.4
0.6
1970
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
$1 - SE(\hat{r_y})^2/\sigma_R^2$
1.0
0.5
0.0
−0.5
1970
1980
1990
2000
2010
2020
Year


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
0
200
600
1000
1400
0
2
4
6
8
Spawning biomass (mt)


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
1984
1985
1967
2021
Recruitment (1,000s)
0
200
600
1000
1400
0
2
4
6
8
Spawning biomass (mt)


1984
1985
1967
2021
0.6
0.4
0.2
0.0
−0.2
−0.4
−0.6
Log recruitment deviation
0.0
0.2
0.4
0.6
0.8
1.0
1.2
1.4
Spawning output (relative to B_0)


Landings (mt)
0
2
4
6
8
10
1970
1980
1990
2000
2010
2020
Year


Landings (mt)
0
2
4
6
8
1970
1980
1990
2000
2010
2020
Year


FISHERY
FISHERY obs.
Observed and expected Landings (mt)
0
2
4
6
8
10
12
1970
1980
1990
2000
2010
2020
Year


Continuous F
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0
1
2
3
4
5
Relative spawning output: B/B_MSY
2010
1980
2000
1990


Index
0
50
100
150
1990
1995
2000
2005
2010
2015
Year


Index
0
50
100
150
1990
1995
2000
2005
2010
2015
Year


Expected index
0
2
4
6
8
10
0
10
20
30
Observed index


Log index
5
4
3
2
1
0
−1
1990
1995
2000
2005
2010
2015
Year


Log index
5
4
3
2
1
0
−1
1990
1995
2000
2005
2010
2015
Year


Log expected index
1.7
1.8
1.9
2.0
2.1
2.2
2.3
2.4
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
Log observed index


Residual
2
1
0
−1
−2
1990
1995
2000
2005
2010
2015
Year


Deviation
1.5
1.0
0.5
−0.5
−1.5
1990
1995
2000
2005
2010
2015
Year


FISHERY
Standardized index
0
1
2
3
1990
1995
2000
2005
2010
2015
Year


0.1
2
Age
8
7
6
5
4
3
2
1
0
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


200
600
1000
Age
8
7
6
5
4
3
2
1
0
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
2
35
30
25
20
15
10
5
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


100
300
500
700
Length
Year
0
5
10
15
20
25
30
35
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021


Numbers at age at equilibrium
0
100
200
300
400
500
0
2
4
6
8
Age


FISHERY
Sum of N input=9502
Proportion
0.0
0.1
0.2
0.3
0.4
14
16
18
20
22
24
26
28
Length (cm)


FISHERY
0.01
0.4
Length (cm)
28
26
24
22
20
18
16
14
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


2004
N input=235
2010
N input=144
2016
N input=369
2005
N input=147
2011
N input=782
2017
N input=204
2006
N input=147
2012
N input=1255
2018
N input=170
2007
N input=292
2013
N input=2416
2019
N input=212
2008
N input=346
2014
N input=1062
2020
N input=155
2009
N input=456
2015
N input=1023
2021
N input=87
Proportion
0.0
0.1
0.2
0.3
0.4
14
18
22
26
Length (cm)


0.01
0.4
Length (cm)
28
26
24
22
20
18
16
14
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


FISHERY (whole catch)
22.5
22.0
21.5
21.0
20.5
Mean length
2005
2010
2015
2020
Year


FISHERY
Sum of N input=9502
Sum of N adj.=2677.2
Proportion
0.0
0.1
0.2
0.3
0.4
14
16
18
20
22
24
26
28
Length (cm)


FISHERY
−10
−6
2
6
10
Length (cm)
28
26
24
22
20
18
16
14
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


2004
N input=235
N adj.=66.6
2010
N input=144
N adj.=41.1
2016
N input=369
N adj.=104.2
2005
N input=147
N adj.=41.9
2011
N input=782
N adj.=220
2017
N input=204
N adj.=57.9
2006
N input=147
N adj.=41.9
2012
N input=1255
N adj.=352.6
2018
N input=170
N adj.=48.4
2007
N input=292
N adj.=82.6
2013
N input=2416
N adj.=678.1
2019
N input=212
N adj.=60.2
2008
N input=346
N adj.=97.7
2014
N input=1062
N adj.=298.5
2020
N input=155
N adj.=44.2
2009
N input=456
N adj.=128.6
2015
N input=1023
N adj.=287.6
2021
N input=87
N adj.=25.1
Proportion
0.0
0.1
0.2
0.3
0.4
14
18
22
26
Length (cm)


−10
−6
2
6
10
Length (cm)
28
26
24
22
20
18
16
14
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


FISHERY (whole catch)
22.5
22.0
21.5
21.0
20.5
Mean length
2005
2010
2015
2020
Year


Equilibrium yield (mt)
0
2
4
6
8
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0
2
4
6
8
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
0
5
10
0
10
20
30
40
50
60
70
Total biomass (mt)


Yield per recruit (kg)
0.000
0.004
0.008
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


SR_LN(R0)
prior
max. likelihood
3
4
5
6
7
8
Size_inflection_FISHERY(1)
20
21
22
23
24
InitF_seas_1_flt_1FISHERY
0.0
0.1
0.2
0.3
0.4
0.5
0.6
Size_95%width_FISHERY(1)
2.0
2.5
3.0
3.5
4.0
LnQ_base_FISHERY(1)
−4
−3
−2
−1
0
1
2
ln(DM_theta)_1
−4
−2
0
2
4
6
Q_extraSD_FISHERY(1)
0.0
0.5
1.0
1.5
2.0
Density
Parameter value