Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Fri Jul 01 13:26:47 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
70
60
50
40
30
20
10
0
0
5
10
15
20
25
Age (yr)


Age (yr)
0
5
10
15
20
25
Length (cm, beginning of the year)
70
60
50
40
30
20
10
0
CV of lengths
0.08
0.04
0
SD of lengths
4
2
0
CV of lengths
0.04
0.08
0
1
2
3
4
SD of lengths
Length (cm, beginning of the year)
Age (yr)


Age (yr)
Mean weight
Length (cm, beginning of the year)
Mean weight
Frac. mature
Age (yr)
Frac. mature


Length
70
60
50
40
30
20
10
0
0
5
10
15
20
25
30
Age


Length
70
60
50
40
30
20
10
0
0
5
10
15
20
25
30
Age


Mean weight (kg) in last year
0
1
2
3
4
5
0
10
20
30
40
50
60
Length (cm)


Maturity
0.0
0.2
0.4
0.6
0.8
0
10
20
30
40
50
60
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0
1
2
3
4
Female weight (kg)


Fecundity
0
1
2
3
4
5
0
1
2
3
4
Female weight (kg)


Fecundity
0
1
2
3
4
5
0
10
20
30
40
50
60
Female length (cm)


Spawning output
0
1
2
3
4
5
0
10
20
30
40
50
60
Length (cm)


Spawning output
0.0
0.2
0.4
0.6
0.8
1.0
1.2
0
5
10
15
20
25
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
50
60
Length (cm)
FISHERY
SURVEY2


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
5
10
15
20
25
Age (yr)
FISHERY
SURVEY2


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
50
60
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
50
60
Length (cm)


Length (cm)
0
10
20
30
40
50
60
1
0.8
0.6
0.4
0.2
0
5
10
15
20
25
Age (yr)


Length (cm)
0
10
20
30
40
50
60
1
0.8
0.6
0.4
0.2
0
5
10
15
20
25
Age (yr)


Spawning biomass (mt)
0.0
0.5
1.0
1.5
2.0
2.5
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
1
2
3
4
5
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
1
2
3
4
1990
2000
2010
2020
Year


Total biomass (mt)
0
1
2
3
4
5
6
1990
2000
2010
2020
Year


Summary biomass (mt)
0
1
2
3
4
5
6
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
1
2
3
4
5
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
5
10
15
20
1990
2000
2010
2020
Year


Summary Fishing Mortality
0.0
0.2
0.4
0.6
0.8
1985
1990
1995
2000
2005
2010
2015
2020
Year


Spawning biomass (mt)
0
1
2
3
4
1985
1990
1995
2000
2005
2010
2015
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
0.6
0.2
−0.2
−0.6
1970
1980
1990
2000
2010
2020
Year


Log recruitment deviation
2
1
0
−1
1970
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.2
0.4
0.6
0.8
1970
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
$1 - SE(\hat{r}_y)^2/\sigma_R^2$
1.0
0.5
0.0
−0.5
−1.0
1970
1980
1990
2000
2010
2020
Year


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
0
1
2
3
4
5
0.0
0.2
0.4
0.6
0.8
1.0
1.2
Spawning biomass (mt)


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
1997
1998
2001
2000
2021
1986
2013
Recruitment (1,000s)
0
1
2
3
4
5
0.0
0.2
0.4
0.6
0.8
1.0
1.2
Spawning biomass (mt)


1997
2001
1998
2000
2021
1986
2013
0.5
0.0
−0.5
0.0
0.2
0.4
0.6
0.8
1.0
Log recruitment deviation
Spawning output (relative to B_0)


Landings (mt)
0.0
0.2
0.4
0.6
0.8
1.0
1.2
1985
1990
1995
2000
2005
2010
2015
2020
Year


Landings (mt)
0.0
0.2
0.4
0.6
0.8
1.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


Continuous F
0.00
0.10
0.20
0.30
1985
1990
1995
2000
2005
2010
2015
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0.0
0.5
1.0
1.5
2.0
2.5
3.0
Relative spawning output: B/B_MSY
1987
2021


Index
0
5
10
15
20
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
5
10
15
20
1990
1995
2000
2005
2010
2015
2020
Year


Expected index
0.0
0.2
0.4
0.6
0.8
1.0
1.2
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
Observed index


Log index
2
0
−2
−4
−6
1990
1995
2000
2005
2010
2015
2020
Year


Log index
2
0
−2
−4
−6
1990
1995
2000
2005
2010
2015
2020
Year


Log expected index
0.2
0.1
0.0
−0.2
−0.4
−2
−1
0
1
Log observed index


Residual
2
1
0
−1
−2
1990
1995
2000
2005
2010
2015
2020
Year


Deviation
2
1
0
−1
−2
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
5
10
15
20
25
30
35
1990
1995
2000
2005
Year


Index
0
5
10
15
20
25
30
35
1990
1995
2000
2005
Year


Expected index
0.0
0.5
1.0
1.5
2.0
2.5
0
1
2
3
4
5
Observed index


Log index
2
0
−2
−4
−6
−8
1990
1995
2000
2005
Year


Log index
2
0
−2
−4
−6
−8
1990
1995
2000
2005
Year


Log expected index
1.0
0.9
0.8
0.7
0.6
0.5
−4
−2
0
Log observed index


Residual
4
2
0
−2
−4
1990
1995
2000
2005
Year


Deviation
6
4
2
0
−2
−4
−6
1990
1995
2000
2005
Year


FISHERY
SURVEY2
Standardized index
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
1990
1995
2000
2005
2010
2015
2020
Year


0.1
4
30
25
20
15
10
5
0
Age
1985
1989
1993
1997
2001
2005
2009
2013
2017
2021
Year


0.1
2
30
25
20
15
10
5
0
Age
1985
1989
1993
1997
2001
2005
2009
2013
2017
2021
Year


0.1
4
70
60
50
40
30
20
10
0
Length
1986
1990
1994
1998
2002
2006
2010
2014
2018
2022
Year


0.1
2
70
60
50
40
30
20
10
0
Length
1986
1990
1994
1998
2002
2006
2010
2014
2018
2022
Year


Numbers at age at equilibrium
0.0
0.5
1.0
1.5
2.0
2.5
0
5
10
15
20
25
Age


FISHERY
Sum of N adj.=217.6
Proportion
0.35
0.30
0.25
0.20
0.15
0.10
0.05
0.00
20
25
30
35
40
45
50
55
Length (cm)


FISHERY
0.01
0.2
0.4
55
50
45
40
35
30
25
20
Length (cm)
2013
2015
Year


2012
N adj.=84.8
2013
N adj.=89.9
2015
N adj.=42.8
0.0
0.1
0.2
0.3
0.4
20
30
40
50
Proportion
Length (cm)


0.01
0.2
0.4
55
50
45
40
35
30
25
20
Length (cm)
2012
2013
2015
Year


FISHERY (whole catch)
35
34
33
32
31
Mean length
2012.0
2012.5
2013.0
2013.5
2014.0
2014.5
2015.0
Year


FISHERY
Sum of N adj.=217.6
Sum of N eff.=407.8
Proportion
0.35
0.30
0.25
0.20
0.15
0.10
0.05
0.00
20
25
30
35
40
45
50
55
Length (cm)


FISHERY
−25
−15
5
15
25
55
50
45
40
35
30
25
20
Length (cm)
2013
2015
Year


2012
N adj.=84.8
N eff.=137.9
2013
N adj.=89.9
N eff.=128.5
2015
N adj.=42.8
N eff.=141.4
0.0
0.1
0.2
0.3
0.4
20
30
40
50
Proportion
Length (cm)


−25
−15
5
15
25
55
50
45
40
35
30
25
20
Length (cm)
2012
2013
2015
Year


harmonic mean
2015
2012
2013
arithmetic mean
Effective sample size
Observed sample size
0
50
100
150
0
20
40
60
80
100


FISHERY (whole catch)
Mean length
35
34
33
32
31
2012.0
2012.5
2013.0
2013.5
2014.0
2014.5
2015.0
Year


Equilibrium yield (mt)
0.0
0.1
0.2
0.3
0.4
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0.0
0.1
0.2
0.3
0.4
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
0.6
0.2
−0.2
0
1
2
3
4
5
6
Total biomass (mt)


Yield per recruit (kg)
0.00
0.10
0.20
1985
1990
1995
2000
2005
2010
2015
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
SURVEY2
Length compositions
FISHERY
1985
1990
1995
2000
2005
2010
2015
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
SURVEY2
Length compositions
FISHERY
1985
1990
1995
2000
2005
2010
2015
2020
Year


SR_LN(R0)
prior
max. likelihood
5
10
15
20
25
30
Size_95%width_FISHERY(1)
0
2
4
6
8
10
12
14
LnQ_base_FISHERY(1)
−6
−4
−2
0
2
4
LnQ_base_SURVEY2(2)
−6
−4
−2
0
2
4
Size_inflection_FISHERY(1)
15
20
25
30
35
40
45
50
Density
Parameter value