Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Wed Jul 20 08:40:59 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
50
40
30
20
10
0
0
2
4
6
8
10
12
14
Age (yr)


Age (yr)
0
2
4
6
8
10
12
14
Length (cm, beginning of the year)
0
10
20
30
40
50
CV of lengths
0
0.02
0.04
0.06
SD of lengths
0.0
1.0
2.0
Length (cm, beginning of the year)
CV of lengths
SD of lengths
Age (yr)


Age (yr)
Mean weight
Length (cm, beginning of the year)
Mean weight
Frac. mature
Age (yr)


Length
50
40
30
20
10
0
0
2
4
6
8
10
12
14
Age


Length
50
40
30
20
10
0
0
2
4
6
8
10
12
14
Age


Mean weight (kg) in last year
0.0
0.5
1.0
1.5
2.0
0
10
20
30
40
50
Length (cm)


1.0
0.8
0.6
0.4
0.2
0.0
Maturity
0
10
20
30
40
50
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0.0
0.5
1.0
1.5
2.0
Female weight (kg)


Fecundity
0.0
0.5
1.0
1.5
2.0
0.0
0.5
1.0
1.5
2.0
Female weight (kg)


Fecundity
2.5
2.0
1.5
1.0
0.5
0.0
0
10
20
30
40
50
Female length (cm)


Spawning output
0.0
0.5
1.0
1.5
2.0
0
10
20
30
40
50
Length (cm)


Spawning output
1.5
1.0
0.5
0.0
0
2
4
6
8
10
12
14
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
50
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
2
4
6
8
10
12
14
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
50
Length (cm)


Length (cm)
0
10
20
30
40
0.8
0.6
0.4
0.2
0
2
4
6
8
10
12
14
Age (yr)


Spawning biomass (mt)
0.0
0.4
0.8
1.2
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0.0
0.5
1.0
1.5
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
1
2
3
1970
1980
1990
2000
2010
2020
Year


Total biomass (mt)
0.0
1.0
2.0
3.0
1970
1980
1990
2000
2010
2020
Year


Summary biomass (mt)
0.0
1.0
2.0
3.0
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
1
2
3
4
5
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
2
4
6
8
10
12
1970
1980
1990
2000
2010
2020
Year


Summary Fishing Mortality
0
1
2
3
4
5
6
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0.0
0.5
1.0
1.5
1970
1980
1990
2000
2010
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
0.3
0.1
−0.1
−0.3
1970
1980
1990
2000
2010
2020
Year


Log recruitment deviation
1.0
0.5
0.0
−0.5
−1.0
1970
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.1
0.2
0.3
0.4
0.5
1970
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
1.0
0.5
0.0
$1-SE(\hat{r}_y)^2/\sigma_R{}^2$
1970
1980
1990
2000
2010
2020
Year


Recruitment (1,000s)
0
1
2
3
4
5
0.0
0.1
0.2
0.3
0.4
0.5
0.6
0.7
Spawning biomass (mt)
Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium


Recruitment (1,000s)
0
1
2
3
4
5
0.0
0.1
0.2
0.3
0.4
0.5
0.6
0.7
Spawning biomass (mt)
1967
2021
Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium


0.4
0.2
0.0
−0.2
−0.4
Log recruitment deviation
2021
1967
0.0
0.2
0.4
0.6
0.8
Spawning output (relative to B_0)


Landings (mt)
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Landings (mt)
0.0
0.2
0.4
0.6
0.8
1970
1980
1990
2000
2010
2020
Year


FISHERY
FISHERY obs.
Observed and expected Landings (mt)
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Continuous F
0.0
0.5
1.0
1.5
1970
1980
1990
2000
2010
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1.2
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0.0
0.5
1.0
1.5
2.0
2.5
3.0
Relative spawning output: B/B_MSY
2021
1967


Index
0
1
2
3
4
5
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
1
2
3
4
5
1990
1995
2000
2005
2010
2015
2020
Year


Expected index
0.0
0.2
0.4
0.6
0.0
0.5
1.0
1.5
Observed index


Log index
0
−2
−4
−6
1990
1995
2000
2005
2010
2015
2020
Year


Log index
0
−2
−4
−6
1990
1995
2000
2005
2010
2015
2020
Year


Log expected index
−0.5
−1.0
−1.5
−2.0
−2.5
−3
−2
−1
0
Log observed index


Effective catchability
0.3568690
0.3568680
0.3568670
0.5
1.0
1.5
2.0
Vulnerable biomass


Residual
2
1
0
−1
−2
1990
1995
2000
2005
2010
2015
2020
Year


Deviation
2
1
0
−1
−2
1990
1995
2000
2005
2010
2015
2020
Year


FISHERY
Standardized index
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1990
1995
2000
2005
2010
2015
2020
Year


0.1
4
Age
14
13
12
11
10
9
8
7
6
5
4
3
2
1
0
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
2
Age
14
13
12
11
10
9
8
7
6
5
4
3
2
1
0
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
2
50
40
30
20
10
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


0.1
2
50
40
30
20
10
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


Numbers at age at equilibrium
0
1
2
3
4
0
2
4
6
8
10
12
14
Age


FISHERY
Sum of N adj.=36.6
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
15
20
25
30
35
40
45
50
Length (cm)


FISHERY
0.01
0.2
0.4
Length (cm)
48
45
42
39
36
33
30
27
24
21
18
15
2012
2013
2014
2015
Year


2011
N adj.=6.8
2012
N adj.=7.9
2013
N adj.=7.3
2014
N adj.=5.4
2015
N adj.=9.2
0.00
0.10
0.20
0.30
15
25
35
45
Proportion
Length (cm)


0.01
0.2
0.4
48
45
42
39
36
33
30
27
24
21
18
15
Length (cm)
2011
2012
2013
2014
2015
Year


FISHERY (whole catch)
Mean length
35
30
25
20
2011
2012
2013
2014
2015
Year


FISHERY
Sum of N adj.=36.6
Sum of N eff.=160.4
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
15
20
25
30
35
40
45
50
Length (cm)


FISHERY
−10
−5
0.1
5
10
Length (cm)
48
45
42
39
36
33
30
27
24
21
18
15
2012
2013
2014
2015
Year


2011
N adj.=6.8
N eff.=7
2012
N adj.=7.9
N eff.=42.7
2013
N adj.=7.3
N eff.=42
2014
N adj.=5.4
N eff.=46.9
2015
N adj.=9.2
N eff.=21.8
0.00
0.10
0.20
0.30
15
25
35
45
Proportion
Length (cm)


−10
−5
0.1
5
10
48
45
42
39
36
33
30
27
24
21
18
15
Length (cm)
2011
2012
2013
2014
2015
Year


Effective sample size
0
10
20
30
40
50
0
2
4
6
8
10
Observed sample size
2014
2013
2012
2015
2011
harmonic mean
arithmetic mean


FISHERY (whole catch)
Mean length
35
30
25
20
2011
2012
2013
2014
2015
Year


Equilibrium yield (mt)
0.0
0.1
0.2
0.3
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0.0
0.1
0.2
0.3
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
0.0
0.2
0.4
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
Total biomass (mt)


Yield per recruit (kg)
0.00
0.04
0.08
0.12
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


SR_LN(R0)
prior
max. likelihood
1
2
3
4
5
Size_95%width_FISHERY(1)
5
10
15
20
InitF_seas_1_flt_1FISHERY
0.00
0.05
0.10
0.15
0.20
0.25
0.30
LnQ_base_FISHERY(1)
−4
−3
−2
−1
0
1
2
Size_inflection_FISHERY(1)
20
25
30
35
40
45
50
Density
Parameter value