


American Samoa Model Checks


2022-09-22

This is a summary report for the LUKA base model run.

Model Output

Input Data





Convergence Check

```

Converged      MaxGrad
1      TRUE 4.42361e-06

```



- [1] "1 NOTE: Max data length bin: 28 < max pop len bins: 31; so will accumulate larger pop
- [2] "2 warning: poor convergence in Fspr search 0.4 0.452066"
- [3] "3 warning: Fmult = 40 cannot get high enough to achieve low SPR target: 0.4; SPR achiev
- [4] "4 warning: poor convergence in Btarget search 5.61753 7.17708"
- [5] "5 warning: poor convergence in Fmsy, final dy/dy2= -0.0152624"
- [6] "6 Forecast F capped by max possible F from control file: 2.9"
- [7] "7 Forecast F capped by max possible F from control file: 2.9"
- [8] "N warnings: 7"

Fit to Model

CPUE

Residual Runs Test (/w plot) stats by Index:

RMSE stats by Index:




Length Comp

Residual Runs Test (/w plot) stats by Mean length:



```
Index runs.p  test  sigma3.lo  sigma3.hi  type
1 FISHERY    0.024 Failed -0.02803011 0.02803011  len
```


RMSE stats by Index:

```
# A tibble: 2 x 3
  Fleet    RMSE.perc  Nobs
  <chr>      <dbl> <int>
1 FISHERY      1.2     18
2 Combined      1.2     18
```




Retrospective and Hindcasting





Retrospective

Mohn's Rho stats, including one step ahead forecasts:



Mohn's Rho stats, including one step ahead forecasts:


	type	peel	Rho	ForecastRho
1	F	2020	-0.0002433336	-0.0002272745
2	F	2019	-0.0154574401	-0.0151827685
3	F	2018	-0.0090489327	-0.0087113446
4	F Combined		-0.0082499021	-0.0080404625

Hindcasting

Plotting Hindcast Cross-Validation (one-step-ahead)

Computing MASE with only 2 of 3 prediction residuals for Index FISHERY

Warning: Unequal spacing of naive predictions residuals may influence the interpretation of




MASE stats by Index:

Plotting Hindcast Cross-Validation (one-step-ahead)

Computing MASE with only 2 of 3 prediction residuals for Index FISHERY

Warning: Unequal spacing of naive predictions residuals may influence the interpretation of

MASE stats by Index:




Recruitment Deviations

Likelihood Profile


[1] "SR_LN"

	frac_change	include	label
TOTAL	1.0000	TRUE	Total
Catch	0.7217	TRUE	Catch
Equil_catch	0.0105	TRUE	Equilibrium catch
Survey	0.0510	TRUE	Index data
Length_comp	0.3381	TRUE	Length data
Recruitment	0.0000	FALSE	Recruitment
InitEQ_Regime	0.0000	FALSE	Initital equilibrium recruitment
Forecast_Recruitment	0.0000	FALSE	Forecast recruitment
Parm_priors	0.0011	FALSE	Priors

Parm_softbounds	0.0000	FALSE	Soft bounds
Parm_devs	0.0000	FALSE	Parameter deviations
Crash_Pen	0.0000	FALSE	Crash penalty






Changes in survey likelihood by fleet





Management Quantities

starter.sso with Bratio: SSB/SSB_{MSY} and F: `_abs_F`







null device
1

Jitter

