Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Tue Feb 14 14:18:42 2023
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
0
20
40
60
0
2
4
6
8
10
12
Age (yr)


Age (yr)
0
2
4
6
8
10
12
Length (cm, beginning of the year)
0
20
40
60
CV of lengths
0
0.02
0.04
SD of lengths
0.0
1.0
2.0
3.0
1.5


Age (yr)
Mean weight
Length (cm, beginning of the year)
Mean weight
Frac. mature
Age (yr)
Frac. mature
Length (cm, beginning of the year)


80
60
40
20
0
Length
0
2
4
6
8
10
12
Age


80
60
40
20
0
Length
0
2
4
6
8
10
12
Age


80
60
40
20
0
Length
0
2
4
6
8
10
12
Age


80
60
40
20
0
Length
0
2
4
6
8
10
12
Age


80
60
40
20
0
Length
0
2
4
6
8
10
12
Age


80
60
40
20
0
Length
0
2
4
6
8
10
12
Age


Mean weight (kg) in last year
0
2
4
6
8
0
10
20
30
40
50
60
70
Length (cm)


1.0
0.8
0.6
0.4
0.2
0.0
Maturity
0
10
20
30
40
50
60
70
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0
2
4
6
8
Female weight (kg)


Fecundity
0
2
4
6
8
0
2
4
6
8
Female weight (kg)


Fecundity
0
2
4
6
8
10
0
10
20
30
40
50
60
70
Female length (cm)


Spawning output
0
2
4
6
8
0
10
20
30
40
50
60
70
Length (cm)


Spawning output
0
2
4
6
8
0
2
4
6
8
10
12
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
50
60
70
Length (cm)
FISHERY
SURVEY2
SURVEY3


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
2
4
6
8
10
12
Age (yr)
FISHERY
SURVEY2
SURVEY3
FISHERY
SURVEY2
SURVEY3
FISHERY
SURVEY2
SURVEY3


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
50
60
70
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
50
60
70
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
50
60
70
Length (cm)


Length (cm)
0
10
20
30
40
50
60
70
0.8
0.6
0.4
0.2
0
2
4
6
8
10
12
Age (yr)


Length (cm)
0
10
20
30
40
50
60
70
0.8
0.6
0.4
0.2
0
2
4
6
8
10
12
Age (yr)


Length (cm)
0
10
20
30
40
50
60
70
0.8
0.6
0.4
0.2
0
2
4
6
8
10
12
Age (yr)


Spawning biomass (mt)
0.0
1.0
2.0
3.0
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
1
2
3
4
5
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
1.0
2.0
3.0
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Total biomass (mt)
0
2
4
6
8
1970
1980
1990
2000
2010
2020
Year


Summary biomass (mt)
0
2
4
6
8
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
1
2
3
4
5
6
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
5
10
15
1970
1980
1990
2000
2010
2020
Year


Summary Fishing Mortality
0.0
0.2
0.4
0.6
0.8
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
1
2
3
4
5
1970
1980
1990
2000
2010
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
0.4
0.2
0.0
−0.2
1970
1980
1990
2000
2010
2020
Year


Log recruitment deviation
1.0
0.5
0.0
−0.5
−1.0
1970
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.1
0.2
0.3
0.4
0.5
1970
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
1.0
0.5
0.0
$1-SE(\hat{r}_y)^2/\sigma_R^2$
1970
1980
1990
2000
2010
2020
Year


Recruitment (1,000s)
0
1
2
3
4
5
6
0.0
0.5
1.0
1.5
Spawning biomass (mt)
Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium


Recruitment (1,000s)
0
1
2
3
4
5
6
0.0
0.5
1.0
1.5
Spawning biomass (mt)
2021
1967
Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium


Log recruitment deviation
0.4
0.2
0.0
−0.2
−0.4
0.0
0.2
0.4
0.6
0.8
1.0
Spawning output (relative to B_0)
2021
1967


Landings (mt)
0.0
0.5
1.0
1.5
2.0
1970
1980
1990
2000
2010
2020
Year


Landings (mt)
0.0
0.5
1.0
1.5
2.0
1970
1980
1990
2000
2010
2020
Year


Continuous F
0.0
0.1
0.2
0.3
0.4
0.5
0.6
0.7
1970
1980
1990
2000
2010
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0
1
2
3
Relative spawning output: B/B_MSY
2010
2000
1990
2020
1980
1968
2021
1970


Index
0
5
10
15
20
25
30
1990
1995
2000
2005
2010
2015
Year


Index
0
5
10
15
20
25
30
1990
1995
2000
2005
2010
2015
Year


Expected index
Observed index
0
1
2
3
0
2
4
6
8


Log index
3
2
1
0
−1
1990
1995
2000
2005
2010
2015
Year


Log index
3
2
1
0
−1
1990
1995
2000
2005
2010
2015
Year


Log expected index
0.8
0.9
1.0
1.1
1.2
1.3
0.5
1.0
1.5
2.0
Log observed index


Residual
1
0
−1
1990
1995
2000
2005
2010
2015
Year


Deviation
1.0
0.5
0.0
−0.5
−1.0
1990
1995
2000
2005
2010
2015
Year


Index
0
1
2
3
4
5
2016
2017
2018
2019
2020
Year


Index
0
1
2
3
4
5
2016
2017
2018
2019
2020
Year


Expected index
0.0
0.5
1.0
1.5
2.0
0.0
0.5
1.0
1.5
2.0
2.5
3.0
Observed index


Log index
1.5
1.0
0.5
0.0
−0.5
−1.0
2016
2017
2018
2019
2020
Year


Log index
1.5
1.0
0.5
0.0
−0.5
−1.0
2016
2017
2018
2019
2020
Year


Log expected index
0.85
0.80
0.75
0.70
0.2
0.4
0.6
0.8
1.0
1.2
Log observed index


Residual
1.5
1.0
0.5
0.0
−0.5
−1.5
2016
2017
2018
2019
2020
Year


Deviation
0.5
0.0
−0.5
2016
2017
2018
2019
2020
Year


SURVEY2
SURVEY3
Standardized index
0.0
0.5
1.0
1.5
2.0
2.5
1990
1995
2000
2005
2010
2015
2020
Year


0.1
4
8
Age
12
11
10
9
8
7
6
5
4
3
2
1
0
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
2
12
11
10
9
8
7
6
5
4
3
2
1
0
Age
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
2
80
60
40
20
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


0.1
2
80
60
40
20
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


Numbers at age at equilibrium
0.0
0.5
1.0
1.5
2.0
2.5
3.0
0
2
4
6
8
10
12
Age


FISHERY
Sum of N input=559
Proportion
0.00
0.05
0.10
0.15
0.20
0.25
10
20
30
40
50
60
70
Length (cm)


FISHERY
0.01
0.2
Length (cm)
65
60
55
50
45
40
35
30
25
20
15
2009
2012
2014
2016
2018
Year


2007
N input=71
2016
N input=80
2009
N input=113
2018
N input=70
2012
N input=76
2013
N input=47
2014
N input=45
2015
N input=57
0.30
0.20
0.10
0.00
10
20
30
40
50
60
70
Proportion
Length (cm)


0.01
0.2
65
60
55
50
45
40
35
30
25
20
15
Length (cm)
2009
2012
2014
2016
2018
Year


FISHERY (whole catch)
Mean length
45
40
35
2008
2010
2012
2014
2016
2018
Year


FISHERY
Sum of N input=559
Sum of N adj.=367.3
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
10
20
30
40
50
60
70
Length (cm)


FISHERY
−2
0.1
2
Length (cm)
65
60
55
50
45
40
35
30
25
20
15
2009
2012
2014
2016
2018
Year


2007
N input=71
N adj.=46.6
2016
N input=80
N adj.=52.5
2009
N input=113
N adj.=74
2018
N input=70
N adj.=46
2012
N input=76
N adj.=49.9
2013
N input=47
N adj.=31
2014
N input=45
N adj.=29.7
2015
N input=57
N adj.=37.5
0.00
0.10
0.20
0.30
10
20
30
40
50
60
70
Proportion
Length (cm)


−2
0.1
2
65
60
55
50
45
40
35
30
25
20
15
Length (cm)
2009
2012
2014
2016
2018
Year


FISHERY (whole catch)
Mean length
45
40
35
2008
2010
2012
2014
2016
2018
Year


Equilibrium yield (mt)
0.0
0.2
0.4
0.6
0.8
1.0
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0.0
0.2
0.4
0.6
0.8
1.0
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
0.0
0.4
0.8
1.2
0
2
4
6
8
Total biomass (mt)


Yield per recruit (kg)
0.00
0.10
0.20
0.30
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
SURVEY2
SURVEY3
Length compositions
FISHERY
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
SURVEY2
SURVEY3
Length compositions
FISHERY
1970
1980
1990
2000
2010
2020
Year


SR_LN(R0)
prior
max. likelihood
2
4
6
8
10
12
14
Q_extraSD_SURVEY3(3)
0.0
0.5
1.0
1.5
2.0
LnQ_base_SURVEY2(2)
−4
−2
0
2
4
Size_inflection_FISHERY(1)
30
35
40
45
50
55
60
Q_extraSD_SURVEY2(2)
0.0
0.5
1.0
1.5
2.0
Size_95%width_FISHERY(1)
5
10
15
20
25
LnQ_base_SURVEY3(3)
−4
−2
0
2
4
ln(DM_theta)_1
−4
−2
0
2
4
6
8
10
Density
Parameter value