Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Fri Jul 01 08:34:10 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
0
10
20
30
40
50
60
0
2
4
6
8
10
12
14
Age (yr)


Age (yr)
0
2
4
6
8
10
12
14
Length (cm, beginning of the year)
0
10
20
30
40
50
60
CV of lengths
0
0.04
0.08
SD of lengths
0
1
2
3
4
Age (yr)


Age (yr)
Mean weight
Length (cm, beginning of the year)
Mean weight
Frac. mature
Age (yr)


Length
Age
0
10
20
30
40
50
60
0
2
4
6
8
10
12
14


Length
Age
0
10
20
30
40
50
60
0
2
4
6
8
10
12
14


Mean weight (kg) in last year
0
1
2
3
4
0
10
20
30
40
50
60
Length (cm)


Maturity
0.0
0.2
0.4
0.6
0.8
0
10
20
30
40
50
60
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0
1
2
3
Female weight (kg)


Fecundity
0
1
2
3
4
0
1
2
3
Female weight (kg)


Fecundity
0
1
2
3
4
0
10
20
30
40
50
60
Female length (cm)


Spawning output
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
0
10
20
30
40
50
60
Length (cm)


0.5
0.4
0.3
0.2
0.1
0.0
Spawning output
0
2
4
6
8
10
12
14
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
50
60
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
2
4
6
8
10
12
14
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
50
60
Length (cm)


Length (cm)
0
10
20
30
40
50
60
Age (yr)
0
2
4
6
8
10
12
14
1
0.8
0.6
0.2


Spawning biomass (mt)
0.00
0.05
0.10
0.15
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0.00
0.10
0.20
0.30
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
1
2
3
4
5
1970
1980
1990
2000
2010
2020
Year


Total biomass (mt)
0
1
2
3
4
5
6
1970
1980
1990
2000
2010
2020
Year


Summary biomass (mt)
0
1
2
3
4
5
6
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
5
10
15
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
10
20
30
40
50
1970
1980
1990
2000
2010
2020
Year


Summary Fishing Mortality
0.0
0.2
0.4
0.6
0.8
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0.00
0.05
0.10
0.15
0.20
0.25
1970
1980
1990
2000
2010
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
0.5
0.0
−0.5
−1.0
1970
1980
1990
2000
2010
2020
Year


Log recruitment deviation
1.5
1.0
0.5
−0.5
−1.5
1970
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.1
0.2
0.3
0.4
0.5
1970
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
1.0
0.5
0.0
$1-SE(\hat{r}_y)^2/\sigma_R{}^2$
1970
1980
1990
2000
2010
2020
Year


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
0
5
10
15
20
0.00
0.02
0.04
0.06
0.08
0.10
Spawning biomass (mt)


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
Spawning biomass (mt)
0
5
10
15
20
0.00
0.02
0.04
0.06
0.08
0.10
2021
2009
2010


1.0
0.5
0.0
−0.5
−1.0
Log recruitment deviation
0.0
0.2
0.4
0.6
0.8
1.0
Spawning output (relative to B_0)
2021
2012
2013
2011
2010
2009


Landings (mt)
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Landings (mt)
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Continuous F
0.0
0.1
0.2
0.3
0.4
1970
1980
1990
2000
2010
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0
1
2
3
4
Relative spawning output: B/B_MSY
1969
2021


Index
0
1
2
3
4
5
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
1
2
3
4
5
1990
1995
2000
2005
2010
2015
2020
Year


Expected index
0.0
0.1
0.2
0.3
0.4
0.5
0.6
0.0
0.5
1.0
1.5
Observed index


Log index
0
−2
−4
−6
1990
1995
2000
2005
2010
2015
2020
Year


Log index
0
−2
−4
−6
1990
1995
2000
2005
2010
2015
2020
Year


Log expected index
−1.6
−1.4
−1.2
−1.0
−0.8
−0.6
−3
−2
−1
0
Log observed index


Residual
2
1
0
−1
−2
1990
1995
2000
2005
2010
2015
2020
Year


Deviation
2
1
0
−1
−2
1990
1995
2000
2005
2010
2015
2020
Year


FISHERY
Standardized index
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1990
1995
2000
2005
2010
2015
2020
Year


5
15
Age
14
13
12
11
10
9
8
7
6
5
4
3
2
1
0
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


5
15
Age
14
13
12
11
10
9
8
7
6
5
4
3
2
1
0
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


5
15
60
50
40
30
20
10
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


0.1
5
10
Length
0
10
20
30
40
50
60
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


Numbers at age at equilibrium
0
2
4
6
8
10
12
0
2
4
6
8
10
12
14
Age


FISHERY
Sum of N adj.=334.3
Proportion
0.00
0.05
0.10
0.15
0.20
15
20
25
30
35
40
45
50
Length (cm)


FISHERY
0.01
0.2
0.4
Length (cm)
48
45
42
39
36
33
30
27
24
21
18
15
2012
2013
2014
2015
Year


2011
N adj.=62.2
2012
N adj.=72.2
2013
N adj.=66.4
2014
N adj.=49.6
2015
N adj.=84
0.00
0.10
0.20
0.30
15
25
35
45
Proportion
Length (cm)


0.01
0.2
0.4
48
45
42
39
36
33
30
27
24
21
18
15
Length (cm)
2011
2012
2013
2014
2015
Year


FISHERY (whole catch)
Mean length
36
34
32
30
28
26
24
22
2011
2012
2013
2014
2015
Year


FISHERY
Sum of N adj.=334.3
Sum of N eff.=322.5
Proportion
0.00
0.05
0.10
0.15
0.20
15
20
25
30
35
40
45
50
Length (cm)


FISHERY
−2
0.1
2
Length (cm)
48
45
42
39
36
33
30
27
24
21
18
15
2012
2013
2014
2015
Year


2011
N adj.=62.2
N eff.=18.4
2012
N adj.=72.2
N eff.=28.9
2013
N adj.=66.4
N eff.=61.8
2014
N adj.=49.6
N eff.=60.6
2015
N adj.=84
N eff.=152.8
0.00
0.10
0.20
0.30
15
25
35
45
Proportion
Length (cm)


−2
0.1
2
48
45
42
39
36
33
30
27
24
21
18
15
Length (cm)
2011
2012
2013
2014
2015
Year


Effective sample size
0
50
100
150
0
20
40
60
80
Observed sample size
2015
2014
2013
2012
2011
harmonic mean
arithmetic mean


FISHERY (whole catch)
Mean length
36
34
32
30
28
26
24
22
2011
2012
2013
2014
2015
Year


Equilibrium yield (mt)
0.0
0.1
0.2
0.3
0.4
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


MSY
B target
SPR target
Current
Equilibrium yield (mt)
0.0
0.1
0.2
0.3
0.4
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Surplus production (mt)
0.0
0.2
0.4
0.6
0.8
0
1
2
3
4
5
6
Total biomass (mt)


Yield per recruit (kg)
0.00
0.02
0.04
0.06
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1970
1980
1990
2000
2010
2020
Year


Catches
Abundance indices
Length compositions
FISHERY
FISHERY
FISHERY
1970
1980
1990
2000
2010
2020
Year


SR_LN(R0)
prior
max. likelihood
5
10
15
20
25
30
LnQ_base_FISHERY(1)
−6
−4
−2
0
2
4
Size_inflection_FISHERY(1)
15
20
25
30
35
40
Size_95%width_FISHERY(1)
0
1
2
3
4
5
Density
Parameter value