Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Wed Oct 19 12:47:39 2022
Data_File: data.ss
Control_File: control.ss


Females
Males
Length (cm, beginning of the year)
Age (yr)
0
10
20
30
40
50
60
0
2
4
6
8
10
12
14


Age (yr)
CV of lengths
Length (cm, beginning of the year)
Females
Males
CV of lengths
SD of lengths
Age (yr)


Age (yr)
Mean weight
Length (cm, beginning of the year)
Females
Males
Mean weight
Frac. mature
Age (yr)


Length
60
50
40
30
20
10
0
0
5
10
15
Age


Length
60
50
40
30
20
10
0
0
5
10
15
Age


Length
60
50
40
30
20
10
0
0
5
10
15
Age


Length
60
50
40
30
20
10
0
0
5
10
15
Age


Length
60
50
40
30
20
10
0
0
5
10
15
Age


Length
60
50
40
30
20
10
0
0
5
10
15
Age


Length
60
50
40
30
20
10
0
0
5
10
15
Age


Length
60
50
40
30
20
10
0
0
5
10
15
Age


Length
60
50
40
30
20
10
0
0
5
10
15
Age


Length
60
50
40
30
20
10
0
0
5
10
15
Age


Length
60
50
40
30
20
10
0
0
5
10
15
Age


Length
60
50
40
30
20
10
0
0
5
10
15
Age


Females
Males
Mean weight (kg) in last year
0
1
2
3
0
10
20
30
40
50
Length (cm)


Females
Males
Mean weight (kg) in last year
0
1
2
3
0
10
20
30
40
50
Length (cm)


1.0
0.8
0.6
0.4
0.2
0.0
Maturity
0
10
20
30
40
50
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
Female weight (kg)


Fecundity
0
1
2
3
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
Female weight (kg)


Fecundity
0
1
2
3
4
0
10
20
30
40
50
Female length (cm)


Spawning output
0
1
2
3
0
10
20
30
40
50
Length (cm)


Spawning output
2.0
1.5
1.0
0.5
0.0
0
5
10
15
Age (yr)


Hermaphroditism transition rate
1.0
0.8
0.6
0.4
0.2
0.0
0
2
4
6
8
10
12
14
Year


Fraction females by age at equilibrium
1.0
0.8
0.6
0.4
0.2
0.0
0
2
4
6
8
10
12
14
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
50
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
5
10
15
Age (yr)
FISHERY
FISHERY
FISHERY


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
50
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
50
Length (cm)


Length (cm)
0
10
20
30
40
50
0
5
10
15
Age (yr)
1
0.8
0.6
0.4
0.2


Length (cm)
0
10
20
30
40
50
0
5
10
15
Age (yr)
1
0.8
0.6
0.4
0.2


Spawning biomass (mt)
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
2
4
6
8
10
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
1.0
2.0
3.0
1970
1980
1990
2000
2010
2020
Year


Total biomass (mt)
0
2
4
6
8
10
12
1970
1980
1990
2000
2010
2020
Year


Summary biomass (mt)
0
2
4
6
8
10
12
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
2
4
6
8
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
5
10
15
20
25
30
1970
1980
1990
2000
2010
2020
Year


Summary Fishing Mortality
0.0
0.1
0.2
0.3
0.4
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
1
2
3
4
5
equilibrium
without fishing
with fishing
1970
1980
1990
2000
2010
2020
Year


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
0
2
4
6
8
0
1
2
3
4
5
Spawning biomass (mt)


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
0
2
4
6
8
0
1
2
3
4
5
Spawning biomass (mt)


Landings (mt)
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Landings (mt)
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Continuous F
0.00
0.05
0.10
0.15
1970
1980
1990
2000
2010
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0.0
0.5
1.0
1.5
2.0
2.5
3.0
Relative spawning output: B/B_MSY
1980
2000
1968
1970


Index
0.0
0.1
0.2
0.3
0.4
0.5
0.6
2016
2017
2018
2019
2020
Year


Index
0.0
0.1
0.2
0.3
0.4
0.5
0.6
2016
2017
2018
2019
2020
Year


Expected index
0.00
0.02
0.04
0.06
0.08
0.10
0.00
0.05
0.10
0.15
Observed index


Log index
−1
−2
−3
−4
−5
2016
2017
2018
2019
2020
Year


Log index
−1
−2
−3
−4
−5
2016
2017
2018
2019
2020
Year


Log expected index
−2.21
−2.23
−2.25
−3.0
−2.5
−2.0
Log observed index


Residual
1.0
0.5
0.0
−0.5
−1.0
2016
2017
2018
2019
2020
Year


Deviation
1.0
0.5
0.0
−0.5
−1.0
2016
2017
2018
2019
2020
Year


FISHERY
Standardized index
0.0
0.5
1.0
1.5
2015
2016
2017
2018
2019
2020
2021
Year


0.1
5
10
Age
15
14
13
12
11
10
9
8
7
6
5
4
3
2
1
0
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


200
600
1000
Age
15
14
13
12
11
10
9
8
7
6
5
4
3
2
1
0
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


Mean age (yr)
0
2
4
6
8
1970
1980
1990
2000
2010
2020
Year
Females
Males


0.1
5
10
Age
15
14
13
12
11
10
9
8
7
6
5
4
3
2
1
0
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


100
300
500
700
Age
15
14
13
12
11
10
9
8
7
6
5
4
3
2
1
0
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


Mean age (yr)
0
2
4
6
8
1970
1980
1990
2000
2010
2020
Year
Females
Males


Age
0
2
4
6
8
10
12
14
1970
1980
1990
2000
2010
2020
Year
0.1
0.2
0.3
0.4
0.5
0.6
0.7
0.8
0.9
1


0.1
2
60
50
40
30
20
10
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


50
150
250
Length
0
10
20
30
40
50
60
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


Mean length (cm)
0
10
20
30
40
1970
1980
1990
2000
2010
2020
Year
Females
Males


0.1
2
60
50
40
30
20
10
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


50
150
250
Length
0
10
20
30
40
50
60
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


Mean length (cm)
0
10
20
30
40
1970
1980
1990
2000
2010
2020
Year
Females
Males


0.1
0.2
0.3
0.4
0.5
0.6
0.7
0.8
0.9
Length
10
20
30
40
50
1970
1980
1990
2000
2010
2020
Year


Numbers at age at equilibrium
0
1
2
3
4
5
6
0
2
4
6
8
10
12
14
Age
Females
Males


FISHERY
Sum of N input=368.8
0.20
0.15
0.10
0.05
0.00
Proportion
20
30
40
50
Length (cm)


FISHERY
0.01
0.2
0.4
Length (cm)
51
48
45
42
39
36
33
30
27
24
21
18
15
2012
2013
2014
2015
2016
Year


2011
N input=62.2
2012
N input=72.2
2013
N input=66.4
2014
N input=49.6
2015
N input=84
2016
N input=34.4
0.00
0.10
0.20
0.30
20
30
40
50
Proportion
Length (cm)


0.01
0.2
0.4
51
45
39
33
27
21
15
Length (cm)
2011
2012
2013
2014
2015
2016
Year


FISHERY (whole catch)
Mean length
35
30
25
2011
2012
2013
2014
2015
2016
Year


FISHERY
Sum of N input=368.8
Sum of N adj.=260.5
Proportion
0.00
0.05
0.10
0.15
0.20
20
30
40
50
Length (cm)


FISHERY
−4
0.1
4
Length (cm)
51
48
45
42
39
36
33
30
27
24
21
18
15
2012
2013
2014
2015
2016
Year


2011
N input=62.2
N adj.=43.9
2012
N input=72.2
N adj.=51
2013
N input=66.4
N adj.=46.9
2014
N input=49.6
N adj.=35.1
2015
N input=84
N adj.=59.2
2016
N input=34.4
N adj.=24.5
0.00
0.10
0.20
0.30
20
30
40
50
Proportion
Length (cm)


−4
0.1
4
51
45
39
33
27
21
15
Length (cm)
2011
2012
2013
2014
2015
2016
Year


FISHERY (whole catch)
Mean length
35
30
25
2011
2012
2013
2014
2015
2016
Year


Equilibrium yield (mt)
0.0
0.2
0.4
0.6
0.8
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0.0
0.2
0.4
0.6
0.8
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
0.5
0.3
0.1
−0.1
0
2
4
6
8
10
Total biomass (mt)


Yield per recruit (kg)
0.00
0.04
0.08
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1970
1980
1990
2000
2010
2020
Year


SR_LN(R0)
prior
max. likelihood
1
2
3
4
5
ln(DM_theta)_1
−4
−2
0
2
4
6
8
10
LnQ_base_FISHERY(1)
−6
−4
−2
0
2
Size_inflection_FISHERY(1)
20
25
30
35
40
45
50
Size_95%width_FISHERY(1)
5
10
15
20
Density
Parameter value