Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Fri Jul 29 14:07:56 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
0
20
40
60
80
0
2
4
6
8
10
12
Age (yr)


Age (yr)
0
2
4
6
8
10
12
Length (cm, beginning of the year)
0
20
40
60
80
CV of lengths
0
0.02
0.04
0.06
SD of lengths
0
1
2
3
4
Length (cm, beginning of the year)
CV of lengths
0
0.02
0.04
0.06
SD of lengths
0
2
4
Age (yr)
0
2
4
6
8
10
12


Age (yr)
Mean weight
Length (cm, beginning of the year)
Mean weight
Frac. mature
Age (yr)
Frac. mature
Length (cm, beginning of the year)


Length
0
20
40
60
80
0
2
4
6
8
10
12
Age


Length
0
20
40
60
80
0
2
4
6
8
10
12
Age


Mean weight (kg) in last year
14
12
10
8
6
4
2
0
0
20
40
60
80
Length (cm)


1.0
0.8
0.6
0.4
0.2
0.0
Maturity
0
20
40
60
80
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0
2
4
6
8
10
12
Female weight (kg)


Fecundity
0
2
4
6
8
10
12
14
0
2
4
6
8
10
12
Female weight (kg)


Fecundity
0
2
4
6
8
10
12
14
0
20
40
60
80
Female length (cm)


Spawning output
14
12
10
8
6
4
2
0
0
20
40
60
80
Length (cm)


Spawning output
10
8
6
4
2
0
0
2
4
6
8
10
12
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
2
4
6
8
10
12
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
Length (cm)


Length (cm)
0
20
40
60
80
0
2
4
6
8
10
12
Age (yr)
0.8
0.6
0.4
0.2


Spawning biomass (mt)
0
50
100
150
200
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
50
100
150
200
250
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
1
2
3
4
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
1
2
3
4
5
6
1990
2000
2010
2020
Year


Total biomass (mt)
0
100
200
300
400
500
1990
2000
2010
2020
Year


Summary biomass (mt)
0
100
200
300
400
500
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
50
100
150
200
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
100
200
300
400
1990
2000
2010
2020
Year


Summary Fishing Mortality
0.000
0.004
0.008
1985
1990
1995
2000
2005
2010
2015
2020
Year


Spawning biomass (mt)
0
50
100
150
200
1985
1990
1995
2000
2005
2010
2015
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
0.2
0.0
−0.2
−0.4
1980
1990
2000
2010
2020
Year


Log recruitment deviation
1.0
0.5
0.0
−0.5
−1.0
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.1
0.2
0.3
0.4
0.5
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
1.0
0.5
0.0
$1 - SE(\hat{r_y})^2/\sigma_R^2$
1980
1990
2000
2010
2020
Year


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
0
50
100
150
200
0
20
40
60
80
100
Spawning biomass (mt)


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
1986
2021
Recruitment (1,000s)
0
50
100
150
200
0
20
40
60
80
100
Spawning biomass (mt)


Log recruitment deviation
0.4
0.2
0.0
−0.2
−0.4
1986
2021
0.0
0.2
0.4
0.6
0.8
1.0
1.2
Spawning output (relative to B_0)


Landings (mt)
0.0
0.5
1.0
1.5
2.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


Landings (mt)
0.0
0.5
1.0
1.5
2.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


FISHERY
FISHERY obs.
Observed and expected Landings (mt)
0.0
0.5
1.0
1.5
2.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


Continuous F
0.000
0.002
0.004
0.006
1985
1990
1995
2000
2005
2010
2015
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0
1
2
3
4
Relative spawning output: B/B_MSY
1986
1990


Index
0
5
10
15
20
25
30
35
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
5
10
15
20
25
30
35
1990
1995
2000
2005
2010
2015
2020
Year


Expected index
0.0
0.5
1.0
1.5
2.0
2.5
3.0
0
2
4
6
8
10
Observed index


Log index
3
2
1
0
−1
−2
1990
1995
2000
2005
2010
2015
2020
Year


Log index
3
2
1
0
−1
−2
1990
1995
2000
2005
2010
2015
2020
Year


Log expected index
0.80
0.90
1.00
1.10
0.0
0.5
1.0
1.5
2.0
Log observed index


Residual
2
1
0
−1
−2
1990
1995
2000
2005
2010
2015
2020
Year


Deviation
1.0
0.5
0.0
−0.5
−1.0
1990
1995
2000
2005
2010
2015
2020
Year


FISHERY
Standardized index
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1990
1995
2000
2005
2010
2015
2020
Year


50
150
250
Age
12
11
10
9
8
7
6
5
4
3
2
1
0
1985
1989
1993
1997
2001
2005
2009
2013
2017
2021
Year


50
150
Age
12
11
10
9
8
7
6
5
4
3
2
1
0
1985
1989
1993
1997
2001
2005
2009
2013
2017
2021
Year


20
60
100
140
Length
0
20
40
60
80
1986
1990
1994
1998
2002
2006
2010
2014
2018
2022
Year


10
30
50
70
Length
0
20
40
60
80
1986
1990
1994
1998
2002
2006
2010
2014
2018
2022
Year


Numbers at age at equilibrium
0
50
100
150
0
2
4
6
8
10
12
Age


FISHERY
Sum of N adj.=132.7
Proportion
0.30
0.25
0.20
0.15
0.10
0.05
0.00
10
20
30
40
50
60
70
80
Length (cm)


FISHERY
0.01
0.2
0.4
Length (cm)
75
70
65
60
55
50
45
40
35
30
25
20
15
2009
2015
2017
Year


2007
N adj.=37.3
2009
N adj.=34.8
2015
N adj.=37.7
2017
N adj.=22.8
0.00
0.10
0.20
0.30
10
30
50
70
Proportion
Length (cm)


0.01
0.2
0.4
75
65
55
45
35
25
15
Length (cm)
2007
2009
2015
2017
Year


FISHERY (whole catch)
50
45
40
Mean length
2008
2010
2012
2014
2016
Year


FISHERY
Sum of N adj.=132.7
Sum of N eff.=141
Proportion
0.00
0.05
0.10
0.15
0.20
0.25
0.30
10
20
30
40
50
60
70
80
Length (cm)


FISHERY
−2
0.1
2
Length (cm)
75
70
65
60
55
50
45
40
35
30
25
20
15
2009
2015
2017
Year


2007
N adj.=37.3
N eff.=39
2009
N adj.=34.8
N eff.=20.7
2015
N adj.=37.7
N eff.=22.1
2017
N adj.=22.8
N eff.=59.2
0.30
0.20
0.10
0.00
10
30
50
70
Proportion
Length (cm)


−2
0.1
2
75
65
55
45
35
25
15
Length (cm)
2007
2009
2015
2017
Year


Effective sample size
0
10
30
50
0
10
20
30
40
Observed sample size
2017
2007
2015
2009
harmonic mean
arithmetic mean


FISHERY (whole catch)
50
45
40
Mean length
2008
2010
2012
2014
2016
Year


Equilibrium yield (mt)
0
10
20
30
40
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0
10
20
30
40
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
60
40
20
0
−20
0
100
200
300
400
500
Total biomass (mt)


Yield per recruit (kg)
0.000
0.004
0.008
0.012
1985
1990
1995
2000
2005
2010
2015
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1985
1990
1995
2000
2005
2010
2015
2020
Year


Catches
Abundance indices
Length compositions
FISHERY
FISHERY
FISHERY
1985
1990
1995
2000
2005
2010
2015
2020
Year


SR_LN(R0)
prior
max. likelihood
2
4
6
8
10
12
14
Size_inflection_FISHERY(1)
30
35
40
45
50
55
60
InitF_seas_1_flt_1FISHERY
0.0
0.1
0.2
0.3
0.4
0.5
Size_95%width_FISHERY(1)
5
10
15
20
25
LnQ_base_FISHERY(1)
−4
−2
0
2
4
Q_extraSD_FISHERY(1)
0.0
0.5
1.0
1.5
2.0
Density
Parameter value