American Samoa Model Checks

2022-08-25

This is a summary report for the LERU base model run.

Model Output

Input Data





Convergence Check

```
  Converged     MaxGrad
1      TRUE 1.95184e-05
```

```
[1] "1 NOTE:  Max data length bin: 38.5  < max pop len bins: 43; so will accumulate larger po
[2] "2 parameter init value is greater than parameter max 0.8 > 0.6 for parm: 14 ; search for
[3] "N warnings: 2"
```

Fit to Model

CPUE

```
Residual Runs Test (/w plot) stats by Index:
```

```
RMSE stats by Index:
```


Index
0
2
4
6
8
10
12
2016
2018
2020
Year
Index residuals
−0.6
−0.2
0.2
0.4
0.6
FISHERY
2016
2018
2020
Year


Index residuals
−0.4
−0.2
0.0
0.2
0.4
RMSE = 24.6%
FISHERY
Loess
2017
2018
2019
2020
2021
Year

Length Comp

#Factor	Fleet	New_Var_adj	Type	Name
4	1	0.216751	len	FISHERY

```
Residual Runs Test (/w plot) stats by Mean length:

    Index runs.p   test   sigma3.lo  sigma3.hi type
1 FISHERY  0.176 Passed -0.03666048 0.03666048  len

RMSE stats by Index:

# A tibble: 2 x 3
  Fleet    RMSE.perc  Nobs
  <chr>        <dbl> <int>
1 FISHERY          2    18
2 Combined         2    18
```



Retrospective and Hindcasting


FISHERY (whole catch)
Mean length
29
28
27
26
25
24
2005
2010
2015
2020
Year


FISHERY
Sum of N adj.=1180.7
Sum of N eff.=1359.7
Proportion
0.4
0.3
0.2
0.1
0.0
15
20
25
30
35
40
Length (cm)



Retrospective

`Mohn's Rho stats, including one step ahead forecasts:`



```
Mohn's Rho stats, including one step ahead forecasts:

  type     peel          Rho  ForecastRho
1    F     2020  0.004520378  0.010143632
2    F     2019 -0.032900248 -0.012529282
3    F     2018  0.015369773  0.016169683
4    F Combined -0.004336699  0.004594677
```

Hindcasting

```
Plotting Hindcast Cross-Validation (one-step-ahead)

 Computing MASE with only 2 of 3  prediction residuals for Index FISHERY

 Warning:  Unequal spacing of naive predictions residuals may influence the interpretation o
```




```
MASE stats by Index:
Plotting Hindcast Cross-Validation (one-step-ahead)

 Computing MASE with only 2 of 3  prediction residuals for Index FISHERY
```

```
 Warning:  Unequal spacing of naive predictions residuals may influence the interpretation o
```

```
MASE stats by Index:
```



Recruitment Deviations

Likelihood Profile

```
[1] "SR_LN"
```

	frac_change	include	label
TOTAL	1.0000	TRUE	Total
Catch	0.9133	TRUE	Catch
Equil_catch	0.0000	FALSE	Equilibrium catch
Survey	0.0220	TRUE	Index data
Length_comp	0.0883	TRUE	Length data
Recruitment	0.0000	FALSE	Recruitment
InitEQ_Regime	0.0000	FALSE	Initital equilibrium recruitment
Forecast_Recruitment	0.0000	FALSE	Forecast recruitment
Parm_priors	0.0000	FALSE	Priors

```
Parm_softbounds          0.0000   FALSE                    Soft bounds
Parm_devs                0.0000   FALSE          Parameter deviations
Crash_Pen                0.0000   FALSE                  Crash penalty
```





Management Quantities

starter.sso with Bratio: SSB/SSBMSY and F: _abs_F




MVLN
SSB/SSB$_{MSY}$
0.0
1.0
2.0
3.0
1970
1980
1990
2000
2010
2020
Year


MVLN
F/F$_{MSY}$
0
2
4
6
8
10
1970
1980
1990
2000
2010
2020
Year






```
null device
          1
```

Jitter


93.400
93.375
93.350
93.325
Likelihood
1.0
1.5
2.0
2.5
3.0
Run


Spawning biomass (t)
0
5
10
15
20
1970
1980
1990
2000
2010
2020
Year
model 1
model 2
model 3


abs F; with F=sum(full Fs)
0.0
0.5
1.0
1.5
2.0
2.5
1970
1980
1990
2000
2010
2020
Year
model 1
model 2
model 3


Age−0 recruits (1,000s)
0
10
20
30
40
1970
1980
1990
2000
2010
2020
Year
model 1
model 2
model 3