Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Mon Aug 08 10:03:11 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
0
10
20
30
40
0
2
4
6
8
10
12
14
Age (yr)


Age (yr)
0
2
4
6
8
10
12
14
Length (cm, beginning of the year)
0
10
20
30
40
CV of lengths
0
0.04
0.08
SD of lengths
0.0
1.0
2.0
3.0


Age (yr)
Mean weight
Length (cm, beginning of the year)
Mean weight
Frac. mature
Age (yr)
0
2
4
6
8
10
12
14
0.5
1
20
30
40
0.0


Length
0
10
20
30
40
0
5
10
15
Age


Length
0
10
20
30
40
0
5
10
15
Age


Mean weight (kg) in last year
1.5
1.0
0.5
0.0
0
10
20
30
40
Length (cm)


1.0
0.8
0.6
0.4
0.2
0.0
Maturity
0
10
20
30
40
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0.0
0.5
1.0
1.5
Female weight (kg)


Fecundity
0.0
0.5
1.0
1.5
0.0
0.5
1.0
1.5
Female weight (kg)


Fecundity
0.0
0.5
1.0
1.5
0
10
20
30
40
Female length (cm)


Spawning output
1.5
1.0
0.5
0.0
0
10
20
30
40
Length (cm)


0.8
0.6
0.4
0.2
0.0
Spawning output
0
5
10
15
Age (yr)


Hermaphroditism transition rate
1.0
0.8
0.6
0.4
0.2
0.0
0
2
4
6
8
10
12
14
Year


Fraction females by age at equilibrium
1.0
0.8
0.6
0.4
0.2
0.0
2
4
6
8
10
12
14
16
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
5
10
15
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
Length (cm)


Length (cm)
0
10
20
30
40
1
0.8
0.6
0.4
0.2
0
5
10
15
Age (yr)


Spawning biomass (mt)
0
5
10
15
20
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
5
10
15
20
25
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Total biomass (mt)
0
5
10
15
20
1970
1980
1990
2000
2010
2020
Year


Summary biomass (mt)
0
5
10
15
20
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
10
20
30
40
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
20
40
60
80
100
1970
1980
1990
2000
2010
2020
Year


Summary Fishing Mortality
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
5
10
15
20
25
1970
1980
1990
2000
2010
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
0.5
0.0
−0.5
1970
1980
1990
2000
2010
2020
Year


Log recruitment deviation
1.5
1.0
0.5
−0.5
−1.5
1970
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.2
0.4
0.6
1970
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
$1 - SE(\hat{r_y})^2/\sigma_R^2$
1.0
0.6
0.2
−0.2
1970
1980
1990
2000
2010
2020
Year


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
0
10
20
30
40
50
0
2
4
6
8
10
Spawning biomass (mt)


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
2005
2018
2021
1967
2002
Recruitment (1,000s)
Spawning biomass (mt)
0
10
20
30
40
50
0
2
4
6
8
10


Log recruitment deviation
0.5
0.0
−0.5
2005
2018
2021
1967
2002
0.0
0.2
0.4
0.6
0.8
1.0
Spawning output (relative to B_0)


Landings (mt)
0
1
2
3
4
5
6
7
1970
1980
1990
2000
2010
2020
Year


Landings (mt)
0
1
2
3
4
5
6
7
1970
1980
1990
2000
2010
2020
Year


Continuous F
0.0
0.5
1.0
1.5
1970
1980
1990
2000
2010
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
Relative spawning output: B/B_MSY
2010
1990
1980
2000
2020
2021
1967
1970


Index
0
2
4
6
8
10
12
14
2016
2017
2018
2019
2020
2021
Year


Index
0
2
4
6
8
10
12
14
2016
2017
2018
2019
2020
2021
Year


Expected index
Observed index
0
1
2
3
4
5


Log index
0.5
1.0
1.5
2.0
2.5
2016
2017
2018
2019
2020
2021
Year


Log index
0.5
1.0
1.5
2.0
2.5
2016
2017
2018
2019
2020
2021
Year


Log expected index
0.6
0.8
1.0
1.2
1.4
1.6
1.0
1.2
1.4
1.6
Log observed index


Residual
1
0
−1
2016
2017
2018
2019
2020
2021
Year


Deviation
0.4
0.2
0.0
−0.2
−0.4
2016
2017
2018
2019
2020
2021
Year


FISHERY
Standardized index
0.0
0.5
1.0
1.5
2015
2016
2017
2018
2019
2020
2021
2022
Year


10
30
50
Age
15
14
13
12
11
10
9
8
7
6
5
4
3
2
1
0
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


10
30
Age
15
14
13
12
11
10
9
8
7
6
5
4
3
2
1
0
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


5
15
25
35
Length
0
10
20
30
40
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


2
6
10
14
Length
0
10
20
30
40
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


Numbers at age at equilibrium
0
5
10
15
20
25
30
0
2
4
6
8
10
12
14
Age


FISHERY
Sum of N adj.=6477
Proportion
0.0
0.1
0.2
0.3
0.4
15
20
25
30
35
40
Length (cm)


FISHERY
0.01
0.5
1
38.5
35.0
31.5
28.0
24.5
21.0
17.5
14.0
Length (cm)
1992
2004
2008
2012
2016
2020
Year


1992
N adj.=1
2009
N adj.=331.8
2015
N adj.=542.6
2021
N adj.=79.8
2004
N adj.=149
2010
N adj.=104.2
2016
N adj.=247
2005
N adj.=125.2
2011
N adj.=782
2017
N adj.=110.9
2006
N adj.=99.1
2012
N adj.=1329.7
2018
N adj.=143.6
2007
N adj.=81.5
2013
N adj.=1564.9
2019
N adj.=189.8
2008
N adj.=250.5
2014
N adj.=190.7
2020
N adj.=153.7
Proportion
0.0
0.1
0.2
0.3
0.4
15
20
25
30
35
40
Length (cm)


0.01
0.5
1
38.5
35.0
31.5
28.0
24.5
21.0
17.5
14.0
Length (cm)
1992
2004
2008
2012
2016
2020
Year


FISHERY (whole catch)
Mean length
35
30
25
20
15
1995
2000
2005
2010
2015
2020
Year


FISHERY
Sum of N adj.=6477
Sum of N eff.=1625.3
Proportion
0.0
0.1
0.2
0.3
0.4
15
20
25
30
35
40
Length (cm)


FISHERY
−25
−15
5
15
25
38.5
35.0
31.5
28.0
24.5
21.0
17.5
14.0
Length (cm)
1992
2004
2008
2012
2016
2020
Year


1992
N adj.=1
N eff.=1.5
2009
N adj.=331.8
N eff.=76
2015
N adj.=542.6
N eff.=173.2
2021
N adj.=79.8
N eff.=79.5
2004
N adj.=149
N eff.=18.6
2010
N adj.=104.2
N eff.=43.2
2016
N adj.=247
N eff.=126.4
2005
N adj.=125.2
N eff.=23.4
2011
N adj.=782
N eff.=51.4
2017
N adj.=110.9
N eff.=280.2
2006
N adj.=99.1
N eff.=22.2
2012
N adj.=1329.7
N eff.=158.6
2018
N adj.=143.6
N eff.=215.9
2007
N adj.=81.5
N eff.=36.4
2013
N adj.=1564.9
N eff.=132.2
2019
N adj.=189.8
N eff.=32.9
2008
N adj.=250.5
N eff.=35.8
2014
N adj.=190.7
N eff.=54.7
2020
N adj.=153.7
N eff.=63.3
0.0
0.1
0.2
0.3
0.4
15
20
25
30
35
40
Proportion
Length (cm)


−25
−15
5
15
25
38.5
35.0
31.5
28.0
24.5
21.0
17.5
14.0
Length (cm)
1992
2004
2008
2012
2016
2020
Year


Effective sample size
0
50
150
250
0
500
1000
1500
Observed sample size
2017
2018
2015
2012
2013
2016
2021
2009
2020
2014
2011
2019
2008
2004
1992
arithmetic mean
harmonic mean


FISHERY (whole catch)
Mean length
35
30
25
20
15
1995
2000
2005
2010
2015
2020
Year


Equilibrium yield (mt)
0.0
0.5
1.0
1.5
2.0
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0.0
0.5
1.0
1.5
2.0
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
0.0
1.0
2.0
3.0
0
5
10
15
20
Total biomass (mt)


Yield per recruit (kg)
0.00
0.04
0.08
0.12
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


SR_LN(R0)
prior
max. likelihood
2
3
4
5
6
7
8
Size_inflection_FISHERY(1)
16
18
20
22
24
InitF_seas_1_flt_1FISHERY
0.00
0.05
0.10
0.15
0.20
0.25
0.30
Size_95%width_FISHERY(1)
1
2
3
4
5
6
7
8
Density
LnQ_base_FISHERY(1)
−2
−1
0
1
2
Q_extraSD_FISHERY(1)
0.0
0.5
1.0
1.5
2.0
Parameter value