

Internship AWI

Author: Leonhard Schatt
Supervisor: Felix Pithan

September 15, 2023

Contents

1	Introduction	5
2	Theory	6
3	Discussion	7
3.1	Diurnal temperature range	7
3.2	Robustness	8
4	Summary	11
A	Append	12
A.1	Robustness	12
	Bibliography	15

1 Introduction

The diurnal temperature range (DTR), the difference between daily minimum temperatures (T_{\min}) and daily maximum temperatures (T_{\max}), decreases on a global scale and in many regional scales since the 1990s [7, 9, 2]. This decrease is caused by the mean surface minimum temperatures rising faster than the maximum temperatures [9]. The reasons for this dynamic are quantitatively not fully understood yet, but the asymmetry in warming has been linked to factors of external forcing such as greenhouse gases, aerosols, and changes in landscape due to land use, as well as secondary effects such as changes in cloudiness and relative humidity [16, 10, 11, 5, text]. The extent of change in DTR is of magnitudes smaller than the trends in mean surface temperatures (T_{mean}). The uncertainty in these trends surpasses that of surface mean temperature, and they were only assessed with medium confidence in the fourth and fifth IPCC reports [1].

In this work, we examine the DTR for the Arctic region. The mean surface temperatures in the Arctic warm more rapidly than in any other region on Earth due to Arctic amplification [12]. At the same time, the Arctic shows great climate variability, which decreases the signal-to-noise ratio (SNR) of the trends in mean surface temperature [8]. As there is no linear coupling between the diurnal temperature range and the mean temperature, the DTR might be less influenced by the high climate variability. In the following, we examine whether the trends in the diurnal temperature range show a better SNR in the Arctic region than the mean surface temperature. This could lead to a more robust parameter to track changes in the polar regions.

In section (sec1), we investigate factors causing the trends in the Diurnal Temperature Range (DTR) at the AWIPEV meteorological station in Spitsbergen. Since the average temperatures in Antarctica have not changed despite climate change, in section (ref), we address the question of whether there have been shifts in the daily temperature range.

2 Theory

3 Discussion

3.1 Diurnal temperature range

In the following section, the diurnal temperature range (DTR) is investigated. The CRU TS data is analyzed by month and latitude to determine trends.




Figure 3.1: Diurnal temperature range change (a) over 50 years for the entire polar region (66-90°). (b) shows, for comparison, the maximum and minimum temperature trends. The variance for these temperatures is much larger compared to (a). (c) presents the signal-to-noise ratio for the two previous plots.

After taking the over all average of the weighted data¹, we investigate the temporal development of DTR monthly average using linear regression. To examine the significance of the trend, the empirical variance in relation to the fit is calculated.

¹The data are weighted by area. This leads to a stronger emphasis on the "lower" latitudes. The effect of the weighting is decreasingly small at the intervals of 5° chosen later on.

3 Discussion

As shown in Fig. 3.1, the changes in DTR are significant. But this is also true for the changes in the minimum temperature T_{\min} as for the maximum temperature T_{\max} . Compared to the changes in mean min./max. temperature the DTR trend does have a bigger SNR, as shown in Fig. 3.1c. This can be explained through the strong correlation between maximum and minimum temperature (see Fig. 3.1d). Due to this higher SNR, the changes in DTR could be an advantageous indicator for climatic changes in regions with high climate variability. Therefore, the robustness of this parameter and trends are investigated in Section 3.2.

3.2 Robustness of the DTR for subgroups of data

The mathematical derivation of uncertainty is too complex for this internship. To gain a sense of how reliable the effect of DTR change is, we compare the results of the left and right hemisphere – the hemisphere from -180 to 0 and 0 to 180 degrees – and further latitude subsets.

As depicted in Figure 3.2, the Diurnal Temperature Range (DTR) exhibits significant variations, particularly at higher latitudes. This variability may be attributed to the data selection process. The dataset encompasses the geographic region illustrated in Figure 3.3. When segregating the data into left and right hemispheres, it becomes apparent that the distribution of landmass is uneven. The left half of the Earth predominantly comprises mainland, whereas the right hemisphere consists mainly of islands. Consequently, temperature variations are influenced by the geographical circumstances, with islands experiencing unique factors such as ice formation, which influence their climate differently.

As shown in Figs. A.1 and A.2, the temperatures and their trends differ significantly for the two areas. The mainland experiences lower temperatures than the island dominated areas, especially for the areas of higher latitude. To check whether the data is robust, we have to compare two equivalent areas. Therefore, Greenland is divided into two halves and the process is repeated.

3.2 Robustness




Figure 3.2: Subfigure with Two Columns






Figure 3.3: Data coverage of the CRU data.



4 Summary

A Append

A.1 Robustness





(a) T_{min} for the left hemisphere between 66 and 70° (b) T_{min} for the right hemisphere between 66 and 70°





(c) T_{min} between 70 and 75°

(d) T_{min} between 70 and 75°



(e) T_{min} between 75 and 80°

(f) T_{min} between 75 and 80°




(g) T_{min} between 80 and 82°


(h) T_{min} between 80 and 82°

Figure A.1: Temperature for left and right hemisphere

A Append




(a) T_{max} for the left hemisphere between 66 and 70° (b) T_{max} for the right hemisphere between 66 and 70°




(c) T_{max} between 70 and 75°

(d) T_{max} between 70 and 75°



(e) T_{max} between 75 and 80°

(f) T_{max} between 75 and 80°



(g) T_{max} between 80 and 82°

(h) T_{max} between 80 and 82°

Bibliography

- [1] Observations: Atmosphere and Surface. In Intergovernmental Panel on Climate Change, editor, *Climate Change 2013 – The Physical Science Basis*, pages 159–254. Cambridge University Press, 2014.
- [2] L. V. Alexander, X. Zhang, T. C. Peterson, J. Caesar, B. Gleason, A. M. G. Klein Tank, M. Haylock, D. Collins, B. Trewin, F. Rahimzadeh, A. Tagipour, K. Rupa Kumar, J. Revadekar, G. Griffiths, L. Vincent, D. B. Stephenson, J. Burn, E. Aguilar, M. Brunet, M. Taylor, M. New, P. Zhai, M. Rusticucci, and J. L. Vazquez-Aguirre. Global observed changes in daily climate extremes of temperature and precipitation. *Journal of Geophysical Research*, 111(D5), 2006.
- [3] Randall S. Cerveny and Robert C. Balling. The impact of snow cover on diurnal temperature range. *Geophysical Research Letters*, 19(8):797–800, 1992.
- [4] Intergovernmental Panel on Climate Change, editor. *Climate Change 2013 – The Physical Science Basis*. Cambridge University Press, 2014.
- [5] Nikolaos Christidis and Peter A. Stott. Attribution analyses of temperature extremes using a set of 16 indices. *Weather and Climate Extremes*, 14:24–35, 2016.
- [6] C. Derksen and R. Brown. Spring snow cover extent reductions in the 2008-2012 period exceeding climate model projections. *Geophysical Research Letters*, 39(19):n/a–n/a, 2012.
- [7] David R. Easterling, Briony Horton, Philip D. Jones, Thomas C. Peterson, Thomas R. Karl, David E. Parker, M. James Salinger, Vyacheslav Razuvayev, Neil Plummer, Paul Jamason, and Christopher K. Folland. Maximum and Minimum Temperature Trends for the Globe. *Science*, 277(5324):364–367, 1997.
- [8] Katharina Hartmuth, Lukas Papritz, Maxi Boettcher, and Heini Wernli. Arctic Seasonal Variability and Extremes, and the Role of Weather Systems in a Changing Climate. *Geophysical Research Letters*, 50(8):e2022GL102349, 2023.
- [9] Thomas R. Karl, Richard W. Knight, Kevin P. Gallo, Thomas C. Peterson, Philip D. Jones, George Kukla, Neil Plummer, Vyacheslav Razuvayev, Janette Lindseay, and Robert J. Charlson. A New Perspective on Recent Global Warming: Asymmetric Trends of Daily Maximum and Minimum Temperature. *Bulletin of the American Meteorological Society*, 74(6):1007–1023, 1993.

Bibliography

- [10] Sophie C. Lewis and David J. Karoly. Evaluation of Historical Diurnal Temperature Range Trends in CMIP5 Models. *Journal of Climate*, 26(22):9077–9089, 2013.
- [11] Lin Liu, Zhanqing Li, Xin Yang, Hainan Gong, Chao Li, and Anyuan Xiong. The long-term trend in the diurnal temperature range over Asia and its natural and anthropogenic causes. *Journal of Geophysical Research: Atmospheres*, 121(7):3519–3533, 2016.
- [12] Felix Pithan and Thorsten Mauritsen. Arctic amplification dominated by temperature feedbacks in contemporary climate models. *Nature Geoscience*, 7(3):181–184, 2014.
- [13] Michael Qu, Joe Wan, and Xianjun Hao. Analysis of diurnal air temperature range change in the continental United States. *Weather and Climate Extremes*, 4:86–95, 2014.
- [14] Susan Solomon. *CLIMATE change 2007. The physical science Basis*. Cambridge University Press, New York, 2007.
- [15] D. Stone and A. Weaver. Factors contributing to diurnal temperature range trends in twentieth and twenty-first century simulations of the CCCma coupled model. *Climate Dynamics*, 20(5):435–445, 2003.
- [16] Liming Zhou, Robert E. Dickinson, Aiguo Dai, and Paul Dirmeyer. Detection and attribution of anthropogenic forcing to diurnal temperature range changes from 1950 to 1999: comparing multi-model simulations with observations. *Climate Dynamics*, 35(7-8):1289–1307, 2010.