
Brouillon 4

Léo Burgund

May 23, 2025

1 Nomenclature

1.1 Dimensions

- b Mini-batch size
- d_e Embedding dimension
- d_s Sequence length
- d_k Query/Keys dimension
- d_v Value dimension
- h Number of heads

We make the hypothesis that $d_k < d_e < d_s$.

1.2 Matrix operations in a self-attention block

In the case of multi head attention, for each head $i = 1, \dots, h$, we have:

- Input $X \in \mathbb{R}^{d_s \times d_e}$
- $W_{Q_i} \in \mathbb{R}^{d_e \times \frac{d_k}{h}}$, $Q_i := XW_{Q_i} \in \mathbb{R}^{d_s \times \frac{d_k}{h}}$
- $W_{K_i} \in \mathbb{R}^{d_e \times \frac{d_k}{h}}$, $K_i := XW_{K_i} \in \mathbb{R}^{d_s \times \frac{d_k}{h}}$
- $S_i := \frac{Q_i K_i^\top}{\sqrt{\frac{d_k}{h}}} \in \mathbb{R}^{d_s \times d_s}$
- $A_i := \text{softmax}_{\text{row}}(S)$
- $W_{V_i} \in \mathbb{R}^{d_e \times \frac{d_v}{h}}$, $V_i := XW_{V_i} \in \mathbb{R}^{d_s \times \frac{d_v}{h}}$
- $H_i := A_i V_i \in \mathbb{R}^{d_s \times \frac{d_v}{h}}$, $H = [H_1, \dots, H_h] \in \mathbb{R}^{d_s \times d_v}$
- $W_O \in \mathbb{R}^{d_v \times d_e}$
- Output $Y := HW_O + X \in \mathbb{R}^{d_s \times d_e}$

For now, we study the case $h = 1$.

🔥 We omit the $\frac{1}{\sqrt{d_k}}$ scaling for the S matrix, it can cause problem with growing, so for growing we will make it a learnable parameter (and initialize it at $\frac{1}{\sqrt{d_{k_{\text{initial}}}}}$?). Or maybe scale with $\left(\frac{\sqrt{d_k}}{\sqrt{d_k+p}}\right)$?

2 Problem

Goal:

$$\min_f \mathcal{L}(f).$$

We will study the variations of the loss made by the variations of S , with other parameters fixed. Hence we will study

$$\arg \min_S \mathcal{L}(S)$$

Let $G := \nabla_S \mathcal{L}(S)$.

We have

$$\text{rk}(G) \leq d_s, \quad \text{rk}(S) \leq d_k$$

We have the first order approximation, with the introduction of $\gamma \in \mathbb{R}_+^*$, similar to a step size:

$$\mathcal{L}(S + \gamma dS) = \mathcal{L}(S) + \gamma \langle G, dS \rangle_F + o(\|dS\|_F)$$

We consider the problem

$$\arg \min_{dS} \langle G, dS \rangle_F \quad \text{s.t.} \quad \|dS\|_F \leq \gamma \wedge \text{rk}(dS) \leq d_k < \text{rk}(G)$$

Let $Z = W_Q W_K^\top$, with $\text{rk}(Z) \leq d_k$, we then have

$$\begin{aligned} S &= X W_Q W_K^\top X^\top \\ &= X Z X^\top \end{aligned}$$

and

$$\begin{aligned} dS &= X (W_{Q_{+1}} W_{K_{+1}}) X^\top - X W_Q W_K^\top X^\top \\ &= X (Z + dZ) X^\top - X Z X^\top \\ &= X dZ X^\top \end{aligned}$$

Moreover,

$$\text{rk}(dS) = \text{rk}(X dZ X^\top) = \text{rk}(dZ) \leq d_k < d_e < d_s$$

Hence, the problem becomes

$$\begin{aligned} dZ^* &= \arg \min_{dZ} \langle G, X dZ X^\top \rangle_F \quad \text{s.t.} \quad \|X dZ X^\top\|_F \leq \gamma \\ &= - \arg \max_{dZ} \langle G, X dZ X^\top \rangle_F \quad \text{s.t.} \quad \|X dZ X^\top\|_F \leq \gamma \\ &= - \arg \max_{dZ} \langle G, X dZ X^\top \rangle_F \quad \text{s.t.} \quad \|X dZ X^\top\|_F = \gamma \\ &= -\gamma \arg \max_{dZ} \langle G, X dZ X^\top \rangle_F \quad \text{s.t.} \quad \|X dZ X^\top\|_F = 1 \\ &= -\frac{\gamma}{\alpha} \arg \min_{dZ} \|G - X dZ X^\top\|_F^2 \end{aligned}$$

(*) We make the hypothesis that we can always find a $\langle G, X dZ X^\top \rangle_F > 0$.

🔥 Justification for α , $\alpha = \|\cdot\|_F$

Let $\lambda := \frac{\gamma}{\alpha}$, $P := \arg \min_{dZ} \|G - X dZ X^\top\|_F^2$.

We will first search P , then λ with a line search.

2.1 Find P

Let

$$f(P) = \|G - X P X^\top\|_F^2$$

f is convex.

We have

$$\nabla_P f = -2X^\top (G - X P X^\top) X$$

so

$$\nabla_P f = 0 \iff X^\top X P X^\top X = X^\top G X$$

🔥 Better way to find P by considering $X^\top (X P X^\top - G) X = 0$?

Under the hypothesis that $X^\top X$ is invertible ($\text{rk}(X) = d_e, d_e < d_s$), we have the solution

$$\begin{aligned} P^\star &= (X^\top X)^{-1} X^\top G X (X^\top X)^{-1} \\ &= (X^\top X)^+ X^\top G X (X^\top X)^+ \\ &= X^+ G (X^+)^{\top} \end{aligned}$$

🔥 Which formula to implement for P^\star ?

To account for the batch, there are several ways to average:

$$\begin{aligned} \mathbb{E}_X \left[(X^\top X)^+ X^\top G X (X^\top X)^+ \right] &= \mathbb{E}_X \left[X^+ G (X^+)^{\top} \right] \\ &= \mathbb{E}_X [X^\top X]^+ \mathbb{E}_X [X^\top G X] \mathbb{E}_X [X^\top X]^+ \\ &= \mathbb{E}_X [X]^+ \mathbb{E}_X [G] (\mathbb{E}_X [X])^+{}^{\top} \\ &= \mathbb{E}_X [X^\top X]^{-1} \mathbb{E}_X [X^\top G X] \mathbb{E}_X [X^\top X]^{-1} \end{aligned}$$

🔥 Which batch average?

2.2 Line search

Do a line search to find λ

2.2.1 “Normal” way

Line search on

$$\mathcal{L} \left(X \text{ SVD}_{d_k+p} \left(W_Q^t W_K^{t^\top} - \lambda P \right) X^\top \right)$$

2.2.2 Testing fast way

Lose the rank constraint, but lose the SVD so faster. Get a good approximation of λ ?

$$\mathcal{L} \left(X \left(W_Q^t W_K^{t^\top} - \lambda P \right) X^\top \right) = \mathcal{L} \left(X W_Q^t W_K^{t^\top} X^\top - \lambda X P X^\top \right)$$