

---

# Brouillon 4

---

Léo Burgund

May 23, 2025

## 1 Nomenclature

### 1.1 Dimensions

- $b$  Mini-batch size
- $d_e$  Embedding dimension
- $d_s$  Sequence length
- $d_k$  Query/Keys dimension
- $d_v$  Value dimension
- $h$  Number of heads

We make the hypothesis that  $d_k < d_e < d_s$ .

### 1.2 Matrix operations in a self-attention block

In the case of multi head attention, for each head  $i = 1, \dots, h$ , we have:

- Input  $X \in \mathbb{R}^{d_s \times d_e}$
- $W_{Q_i} \in \mathbb{R}^{d_e \times \frac{d_k}{h}}$ ,  $Q_i := XW_{Q_i} \in \mathbb{R}^{d_s \times \frac{d_k}{h}}$
- $W_{K_i} \in \mathbb{R}^{d_e \times \frac{d_k}{h}}$ ,  $K_i := XW_{K_i} \in \mathbb{R}^{d_s \times \frac{d_k}{h}}$
- $S_i := \frac{Q_i K_i^\top}{\sqrt{\frac{d_k}{h}}} \in \mathbb{R}^{d_s \times d_s}$
- $A_i := \text{softmax}_{\text{row}}(S)$
- $W_{V_i} \in \mathbb{R}^{d_e \times \frac{d_v}{h}}$ ,  $V_i := XW_{V_i} \in \mathbb{R}^{d_s \times \frac{d_v}{h}}$
- $H_i := A_i V_i \in \mathbb{R}^{d_s \times \frac{d_v}{h}}$ ,  $H = [H_1, \dots, H_h] \in \mathbb{R}^{d_s \times d_v}$
- $W_O \in \mathbb{R}^{d_v \times d_e}$
- Output  $Y := HW_O + X \in \mathbb{R}^{d_s \times d_e}$

For now, we study the case  $h = 1$ .

🔥 We omit the  $\frac{1}{\sqrt{d_k}}$  scaling for the  $S$  matrix, it can cause problem with growing, so for growing we will make it a learnable parameter (and initialize it at  $\frac{1}{\sqrt{d_{k_{\text{initial}}}}}$  ?). Or maybe scale with  $\left(\frac{\sqrt{d_k}}{\sqrt{d_k+p}}\right)$ ?

## 2 Problem

Goal:

$$\min_f \mathcal{L}(f).$$

We will study the variations of the loss made by the variations of  $S$ , with other parameters fixed. Hence we will study

$$\arg \min_S \mathcal{L}(S)$$

Let  $G := \nabla_S \mathcal{L}(S)$ .

We have

$$\text{rk}(G) \leq d_s, \quad \text{rk}(S) \leq d_k$$

We have the first order approximation, with the introduction of  $\gamma \in \mathbb{R}_+^*$ , similar to a step size:

$$\mathcal{L}(S + \gamma dS) = \mathcal{L}(S) + \gamma \langle G, dS \rangle_F + o(\|dS\|_F)$$

We consider the problem

$$\arg \min_{dS} \langle G, dS \rangle_F \quad \text{s.t.} \quad \|dS\|_F \leq \gamma \wedge \text{rk}(dS) \leq d_k < \text{rk}(G)$$

Let  $Z = W_Q W_K^\top$ , with  $\text{rk}(Z) \leq d_k$ , we then have

$$\begin{aligned} S &= X W_Q W_K^\top X^\top \\ &= X Z X^\top \end{aligned}$$

and

$$\begin{aligned} dS &= X (W_{Q_{+1}} W_{K_{+1}}) X^\top - X W_Q W_K^\top X^\top \\ &= X (Z + dZ) X^\top - X Z X^\top \\ &= X dZ X^\top \end{aligned}$$

Moreover,

$$\text{rk}(dS) = \text{rk}(X dZ X^\top) = \text{rk}(dZ) \leq d_k < d_e < d_s$$

Hence, the problem becomes

$$\begin{aligned} dZ^* &= \arg \min_{dZ} \langle G, X dZ X^\top \rangle_F \quad \text{s.t.} \quad \|X dZ X^\top\|_F \leq \gamma \\ &= - \arg \max_{dZ} \langle G, X dZ X^\top \rangle_F \quad \text{s.t.} \quad \|X dZ X^\top\|_F \leq \gamma \\ &= - \arg \max_{dZ} \langle G, X dZ X^\top \rangle_F \quad \text{s.t.} \quad \|X dZ X^\top\|_F = \gamma \\ &= -\gamma \arg \max_{dZ} \langle G, X dZ X^\top \rangle_F \quad \text{s.t.} \quad \|X dZ X^\top\|_F = 1 \\ &= -\frac{\gamma}{\alpha} \arg \min_{dZ} \|G - X dZ X^\top\|_F^2 \end{aligned}$$

(\*) We make the hypothesis that we can always find a  $\langle G, X dZ X^\top \rangle_F > 0$ .

🔥 Justification for  $\alpha$ ,  $\alpha = \|\cdot\|_F$

Let  $\lambda := \frac{\gamma}{\alpha}$ ,  $P := \arg \min_{dZ} \|G - X dZ X^\top\|_F^2$ .

We will first search  $P$ , then  $\lambda$  with a line search.

## 2.1 Find $P$

Let

$$f(P) = \|G - X P X^\top\|_F^2$$

$f$  is convex.

We have

$$\nabla_P f = -2X^\top (G - X P X^\top) X$$

so

$$\nabla_P f = 0 \iff X^\top X P X^\top X = X^\top G X$$

🔥 Better way to find  $P$  by considering  $X^\top (X P X^\top - G) X = 0$  ?

Under the hypothesis that  $X^\top X$  is invertible ( $\text{rk}(X) = d_e, d_e < d_s$ ), we have the solution

$$\begin{aligned} P^\star &= (X^\top X)^{-1} X^\top G X (X^\top X)^{-1} \\ &= (X^\top X)^+ X^\top G X (X^\top X)^+ \\ &= X^+ G (X^+)^{\top} \end{aligned}$$

🔥 Which formula to implement?

To account for the batch, there are several ways to average:

$$\begin{aligned} \mathbb{E}_X \left[ (X^\top X)^+ X^\top G X (X^\top X)^+ \right] &= \mathbb{E}_X \left[ X^+ G (X^+)^{\top} \right] \\ \mathbb{E}_X [X^\top X]^+ \mathbb{E}_X [X^\top G X] \mathbb{E}_X [X^\top X]^+ & \\ \mathbb{E}_X [X]^+ \mathbb{E}_X [G] (\mathbb{E}_X [X]^+)^{\top} & \end{aligned}$$

## 2.2 Line search

Do a line search to find  $\lambda$

### 2.2.1 “Normal” way

Line search on

$$\mathcal{L} \left( X \text{ SVD}_{d_k+p} \left( W_Q^t W_K^{t^\top} - \lambda P \right) X^\top \right)$$

### 2.2.2 Testing fast way

Lose the rank constraint, but lose the SVD so faster. Get a good approximation of  $\lambda$  ?

$$\mathcal{L} \left( X \left( W_Q^t W_K^{t^\top} - \lambda P \right) X^\top \right)$$