هـيـئـة الـبـيـئـة - أبـوظـبـي
Environment Agency - ABU DHABI

AIR QUALITY
ANNUAL SUMMARY REPORT
ABU DHABI
2019

01

//INTRODUCTION

The objective of EAD's air quality priority is to ensure that ambient air quality in Abu Dhabi protects human health and the environment. EAD will focus on improving the comprehensiveness of ambient air quality monitoring across the emirate, advancing capabilities for analysing, modelling and reporting air quality information, and ensuring that emission standards, regulations and enforcement regimes are in place for the key sectors and pollutants that pose the greatest threats to public health, wildlife, and quality of life in Abu Dhabi.

The Environment Agency – Abu Dhabi (EAD) started monitoring air quality in 2007. Quality Assurance/Quality Control (QA/QC) methods and procedures are implemented with full documentation and are validated through an international certified calibration reference laboratory. Forms and log sheets document every activity in the air monitoring stations and document all maintenance, calibration, operation and other activities such as all visits to the stations.

This annual report provides an overview and analysis of air quality monitoring data in Abu Dhabi for the year 2019, and a short comparison of monitoring results with earlier years. The analysis covers the three regions in Abu Dhabi Al Ain Region (Eastern Region), Al Dhafra Region (Western Region) and Central Region (Greater Abu Dhabi and its surrounding).

The report summarizes the data available at the twenty fixed stations in Abu Dhabi Emirate, in addition two mobile stations. The network monitors up to 17 parameters.

//AIR POLLUTION SOURCES AND HEALTH EFFECTS

POLLUTANT

SULFUR DIOXIDE SO_2

- Traffic pollution
- Fuel Combustion
- Electric Utilities
- Industrial Processes
- Oil and gas activities

CARBON MONOXIDE CO

- Traffic Pollution
- Fuel Combustion

NITROGEN DIOXIDE NO_2

- Traffic Pollution
- Fuel Combustion
- Electric Utilities
- Industrial Boilers

PARTICULATE MATTER PM

- Arid climate
- Regional dust clouds
- Fuel Combustion
- Industrial processes
- Traffic Pollution
- Construction Activities

OZONE O_3

- Secondary pollutant typically formed by photochemical reaction of volatile organic compounds (VOCs) and NOx in the presence of sunlight.

HYDROGEN SULFIDE H_2S

- Sewage Network
- Oil and Gas industrial activities
- Waste-water treatment plants

Respiratory diseases Cardiovascular diseases Odor nuisance

03

//AMBIENT AIR QUALITY LIMITS (AQL)

The air pollution levels have been compared to the UAE Air Quality Limits defined in the Cabinet of Ministers Decree No. 12 for 2006 concerning Protection of Air from Pollution. A summary of these Ambient Air Quality Limit (AQL) concentration levels is presented bellow.



*TSP = Total Suspended Particles

04

//AIR QUALITY INDEX

This report establishes the Air Quality Index (AQI) to evaluate air pollution. EAD simplifies the Ambient Air Quality State by calculating the AQI Range based on Air Quality National Standards for the major five parameters; Particulate matter, Ground level ozone, Sulfur dioxide, Nitrogen dioxide and Carbon monoxide.

AQI values that are below 100 are compliant with the air quality limits and are therefore considered to be acceptable.

US EPA AQI Classifications

AQI RANGE	CLASSIFICATIONS	CONDITIONS
0 to 50	Good	Considered satisfactory
51 to 100	Moderate	Air quality is acceptable
101 to 150	Unhealthy for Sensitive Groups	Members of sensitive groups may experience health effects
151to 200	Unhealthy	Everyone may begin to experience health effects
201to 300	Very Unhealthy	Health alert, meaning everyone may experience more serious health effects.
301 to 500	Hazardous	Health warnings of emergency conditions.

05 //AIR QUALITY MONITORING STATIONS

STATION NAME	STATION REPRESENTATIVITY (AREA TYPE)	MAIN PARAMETERS						
		SO_2	NO_2	CO	O_3	PM_{10}	$PM_{2.5}$	H_2S
CENTRAL ABU DHABI AREA								
HAMDAN STREET	URBAN TRAFFIC	●	●	●		●	●	
KHADEJAH SCHOOL	URBAN BACKGROUND	●	●		●	●	●	●
KHALIFA SCHOOL	SUBURBAN BACKGROUND	●	●		●	●	●	●
AL MAQTA	SUBURBAN BACKGROUND	●	●	●	●	●	●	●
KHALIFA CITY A	SUBURBAN BACKGROUND	●	●		●	●	●	●
BANIYAS SCHOOL	SUBURBAN BACKGROUND	●	●		●	●	●	●
MUSSFAH	SUBURBAN INDUSTRIAL	●	●			●	●	●
AL MAFRAQ	SUBURBAN INDUSTRIAL	●	●			●	●	●
AL AIN REGION								
AL AIN STREET	URBAN TRAFFIC	●	●	●		●	●	
AL AIN ISLAMIC INSTITUTE	URBAN BACKGROUND	●	●		●	●	●	●
AL TAWIA	SUBURBAN BACKGROUND	●	●		●	●	●	●
ZAKHER	URBAN BACKGROUND	●	●			●	●	●
SWEIHAN	SUBURBAN BACKGROUND	●	●	●	●	●	●	
AL QUA'A	REGIONAL RURAL BACKGROUND	●	●	●	●	●	●	
AL DHAFRA REGION								
E11 ROAD*	RURAL TRAFFIC	●	●	●		●	●	
BIDA ZAYED	SUBURBAN BACKGROUND	●	●		●	●	●	●
HABSHAN SOUTH	RURAL INDUSTRIAL	●	●		●	●	●	●
RUWAIS	SUBURBAN INDUSTRIAL	●	●	●	●	●	●	●
GAYATHI SCHOOL	SUBURBAN BACKGROUND	●	●		●	●	●	●
LIWA OASIS	REGIONAL RURAL BACKGROUND	●	●		●	●	●	

* E11 Road Station currently under relocation.

● Measured Parameters

06 //AIR QUALITY STATUS



PM_{10}
O_3

The percentage of compliant days present AQI values below 100, which are normally expected to be satisfactory

THE PERCENTAGE OF COMPLIANT DAYS WITHIN THE YEAR IN ABU DHABI MONITORING STATIONS



//METEOROLOGICAL DATA

07

All EAD air quality monitoring stations are equipped with sensors to record meteorological parameters, which are essential to understand the ambient air quality patterns and local meteorological conditions. The meteorological parameters measured are: wind speed, wind direction, temperature, relative humidity, net radiation and barometric pressure.

The prevailing winds over Abu Dhabi are northwesterly winds, though differences in wind speed and wind direction may occur in specific locations due to local conditions and patterns. During the night, wind blowing from south-southeast also has a high percentage of occurrence.

The below map shows the wind roses in all of Abu Dhabi stations from 2007 to 2019, and the wind rose is a graphic tool used to describe the distribution of the wind speed and wind direction in a particular location. The frequency of the winds is plotted by wind direction, with colour bands showing wind speed ranges. The direction of the longest spoke shows the predominant wind direction in that location.

//WIND SPEED/DIRECTION DATA IN ABU DHABI STATIONS FROM 2007 TO 2019



PARTICULATE MATTER PM_{10}

Figure 1 shows the annual average PM_{10} concentrations by region together with the linear regression of PM_{10} annual means measured at all stations from the beginning of 2007 until the end of 2019.

Overall, there was a slight decrease in the trend of PM_{10} concentrations from the beginning of 2007 until the end of 2019. During 2019, PM_{10} concentration decreased in all region. In Abu Dhabi Region the lowest concentration of PM_{10} recorded by Hamdan Street station, in Al Ain Region recorded by Sweihan station, and in Al Dhafra Region recorded by Ruwais station and Gayathi School station.

ANNUAL AVERAGE PM_{10} CONCENTRATIONS



Figure1:

Annual average concentrations for PM_{10} given for the sites in the Al Dhafra Region, Al Ain Region, Abu Dhabi Industrial Areas and Abu Dhabi Emirate PM_{10} linear regression from 2007 to 2019
The number of days where daily averages were above 350 µg/m³ are:



PARTICULATE MATTER $PM_{2.5}$

Figure 2 shows the annual average $PM_{2.5}$ concentrations by region together with the linear regression of $PM_{2.5}$ annual means measured at all stations from the beginning of 2012 until the end of 2019. The monitored $PM_{2.5}$ levels in Abu Dhabi.

Overall, there was a slight increase in the trend of $PM_{2.5}$ concentrations from the beginning of 2012 until the end of 2018. During 2019, $PM_{2.5}$ concentration slightly decreased in all region except in Al Ain Region which extremely decreased. In Abu Dhabi Region the lowest concentration of $PM_{2.5}$ recorded by Al Maqta station, in Al Ain Region recorded by Al Tawia station, and in Al Dhafra Region recorded by three different stations.

ANNUAL AVERAGE $PM_{2.5}$ CONCENTRATIONS



Figure2:

Annual average concentrations for $PM_{2.5}$ given for the sites in the Al Dhafra Region, Al Ain Region, Abu Dhabi Industrial Areas and Abu Dhabi Emirate $PM_{2.5}$ linear regression from 2012 to 2019

12

OZONE O_3

Figure 3 shows the annual average O_3 concentrations by region together with the linear regression of O_3 annual means measured at all stations from the beginning of 2007 until the end of 2019.

Overall, there was a notably increasing in the trend of O_3 concentrations from the beginning of 2007 until the end of 2019. During 2019, O_3 concentration decreased in all regions. In Abu Dhabi Region the lowest concentration of O_3 recorded by Khalifa School station, in Al Ain Region recorded by Al Ain Islamic Institute station, and in Al Dhafra Region recorded by Bida Zayed Station.

ANNUAL AVERAGE O_3 CONCENTRATIONS



Figure3:

Annual average concentrations for O_3 given for the sites in the Al Dhafra Region, Al Ain Region, Abu Dhabi Industrial Areas and Abu Dhabi Emirate O_3 linear regression from 2007 to 2019

O_3 TREND

Ozone trend can be related to the following factors:

O_3

DECREASE

Expand the use of renewable energy(*e.g.* solar energy) to reduce greenhouse gases.

Apply the best available technologies to control emissions.

Encourage the use of international best practices to minimise emissions.

INCREASE

An Increase of traffic and industrial activities.

Effect of global atmospheric circulation.

Climate change increasing global temperature leading to higher potential for O_3 generation

13

SULFUR DIOXIDE SO_2

Figure 4 shows the annual average SO_2 concentrations by region together with the linear regression of SO_2 annual means measured at all stations from the beginning of 2007 until the end of 2019.

Overall, there was a slight increase in the trend of SO_2 concentrations from the beginning of 2007 until the end of 2019. During 2019, SO_2 concentration decreased in all regions except Al Ain Region. In Abu Dhabi Region the lowest concentration of SO_2 recorded by Khalifa City A Station, in Al Ain Region recorded by Zakher Station and in Al Dhafra Region recorded by Liwa Oasis Station.

ANNUAL AVERAGE SO_2 CONCENTRATIONS



Figure 4:

Annual average concentrations for SO_2 given for the sites in the Al Dhafra Region, Al Ain Region, Abu Dhabi Industrial Areas and Abu Dhabi Emirate SO_2 linear regression from 2007 to 2019

14

NITROGEN DIOXIDE NO_2

Figure 5 shows the annual average NO_2 concentrations by region together with the linear regression of NO_2 annual means measured at all stations from the beginning of 2007 until the end of 2019.

Overall, NO_2 concentrations trend almost constant from the beginning of 2007 until the end of 2019. During 2019, NO_2 concentration slight decreased in all regions except Al Dhafra Region which constant. In Abu Dhabi Region the lowest concentration of NO_2 recorded by Baniyas School Station, in Al Ain Region recorded by Al Qua`a Station, and in Al Dhafra Region recorded by Liwa Oasis Station.

The annual average NO_2 concentrations reached 157%, 103%, 123%, 127% and 127% of the annual WHO guideline value at Hamdan Street, Khalifa City A, Mussafah, Al Mafraq and Al Ain Stree stations respectively.

ANNUAL AVERAGE NO_2 CONCENTRATIONS



Figure 5:

Annual average concentrations for NO_2 given for the sites in the Al Dhafra Region, Al Ain Region, Abu Dhabi Industrial Areas and Abu Dhabi Emirate NO_2 linear regression from 2007 to 2019

15

CARBON MONOXIDE CO

Figure 6 shows the annual average CO concentrations by region together with the linear regression of CO annual means measured at all stations from the beginning of 2007 until the end of 2019.

Overall, there was a decrease in the trend of CO concentrations from the beginning of 2007 until the end of 2019. During 2019, CO concentration slightly increased in all regions except Al Ain Region which constant and CO concentrations never exceeded any of the air quality limit value in the same year.



Figure 6 :

Annual average concentrations for CO given for the sites in the Al Dhafra Region, Al Ain Region, Abu Dhabi Industrial Areas and Abu Dhabi Emirate CO linear regression from 2007 to 2019

HYDROGEN SULFIDE H_2S

There is no air quality limit value for H_2S in UAE. H_2S is not one of the criteria pollutants, but may cause odour nuisance at concentrations far below those that cause health hazards. The World Health Organization has presented a 24 hour average guideline value of 150 µg/m^3. Half-hour average concentrations exceeding 7 µg/m^3 are likely to produce odor problems and complaints among persons exposed.

The WHO guideline given for 24-hour average concentration of H_2S at 150 µg/m^3 was never exceeded.

In Abu Dhabi Emirate H_2S concentrations are well within the health threshold recommended by WHO. However, there has been as increase in H_2S concentrations in some specific area, which may cause odour nuisances.

16

//CONCLUSION

Natural and man-made activities are the main contributors to the air quality of Abu Dhabi. Our desert environment plays an important role in quality of the air as well. As our population grows, our contribution to the level of air quality will get affected.

Together with the efforts of the Environment Agency - Abu Dhabi and all government entities who do their best in monitoring and minimising the effects of these man made activities in order to provide us with a clean environment.

To sum up, compared to the previous year, 2019 show a decreased in the average of O_3, $PM_{2.5}$ and PM_{10} concentration at all regions. However SO_2, NO_2 and CO were compliant in all the stations.

Photo credit: **Environment Agency - Abu Dhabi**

CONTACT US

Environment Agency - Abu Dhabi
Call us: **(+971) 2 4454777**
E-mail us: **customerhappiness@ead.gov.ae**
Know more about us: **www.ead.gov.ae**

@EADTweets
Environment Agency -Abu Dhabi
Environment Abu Dhabi
Environment Agency -Abu Dhabi