

Table 1: Standard error, skewness, and 95% confidence limits for three functionals in the worked example, by interval procedure (a single data set with $N = 500$; $B = 5,000$ weight vectors).

Functional	Method	SE	Skewness	2.5%	97.5%
ab	Wald (Expected)	0.010	0.000	-0.005	0.033
	Wald (HW)	0.010	0.000	-0.005	0.033
	Monte Carlo (HW)	0.010	0.694	-0.003	0.037
	IJ1 percentile	0.010	0.610	-0.004	0.037
	HOLJ-2 percentile	0.010	0.662	-0.004	0.037
	Bootstrap percentile	0.011	0.672	-0.004	0.037
ψ_{speed}	Wald (Expected)	0.068	0.000	0.252	0.516
	Wald (HW)	0.069	0.000	0.249	0.519
	Monte Carlo (HW)	0.069	-0.030	0.248	0.517
	IJ1 percentile	0.069	0.009	0.250	0.517
	HOLJ-2 percentile	0.069	0.239	0.257	0.524
	Bootstrap percentile	0.069	0.240	0.257	0.525
ω_{speed}	Wald (Expected)	0.025	0.000	0.627	0.725
	Wald (HW)	0.026	0.000	0.626	0.727
	Monte Carlo (HW)	0.028	-0.676	0.609	0.720
	IJ1 percentile	0.028	-0.562	0.616	0.722
	HOLJ-2 percentile	0.027	-0.276	0.622	0.725
	Bootstrap percentile	0.026	-0.245	0.624	0.725




Figure 1: Replicate distributions in the lower (top row) and upper (bottom row) tail of three functionals, for a single data set. The shaded region is the bootstrap; the vertical line marks the corresponding bootstrap percentile limit (2.5% above, 97.5% below).