

Graph Theory and Optimization

Integer Linear Programming

Nicolas Nisse

Université Côte d'Azur, Inria, CNRS, I3S, France

October 2018

Outline

- 1 Integer Linear Programme
- 2 Some examples
- 3 Integrality gap
- 4 Polynomial Cases
- 5 More Examples

Linear Programme (reminder)

Linear programmes can be written under the **standard form**:

$$\begin{array}{ll}\text{Maximize} & \sum_{j=1}^n c_j x_j \\ \text{Subject to:} & \sum_{j=1}^n a_{ij} x_j \leq b_i \quad \text{for all } 1 \leq i \leq m \\ & x_j \geq 0 \quad \text{for all } 1 \leq j \leq n.\end{array}$$

- the problem is a **maximization**;
- all constraints are **inequalities** (and not equations);
- all variables x_1, \dots, x_n are **non-negative**.

Linear Programme (Real variables) can be solved in polynomial-time in the number of variables and constraints (e.g., ellipsoid method)

Linear Programme (reminder)

Linear programmes can be written under the **standard form**:

$$\begin{aligned}
 &\text{Maximize} && \sum_{j=1}^n c_j x_j \\
 &\text{Subject to:} && \sum_{j=1}^n a_{ij} x_j \leq b_i && \text{for all } 1 \leq i \leq m \\
 &&& x_j \geq 0 && \text{for all } 1 \leq j \leq n.
 \end{aligned}$$

- the problem is a **maximization**;
- all constraints are **inequalities** (and not equations);
- all variables x_1, \dots, x_n are **non-negative**.

Linear Programme (Real variables) can be solved in polynomial-time in the number of variables and constraints (e.g., ellipsoid method)

Integer Linear Programme

Integer Linear programmes:

$$\begin{aligned}
 &\text{Maximize} && \sum_{j=1}^n c_j x_j \\
 &\text{Subject to:} && \sum_{j=1}^n a_{ij} x_j \leq b_i && \text{for all } 1 \leq i \leq m \\
 &&& x_j \in \mathbb{N} && \text{for all } 1 \leq j \leq n.
 \end{aligned}$$

- the problem is a **maximization**;
- all constraints are **inequalities** (and not equations);
- all variables x_1, \dots, x_n are **Integers**.

Integer Linear Programme is NP-complete in general!


Outline

- 1 Integer Linear Programme
- 2 Some examples
- 3 Integrality gap
- 4 Polynomial Cases
- 5 More Examples

Knapsack Problem

(Weakly NP-hard)


- **Data:**
 - a knapsack with maximum weight 15 Kg
 - 12 objects with
 - a weight w_i
 - a value v_i
- **Objective:** which objects should be chosen to maximize the value carried while not exceeding 15 Kg?



Knapsack Problem

(Weakly NP-hard)

- **Data:**
 - a knapsack with maximum weight 15 Kg
 - 12 objects with
 - a weight w_i
 - a value v_i
- **Objective:** which objects should be chosen to maximize the value carried while not exceeding 15 Kg?



$$\begin{array}{ll}
 \max & \sum_{1 \leq i \leq 12} v_i x_i \\
 \text{subject to} & \sum_{1 \leq i \leq 12} w_i x_i \leq 15 \\
 & x_i \in \{0, 1\}
 \end{array}$$

Minimum Vertex Cover

(NP-hard)

Let $G = (V, E)$ be a graph

Vertex Cover: set $K \subseteq V$ such that $\forall e \in E, e \cap K \neq \emptyset$

set of vertices that "touch" every edge

Minimum Vertex Cover

(NP-hard)

Let $G = (V, E)$ be a graph

Vertex Cover: set $K \subseteq V$ such that $\forall e \in E, e \cap K \neq \emptyset$
set of vertices that "touch" every edge

Solution: $K \subseteq V$

\Rightarrow variables x_v , for each $v \in V$
 $x_v = 1$ if $v \in K$, $x_v = 0$ otherwise.

Minimum Vertex Cover

(NP-hard)

Let $G = (V, E)$ be a graph

Vertex Cover: set $K \subseteq V$ such that $\forall e \in E, e \cap K \neq \emptyset$
set of vertices that "touch" every edge

Solution: $K \subseteq V$

\Rightarrow variables x_v , for each $v \in V$

$x_v = 1$ if $v \in K$, $x_v = 0$ otherwise.

Objective function: minimize $|K|$

minimize $\sum_{v \in V} x_v$

Minimum Vertex Cover

(NP-hard)

Let $G = (V, E)$ be a graph

Vertex Cover: set $K \subseteq V$ such that $\forall e \in E, e \cap K \neq \emptyset$

set of vertices that "touch" every edge

Solution: $K \subseteq V$

\Rightarrow variables x_v , for each $v \in V$

$x_v = 1$ if $v \in K$, $x_v = 0$ otherwise.

Objective function: minimize $|K|$

minimize $\sum_{v \in V} x_v$

Constraint: $\forall \{u, v\} \in E, u \in K \text{ or } v \in K$

$x_u + x_v \geq 1$

Minimum Vertex Cover

(NP-hard)

Let $G = (V, E)$ be a graph

Vertex Cover: set $K \subseteq V$ such that $\forall e \in E, e \cap K \neq \emptyset$

set of vertices that "touch" every edge

Solution: $K \subseteq V$

\Rightarrow variables x_v , for each $v \in V$

$x_v = 1$ if $v \in K$, $x_v = 0$ otherwise.

Objective function: minimize $|K|$

minimize $\sum_{v \in V} x_v$

Constraint: $\forall \{u, v\} \in E, u \in K \text{ or } v \in K$

$x_u + x_v \geq 1$

$$\begin{array}{ll} \text{Minimize} & \sum_{v \in V} x_v \\ \text{Subject to:} & x_v + x_u \geq 1 \quad \text{for all } \{u, v\} \in E \\ & x_v \in \{0, 1\} \quad \text{for all } v \in V \end{array}$$

Vertex Coloring

(NP-hard)

Let $G = (V, E)$ be a graph

k -Proper coloring: $c : V \rightarrow \{1, \dots, k\}$ s.t. $c(u) \neq c(v)$ for all $\{u, v\} \in E$.
color the vertices s ($\leq k$ colors) s.t. adjacent vertices receive \neq colors

Vertex Coloring

(NP-hard)

Let $G = (V, E)$ be a graph

k -Proper coloring: $c : V \rightarrow \{1, \dots, k\}$ s.t. $c(u) \neq c(v)$ for all $\{u, v\} \in E$.
color the vertices s ($\leq k$ colors) s.t. adjacent vertices receive \neq colors

Solution: $c : V \rightarrow \{1, \dots, n\} \Rightarrow$ variables y_j , is color $j \in \{1, \dots, n\}$ used?
 variable c_v^j for color j and vertex v : $c_v^j = 1$ if v colored j , $c_v^j = 0$ otherwise

Vertex Coloring

(NP-hard)

Let $G = (V, E)$ be a graph

k -Proper coloring: $c : V \rightarrow \{1, \dots, k\}$ s.t. $c(u) \neq c(v)$ for all $\{u, v\} \in E$.
color the vertices s ($\leq k$ colors) s.t. adjacent vertices receive \neq colors

Solution: $c : V \rightarrow \{1, \dots, n\} \Rightarrow$ variables y_j , is color $j \in \{1, \dots, n\}$ used?
 variable c_v^j for color j and vertex v : $c_v^j = 1$ if v colored j , $c_v^j = 0$ otherwise

Objective function: minimize # of used colors minimize $\sum_{1 \leq j \leq n} y_j$

Vertex Coloring

(NP-hard)

Let $G = (V, E)$ be a graph

k -Proper coloring: $c : V \rightarrow \{1, \dots, k\}$ s.t. $c(u) \neq c(v)$ for all $\{u, v\} \in E$.
color the vertices s ($\leq k$ colors) s.t. adjacent vertices receive \neq colors

Solution: $c : V \rightarrow \{1, \dots, n\} \Rightarrow$ variables y_j , is color $j \in \{1, \dots, n\}$ used?
 variable c_v^j for color j and vertex v : $c_v^j = 1$ if v colored j , $c_v^j = 0$ otherwise

Objective function: minimize # of used colors minimize $\sum_{1 \leq j \leq n} y_j$

Constraints: each vertex v has 1 color $\sum_{1 \leq j \leq n} c_v^j = 1$

ends of each edge $\{u, v\} \in E$ have \neq colors: $c_v^j + c_u^j \leq 1$ for all $j \in \{1, \dots, n\}$
 color j used if ≥ 1 vertex colored with j $c_v^j \leq y_j$ for all $v \in V$

Vertex Coloring

(NP-hard)

Let $G = (V, E)$ be a graph

k -Proper coloring: $c : V \rightarrow \{1, \dots, k\}$ s.t. $c(u) \neq c(v)$ for all $\{u, v\} \in E$.
color the vertices s ($\leq k$ colors) s.t. adjacent vertices receive \neq colors

$$\begin{array}{ll}
 \text{Minimize} & \sum_{1 \leq j \leq n} y_j \\
 \text{Subject to:} & \sum_{1 \leq j \leq n} c_v^j = 1 \quad \text{for all } v \in V \\
 & c_v^j + c_u^j \leq 1 \quad \text{for all } j \in \{1, \dots, n\}, \{u, v\} \in E \\
 & c_v^j \leq y_j \quad \text{for all } j \in \{1, \dots, n\}, v \in V \\
 & y_j \in \{0, 1\} \quad \text{for all } j \in \{1, \dots, n\} \\
 & c_v^j \in \{0, 1\} \quad \text{for all } j \in \{1, \dots, n\}, v \in V
 \end{array}$$

Outline

- 1 Integer Linear Programme
- 2 Some examples
- 3 Integrality gap
- 4 Polynomial Cases
- 5 More Examples

Integer Programme vs. Linear Programme

Integer Linear programme (ILP):

$$\begin{array}{ll}
 \text{Max.} & \sum_{j=1}^n c_j x_j \\
 \text{s.t.:} & \sum_{j=1}^n a_{ij} x_j \leq b_i \quad \forall 1 \leq i \leq m \\
 & x_j \in \mathbb{N} \quad \forall 1 \leq j \leq n.
 \end{array}$$

NP-hard in general

Integer Programme vs. Linear Programme

Integer Linear programme (ILP):

$$\begin{array}{ll} \text{Max.} & \sum_{j=1}^n c_j x_j \\ \text{s.t.:} & \sum_{j=1}^n a_{ij} x_j \leq b_i \quad \forall 1 \leq i \leq m \\ & x_j \in \mathbb{N} \quad \forall 1 \leq j \leq n. \end{array}$$

NP-hard in general

Fractional Relaxation: Linear Programme

$$\begin{array}{ll} \text{Max.} & \sum_{j=1}^n c_j x_j \\ \text{s.t.:} & \sum_{j=1}^n a_{ij} x_j \leq b_i \quad \forall 1 \leq i \leq m \\ & x_j \geq 0 \quad \forall 1 \leq j \leq n. \end{array}$$

Polynomial-time solvable

Integer Programme vs. Linear Programme

Integer Linear programme (ILP):

$$\begin{array}{ll} \text{Max.} & \sum_{j=1}^n c_j x_j \\ \text{s.t.:} & \sum_{j=1}^n a_{ij} x_j \leq b_i \quad \forall 1 \leq i \leq m \\ & x_j \in \mathbb{N} \quad \forall 1 \leq j \leq n. \end{array}$$

NP-hard in general

Fractional Relaxation: Linear Programme

$$\begin{array}{ll} \text{Max.} & \sum_{j=1}^n c_j x_j \\ \text{s.t.:} & \sum_{j=1}^n a_{ij} x_j \leq b_i \quad \forall 1 \leq i \leq m \\ & x_j \geq 0 \quad \forall 1 \leq j \leq n. \end{array}$$

Polynomial-time solvable

What is the difference between Optimal solutions of *LP* and of *ILP*?

Integer Programme vs. Linear Programme

Integer Linear programme (ILP):

$$\begin{array}{ll} \text{Max.} & \sum_{j=1}^n c_j x_j \\ \text{s.t.:} & \sum_{j=1}^n a_{ij} x_j \leq b_i \quad \forall 1 \leq i \leq m \\ & x_j \in \mathbb{N} \quad \forall 1 \leq j \leq n. \end{array}$$

NP-hard in general

Fractional Relaxation: Linear Programme

$$\begin{array}{ll} \text{Max.} & \sum_{j=1}^n c_j x_j \\ \text{s.t.:} & \sum_{j=1}^n a_{ij} x_j \leq b_i \quad \forall 1 \leq i \leq m \\ & x_j \geq 0 \quad \forall 1 \leq j \leq n. \end{array}$$

Polynomial-time solvable

What is the difference between Optimal solutions of LP and of ILP?

$$OPT(LP) \geq OPT(ILP) \quad (\text{for a maximization problem})$$

$$OPT(LP) \leq OPT(ILP) \quad (\text{for a minimization problem})$$

If $OPT(LP)$ is "closed" to $OPT(ILP)$, then solving the **Fractional Relaxation** (in polynomial-time) gives a good bound for the ILP

Fractional Relaxation of Vertex Coloring

Integer Linear programme (ILP):

$$\begin{array}{ll}
 \text{Minimize} & \sum_{1 \leq j \leq n} y_j \\
 \text{Subject to:} & \sum_{1 \leq j \leq n} c_v^j = 1 \\
 & c_v^j + c_u^j \leq 1 \\
 & c_v^j \leq y_j \\
 & y_j \in \{0, 1\} \\
 & c_v^j \in \{0, 1\}
 \end{array}$$

Fractional Relaxation (LP):

$$\begin{array}{ll}
 \text{Minimize} & \sum_{1 \leq j \leq n} y_j \\
 \text{Subject to:} & \sum_{1 \leq j \leq n} c_v^j = 1 \\
 & c_v^j + c_u^j \leq 1 \\
 & c_v^j \leq y_j \\
 & y_j \geq 0 \\
 & c_v^j \geq 0
 \end{array}$$


$$\begin{array}{l}
 y_{\text{red}} = y_{\text{blue}} = y_{\text{green}} = 1 \\
 c_a^{\text{red}} = c_b^{\text{blue}} = c_c^{\text{green}} = 1 \\
 \text{OPT(ILP)} = \sum_c y_c = 3
 \end{array}$$

$$\begin{array}{l}
 y_{\text{red}} = y_{\text{blue}} = 1/2, y_{\text{green}} = 0 \\
 c_a^{\text{red}} = c_b^{\text{red}} = c_c^{\text{red}} = 1/2 \\
 c_a^{\text{blue}} = c_b^{\text{blue}} = c_c^{\text{blue}} = 1/2 \\
 \text{OPT(LP)} = \sum_c y_c = 1
 \end{array}$$

Fractional Relaxation of Vertex Coloring

Integer Linear programme (ILP):


$$\begin{aligned}
 &\text{Minimize} && \sum_{1 \leq j \leq n} y_j \\
 &\text{Subject to:} && \sum_{1 \leq j \leq n} c_v^j = 1 \\
 & && c_v^j + c_u^j \leq 1 \\
 & && c_v^j \leq y_j \\
 & && y_j \in \{0, 1\} \\
 & && c_v^j \in \{0, 1\}
 \end{aligned}$$



$$\begin{aligned}
 y_{\text{red}} &= y_{\text{blue}} = y_{\text{green}} = 1 \\
 c_a^{\text{red}} &= c_b^{\text{blue}} = c_c^{\text{green}} = 1 \\
 \text{OPT(ILP)} &= \sum_c y_c = 3
 \end{aligned}$$

Fractional Relaxation (LP):

$$\begin{aligned}
 &\text{Minimize} && \sum_{1 \leq j \leq n} y_j \\
 &\text{Subject to:} && \sum_{1 \leq j \leq n} c_v^j = 1 \\
 & && c_v^j + c_u^j \leq 1 \\
 & && c_v^j \leq y_j \\
 & && y_j \geq 0 \\
 & && c_v^j \geq 0
 \end{aligned}$$



$$\begin{aligned}
 y_{\text{red}} &= y_{\text{blue}} = 1/2, y_{\text{green}} = 0 \\
 c_a^{\text{red}} &= c_b^{\text{red}} = c_c^{\text{red}} = 1/2 \\
 c_a^{\text{blue}} &= c_b^{\text{blue}} = c_c^{\text{blue}} = 1/2 \\
 \text{OPT(LP)} &= \sum_c y_c = 1
 \end{aligned}$$

Fractional Relaxation of Knapsac

Integer Linear programme (ILP):

$$\begin{array}{ll}\max & \sum_{1 \leq i \leq n} v_i x_i \\ \text{subject to} & \sum_{1 \leq i \leq n} w_i x_i \leq W \\ & x_i \in \{0, 1\}\end{array}$$

Fractional Relaxation (LP):

$$\begin{array}{ll}\max & \sum_{1 \leq i \leq n} v_i x_i \\ \text{subject to} & \sum_{1 \leq i \leq n} w_i x_i \leq W \\ & x_i \geq 0\end{array}$$

Example:

- Sac: $W = n$
- Objects:
 - one object (O_1) of weight $n + 0,1$ and value n
 - $n - 1$ objects (O_2, \dots, O_n) of weight 1 and value $1/n$

$$x_1 = 0, x_2 = \dots = x_n = 1$$

$$OPT(ILP) = \sum_c v_i x_i = (n-1)/n$$

$$x_1 = \frac{n}{n+0,1}, x_2 = \dots = x_n = 0$$

$$OPT(LP) = \sum_c v_i x_i = \frac{n^2}{n+0,1}$$

⇒ the ratio between the LP optimal solution and the Integral opt. solution may be arbitrary large

Fractional Relaxation of Knapsac

Integer Linear programme (ILP):

$$\begin{array}{ll}\max & \sum_{1 \leq i \leq n} v_i x_i \\ \text{subject to} & \sum_{1 \leq i \leq n} w_i x_i \leq W \\ & x_i \in \{0, 1\}\end{array}$$

Fractional Relaxation (LP):

$$\begin{array}{ll}\max & \sum_{1 \leq i \leq n} v_i x_i \\ \text{subject to} & \sum_{1 \leq i \leq n} w_i x_i \leq W \\ & x_i \geq 0\end{array}$$

Example:

- Sac: $W = n$
- Objects:
 - one object (O_1) of weight $n+0,1$ and value n
 - $n-1$ objects (O_2, \dots, O_n) of weight 1 and value $1/n$

$$x_1 = 0, x_2 = \dots = x_n = 1$$

$$OPT(ILP) = \sum_c v_i x_i = (n-1)/n$$

$$x_1 = \frac{n}{n+0,1}, x_2 = \dots = x_n = 0$$

$$OPT(LP) = \sum_c v_i x_i = \frac{n^2}{n+0,1}$$

⇒ the ratio between the LP optimal solution and the Integral opt. solution may be arbitrary large

Fractional Relaxation of Knapsac

Integer Linear programme (ILP):

$$\begin{array}{ll}\max & \sum_{1 \leq i \leq n} v_i x_i \\ \text{subject to} & \sum_{1 \leq i \leq n} w_i x_i \leq W \\ & x_i \in \{0, 1\}\end{array}$$

Fractional Relaxation (LP):

$$\begin{array}{ll}\max & \sum_{1 \leq i \leq n} v_i x_i \\ \text{subject to} & \sum_{1 \leq i \leq n} w_i x_i \leq W \\ & x_i \geq 0\end{array}$$

Example:

- Sac: $W = n$
- Objects:
 - one object (O_1) of weight $n+0,1$ and value n
 - $n-1$ objects (O_2, \dots, O_n) of weight 1 and value $1/n$

$$x_1 = 0, x_2 = \dots = x_n = 1$$

$$OPT(ILP) = \sum_c v_i x_i = (n-1)/n$$

$$x_1 = \frac{n}{n+0,1}, x_2 = \dots = x_n = 0$$

$$OPT(LP) = \sum_c v_i x_i = \frac{n^2}{n+0,1}$$

\Rightarrow the ratio between the LP optimal solution and the Integral opt. solution may be arbitrary large

Outline

- 1 Integer Linear Programme
- 2 Some examples
- 3 Integrality gap
- 4 Polynomial Cases
- 5 More Examples

No integrality gap

Integer Linear programme (ILP):

$$\begin{aligned} \text{Max.} \quad & \sum_{j=1}^n c_j x_j \\ \text{s.t.:} \quad & \sum_{j=1}^n a_{ij} x_j \leq b_i \quad \forall 1 \leq i \leq m \\ & x_j \in \mathbb{N} \quad \forall 1 \leq j \leq n. \end{aligned}$$

NP-hard in general

Fractional Relaxation: Linear Programme

$$\begin{aligned} \text{Max.} \quad & \sum_{j=1}^n c_j x_j \\ \text{s.t.:} \quad & \sum_{j=1}^n a_{ij} x_j \leq b_i \quad \forall 1 \leq i \leq m \\ & x_j \geq 0 \quad \forall 1 \leq j \leq n. \end{aligned}$$

Polynomial-time solvable

In some cases: $OPT(ILP) = OPT(LP)$.

\Leftrightarrow there exists an integral solution with value $OPT(LP)$.

In such case, Polynomial-time solvable: solve the Fractional Relaxation

Integer Programme Example: Shortest path

$D = (V, A)$ be a digraph with length $\ell : A \rightarrow \mathbb{R}^+$, and $s, t \in V$.

Problem: Compute a shortest directed path from s to t .

Integer Programme Example: Shortest path

$D = (V, A)$ be a digraph with length $\ell : A \rightarrow \mathbb{R}^+$, and $s, t \in V$.

Problem: Compute a shortest directed path from s to t .

Solution: A path P from s to t \Rightarrow variables x_a for each $a \in A$
 $x_a = 1$ if $a \in A(P)$, $x_a = 0$ otherwise.

Integer Programme Example: Shortest path

$D = (V, A)$ be a digraph with length $\ell : A \rightarrow \mathbb{R}^+$, and $s, t \in V$.


Problem: Compute a shortest directed path from s to t .

Solution: A path P from s to t \Rightarrow variables x_a for each $a \in A$
 $x_a = 1$ if $a \in A(P)$, $x_a = 0$ otherwise.


$$\begin{array}{ll}
 \text{Minimize} & \sum_{a \in A} \ell(a) x_a \\
 \text{Subject to:} & \sum_{u \in N^+(s)} x(su) = 1 \\
 & \sum_{u \in N^-(t)} x(tu) = 1 \\
 & \sum_{u \in N^+(v)} x(uv) = \sum_{u \in N^-(v)} x(vu) \quad \text{for all } v \in V \setminus \{s, t\} \\
 & x(a) \in \{0, 1\} \quad \text{for all } a \in A
 \end{array}$$

Integer Programme Example: Shortest path

$$\begin{array}{ll}
 \text{Minimize} & \sum_{a \in A} \ell(a) x_a \\
 \text{Subject to:} & \sum_{u \in N^+(s)} x(su) = 1 \\
 & \sum_{u \in N^-(t)} x(tu) = 1 \\
 & \sum_{u \in N^+(v)} x(uv) = \sum_{u \in N^-(v)} x(vu) \quad \text{for all } v \in V \setminus \{s, t\} \\
 & x(a) \geq 0 \quad \text{for all } a \in A
 \end{array}$$



Example: all edges
have length 1



Exercise: Prove that this LP always admits an integral optimal solution

Integer Programme Example: Maximum Matching

$G = (V, E)$ be a graph

Problem: Compute a maximum matching

Solution: a set $M \subseteq E$ of pairwise disjoint edges

\Rightarrow variables x_e for each $e \in E$

$x_e = 1$ if $e \in M$, $x_e = 0$ otherwise.

Maximize $\sum_{e \in E} x_e$

Subject to: $\sum_{e \in E, v \in e} x_e \leq 1$ for all $v \in V$

$x_e \in \{0, 1\}$ for all $e \in E$

Exercise: Prove that the fractional relaxation of this ILP always admits an integral optimal solution

Totally unimodular matrices

unimodular matrix: square matrix with determinant $+1$ or -1

totally unimodular matrix: every square non-singular submatrix is unimodular

Integer Linear programme (ILP):

$$\begin{aligned} \text{Max.} \quad & \sum_{j=1}^n c_j x_j \\ \text{s.t.:} \quad & \sum_{j=1}^n a_{ij} x_j \leq b_i \quad \forall 1 \leq i \leq m \\ & x_j \in \mathbb{N} \quad \forall 1 \leq j \leq n. \end{aligned}$$

NP-hard in general

Fractional Relaxation: Linear Programme

$$\begin{aligned} \text{Max.} \quad & \sum_{j=1}^n c_j x_j \\ \text{s.t.:} \quad & \sum_{j=1}^n a_{ij} x_j \leq b_i \quad \forall 1 \leq i \leq m \\ & x_j \geq 0 \quad \forall 1 \leq j \leq n. \end{aligned}$$

Polynomial-time solvable

Theorem

[Hoffman, Kruskal, 1956]

If the matrix $A = [a_{ij}]$ is totally unimodular then every *basic* feasible solution (the "corner" of the polytope) is integral

\Rightarrow exist integral optimal solution of the LP

$\Rightarrow \text{OPT(ILP)}$ can be computed by solving the Fractional relaxation

Outline

- 1 Integer Linear Programme
- 2 Some examples
- 3 Integrality gap
- 4 Polynomial Cases
- 5 More Examples

Integer Programme Example: Minimum Spanning Tree

$G = (V, E)$ be a graph with weight $w : E \rightarrow \mathbb{R}^+$, and $s, t \in V$.

Problem: Compute a minimum spanning tree


Solution: A spanning tree $T \Rightarrow$ variables x_e for each $e \in E$
 $x_e = 1$ if $e \in E(T)$, $x_e = 0$ otherwise.

$$\begin{array}{ll}
 \text{Minimize} & \sum_{e \in E} w(e)x_e \\
 \text{Subject to:} & \sum_{e=\{u,v\} \in E, u \in S, v \notin S} x_e \geq 1 \quad \text{for all } S \subseteq V \\
 & x_e \in \{0, 1\} \quad \text{for all } e \in E
 \end{array}$$

Remark: The number of constraints is exponential

Optical Networks (WDM)

Optical network: optical fiber connecting e.g. routers




Wavelength-division Multiplexing (WDM): technology which multiplexes a number of optical carrier signals onto a single optical fiber by using different wavelengths (i.e., colors) of laser light [Wikipedia]

⇒: **different signals** on the same link must have **different wavelengths** (colors)


Optical Networks (WDM)

Optical network: optical fiber connecting e.g. routers



Wavelength-division Multiplexing (WDM): technology which multiplexes a number of optical carrier signals onto a single optical fiber by using different wavelengths (i.e., colors) of laser light [Wikipedia]

⇒: **different signals** on the same link must have **different wavelengths** (colors)



Optical Networks (WDM)

RWA problem


RWA: Routing and Wavelength Assignment Problem

Given a graph $G = (V, E)$ with capacity on links, and a traffic-demand matrix T , where $T[u, v]$ is the amount of traffic that must transit from u to v , for any $u, v \in V$. Find a set of paths and one wavelength assignment for each path such that:

- all demands are routed
- capacity of each link cannot be exceeded
- total number of wavelength is as small as possible

demand matrix:

	a	b	c	d	e	f	g
a	0	0	1	1	0	1	0
b	0	0	0	0	0	2	0
c	0	0	0	0	0	0	0
d	0	0	0	0	0	0	5
e	0	0	0	0	0	0	0
f	0	0	0	0	0	0	0
g	0	1	0	0	0	0	0



Optical Networks (WDM)

RWA problem


RWA: Routing and Wavelength Assignment Problem

Given a graph $G = (V, E)$ with capacity on links, and a traffic-demand matrix T , where $T[u, v]$ is the amount of traffic that must transit from u to v , for any $u, v \in V$. Find a set of paths and one wavelength assignment for each path such that:

- all demands are routed
- capacity of each link cannot be exceeded
- total number of wavelength is as small as possible

demand matrix:

	a	b	c	d	e	f	g
a	0	0	1	1	0	1	0
b	0	0	0	0	0	2	0
c	0	0	0	0	0	0	0
d	0	0	0	0	0	0	5
e	0	0	0	0	0	0	0
f	0	0	0	0	0	0	0
g	0	1	0	0	0	0	0



Optical Networks (WDM)

RWA problem

Let us simplify the problem


⇒ consider only Wavelength assignment

WA: Wavelength Assignment Problem

Given a graph $G = (V, E)$ with capacity on links, and a set of paths

Give One color to each path s.t. no two paths with the same color cross a same link

Minimize the number of colors



Optical Networks (WDM)

RWA problem

Let us simplify the problem



⇒ consider only Wavelength assignment

WA: Wavelength Assignment Problem

Given a graph $G = (V, E)$ with capacity on links, and a set of paths

Give One color to each path s.t. no two paths with the same color cross a same link

Minimize the number of colors



It is the problem of PROPER COLORING in graphs !!

the “simplified” problem is already NP-complete :(

Optical Networks (WDM)

RWA problem

Let us simplify the problem



⇒ consider only Wavelength assignment

WA: Wavelength Assignment Problem

Given a graph $G = (V, E)$ with capacity on links, and a set of paths

Give One color to each path s.t. no two paths with the same color cross a same link

Minimize the number of colors



It is the problem of PROPER COLORING in graphs !!

the "simplified" problem is already NP-complete :(

Optical Networks (WDM)

RWA problem

Let us simplify the problem



⇒ consider only Wavelength assignment

WA: Wavelength Assignment Problem

Given a graph $G = (V, E)$ with capacity on links, and a set of paths

Give One color to each path s.t. no two paths with the same color cross a same link

Minimize the number of colors



Exercise: Write a (Integer) Linear Programme that solves the RWA problem

Summary: To be remembered

- ILP allow to model many problems
- there may be a huge **integrality gap**
(between $OPT(LP)$ and $OPT(ILP)$).
- if no integrality gap (e.g., **totally unimodular matrices**)
⇒ Fractional Relaxation gives Optimal Integral Solution