

Assessing Planetary Boundaries through an Interactive Accessible Dashboard (WIP)

Author: Leonardo Torres

Date: November 4, 2025

Introduction Section : Accessible analysis on visualizing Planetary Boundaries

In the Anthropocene era, the scale of human influence on the Earth natural system has become so significant that it threatens the overall stability and maintenance of the Holocene like state of the world which has supported the development of human societies for the past millennia. The constant change in global biogeochemical flows, climate change, and biosphere integrity throughout the past few decades are reason why we need to implement some scientific framework to quantify and mitigate the risks our planet is facing. The Planetary Boundaries (PB) framework was established to address this need, proposing quantitative limits for nine critical Earth system processes to define a “safe operating space” for humanity (Rockstrom, 2009). However, a significant implementation gap persists between this high-level conceptual framework and the ability to perform consistent, transparent, and reproducible assessments of boundary status.

Current assessments indicate that multiple planetary boundaries have already been transgressed, moving humanity into a zone of increasing risk for systemic environmental destabilization. Understanding the precise timing, magnitude, and spatial patterns of these transgressions is crucial for informing effective responses and sustainability changes to ensure we don’t reach a high risk zone. Yet, the scientific community has lacked a standardized computational tool to facilitate such analyses across different models and datasets. Existing approaches often rely on dispersed, one-off scripts that are difficult to trace, validate, or reproduce, creating barriers to the rigorous and repeated evaluation of Earth system states (Gerten, 2025).

The recent development of the boundaries R package represents a substantial step toward addressing this methodological gap. This open-source software provides a structured environment for calculating and visualizing PB statuses based on outputs from biosphere models such as LPJmL, which are publically available online. While this package offers powerful analytical capabilities for experts, its complex nterface and computational requirements present a significant accessibility barrier for a broader audience of policymakers, educators, and researchers.

This project aims to offer access to planetary boundaries assessment by developing an interactive, dashboard that serves as a user-friendly interface for the boundaries software. The primary objective is to abstract the underlying computational complexity of the R package while preserving its analytical rigor, enabling users to intuitively explore spatial and temporal patterns of planetary boundary transgression. By providing dynamic visualization tools and configurable parameters, this dashboard will empower non-technical users to generate customized PB assessments, thereby enhancing the transparency, accessibility, and utility of planetary boundaries science for global sustainability and education of our current Earth Status and potential solutions to address it.

Background: Planetary Boundaries and “boundaries” R package.

In this section, I will establish the scientific and methodological groundwork for this project. I will first dive into the Planetary Boundaries (PB) framework as a foundation concept for quantifying global environmental system risk. Afterwards, I will go into what the boundaries R package is, the contribution it offers, and overall its analytical functions for the purpose of this project.

Planetary Boundaries Framework

The Planetary Boundaries framework, introduced by Rockstrom (2009), provides a science-based proposal for understanding and quantifying the limits of anthropogenic perturbation on critical Earth system processes, which we will describe within this section. This framework has been established to represent a planetary state throughout the past ~11,000 years, known as the Holocene epoch, which is the only planetary state known to support humans. The drastic human impact on the Earth system throughout the last few decades has put the planet into a risky position that could be out of the scope of the Holocene epoch, putting us into a stage of non-linear, abrupt, and irreversible environmental changes on a planetary scale.

This framework establishes nine interlinked planetary boundaries where each correspond with an important Earth system process: 1. Climate Change 2. Biosphere Integrity 3. Land-system change 4. Freshwater change 5. Biogeochemical Flows (Nitrogen and Phosphorus cycles) 6. Ocean Acidification 7. Atmospheric Aerosol Loading 8. Stratospheric Ozone Depletion 9. Novel Entities

Back in 2015, this framework was built upon by Steffen (2015) by incorporating:

- A Two-Tier approach: Many processes operate at regional scales, boundaries were defined for both global and sub-global levels (such as for fresh water and land-system change)
- Core Boundaries: Boundaries, such as Climate Change and Biosphere Integrity, regulate the overall state of the Earth system and can individually drive the planet into a new state.
- Refining Control Variables: Control variables and their quantitative boundaries are updated based on latest scientific evidence.

Overall, there are several boundaries at the moment that have already transgressed the safety threshold, placing humanity in a zone to enter a new inhabitable state.

Boundaries R Package

Gerten (2025) developed the boundaries R package, an open-source R software designed to facilitate the standardized calculation and visualization of planetary boundary statuses. This package serves as a post-processing tool

that translates raw model output into PB assessments. Some key features of the package: - Model Input: Designed to work with outputs from the LPJmL (LundPotsdam-Jena managed Land) dynamic global vegetation model, which overall simulates ecological, hydrological, and biogeochemical processes which will then be used in relevancy to terrestrial PBs. - PB Coverage: As mentioned previously, this model is only functionable with terrestrial PBs, this includes the following boundaries: Land-System Change, Biosphere Integrity, Biogeochemical Flows, and Freshwater Change. - Core functionality: - Calculation (`calc_status`) Computes the status of a selected PB for user-defined timer periods and spatial scales within the model - Visualization (`plot_status`) Generates temporal trajectories and spatial patterns to illustrate when and where boundaries are transgressed, when they cross the safety threshold. - Validation (`validate_simulation`) Compares the output of the model against independent data sources to ensure plausibility/accuracy.

This important open-source tool, the boundaries package, will allow us to make an advancement in making PB science more accessible, robust, and reproducible. From this, we will build directly upon from this purpose to create a user-friendly dashboard interface that utilizes the package to correctly understand and visualize planetary boundary assessments

Data sources and Processing:

The primary data source for this dashboard is output from the LPJmL biosphere model, which simulates the growth and productivity of natural and agricultural vegetation in coupling with ecological, hydrological, and biogeochemical processes. The boundaries package post-processes these outputs—which include variables like net primary production, streamflow, soil moisture, and nitrogen fluxes—to compute the status of the four terrestrial planetary boundaries. The data is structured in a standardized netCDF format, allowing for consistent spatiotemporal analysis over the period 1901-2017, relative to a pre-industrial reference state.

Implementation: Building the Dashboard Interface & Functionality

The dashboard (will be) is implemented as a Shiny web application within the R environment. Shiny provides a framework for building interactive web apps directly from R, making it the ideal middleware to connect a user-friendly frontend with the computational backend of the boundaries package. The boundaries R package and its dependencies (e.g., `lpjmkkit`), which perform all complex calculations. The Shiny application server, which handles user input, calls the appropriate boundaries functions, and serves the results. The user interface (UI) built with Shiny’s UI components, rendered in a web browser. This includes control widgets (sliders, dropdowns, boolean) and output areas for maps

and plots. Allow user to choose which of the four boundaries to analyze (or multiple) Allow to switch between viewing temporal trends (line graphs) and spatial patterns (interactive maps). Modify key scientific thresholds (e.g., the forest cover percentage defining the Land-System Change boundary) to explore different scenarios.

Visual Demonstrations: Exploring Scenarios and Visualizations

Discussion: Limitations, Future Enhancements, and Applications in the Real World

Ethical Challenges

The development of an interactive dashboard for planetary boundaries assessment presents several important ethical considerations that have been addressed throughout this project's design and implementation.

Data Transparency and Reproducibility

Without a clear documentation of data and calculation methods, there are risks of this data being misinterpreted or misused in policy/academic contexts. This should be addressed by: 1. Follow established scientific methodologies from the boundaries R package. 2. Provide documentation of all data sources and model assumptions within the dashboard. 3. Enable full reproducibility through Github. 4. Keep the project open-source in order to maintain validation and allowing for community input.

Scientific Communication and Risk of Misinterpretation

The visualizations of planetary boundary data in an accessible dashboard carries the risk of oversimplification or misinterpretation. Users without technical backgrounds might draw inappropriate or incorrect conclusions from presented data. To mitigate such a risk we can: 1. Include contextual information and explanatory notes for each planetary boundary metric. 2. Establish the distinction between the zones of increasing risk, high risk, and holocene. 3. Establish what the limitations are of the model, such that we only have access to certain terrestrial planetary boundaries and terrestrial values in the world.

Global Sustainability

The planetary boundaries framework inherently raises questions of global equity and justice. The current rules documented in the dashboard result dis-

proportionately from historical and contemporary resource use by industrialized nations, while the consequences often affect vulnerable populations most severely. This project acknowledges these ethical worries by: 1. Clearly presenting data at multiple scales (global, regional, and biome level) to highlight different impacts. 2. Avoiding framing that suggests equal responsibility or equal consequences across different regions and populations

Open Access and Knowledge

This project is built on the belief that important scientific information should be available to everyone. To achieve this, the dashboard is designed to be free to use and easy to understand, even for people who are not experts. This ensures that students, teachers, and community leaders can all use it to learn about planetary boundaries. The code behind the dashboard is also open for anyone to see and improve, encouraging a community effort to make the tool better for everyone. Some demo or examples will be available as well to ensure people who want to access this knowledge can freely do this.

Responsible Implementation/Use and Impact Considerations

The dashboard is meant to provide reliable information and support discussion surrounding planetary boundaries. It does not tell users what decisions to make or what policies to support as these worlds don't reflect the real one. The data shows the physical limits of our planet, but how we choose to live within those limits involves complex social and ethical choices that are up to society to decide. By focusing on presenting the facts clearly and without bias, the tool aims to empower users to make their own informed choices. In order to ensure this message goes across, users must understand a terms and conditions that shows that show data and visualizations is for the sole purpose of getting a better understanding of planetary boundaries within a scope of another world.

Conclusion & Reflection

The project provides an example of an accessible interface built for complex scientific computational tools. The dashboard built from the boundaries R package, while retaining the technical complexity of the dashboard, democratization of closure planetary boundaries science. This can enable educators, provide transparent, data-driven visualizations, and support policymakers brainstorm ideas towards change.

This dashboard design and implementation showcase the value of user centered and literate programming in the scientific computing. Breaking the analytical bottleneck while integrating an interface with high performance, analytical backend and is user friendly, yet essential, design task in scientific computing.

Future work will focus on enhancing the dashboard by incorporating multi-model ensemble data to better represent uncertainties, expanding the analysis to include more planetary boundaries, and adding features for comparing different climate or policy scenarios. By continuing to lower the barrier to entry for planetary boundaries science, this project contributes to a more informed and engaged global community, which is a fundamental prerequisite for navigating a sustainable path within our planet's safe operating space.

References

- Braun, Johanna, Jannes Breier, Fabian Stenzel, and Caterina Vanelli. 2025. "Boundaries: Planetary Boundary Status Based on LPJmL Simulations." <https://doi.org/10.5281/zenodo.11550559>.
- Rockström, Johan, Will Steffen, Kevin Noone, Åsa Persson, F. Stuart Chapin, Eric Lambin, Timothy M. Lenton, et al. 2009. "Planetary Boundaries: Exploring the Safe Operating Space for Humanity." *Ecology and Society* 14 (2). <http://www.jstor.org/stable/26268316>.
- Running, Steven W. 2012. "A Measurable Planetary Boundary for the Biosphere." *Science* 337 (6101): 1458–59. <https://doi.org/10.1126/science.1227620>.
- Steffen, Will, Katherine Richardson, Johan Rockström, Sarah E. Cornell, Ingo Fetzer, Elena M. Bennett, Reinette Biggs, et al. 2015. "Planetary Boundaries: Guiding Human Development on a Changing Planet." *Science* 347 (6223): 1259855. <https://doi.org/10.1126/science.1259855>.