Mineralium Deposita

Sources, transport, and deposition of epithermal metal(loid)s recorded by sulfide and rock geochemistry: Constraints from a vertical profile through the epithermal Profitis Ilias Au prospect, Milos Island, Greece

Anna Grosche[1]*, Reiner Klemd[1], Katja Denkel[1], Manuel Keith[1], Karsten M. Haase[1], Panagiotis C. Voudouris[2], Dimitrios Alfieris[3], Michael Wiedenbeck[4]

[1]GeoZentrum Nordbayern, Friedrich-Alexander-Universität (FAU) Erlangen-Nürnberg, Germany
[2]Faculty of Geology & Geoenvironment, National and Kapodistrian University of Athens, Greece
[3]Research and Consulting Geoscientist at DALFEX, Athens, Greece
[4]Deutsches GeoForschungsZentrum GFZ, Potsdam, Germany
*anna.grosche@fau.de

ESM1, Figure S1




Fig. S1. Bar chart showing the results of (A) XRF major element oxide and (B) XRD mineralogical analyses of bulk rock samples of the different host rock lithologies. The samples are ordered by depth from high elevations (left) to low elevations (right).

ESM1, Figure S2




Fig. S2. Lead isotope composition of sphalerite from Profitis Ilias (this study) compared to published data of sulfides from Kondaros-Katsimouti (Schaarschmidt et al. 2021a), Triades (Schaarschmidt et al. 2021a), and Triades-Galana (Marschik et al. 2010).