

Drawing Math

Lev Stambler

May 4, 2022

1 Introduction

- Numberphile video - I was bored and this looked fun - I wasn't paying attention in one of my lectures... - This paper is purely for funs... for now

2 Background

- Digital math - Euler's formula - I paid attention in some of my lectures

3 Definitions and questions

3.1 Definitions

Say you (yes you!) had a turtle living in D dimensional Euclidean space and in discrete time. At time step i , where $i \in \mathbb{Z}$ and $i > 0$, the turtle has position $p_i \in \mathbb{R}^D$. Then, let's define $\Delta p_{i+1} = p_{i+1} - p_i$; in other words, Δp_{i+1} is the change in position from time i to $i + 1$.

Now say that the turtle's movement is determined by k seed parameters drawn from the same set. Then, for some state space \mathcal{S} , define $s_i^j \in \mathcal{S}$ to be some arbitrary state associated with timestamp i for the j th seed parameter where $j \in [k]$. Also, define $\mathbf{s}_i = (s_i^1, s_i^2, \dots, s_i^k)$. Next we will define a set of functions $SU^j : \mathcal{S} \rightarrow \mathcal{S}$ (for *State Updater*) such that $s_{i+1}^j = SU^j(s_i^j, i)$. Note that for $j, a \in [m]$ where $j \neq a$, s_{i+1}^j is determined solely by s_i^j and i and not s_i^a .

Now that we have our machinery built up, let's define $Comb : \mathcal{S}^k \rightarrow \mathbb{R}^d$ such that

$$\Delta p_{i+1} = Comb(s_{i+1}^1, s_{i+1}^2, \dots, s_{i+1}^k).$$

In other words, $Comb$ takes in the state of each seed and returns an update to the position of the turtle.

Finally, let us define

$$\Delta P_{a,b} = \sum_{i=a}^b \Delta p_i.$$

In other words, $\Delta P_{a,b}$ is the change in position from timestep a to b .

3.2 The problem

Say we are given, $Comb$, $hhSU^j$, p_0 , and s_0^j for all $j \in [k]$. Informally, the question is whether the turtle draws a "closed" shape or not.

More formally, is there some period T such that

$$p_{i+\ell T} = p_i$$

for $i, \ell \in \mathbb{N}$. Then, note that if there exists a period T such that $\Delta P_{i,i+\ell T} = 0$ for all $i, \ell \in \mathbb{N}$, $p_{i+\ell T} = p_i$ and the turtle forms a closed shape.

3.3 Specifying the task ahead of us

For our case, we consider $Comb, SU^j$ to all be memoryless (i.e. their output is uniquely determined by the current input). So, we can simplify the overall question. If, $\mathbf{s}_i = \mathbf{s}_{i+\ell T}$ for some $T \in \mathbb{N}$ and all $i \in \mathbb{N}$, then $\Delta p_i = \Delta p_{i+\ell T}$. So then, $\Delta P_{i,i+\ell T} = \Delta P_{i,i+\ell' T}$ for all $\ell, \ell' \in \mathbb{N}$. Thus, T is a period of the change in position. We can thus break down our problem into two parts:

1. Finding the period, T , of the state \mathbf{s} .
2. Checking whether $\Delta P_{i,i+T} = 0$.

3.4 Some more restrictions on our problem

We further restrict the problem by only considering the case where

$$Comb((.,.,.,.,\theta^1), (.,.,.,.,\theta^2), ..., (.,.,.,.,\theta^k)) = \left(\prod_{i=1}^k \cos(\theta_j)^{\text{incl}_1^j}, ..., \prod_{i=1}^k \cos(\theta_j)^{\text{incl}_D^j} \right)$$

where $\text{incl}_d^j \in \{0, 1\}$ for $d \in [D]$ indicates whether to include a given $x \in R$ determined by seed j for position update in the d th dimension.

We will also let $\mathcal{S} = N^5$ where for $(n, d, b, i, \theta) \in \mathcal{S}$, n is the numerator of a rational in fraction form, d is the denominator, b is the base (i.e. base 10, base 12, etc.), i is the current time step, and $\frac{2\theta}{b\pi}$ is an “angle” associated with the state. Moreover, we will only consider

$$SU^j(n, b, d, i, \theta) = (n, b, d, i + 1, \theta + \text{digit}(n, b, d, i) \mod b).$$

where $\text{digit}(n, b, d, i)$ gives us the i th digit of the decimal expansion of $\frac{n}{d}$ in base b .

Finally, for simplicity’s sake, assume that $\theta = 0$ for all $(n, b, d, i, \theta) \in \mathbf{s}_0$, $n < d$, and $\frac{n}{d}$ ’s decimal expansion is periodic after some $N \geq 0$ decimal places and does not terminate in base b .

3.4.1 Some intuition

While the restrictions may seem arbitrary, they aptly match our original problem statement. Take the three dimensional case with two seeds. The two seeds can be thought of as the fraction which generates the updates to two angles, θ, α . θ and α can then uniquely determine a direction in 3D space in which the turtle is pointing. So, at each time step $i + 1$, θ_{i+1} equals to θ_i , but rotated by some fraction of 2π determined by the i th digit of the decimal expansion of one fraction and its base.

4 Results

4.1 Finding period T

4.1.1 Finding the period of $\frac{n}{d}$

We will first aim to find period T of the state \mathbf{s} . For some $(n, b, d, i, \theta) \in \mathbf{s}_0$, by [ho], we have that the period of the decimal expansion of $\frac{n}{d}$ can be determined by finding the smallest $T^{j'}$ such that

$$b^{T^{j'}} \equiv 1 \mod d.$$

See, appendix A for more detail.

4.1.2 Finding the digital root

Next, we introduce the idea of a digital root. TODO: introduce. Basically this is the amount you add with every $T^{j'}$. So, period is $T^{j'} \cdot \text{DigRoot}$.

4.1.3 Finding the period of s^k

4.1.4 Finding the period of s

Simply lcm of all other periods

4.2 $\Delta P_{i,i+T} = 0$?

5 Conclusion

6 Open Questions

Acknowledgments

A Finding the period of $\frac{n}{d}$

aaaaa

References

- [ho] Glen O (https://math.stackexchange.com/users/67842/glen_o). Length of period of decimal expansion of a fraction. Mathematics Stack Exchange. URL:https://math.stackexchange.com/q/2611737 (version: 2018-01-19). 4.1.1