

Math 55b Problem Set 2

Lev Kruglyak

February 8, 2022

I collaborated with AJ LaMotta for this problem set, and it took me about 10 hours.

Problem 1. The *Zariski topology* on \mathbb{R}^2 is the topology generated by the basis

$$U_f = \{(x, y) \in \mathbb{R}^2 \mid f(x, y) \neq 0\}$$

where f ranges over all polynomials in $\mathbb{R}[x, y]$.

- (a) Show that the subsets $U_f \subset \mathbb{R}^2$ are indeed a basis for a topology.
- (b) Show that this topology is not Hausdorff.
- (c) What is the closure \bar{I} of the line segment $I = \{(x, 0) \mid x \in [0, 1]\} \subset \mathbb{R}^2$ in the Zariski topology?
- (d) If $\mathbb{R} \subset \mathbb{R}^2$ is the x -axis, show that the subspace topology on \mathbb{R} induced by the Zariski topology is the finite complement topology.

(a) To prove that U_f form a basis set for a topology, we need to show that the basis elements cover the whole space, and that the intersection of two basis elements contains a third basis element. Clearly, the basis elements cover the whole space, just consider $U_1 = \mathbb{R}^2$. Next, suppose $f_1, f_2 \in \mathbb{R}[x, y]$. Then if $U_{f_1} \cap U_{f_2} \neq \emptyset$, it must contain some point (x_0, y_0) . We are done since $(x, y) = U_{(x-x_0)^2+(y-y_0)^2}$.

(b) If the topology were Hausdorff, then any subspace of it would also be Hausdorff. However by (d), $\mathbb{R} \subset \mathbb{R}^2$ has the finite complement topology, which isn't Hausdorff because any two open sets have nontrivial intersection.

(c) By (d), $\bar{I} \cap \mathbb{R}$ is equal to the closure of $I \cap \mathbb{R}$ in \mathbb{R} with the finite complement topology. However in the finite complement topology, the only closed sets are collections of finite sets or the space itself. Thus $\bar{I} \cap \mathbb{R}$ must be \mathbb{R} since it has an infinite number of points. However \mathbb{R} is a closed subset of \mathbb{R}^2 because $\mathbb{R} = \mathbb{R}^2 - U_y$. So $\bar{I} = \mathbb{R} = \{(x, 0) \mid x \in \mathbb{R}\}$.

(d) Clearly for any $f \in \mathbb{R}[x, y]$, $U_f \cap \mathbb{R} = \{x \in \mathbb{R} \mid f(x, 0) \neq 0\}$. $f(x, 0)$ is a polynomial in $\mathbb{R}[x]$, so it either vanishes completely or has a finite number of roots. This is exactly an open set in the finite complement topology. Conversely, if $U = \mathbb{R} - \{x_1, x_2, \dots, x_n\}$ is an open subset in the finite complement topology, note that $U_{(x-x_1)\dots(x-x_n)} \cap \mathbb{R} = U$.

Problem 2. Let x_1, x_2, x_3, \dots be a sequence of points in a product space $\prod_{\alpha} X_{\alpha}$. Show that this sequence converges to a point x if and only if the sequence $\pi_{\alpha}(x_1), \pi_{\alpha}(x_2), \dots$ converges to $\pi_{\alpha}(x)$ for each α . Is this fact true if one uses the box topology instead of the product topology?

Suppose x_1, x_2, x_3, \dots is a sequence of points in $X = \prod_{\alpha} X_{\alpha}$ converging to some $x \in X$. Then since the projection maps π_{α} are continuous in both product and box topologies, so $\pi_{\alpha}(x_i)$ converge to $\pi_{\alpha}(x)$ in X_{α} .

Conversely, suppose $\pi_{\alpha}(x_i)$ converges to $\pi_{\alpha}(x)$ for all α . Let U be some open neighborhood of x , which contains some open set $\prod_{\alpha} U_{\alpha}$. By definition of the box topology, all but finitely many of the U_{α} are not equal to X_{α} , say $U_{\alpha_1}, \dots, U_{\alpha_n}$. Since $\pi_{\alpha_j}(x_i)$ converges to $\pi_{\alpha_j}(x)$, there exist $N_{\alpha_1}, \dots, N_{\alpha_n}$ such that $m \geq N_{\alpha_j}$ implies $\pi_{\alpha_j}(x_m) \in U_{\alpha_j}$. So letting $N = \max\{N_{\alpha_1}, \dots, N_{\alpha_n}\}$, it follows that $x_m \in \prod_{\alpha} U_{\alpha}$ for all $m \geq N$.

This argument fails in the box topology because the maximum of an infinite collection of integers need not exist, so if you had a sequence in an infinite product space which converges slower and slower in each coordinate, it may never converge in the box topology.

Problem 3. Let \mathbb{R}^{∞} be the subset of \mathbb{R}^{ω} consisting of all sequences that are “eventually zero,” that is, all sequences (x_1, x_2, \dots) such that $x_i \neq 0$ for only finitely many values of i . What is the closure of \mathbb{R}^{∞} in \mathbb{R}^{ω} in the box and product topologies? Justify your answer.

First, we claim that \mathbb{R}^{∞} is dense in \mathbb{R}^{ω} in the product topology, i.e. $\overline{\mathbb{R}^{\infty}} = \mathbb{R}^{\omega}$. Recall that any open set in \mathbb{R}^{ω} is of the form $U = U_1 \times U_2 \times \dots \times U_n \times \mathbb{R} \times \mathbb{R} \dots$. Then, the sequence $x = (x_1, x_2, \dots, x_n, 0, 0, \dots)$ is in U for some $x_i \in U_i$. But $x \in \mathbb{R}^{\infty}$, so every open set in \mathbb{R}^{ω} intersects \mathbb{R}^{∞} nontrivially, which implies density.

In the box topology, \mathbb{R}^{∞} is closed, so $\overline{\mathbb{R}^{\infty}} = \mathbb{R}^{\infty}$. To show this suppose $x \notin \mathbb{R}^{\infty}$, so there are an infinite number of terms in x which are nonzero. Then

$$U = \prod_i U_i \quad \text{where } U_i = \begin{cases} (x_i - |x_i|/2, x_i + |x_i|/2) & x_i \neq 0 \\ (-1, 1) & x_i = 0 \end{cases}$$

is an open set containing x which is disjoint from \mathbb{R}^{∞} .

Problem 4. Let A be a proper subset of X , and let B be a proper subset of Y . If X and Y are connected, show that

$$(X \times Y) - (A \times B)$$

is connected.

Suppose $F : (X \times Y) - (A \times B) \rightarrow \{0, 1\}$ is a continuous function. We claim that F must be constant, since this will imply connectedness of $(X \times Y) - (A \times B)$. (If it was disconnected, then F could take on different values for all of the connected components.)

Fix some $a \in X - A$ and $b \in Y - B$. Now consider an arbitrary point $(x, y) \in (X \times Y) - (A \times B)$. We have two cases. If $x \notin A$, the set $\{x\} \times Y$ is connected so $F(x, y) = F(x, b)$. Similarly, $X \times \{b\}$ is connected, so $F(x, b) = F(a, b)$. So $F(x, y) = F(a, b)$. Otherwise, if $x \in A$ then

$y \notin B$, so we can use a similar argument to get $F(x, y) = F(a, b)$. Since $F(a, b)$ was fixed, it follows that F is constant so $(X \times Y) - (A \times B)$ is connected.

Problem 5. Let \mathbb{R}_ℓ denote the real line with the *lower limit topology*, generated by the basis consisting of all intervals $[a, b)$, $a < b$.

- (a) Show that \mathbb{R}_ℓ is *totally disconnected*, i.e. its only (nonempty) connected subspaces are subsets consisting of a single point.
- (b) Say a function $f : \mathbb{R} \rightarrow \mathbb{R}$ is *continuous from the right* (in the usual topology) if, $\forall a \in \mathbb{R}$, $\lim_{x \rightarrow a^+} f(x) = f(a)$, i.e. $\forall \epsilon > 0, \exists \delta > 0$ s.t. $a < x < a + \delta \Rightarrow |f(x) - f(a)| < \epsilon$. Show that $f : \mathbb{R} \rightarrow \mathbb{R}$ is continuous from the right if and only if it is continuous when considered as a function from \mathbb{R}_ℓ to \mathbb{R} .
- (c) What functions $f : \mathbb{R} \rightarrow \mathbb{R}$ are continuous when considered as maps from \mathbb{R} to \mathbb{R}_ℓ ?
- (d) What can you say about functions which are continuous as maps from \mathbb{R}_ℓ to \mathbb{R}_ℓ ?

(a) Note that we have open sets for any $b \in \mathbb{R}$:

$$(-\infty, b) = \bigcup_{a \leq b} [a, b), \quad [b, \infty) = \bigcup_{b \leq a} [b, a).$$

Then any set X containing more than a single element, say $a < b$ can be separated by disjoint open sets of the form $a \in (-\infty, b)$ and $b \in [b, \infty)$.

(b) First suppose $f : \mathbb{R} \rightarrow \mathbb{R}$ is continuous from the right, and fix $a \in \mathbb{R}$. For each $\epsilon > 0$ there exists a $\delta > 0$ such that for all $x \in \mathbb{R}$ with $a < x < a + \delta$, we have $|f(x) - f(a)| < \epsilon$. This implies that $f([a, a + \delta)) \subset B_\epsilon(f(a))$, so $f^{-1}((a, b))$ is open in the lower limit topology. Conversely suppose $f : \mathbb{R}_\ell \rightarrow \mathbb{R}$ is continuous. Then for any $a \in \mathbb{R}$ and $\epsilon > 0$, by continuity we have $f^{-1}((a - \epsilon, a + \epsilon))$ is open in \mathbb{R}_ℓ . Pick some basis element $[x, y) \subset f^{-1}((a - \epsilon, a + \epsilon))$ containing a . Then we can find some δ such that $[a, a + \delta) \subset [x, y)$, so for any $x \in [a, a + \delta)$, $|f(x) - f(a)| < \epsilon$ by construction. This is exactly a right continuous function.

(c) Note that any continuous map $f : \mathbb{R} \rightarrow \mathbb{R}_\ell$ must map connected sets to connected sets, so $f(\mathbb{R})$ is a connected set in \mathbb{R}_ℓ . However the only connected sets in \mathbb{R}_ℓ are single points, so f is a constant function.

(d) Suppose $f : \mathbb{R}_\ell \rightarrow \mathbb{R}_\ell$ is some continuous function. By (b), since \mathbb{R}_ℓ is finer than \mathbb{R} so any continuous function $f : \mathbb{R}_\ell \rightarrow \mathbb{R}_\ell$ is also continuous $f : \mathbb{R}_\ell \rightarrow \mathbb{R}$. So f must be right continuous by (b).

Note that if $f : \mathbb{R}_\ell \rightarrow \mathbb{R}_\ell$ is continuous, then for any $x \in \mathbb{R}$ and $\epsilon > 0$, there exists a $\delta > 0$ such that for any $y \in \mathbb{R}$ satisfying $x \leq y < x + \delta$ we have $f(x) \leq f(y) < f(x) + \epsilon$. So f must be monotonically increasing. This is a necessary but not sufficient condition.

Problem 6.

- (a) Show that no two of the spaces $(0, 1)$, $(0, 1]$, and $[0, 1]$ are homeomorphic.
- (b) Suppose there exist embeddings $f : X \rightarrow Y$ and $g : Y \rightarrow X$. Show by means of an example that X and Y need not be homeomorphic.
- (c) Show that \mathbb{R}^n and \mathbb{R} are not homeomorphic if $n > 1$.

We'll introduce a powerful invariant to help us answer these questions.

Claim. Let X be a topological space. Let

$$n(X) = \{x \in X \mid X - \{x\} \text{ is path connected}\}.$$

Then if two spaces X, Y are homeomorphic, their associated sets $n(X), n(Y)$ are bijective.

Proof. We'll show that there are injections $\iota_1 : n(X) \rightarrow n(Y)$ and $\iota : n(Y) \rightarrow n(X)$. Indeed, suppose $x \in n(X)$. Then $f(X - \{x\}) = f(X) - \{f(x)\} = Y - \{f(x)\}$ since f is a bijection. However f is also continuous so this image is path connected. Thus $f(x) \in n(Y)$. This gives an injection $n(X) \rightarrow n(Y)$ induced by f . We can do the same thing for f^{-1} , and so there must be a bijection $n(X) \xrightarrow{\sim} n(Y)$. \square

(a) Observe that $n(0, 1) = \emptyset$, $n(0, 1] = \{1\}$, and $n[0, 1] = \{0, 1\}$, hence none of them can be homeomorphic because any homeomorphism would induce bijections of their associated sets.

(b) Consider $(0, 1) \hookrightarrow [0, 1]$ embedded in the natural way, and $[0, 1] \hookrightarrow (0, 1)$ embedded by the map sending $[0, 1] \rightarrow [1/3, 2/3] \subset (0, 1)$. However these spaces aren't homeomorphic by (a).

(c) Using our invariant, note that

$$n(\mathbb{R}^n) = \begin{cases} \mathbb{R} & n = 1 \\ \emptyset & n > 1 \end{cases}$$

because any point from \mathbb{R}^n clearly leaves it path connected for $n > 1$, whereas removing a point x from \mathbb{R} leaves it split into two components, $(-\infty, x) \cup (x, \infty)$.

Problem 7. Let $f : S^1 \rightarrow \mathbb{R}$ be a continuous map. Show that there exists a point $x \in S^1$ such that $f(x) = f(-x)$.

Consider the function $g(x) = f(x) - f(-x)$. Now pick some $x \in S^1$. If $g(x) = 0$, we are done since that would mean that $f(x) = f(-x)$. Otherwise, assume without loss of generality that $g(x) > 0$. Then $g(-x) = f(-x) - f(x) = -g(x) < 0$.

Now reparameterizing the input of g so that it can be considered as a function $\bar{g}(t) : [0, 1] \rightarrow \mathbb{R}$ with $\bar{g}(0) = g(x)$ and $\bar{g}(1) = g(-x)$. Recall that $\bar{g}(0) > 0$ and $\bar{g}(1) < 0$. Thus by the intermediate value theorem there must be a point $t \in [0, 1]$ such that $\bar{g}(t) = 0$. This corresponds to a point $y \in S^1$ satisfying $f(y) = f(-y)$ so we are done.

Problem 8. Let $f : X \rightarrow X$ be a continuous map. Show that if $X = [0, 1]$, there is a point $x \in X$ such that $f(x) = x$. The point x is called a *fixed point* of f . What happens if $X = [0, 1)$ or $(0, 1)$?

Let $g(x) = f(x) - x$. Then $g(0) = f(0) \geq 0$ and $g(1) = f(1) - 1 \leq 0$ so by the intermediate value theorem there must be an $t \in [0, 1]$ such that $g(t) = 0$. Then $f(t) = t$ so we are done.

Problem 9. Show that every compact subspace of a metric space is bounded in that metric and is closed. Find a metric space in which not every closed bounded subspace is compact.

Let C be a compact subspace of a metric space (X, d) . It is clearly closed because compact subspaces are closed in Hausdorff spaces, and every metric space is Hausdorff. To prove that it is bounded, write

$$C \subset \bigcup_{x \in C} B_\epsilon(x) \quad \text{for some } \epsilon > 0.$$

Since C is compact and this is an open cover, there must be some finite collection of points $x_1, x_2, \dots, x_n \in C$ such that $C \subset B_\epsilon(x_1) \cup \dots \cup B_\epsilon(x_n)$. Now let $x \in C$ be arbitrary. Then $C \subset B_r(x)$ where $r = \epsilon + \max_{1 \leq i \leq n} d(x, x_i)$ by the triangle inequality. So C is bounded by a ball of radius r .

To show an example of a metric space where every closed bounded subspace is not compact, consider any infinite set X equipped with the discrete metric, so every set is open. Note that every set is closed and bounded in this space. However any infinite subset S cannot be compact since $S = \bigcup_{s \in S} \{s\}$ is an open cover of S with no finite subcover.

Problem 10. Let A and B be disjoint compact subspaces of the Hausdorff space X . Show that there exist disjoint open sets U and V containing A and B , respectively.

First we'll prove that for any point $a \in A$, there exists open disjoint sets $U \ni a$ and $V \supset B$. Indeed, since X is Hausdorff, we have disjoint open sets $U_b \ni a$ and $V_b \ni b$ for every $b \in B$. Considering $\{V_b\}_{b \in B}$ as an open cover of B , there must be some finite sub collection of the sets also covering B by compactness, say $V_{b_1}, V_{b_2}, \dots, V_{b_n}$. Then $U = \bigcap_i U_{b_i}$ and $V = \bigcup_i V_{b_i}$ satisfy the required conditions.

Now we can construct disjoint open $U \supset A$ and $V \supset B$. By the first paragraph, we have disjoint open sets $U_a \ni a$ and $V_a \supset B$ for every $a \in A$. Since $\{U_a\}_{a \in A}$ is an open cover of A , there must be some finite subcover, say $U_{a_1}, U_{a_2}, \dots, U_{a_n}$. Then $U = \bigcup_i U_{a_i}$ and $V = \bigcap_i V_{a_i}$ satisfy the conditions of the problem.

Problem 11. Let X be the union of \mathbb{R}^n and one additional point called ∞ . Consider the topology with basis given by open balls in \mathbb{R}^n plus the sets $U_r = \{\infty\} \cup \{x \in \mathbb{R}^n \mid |x| > r\}$. Show that X is a compact Hausdorff space.

To show that the space is Hausdorff, it suffices to show that ∞ can be separated from any point in \mathbb{R}^n by disjoint open sets, since any two distinct points in \mathbb{R}^n can already be separated by disjoint open sets. Let $x \in \mathbb{R}^n$. Then letting $U_r(x) = \{\infty\} \cup \{x \in \mathbb{R}^n \mid |x| > r\}$ and choosing some $\epsilon > 0$ we have $\infty \in U_\epsilon(x)$ and $x \in B_\epsilon(x)$, and these sets are clearly disjoint.

To prove compactness, let \mathcal{U} is some open cover of X . Pick some $V \in \mathcal{U}$ such that $\infty \in V$. We can assume without loss of generality that V is a basis element (otherwise split it into a union of basis elements), so $V = X - C$ for some compact ball C . So when restricted to \mathbb{R}^n the set $\mathcal{U} - \{V\}$ is an open cover of C . (Clearly any set of the form $\mathbb{R}^n - C_1$ is open since \mathbb{R}^n is a Hausdorff space so any compact C_1 is closed) Since C is compact, there is some finite subcover $\mathcal{V} \subset \mathcal{U} - \{V\}$. The $\mathcal{V} \cup \{V\}$ is an open cover of X .