

MATH 231BR: ADVANCED ALGEBRAIC TOPOLOGY
HOMEWORK 9
DUE: TUESDAY, APRIL 18 AT 12:00AM (MIDNIGHT) ON CANVAS

In the below, I use LAT to refer to Miller's *Lectures on Algebraic Topology*, available at:
<https://math.mit.edu/~hrm/papers/lectures-905-906.pdf>.

1. PROBLEM 1: JAMES SPLITTING OF $\Sigma\Omega S^{2n+1}$ (15 POINTS)

- (a) Given two path-connected pointed spaces X and Y , prove that there is a splitting

$$\Sigma(X \times Y) \simeq \Sigma X \vee \Sigma Y \vee \Sigma(X \wedge Y).$$

(Hint: use the pinch map $\Sigma Z \rightarrow \Sigma Z \vee \Sigma Z$.)¹

- (b) Composition of loops $\Omega S^{2n+1} \times \Omega S^{2n+1} \rightarrow \Omega^{2n+1}$ makes $H_*(\Omega S^{2n+1})$ into a ring. Prove that $H_*(\Omega S^{2n+1}) \cong \mathbb{Z}[x_{2n}]$. (Hint: the product $H_*(\Omega S^{2n+1}) \otimes H_*(\Omega S^{2n+1}) \rightarrow H_*(\Omega S^{2n+1})$ is dual to a coassociative and counital coproduct $H^*(\Omega S^{2n+1}) \rightarrow H^*(\Omega S^{2n+1}) \otimes H^*(\Omega S^{2n+1})$ which itself is a map of rings.)
- (c) Prove that $\Sigma\Omega S^{2n+1} \simeq \bigvee_{k=1}^{\infty} S^{2kn+1}$.²

2. PROBLEM 2: EHP SEQUENCE (15 POINTS)

- (a) Using 1(c), construct a map $H : \Omega S^{2n+1} \rightarrow \Omega S^{4n+1}$ which induces an isomorphism in H_{4n} .
- (b) Let $E : S^{2n} \rightarrow \Omega S^{2n+1}$ denote the adjoint to the identity on S^{2n+1} . Using the Serre spectral sequence, prove that

$$S^{2n} \xrightarrow{E} \Omega S^{2n+1} \xrightarrow{H} \Omega S^{4n+1}$$

is a mod \mathcal{C}_2 -fiber sequence, i.e. that the map $S^{2n} \rightarrow F$ induces a mod \mathcal{C}_2 -isomorphism on homotopy groups.³ The induced long exact sequence of mod \mathcal{C}_2 homotopy groups is called the *EHP sequence*.

3. PROBLEM 3: COHOMOLOGY OF $V_2(\mathbb{R}^n)$ (10 POINTS)

Let $V_2(\mathbb{R}^n)$ denote the space of pairs (x_1, x_2) of orthonormal vectors in \mathbb{R}^n .

- (a) Identify the map $V_2(\mathbb{R}^n) \rightarrow S^{n-1}$ which sends $(x_1, x_2) \mapsto x_1$ with the unit sphere bundle associated to the tangent bundle of S^{n-1} .
- (b) Using the fact that $\langle e(TM), [M] \rangle = \chi(M)$, compute the cohomology rings $H^*(V_2(\mathbb{R}^n); \mathbb{F}_2)$ and $H^*(V_2(\mathbb{R}^n); \mathbb{Z})$.

4. PROBLEM 4: COHOMOLOGY OF G_2 (10 POINTS)

The exceptional Lie group G_2 lies in a fiber sequence

$$S^3 \rightarrow G_2 \rightarrow V_2(\mathbb{R}^6).$$

Compute the integral and mod 2 cohomology groups of G_2 using the Serre spectral sequence. Explain why the Serre spectral sequence is unable to uniquely determine the ring structure without some additional input.

¹The statement is true without the path-connected assumption, but including this assumption simplifies the proof by allowing you to use homology Whitehead.

²The same results are all true for ΩS^{2n} , but the proof requires a little bit more work, since the rationalization of $H^*(\Omega S^{2n+1})$ is no longer generated by a single element.

³Again, the same result is true for ΩS^{2n} . There is also an odd-primary version, which is a bit more complicated.