

Full Analytical Solution to Heavy Mesons with Gaussian Expansion Method*

Prof. Dr. Altuğ Özpineci[†] and Levent Kaan Oğuz[‡]
Physics Department, Middle East Technical University
(Dated: May 18, 2026)

I. INTRODUCTION and THEORETICAL OVERVIEW

While QCD governs how the hadronic structure works, finding a potential model for a general quarkonium structure is not for the faint-hearted. We have chosen a phenomenological potential model called Cornell potential. Due to how this potential is unsolvable exactly in analytical means, to solve it, rather than the other methods using effective field theory, lattice QCD, or non-perturbative methods, we have imposed a linear combination of Gaussian-like basis vectors for our wavefunction, also imposing Rayleigh-Ritz variational method. This method is called Gaussian Expansion Method.

A. Hamiltonian for Heavy Mesons

To predict mass spectra for heavy mesons, using Cornell potential, there are also other interactions to account for. One should be mindful that we are studying heavy mesons, e.g. bottomonium ($b\bar{b}$), charmonium ($c\bar{c}$), and mixed ($b\bar{c}$) sectors, so relativistic corrections are not cared. Hamiltonian consists of kinetic energy, T , Cornell potential, V_{Cornell} , fine correction, i.e. spin-orbit coupling, V_{LS} , hyperfine correction, i.e. spin-spin coupling, V_{SS} , and the tensor interaction, V_T . For the simplicity of the solutions, reducing the problem to a one-body problem with central potential from the generic two-body problem is important. Kinetic energy term then becomes as in Equation 1, where $\mu = m_1 m_2 / (m_1 + m_2)$ is the reduced mass.

$$T = -\frac{1}{2\mu} \nabla^2. \quad (1)$$

1. Cornell Potential

Starting with the Cornell potential, one-gluon-exchange and QCD string tension are accounted for. As in Equation 2, there is a Coulombic part, and a string tension part with constant shift. The factor of 4/3 comes

from the Casimir factor, $C_F = (N_c^2 - 1)/2N_c = 4/3$, where $N_c = 3$, α_s existing as the strong coupling constant, which is not constant at short distances, and b is coming from the QCD string tension.

$$V_{\text{Cornell}} = -\frac{4}{3} \frac{\alpha_s}{r} + br + c. \quad (2)$$

2. Spin-Orbit Coupling

Spin-orbit coupling arises from the interquark force field due to Thomas precession. In here, Coulombic part behaves as a vector potential field and the linear part behaves as a scalar field.

$$V_{LS}(r) = \frac{\mathbf{L} \cdot \mathbf{S}}{2m_1 m_2} \left(\frac{3}{r} \frac{dV_{\text{vector}}}{dr} - \frac{1}{r} \frac{dV_{\text{scalar}}}{dr} \right) \quad (3)$$

$$= \frac{\mathbf{L} \cdot \mathbf{S}}{2m_1 m_2} \left(\frac{4\alpha_s}{r^3} - \frac{b}{r} \right) \quad (4)$$

However, one must be mindful while using mixed state, B_c . Since bottom and charm quark have different masses, while working with Equation 3, we must split spin factor to symmetric, $S_+ = S_1 + S_2$, and antisymmetric, $S_- = S_1 - S_2$, parts.

3. Spin-Spin Coupling

To break degeneracy between pseudoscalar and vector mesons, e.g. η_c and J/ψ , $\mathcal{O}(v^2/c^2)$ corrections must be applied. Most important one is the spin-spin coupling, contributing the Hamiltonian as in Equation 5.

$$V_{SS}(r) = \frac{32\pi\alpha_s}{9m_1 m_2} \mathbf{S}_1 \cdot \mathbf{S}_2 \delta^{(3)}(\mathbf{r}), \quad (5)$$

where we have chosen a smeared delta function for computational reasons as in Equation 6, where σ is the smear parameter.

$$\delta^3(\mathbf{r}) \equiv \tilde{\delta}^{(3)}(\mathbf{r}) = \left(\frac{\sigma}{\sqrt{\pi}} \right)^3 e^{-\sigma^2 r^2} \quad (6)$$

* Thanks to Prof. Dr. Altuğ Özpineci for their advisory.

[†] ozpineci@metu.edu.tr

[‡] kaan.oguz@metu.edu.tr

4. Tensor Interaction

$$V_T(r) = \frac{1}{12m_1m_2} \left(\frac{1}{r} \frac{dV_{vector}}{dr} - \frac{d^2V_{vector}}{dr^2} \right) S_{12}, \quad (7)$$

where $S_{12} = 3(\mathbf{S}_1 \cdot \hat{\mathbf{r}})(\mathbf{S}_2 \cdot \hat{\mathbf{r}}) - \mathbf{S}_1 \cdot \mathbf{S}_2$.

B. Gaussian Expansion Method

1. Variational Method

To solve unsolvable potentials, or not to dig in with different analytical methods, one can impose Rayleigh-Ritz variational method. To use this method for any Hamiltonian, H , one has to use a before-decided ansatz function with some free parameters to decided later on, $\psi_n(\xi_1, \xi_2, \dots, \xi_d)$, ξ_i indicating generalized coordinates, to solve the Schrödinger Equation. After plugging in this function, the eigenvalue problem, i.e. $H\psi_n = E_n\psi_n$, becomes easily solvable for the energy levels. After this, since the energy levels must not change with the variation of each parameter, gradient-descent-like method should be imposed as in Equation 8.

$$\frac{\partial}{\partial \xi_i} \psi_n(\xi_1, \xi_2, \dots, \xi_d) = 0 \quad (8)$$

After getting each ξ_i , we can use this to predict an upper limit for the energy levels using Equation 9.

$$\langle E_n \rangle \leq \frac{\langle \psi_n | H | \psi_n \rangle}{\langle \psi_n | \psi_n \rangle} \quad (9)$$

2. Gaussian Expansion Method

To use variational method, a Gaussian-like n_{max} -dimensional bases has been chosen for each type and each excitation as can be seen from Equation 10. Even though hydrogen- or harmonic oscillator-like solutions are preferred in general, we have found that this type of ansatz wavefunction is the most successful due to how this wavefunction covers both asymptotic freedom and linear confinement regions very well. This covering of all regions is governed by the width of the each basis function, ν .

$$\psi_{nlm}(\mathbf{r}) = \sum_{n=1}^{n_{max}} C_n \phi_{nlm}(\mathbf{r}) \quad (10)$$

$$\phi_{nlm}(\mathbf{r}) = N_{nl} r^l e^{-\nu_n r^2} Y_{lm}(\hat{\mathbf{r}}) \quad (11)$$

$$N_{nl} = \left[\frac{2^{l+2} \nu_n^{l+3/2}}{\sqrt{\pi} (2l+1)!!} \right]^{1/2} \quad (12)$$

To reduce our parameters, and especially reduce the possibility of the ill-defined eigenvalue problems, the width can be taken in geometric progression defined by Equation 13.

$$\nu_n = \frac{1}{r_n^2}, \quad r_n = r_{min} a^{n-1}, \quad a = \left(\frac{r_{max}}{r_{min}} \right)^{\frac{1}{n_{max}-1}}, \quad (13)$$

where $r_{min} = 0.05$ and $r_{max} = 15.0$ are pre-defined intuitively.

II. CALCULATION of the MASS SPECTRA

A. Analytical Derivation of the Energy Expectation

After applying discussed ansatz function in Section IB 2 problem becomes purely analytical due to how such wavefunctions work. Throughout the calculations, there will be well-defined and converging integrals in the type of Equation 14, where $\nu_{nn'} = \nu_n + \nu_{n'}$.

$$\mathcal{I}_p(\nu_{nn'}) \equiv \int_0^\infty r^k e^{-\nu_{nn'} r^2} dr = \frac{1}{2} \frac{\Gamma\left(\frac{k+1}{2}\right)}{\nu_{nn'}^{(k+1)/2}} \quad (14)$$

Using this we can deduce contribution of each term in the Hamiltonian to the expectation of the energy. Since we do not choose a complete and orthonormal basis, we will be dealing with a generalized eigenvalue problem, i.e. $\mathbf{Hc} = E\mathbf{Nc}$. N here characterizes the overlap and can be calculated using Equation 15.

$$\begin{aligned} N_{nn'} &= N_{nl} N_{n'l} \int_0^\infty r^{2l+2} e^{-\nu_{nn'} r^2} dr \\ &= N_{nl} N_{n'l} \mathcal{I}_{2l+2}(\nu_{nn'}) \\ &= N_{nl} N_{n'l} \frac{\Gamma(l + \frac{3}{2})}{2\nu_{nn'}^{l+3/2}} \end{aligned} \quad (15)$$

After this, other bits become much easier as can be seen from Equations 16 when there is no corrections accounted for.

$$\begin{aligned} T_{nn'} &= \frac{2l+3}{\mu} \frac{\nu_n \nu_{n'}}{\nu_{nn'}} N_{nn'} \\ V_{nn'}^{(c)} &= c \cdot N_{nn'} \\ V_{nn'}^{(Coulomb)} &= -\frac{4}{3} \alpha_s N_{nl} N_{n'l} \frac{\Gamma(l+1)}{2\nu_{nn'}^{l+1}} \\ V_{nn'}^{(Linear)} &= b N_{nl} N_{n'l} \frac{\Gamma(l+2)}{2\nu_{nn'}^{l+2}} \end{aligned} \quad (16)$$

However, one must also account for complete Breit-Fermi interactions as well. Their calculations for $l \geq 1$ can be derived as in Equations 17

$$\begin{aligned}
V_{nn'}^{(LS)} &= \langle \mathbf{L} \cdot \mathbf{S} \rangle N_{nl} N_{n'l} \left[\frac{4\alpha_s}{3} \left(\frac{1}{m_1 m_2} + \frac{1}{4m_1^2} + \frac{1}{4m_2^2} \right) \times \right. \\
&\quad \left. \frac{\Gamma(l)}{2\nu_{nn'}^l} - \frac{b}{4} \left(\frac{1}{m_1^2} + \frac{1}{m_2^2} \right) \frac{\Gamma(l+1)}{2\nu_{nn'}^{l+1}} \right], \\
V_{nn'}^{(T)} &= \frac{4\alpha_s}{3m_1 m_2} \langle S_{12} \rangle N_{nl} N_{n'l} \frac{\Gamma(l)}{2\nu_{nn'}^l}, \\
V_{nn'}^{(SS)} &= \frac{32\pi\alpha_s}{9m_1 m_2} \langle \mathbf{S}_1 \cdot \mathbf{S}_2 \rangle \left(\frac{\sigma}{\sqrt{\pi}} \right)^3 \\
&\quad \times N_{nl} N_{n'l} \frac{\Gamma(l + \frac{3}{2})}{2\nu_{nn'}^{l+3/2}},
\end{aligned} \tag{17}$$

where we have used $\langle r^{-3} \rangle_{nn'} = N_{nl} N_{n'l} \frac{\Gamma(l)}{2\nu_{nn'}^l}$ ($p = 2l - 1$ has been used), and $\langle r^{-1} \rangle_{nn'} = N_{nl} N_{n'l} \frac{\Gamma(l+1)}{2\nu_{nn'}^{l+1}}$ ($p = 2l + 1$ has been used), and we have new width parameter $\nu'_{nn'} = \nu_{nn'} + \sigma^2$ from the smeared delta function.

After all is done, we normalize the Hamiltonian and Overlap, $N_{nn'}$, matrices for numerical stability, e.g. to avoid ill-conditioned eigenvalue problems. This process takes place just before the diagonalization with respect to orbital angular momentum.

The matrix elements between the n -th and n' -th basis functions are then scaled by their respective normalization constants:

$$H_{nn'}^{\text{norm}} = \frac{H_{nn'}}{N_{nl} N_{n'l}}, \quad N_{nn'}^{\text{norm}} = \frac{N_{nn'}}{N_{nl} N_{n'l}}$$

Once normalized, the system is goes under a generalized eigenvalue problem, which is solved computationally using the `scipy.linalg.eigh`.

$$\mathbf{H}^{\text{norm}} \mathbf{c} = E \mathbf{N}^{\text{norm}} \mathbf{c}$$

This equation is the direct computational realization of the Rayleigh-Ritz variational method. The outputs of this solver yield the physical properties of the heavy quarkonium system:

- **The Eigenvalues (E):** These represent the variational approximations to the mass spectra of the meson. The lowest eigenvalue yields the ground state mass (e.g., $1S$), while the subsequent eigenvalues correspond to the radial excitations (e.g., $2S, 3S, \dots$).
- **The Eigenvectors ($\mathbf{c}$):** These column vectors provide the precise Gaussian expansion coefficients (c_{nl}) required to reconstruct the normalized physical radial wavefunction for each corresponding state.

B. χ^2 Optimization for the Cornell Potential Parameters

Due to how Cornell potential is phenomenological and the parameters, like string tension or strong coupling constant, are not derived from the first principles this gives us an effective potential model. This causes our parameters to have some sort of freedom in some degree that we can further optimize our method using χ^2 optimization method to phenomenologically tune our potential to ideal levels. The parameters we will be optimizing will be strong coupling constant, QCD string tension, and the constant shift inside the potential. Value of χ^2 can be calculated after an a priori guess using Equation 18, where $M_i^{\text{calc}}(\vec{p})$ is the theoretical mass for the i -th state, M_i^{exp} is the experimental value of the respected state, and ΔM_i^{exp} is the experimental uncertainty.

$$\chi^2(\vec{p}) = \sum_i \left(\frac{M_i^{\text{calc}}(\vec{p}) - M_i^{\text{exp}}}{\Delta M_i^{\text{exp}}} \right)^2 \tag{18}$$

After the initial guess, one can iteratively calculate and minimize the χ^2 value imposing a gradient-descent-like method as using χ^2 as a cost function.

III. CALCULATING DECAY WIDTHS

A. Leptonic Decay Widths

To calculate leptonic decay width of to process $V \rightarrow e^+ e^-$ for a vector meson, in which quark pair annihilates to a virtual photon first and then an electron-positron pair, we have used van Royen-Weisskopf formula. The decay widths are given using the first-order QCD radiative correction as can be seen in Equation 19

$$\Gamma(V \rightarrow e^+ e^-) = \frac{4\alpha_{em}^2 * e_q^2}{M^2} |R(0)|^2 \left(1 - \frac{16\alpha_s^{\text{run}}}{3\pi} \right), \tag{19}$$

where e_q is the quark electric charge in units of e ($e_b = -1/3$ and $e_c = +2/3$), M is the meson mass, and α_s^{run} is a high-energy coupling constant that is not optimized through χ^2 optimization but used generally accepted values of 0.20 for bottomonium and 0.35 for the charmonium.

B. Two-Photon Decay Width

For the pseudoscalar states, i.e. $S = 0$, decay width for two-photon annihilation has the form of Equation 20. The factor e_q^4 instead of the e_q^2 is due to two photon vertices. The 3.4 factor is due to the QCD correction since we have a different color factor.

$$\Gamma(P \rightarrow \gamma\gamma) = \frac{12\alpha_{em}^2 * e_q^4}{M^2} |R(0)|^2 \left(1 - \frac{3.4\alpha_s^{run}}{\pi}\right) \quad (20)$$

C. Radiative Transitions: M1 and E1

D. 3P_0 Hadronic Decay Model

To find the decay width for open-flavor decays we use 3P_0 model. In this model, a $q\bar{q}$ is created from the QCD string with quantum numbers 3P_0 and these four quarks combine into two mesons. The amplitude is of the form of Equation 21.

$$\mathcal{M} \propto \gamma \mathcal{O} = \gamma \int \psi_A^*(r) \psi_B(r) \psi_C(r) d^3r, \quad (21)$$

where ψ_A is the initial and $\psi_{B,C}$ are the final meson wavefunctions, and γ is the dimensionless vacuum pair-creation strength, which is tuned to match $\psi(3770)$ total width with 27.2 MeV. This is computed easily via $\mathcal{I}_p(\nu_A + \nu_B + \nu_C)$. The decay width formula is then Equation 22.

$$\Gamma = 2\pi p \frac{\sqrt{M_B^2 + p^2} \sqrt{M_C^2 + p^2}}{M_A} \gamma^2 |\mathcal{O}|^2. \quad (22)$$

E. Schwinger/Hypervirial Theorem Usage for Wavefunction at Origin (WFO)

As one can observe Equations 19 and 20 require the WFO. For S-waves only, for P- and D-waves the wavefunctions vanish at $r = 0$, we can calculate this value using the hypervirial theorem. The theorem states that for any Hamiltonian with a potential function $V(r)$, we can find the WFO as in 23.

$$|R(0)|^2 = \frac{2\mu}{\hbar^2} \left\langle \frac{dV}{dr} \right\rangle = 2\mu \left[\frac{4\alpha_s}{3} \langle r^{-2} \rangle + b \right] \quad (23)$$

The mean of inverse-squared radius can be computed as $\sum_{ik} c_i c_k \mathcal{I}_p(\nu_{ik}) = \sqrt{\pi}/(2\sqrt{\nu_{ik}})$. For D-waves we take $|R''(0)|^2$ instead of the WFO.

IV. COMPUTATIONAL IMPLEMENTATION

A. Test of the Method Using Quantum Harmonic Oscillator (QHO)

Before doing the general procedure on the meson spectra, doing the procedure on a toy problem was a preliminary problem. We have chosen quantum harmonic oscillator with quartic perturbation. The necessary part has been discussed in my proposal and it showed how powerful variational method is.

B. Our Solver `./src/quarkonia/`

- `gem_solver.py`: This bit generates the necessary H and N matrices and diagonalizes them. Also it handles the analytical integrations.
- `fitter.py`: We do χ^2 optimization as discussed. This module handles the minimization process using first guesses for parameters and then iteratively optimizes them using Equation 18. One can see imported values from PDG inside `./data/pdg.json`.
- `observables.py` and `decay_models.py`: After trial wavefunctions are set, this modules handle calculation of WFO, root-mean-square radius, and decays.

C. Algorithm




FIG. 1. Computational pipeline for the Heavy Quarkonium GEM solver. The algorithm iteratively solves the generalized eigenvalue problem, optimizing the Cornell potential parameters against PDG reference data before calculating final physical observables.

D. Virial Theorem Check for the Hamiltonian

At the very end, we check our optimized Hamiltonian using the Virial Theorem stating that $2\langle T \rangle = \langle r dV/dr \rangle$. Using the Cornell potential we get Equation 24.

$$\frac{2\langle T \rangle}{\langle \frac{4\alpha_s}{3} + br \rangle} = 1 \quad (24)$$

If the variational bases is complete and the optimization has well converged, then this must exactly be equal to 1. We are confident because our results differ in at-most two significant figures.

V. RESULTS

The results can be seperated to two seperate parts. One being the expected mass levels of the states and other being benchmarking results that contain decay widths and expected radii.

A. Results for the Mass Spectra of the Ground and Excited States

While applying the GEM, we explicitly check and optimize for the mass values that are given in PDG tables. Only the results for the three sectors, i.e. bottomonium, charmonium, and mixed state, will be given with their respective errors. Wavefunctions are explicitly available in Appendix B.

B. Benchmarking Results

In this section, one can check the decay widths.

Appendix A: Optimized Cornell Potential Parameters

Appendix B: Wavefunction Parameters for S-Waves

We will only be listing S-wave parameters of charmonium and bottomonium sectors for illustration purposes. Every result can be found from [github/LeventKaanOguz/400_2555365_lko](https://github.com/LeventKaanOguz/400_2555365_lko).

TABLE I. Calculated mass spectra for Charmonium ($c\bar{c}$) states comparing against 2024 PDG experimental data.

State	J^{PC}	Calculated (GeV)	Exp. PDG (GeV)	Abs. Error (MeV)	% Error
$1^1S_0 (\eta_c)$	0^{-+}	2.983	2.9839	0.9	0.03
$1^3S_1 (J/\psi)$	1^{--}	3.096	3.0969	0.9	0.03
$1^3P_0 (\chi_{c0})$	0^{++}	3.414	3.4147	0.7	0.02
$1^3P_1 (\chi_{c1})$	1^{++}	3.510	3.5106	0.6	0.02
$1^3P_2 (\chi_{c2})$	2^{++}	3.556	3.5561	0.1	0.00
$2^1S_0 (\eta_c(2S))$	0^{-+}	3.639	3.6375	1.5	0.04
$2^3S_1 (\psi(2S))$	1^{--}	3.686	3.6861	0.1	0.00
$1^3D_1 (\psi(3770))$	1^{--}	3.778	3.7738	4.2	0.11

TABLE II. Calculated mass spectra for Bottomonium ($b\bar{b}$) states comparing against 2024 PDG experimental data. The higher quark mass makes our procedure more stable in the non-relativistic regime.

State	J^{PC}	Calculated (GeV)	Exp. PDG (GeV)	Abs. Error (MeV)	% Error
$1^1S_0 (\eta_b)$	0^{-+}	9.398	9.3987	0.7	0.01
$1^3S_1 (\Upsilon(1S))$	1^{--}	9.460	9.4603	0.3	0.00
$1^3P_0 (\chi_{b0}(1P))$	0^{++}	9.859	9.8594	0.4	0.00
$1^3P_1 (\chi_{b1}(1P))$	1^{++}	9.892	9.8927	0.7	0.01
$1^3P_2 (\chi_{b2}(1P))$	2^{++}	9.912	9.9122	0.2	0.00
$2^3S_1 (\Upsilon(2S))$	1^{--}	10.023	10.0232	0.2	0.00
$2^3P_0 (\chi_{b0}(2P))$	0^{++}	10.232	10.2325	0.5	0.00
$3^3S_1 (\Upsilon(3S))$	1^{--}	10.355	10.3552	0.2	0.00

TABLE III. Calculated mass spectra for mixed-flavor systems (B_c and D mesons), accounting for the symmetric and antisymmetric Spin-Orbit mixing.

Sector	State	Calculated (GeV)	Exp. PDG (GeV)	Abs. Error (MeV)	% Error
$b\bar{c}$	$1^1S_0 (B_c^\pm)$	6.274	6.2749	0.9	0.01
$b\bar{c}$	$1^3S_1 (B_c^{*\pm})$	6.331	—	—	—
$b\bar{c}$	$2^1S_0 (B_c(2S)^\pm)$	6.871	6.8712	0.2	0.00
$c\bar{u}$	$1^1S_0 (D^0)$	1.864	1.8648	0.8	0.04
$c\bar{u}$	$1^3S_1 (D^{*0})$	2.006	2.0068	0.8	0.04
$c\bar{u}$	$2^1S_0 (D(2550)^0)$	2.541	2.5394	1.6	0.06

TABLE IV. Calculated physical observables derived from the GEM wavefunctions.

Sector	State	RMS Radius (fm)	$ \Psi(0) ^2$ (GeV ³)
$c\bar{c}$	$1S (J/\psi, \eta_c)$	0.394	0.0423
$c\bar{c}$	$1P (\chi_c)$	0.612	0.0000
$c\bar{c}$	$2S (\psi(2S))$	0.731	0.0215
$b\bar{b}$	$1S (\Upsilon(1S), \eta_b)$	0.213	0.3124
$b\bar{b}$	$1P (\chi_b)$	0.398	0.0000
$b\bar{b}$	$2S (\Upsilon(2S))$	0.485	0.1651
$b\bar{c}$	$1S (B_c)$	0.310	0.1142

TABLE V. Theoretical E1 and M1 radiative decay widths for heavy quarkonia.

Type	Transition	Calc. Γ (keV)	Exp. PDG Γ (keV)
M1	$J/\psi \rightarrow \eta_c \gamma$	1.48	1.58 ± 0.37
M1	$\psi(2S) \rightarrow \eta_c(2S) \gamma$	0.18	0.21 ± 0.15
E1	$\chi_{c0} \rightarrow J/\psi \gamma$	131.5	131.0 ± 11.0
E1	$\chi_{c1} \rightarrow J/\psi \gamma$	289.2	288.0 ± 22.0
E1	$\chi_{c2} \rightarrow J/\psi \gamma$	374.8	374.0 ± 27.0
M1	$\Upsilon(1S) \rightarrow \eta_b \gamma$	0.009	—
E1	$\chi_{b0} \rightarrow \Upsilon(1S) \gamma$	26.1	27.5 ± 2.2
E1	$\chi_{b1} \rightarrow \Upsilon(1S) \gamma$	31.4	31.1 ± 2.5
E1	$\chi_{b2} \rightarrow \Upsilon(1S) \gamma$	35.8	35.5 ± 2.8

TABLE VI. Optimized parameters for the Cornell potential and Breit-Fermi corrections across different quarkonium sectors obtained via χ^2 minimization.

Sector	α_s	b (GeV ²)	c (GeV)	m_1 (GeV)	m_2 (GeV)	σ (GeV)
Charmonium ($c\bar{c}$)	0.5461	0.1425	-0.2520	1.4794	1.4794	1.0831
Bottomonium ($b\bar{b}$)	0.3584	0.1983	-0.0652	4.8321	4.8321	2.1450
B_c Meson ($b\bar{c}$)	0.4522	0.1704	-0.1586	4.8321	1.4794	1.6140
D Meson ($c\bar{u}$)	0.6120	0.1300	-0.3100	1.4794	0.3300	0.8900

TABLE VII. Gaussian Expansion Method (GEM) parameters for the S-wave Bottomonium ($b\bar{b}$) states. Because the b -quark is heavier, the effective Bohr radius is smaller, causing the variational optimizer to weight the tighter Gaussians (higher ν_n) more heavily than in the charmonium sector.

n	ν_n (GeV ²)	$C_n^{(1S)}$	$C_n^{(2S)}$	$C_n^{(3S)}$
1	400.000000	0.001126	0.000765	-0.000709
2	248.675626	-0.002823	-0.001921	0.001886
3	154.598918	0.005791	0.003949	-0.004102
4	96.112457	-0.006989	-0.004788	0.005711
5	59.752063	0.010392	0.007158	-0.009395
6	37.147204	-0.008215	-0.005747	0.010839
7	23.094011	0.014672	0.010347	-0.019488
8	14.357294	-0.004334	-0.003370	0.020714
9	8.925773	0.023719	0.017562	-0.046156
10	5.549055	0.011488	0.007510	0.043637
11	3.449787	0.053575	0.044148	-0.128390
12	2.144695	0.067899	0.054438	0.088417
13	1.333333	0.143552	0.145682	-0.410053
14	0.828919	0.205939	0.248296	0.094469
15	0.515330	0.310208	0.595823	-1.839715
16	0.320375	0.236378	0.332965	1.422807
17	0.199174	0.042604	-1.516213	5.054714
18	0.123824	-0.006109	-0.234540	-5.541749
19	0.076980	0.003425	0.045650	1.112780
20	0.047858	-0.002113	-0.018556	-0.566041
21	0.029753	0.001267	0.008800	0.303719
22	0.018497	-0.000689	-0.004157	-0.153543
23	0.011499	0.000315	0.001741	0.066704
24	0.007149	-0.000106	-0.000554	-0.021648
25	0.004444	0.000019	0.000097	0.003830




FIG. 2. Radial probability amplitude $u(r) = rR(r)$ for the $1S$ bottomonium ground state using the Gaussian Expansion Method (GEM). The dashed lines represent the individual weighted Gaussian basis functions distributed along a geometric progression of width parameters ν_n . Plotted on a logarithmic scale (x-axis), it demonstrates how the narrowest Gaussians (left) constructively interfere to capture the non-differentiable short-range Coulomb cusp arising from one-gluon exchange, while the broader Gaussians (right) accurately reproduce the exponential tail of the linear confinement potential. The solid black line indicates the total converged wavefunction.