

Print this Page for Your Records

Close Window

Control/Tracking Number: 24-SA-3774-ACSM **Activity:** Scientific Abstract Current Date/Time: 11/28/2023 9:21:22 PM

EMG Coherence Across Shoulder Muscles During Fatigue In Older Adults With Rotator Cuff Disease

Author Block: Xinyu Li, Huiying Zhu, Lisa Griffin. University of Texas at Austin, Austin, TX.

## Abstract:

The risk of degenerative rotator cuff disease (RCD) increases with age, partly due to changes in neuromuscular control, especially during fatiguing contractions. Intermuscular electromyographic (EMG) coherence, reflecting shared neural inputs between motor neuron pools, is essential for maintaining shoulder joint stability. PURPOSE: To determine differences in EMG coherence within the rotator cuff muscles, and between the rotator cuff and deltoid muscles, during sustained isometric fatiguing contractions in older adults with and without RCD. **METHODS:** Thirteen older adults with RCD (68  $\pm$  8 yrs) and 20 without RCD (69  $\pm$  8yrs) participated. Intramuscular EMG was recorded from the supraspinatus (SS), and surface EMG was recorded from the infraspinatus (IS) and middle deltoid (MD). Participants performed an isometric fatiguing contraction at 30° scaption at 25% maximal voluntary contraction until endurance limit. Endurance time was divided into 2 phases: initial (first 30 seconds) and final (last 30 seconds). Z-transformed pooled coherence of each muscle pair (SS-IS, SS-MD, and IS-MD) in the delta (2-5 Hz) and beta (15-35 Hz) bands in each phase were compared with a twoway repeated-measures ANOVA with Bonferroni post-hoc analysis. RESULTS: For SS-IS, there was a significant main effect of group for both the delta (p=0.03) and beta (p<0.001) bands, with the asymptomatic group showing greater SS-IS coherence than the symptomatic group in both phases. Both groups increased SS-IS coherence with fatigue in both bands. For SS-MD, there was also a significant main effect of group for both the delta (p=0.04) and beta (p<0.001) bands, with the asymptomatic group displaying lower SS-MD coherence than the symptomatic group during both phases. Significant main effects of time were also found for both the delta (p=0.006) and beta (p<0.001) bands, with the SS-MD coherence of both groups decreasing with fatigue. CONCLUSIONS: In both two groups, SS-IS coherence increased, but SS-MD coherence decreased with fatigue in both delta and beta bands. Compared to the asymptomatic group, the symptomatic group exhibited lower SS-IS coherence, but greater SS-MD coherence

in both bands during fatique. This could be a potential risk factor that contributes to humeral head instability in RCD.

Author Disclosure Information:

X. Li: None.

Category (Complete): 1106. Clinical Exercise Physiology and Chronic Disease musculoskeletal/neuromuscular diseases; 301. Skeletal Muscle, Bone and Connective Tissue skeletal muscle physiology

> **Keyword (Complete)**: EMG Coherence; Rotator Cuff Disease; Fatigue **Unlabeled/Disclosure (Complete):**

\*If you disclosed a relationship, has the relationship ended? If the financial relationship existed during the last 24 months, but has now ended, please select YES. CME staff will use this information to determine any mitigation steps.: Nothing to disclose \*If you disclosed a relationship, will any of these relationships impact your ability to present an unbiased presentation? : Nothing to disclose

> : No Please select: Yes

Area of Interest (Complete): Area of Interest: Applied Science Check if this abstract is translational science: True

**Presentation Preference (Complete)**: Poster hall presentation **Payment (Complete)**: Your credit card order has been processed on Tuesday 28 November 2023 at 9:00 PM.

> **Status:** Complete OASIS Helpdesk

## American College of Sports Medicine

401 West Michigan Street Indianapolis, IN 46202-3233 (317) 637-9200

Powered by cOASIS, The Online Abstract Submission and Invitation System SM © 1996 - 2023 CTI Meeting Technology All rights reserved. Privacy Policy