

A harmonised point-extraction tool for heterogeneous paleoclimate NetCDF datasets

Essay or report or something for geoinformatics project seminar

Name: Levin Giersch
Student ID: 1690 0515040
E-Mail: l.giersch@campus.tu-berlin.de
Faculty: Faculty VI - Planning Building Environment
Program: Geodesy and Geoinformation Science, M.Sc.
Date: April 28, 2026
Lecturer: Prof. Dr.-Ing. Martin Kada

Contents

Abstract	1
1 Intro	1
2 The Data	1
2.1 Trace21k	1
2.2 CHELSA-TraCE21k	2
2.3 Beyer 2021 (v4)	2
2.4 PalMod 2	3
3 Homogenization	4
3.1 Variables	4
3.2 Time	4
3.3 Space	4
4 Point Extraction	4
5 Results)	4
6 Future	4
7 Conclusion	5

Abstract

1 Intro

- Paleoclimate models are useful for:
 - Reconstructing Earth's climate before direct measurements existed
 - Understanding how the climate responds to forcings like solar output, volcanic eruptions, orbital cycles, and CO₂
 - Testing and validating modern climate models against known past states
 - Establishing the natural variability range of climate, separate from human influence
 - Constraining climate sensitivity (temperature change per CO₂ doubling)
 - Studying planetary climate evolution — Mars, Venus, and other terrestrial bodies
 - Using past warm periods (e.g. Pliocene) as analogues for near-future climate projections
-
- there are many different paleoclimate models.
 - comparing them is important to detect outliers, irregularities or strong differences eg. to help improve the models

2 The Data

- can be downloaded via `data_downloader.ipynb`

2.1 Trace21k

- Simulation of the Transient Climate of the Last 21,000 Years
- Chosen Version: TraCE Main Land Decadal Average
- developed by: see source
- year of development: see source
- how where they made: fully-coupled, non-accelerated atmosphere-ocean-sea ice-land surface CCSM3 simulation
- what was the goal:
- nr. files: 2
- size (gb): 449,2 MB
- type: netcdf
- used variables: TSA, TSOI

- how to get: downloaded via wget python script
- other info:
- extend time: 22,000 BCE to 1990 CE
- resolution time: decadal averages
- extend space: global land
- resolution space: T31_gx3
- source: Otto-Bliesner and Rosenbloom 2021

2.2 CHELSA-TraCE21k

- Chosen Version: CHELSA-TraCE21k-centennial
- developed by: see source
- year of development: see source
- how where they made: generated with the CHELSA-TraCE21k downscaling model.
- what was the goal:
- nr. files:
- size (gb):
- type: geotiff -> need to be converted to netcdf
- used variables: tasmin, tasmax -> $\text{tasmin} + \text{tasmax} / 2$ to get mean
- how to get: downloaded via wget python script
- other info:
- extend time: Start Time 21k BP, End Time 0 BP
- resolution time: monthly
- extend space: global
- resolution space: kilometer scale
- source: Karger et al. 2023, Karger et al. 2020

2.3 Beyer 2021 (v4)

- Chosen Version: 12293345 version 4 (figshare, published 24 June 2021)
- developed by: Robert M. Beyer, Mario Krapp & Andrea Manica (Department of Zoology, University of Cambridge)
- year of development: original paper published 14 July 2020 (received 03 July 2019); dataset v4 published 24 June 2021; addendum paper published 29 September 2021
- how were they made: Delta-method-based downscaling and bias correction, combining medium-resolution HadCM3 transient simulations with high-resolution HadAM3H snap-

shots and present-day observational data

- what was the goal: Fill the gap between high-temporal-resolution but low-spatial-resolution palaeoclimate simulations and high-spatial-resolution snapshot-only datasets
- nr. files: main NetCDF data file plus R, Python and MATLAB example scripts (verify exact count on the figshare page)
- size (GB): 1.24 GB (main NetCDF data file)
- type: NetCDF (data) + R / Python / MATLAB scripts
- used variables:
- how to get: downloaded via wget Python script
- other info: v4 adds global monthly NPP relative to v3; documented in the 2021 addendum
- extend time: 120,000 BP to pre-industrial modern era (0 BP)
- resolution time: 2000-yr steps from 120000–22000 BP; 1000-yr steps from 22000 BP to pre-industrial (72 snapshots total)
- extend space: global terrestrial (full 360° longitude; latitude coverage 720×300 cells at 0.5°, i.e. 150° of latitude)
- resolution space: 0.5° equirectangular grid (~55 km at the equator)
- source: Beyer, Krapp, and Manica 2020, Beyer, Krapp, and Manica 2021a, Beyer, Krapp, and Manica 2021b

2.4 PalMod 2

- Chosen Version:
- developed by:
- year of development:
- how where they made:
- what was the goal:
- nr. files:
- size (gb):
- type:
- used variables:
- how to get:
- other info:
- extend time:
- resolution time:
- extend space:
- resolution space:

- source:

3 Homogenization

3.1 Variables

- which do i choose, are they directly comparable?

3.2 Time

- time extend
- time resolution

3.3 Space

- grid sizes
- changes in north south
- different resolutions for different times

4 Point Extraction

- do i just take the grid cell the point is in?

5 Results)

- For one example Point
- differences between models

6 Future

- Auto iterate over points and compare
- correlate with eg. north south, climate zones, water vs land etc.

7 Conclusion

- hard to compare due to differences eg. resolutions
- comparison can help to find out where they differ the most -> conclusion to how to improve models?

References

- Beyer, Robert M., Mario Krapp, and Andrea Manica (2021a). “Addendum: High-resolution terrestrial climate, bioclimate and vegetation for the last 120,000 years”. In: *Scientific Data* 8.1, p. 262. DOI: 10.1038/s41597-021-01051-1. URL: <https://doi.org/10.1038/s41597-021-01051-1>.
- (2020). “High-resolution terrestrial climate, bioclimate and vegetation for the last 120,000 years”. In: *Scientific Data* 7.236. DOI: 10.1038/s41597-020-0552-1. URL: <https://doi.org/10.1038/s41597-020-0552-1>.
- (2021b). *Late Quaternary climate, bioclimate and vegetation*. Version v4. Dataset. DOI: 10.6084/m9.figshare.12293345.v4. URL: <https://doi.org/10.6084/m9.figshare.12293345.v4>.
- Karger, Dirk Nikolaus et al. (2023). “CHELSA-TraCE21k – high-resolution (1 km) down-scaled transient temperature and precipitation data since the Last Glacial Maximum”. In: *Climate of the Past* 19, pp. 439–456. DOI: 10.5194/cp-19-439-2023. URL: <https://doi.org/10.5194/cp-19-439-2023> (visited on 04/28/2026).
- (2020). *CHELSA-TraCE21k: Downscaled transient temperature and precipitation data since the last glacial maximum*. DOI: 10.16904/envidat.211. URL: <https://www.doi.org/10.16904/envidat.211> (visited on 04/28/2026).
- Otto-Bliesner, Bette L. and Nan A. Rosenbloom (2021). *Simulation of the Transient Climate of the Last 21,000 Years (TraCE-21ka)*. Boulder, CO. DOI: 10.5065/CXB5-TV56. URL: <https://doi.org/10.5065/CXB5-TV56> (visited on 04/28/2026).