

Sex-equalized high-BMI exposure and potentially avoidable MASLD/NAFLD liver burden


GBD 2023 counterfactual analysis across Southeast Asia, East Asia, and Oceania

1 Evidence base

GBD 2023 estimates for MASLD/NAFLD-related cirrhosis and liver cancer burden.

34 locations

Natural Earth



Scenario A assigns the higher-BMI sex the lower-BMI sex exposure distribution within each location-year-age cell. Scenario B is a high-BMI-at-TMREL upper-bound comparator; the scenarios are nested and not additive. Projection rates are internal scenarios, not IHME official forecasts.

2 2023 burden

Potentially avoidable MASLD/NAFLD-related liver burden under Scenario A.

2,797

potentially avoidable DALYs

107 deaths

0.40% of total DALY burden

11.3% of observed high-BMI DALYs

Effect is small but measurable

3 Risk priority

High BMI was not the dominant regional MASLD/NAFLD-related risk signal.

High FPG 27/34

High BMI 3/34

Smoking 4/34

Glycaemic-risk prevention remains central

2050 internal DALY-rate scenario

Baseline 47.1

Scenario A 46.2

Scenario B 39.3

4 Counterfactual

Higher-BMI sex assigned the lower-BMI sex exposure distribution within each location-year-age cell.

