

# Week Six PHY-480

Lewis

September 2025

## Question 1

Outline how you would structure a python code to carry out MD simulations of equilibrium dynamics at fixed temperature of  $N$  Argon atoms in a periodic box of volume  $V$  using the velocity Verlet algorithm.

### Pseudo-code Solution

1. Initialize system parameters:
  - Number of atoms  $N$
  - Box volume  $V$  and box length  $L = V^{1/3}$
  - Time step  $dt$ , total simulation time, temperature  $T$
  - Lennard-Jones parameters ( $\epsilon$ ,  $\sigma$ )
2. Set up initial positions:
  - Arrange atoms in an fcc lattice inside the box
3. Set up initial velocities:
  - Assign velocities from Maxwell-Boltzmann distribution at temperature  $T$
  - Subtract center of mass velocity to ensure total momentum = 0
4. Apply periodic boundary conditions (PBC):
  - Ensure all positions are mapped into the simulation box
  - When calculating distances, use the minimum image convention
5. Compute forces:
  - For each pair of atoms, compute Lennard-Jones force:  
 $F(r) = -dV(r)/dr$  with  $V(r) = 4\epsilon((\sigma/r)^{12} - (\sigma/r)^6)$
  - Store forces on each atom
6. Velocity Verlet integration loop:  
For each timestep:
  - (a) Update positions:  
 $r_i(t+dt) = r_i(t) + v_i(t)dt + 0.5F_i(t)/m * dt^2$
  - (b) Apply periodic boundary conditions to new positions

- (c) Compute new forces  $F_i(t+dt)$
  - (d) Update velocities:
 
$$v_i(t+dt) = v_i(t) + 0.5*(F_i(t) + F_i(t+dt))/m * dt$$
7. Temperature control:
    - Calculate kinetic energy from velocities
    - Rescale velocities by a factor to match target temperature T
  8. Output and analysis:
    - Record energies (kinetic, potential, total)
    - Record temperature and pressure
    - Save configurations for structural analysis
  9. End simulation when final time is reached

## Question 2

Describe how to treat an atom that moves out of the simulation box.

### Answer

When an atom moves out of the primary simulation box we apply periodic boundary conditions which essentially means the atom is “wrapping” back into the box from the opposite side like folding the ends of a piece of paper. For example, if an atom crosses the right side boundary of the box, it will reenters from the left side at the corresponding position similar to a globe.

## Question 3

Describe how to treat shortest distance interactions with atoms in periodically repeated cells surrounding the primary simulation box.

### Answer

To compute interactions, the minimum image convention is used. When there are two atoms, the distance is calculated not only within the primary box but also considering the nearest periodic images of each atom. The effective distance between two particles is the shortest possible distance between them in all repeated cells.