HW 9

Lewis Collum

Updated: April 21, 2020



$$U_{p} = \frac{c}{n}$$

$$L = \frac{\sqrt{x_{0}^{2} + P_{1}^{2}}}{c/n_{1}} + \frac{\sqrt{(P_{0}x_{0}^{2} - X_{0})^{2} + P_{0}x^{2}}}{c/n_{2}}$$


$$\frac{dt}{dx} = \frac{n_{1} x_{0}}{c\sqrt{x_{0}^{2} + P_{1}x^{2}}} + \frac{n_{1} x_{0}}{c\sqrt{(1-x_{0})^{2} + P_{1}x^{2}}} = 0 \quad \text{Aime}$$

$$\Rightarrow \frac{n_{1} x_{0}}{\sqrt{x_{0}^{2} + P_{1}x^{2}}} = \frac{n_{2}(1-x_{0})}{\sqrt{(1-x_{0})^{2} + P_{1}x^{2}}} = 0$$

$$\Rightarrow \frac{n_{1} x_{0}}{\sqrt{x_{0}^{2} + P_{1}x^{2}}} = \frac{n_{2}(1-x_{0})}{\sqrt{(1-x_{0})^{2} + P_{1}x^{2}}} = 0$$

n, sin O, = nasinoa

2



Nair Sin Gair = Nwoter Sin Gwoter

SinOnir
SinOmir =
$$\sqrt{\frac{1.8}{4062}}$$

SinOmir = $\sqrt{\frac{1.8}{4062}}$
SinOmir = $\sqrt{\frac{1.8}{4062}}$



```
from numpy import *
nAir = 1
nGan = 3.7
criticalAngle = arcsin(nAir/nGan)
print(f'\[\\theta_c ='
      f'{rad2deg(criticalAngle):.3f}^{{\circ}}\]')
solidAngleCone = 2*pi*(1-cos(criticalAngle)) #4*pi
solidAngleHemisphere = 2*pi
coneCount = 2
solidAngleRatio = coneCount * solidAngleCone/solidAngleHemisphere
print(f'\[\\text{{solid Angle Ratio}} = {solidAngleRatio:.3f}\]')
import pint
unit = pint.UnitRegistry()
wavelength = 560
h = 6.63E-34
c = 3E8
E = solidAngleRatio * h*c/wavelength
print(f'\\[\text{{fraction of optical energy}} = \\\text{{E:.2E} J}})]')
```

$$\theta_c=15.680^\circ$$
 solid Angle Ratio $=0.074$ fraction of optical energy $=$ 2.64E-29 J

4


```
from matplotlib import pyplot
import numpy
x = numpy.linspace(2, 6, 20) #wavelenth
#impedance for quartz
eta2 = 1/numpy.sqrt(1 +
               0.6961663*x**2 / (x**2 - 0.0684043**2) +
               0.4079426*x**2 / (x**2 - 0.1162414**2) +
               0.8974794*x**2 / (x**2 - 9.896161**2))
```


```
#impedance for air
eta1 = 1

reflectionCoefficient = (eta2 - eta1)/(eta1 + eta2)
transmissionCoefficient = 1 + reflectionCoefficient

reflectivity = reflectionCoefficient**2
transmissivity = transmissionCoefficient**2 * (eta1/eta2)

pyplot.plot(x, reflectivity)
pyplot.savefig('felectivity')
pyplot.savefig('figure/4-re.png')
pyplot.plot(x, transmissivity)
pyplot.savefig('figure/4-tr.png')
pyplot.savefig('figure/4-tr.png')
pyplot.show()
```









```
from matplotlib import pyplot
import numpy
from numpy import cos, arcsin, sin
```

```
def ganIndex(wavelength):
   return numpy.sqrt(
       0.6961663*wavelength**2 / (wavelength**2 - 0.0684043**2)
        + 0.4079426*wavelength**2 / (wavelength**2 - 0.1162414**2)
        + 0.8974794*wavelength**2 / (wavelength**2 - 9.896161**2)
        + 1).reshape(-1, 1)
def degreeArangeToRadians(start, stop, spacing):
    degreeRange = numpy.arange(start, stop+spacing, spacing)
    radianRange = numpy.deg2rad(degreeRange)
    radianRangeAsColumnVector = radianRange.reshape(-1, 1).T
    return radianRangeAsColumnVector
def snellsTransmissionAngle(eta1, eta2, a1):
   return arcsin(sin(a1)*(eta2/eta1))
def perpendicularReflection(eta1, eta2, a1, a2):
    return ((eta2*cos(a1) - eta1*cos(a2))
            / (eta2*cos(a1) + eta1*cos(a2)))**2
def parallelReflection(eta1, eta2, a1, a2):
    return ((eta2*cos(a2) - eta1*cos(a1))
            / (eta2*cos(a2) + eta1*cos(a1)))**2
wavelength = numpy.linspace(2, 6, 20)
n2 = ganIndex(wavelength)
eta1 = 1 \#Air
eta2 = 1/n2 #by the relationship eta2/eta1 = n1/n2
incidenceAngles = degreeArangeToRadians(
    start = 0,
    stop = 60,
    spacing = 10)
transmissionAngles = snellsTransmissionAngle(
    eta1 = eta1,
    eta2 = eta2,
    a1 = incidenceAngles)
transverseElectricReflectivity = perpendicularReflection(
   eta1 = eta1,
    eta2 = eta2,
    a1 = incidenceAngles,
    a2 = transmissionAngles)
transverseMagneticReflectivity = parallelReflection(
    eta1 = eta1,
    eta2 = eta2,
    a1 = incidenceAngles.
    a2 = transmissionAngles)
pyplot.plot(wavelength, transverseElectricReflectivity)
pyplot.title('TE Polarization Reflectivity at Different Incidence Angles')
pyplot.xlabel('wavelength')
pyplot.ylabel('TE reflectivity')
pyplot.legend(*numpy.round(numpy.rad2deg(incidenceAngles)))
pyplot.savefig('figure/5-te.png')
pyplot.show()
pyplot.plot(wavelength, transverseMagneticReflectivity)
pyplot.title('TM Polarization Reflectivity at Different Incidence Angles')
pyplot.xlabel('wavelength')
pyplot.ylabel('TM reflectivity')
pyplot.legend(*numpy.round(numpy.rad2deg(incidenceAngles)))
pyplot.savefig('figure/5-tm.png')
pyplot.show()
```







6 - Brewster's Angle

```
from numpy import arctan, sqrt, rad2deg

wavelength = 580E-9
schottIndex = 1.5427
airIndex = 1
brewsterAngle = rad2deg(arctan(sqrt(schottIndex/airIndex)))

print(f'\[\\theta_B'
    f' = \\tan^{{-1}}\sqrt{\\dfrac{\\epsilon_2}}{\\epsilon_1}}'
    f' = {\brewsterAngle:.2f}^{\\chickless'}\]')
```

$$\theta_B = \tan^{-1} \sqrt{\frac{\epsilon_2}{\epsilon_1}} = 51.16^{\circ}$$