

Monge–Ampère PDEs And Their Geometry: A Strange Relationship

Lewis Napper (University of Surrey, UK)

La Rochelle University, France
26th September 2024



AMBASSADE
DE FRANCE
AU ROYAUME-UNI

*Liberté
Égalité
Fraternité*

1. Monge–Ampère Equations
2. Differential Geometry Glossary
3. From Forms To Equations (And Back Again)
4. Some Useful (2D) Tools
5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved
Backgrounds and
High-Dimensional
Systems



My Work:

- Final year Ph.D. researcher at the University of Surrey, UK.
- Visiting La Rochelle as part of a French Embassy SSHN fellowship.
- My Website – <https://lewisn3142.github.io>

My Current Projects:

- Monge–Ampère Geometry and Fluid Dynamics.
- Lorentzian Length Spaces and Curvature.
- Cellular Automata on Aperiodic Tilings of Surfaces.

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



Outline of Talk

La Rochelle Seminar

Lewis Napper

1. Monge–Ampère Equations
2. Differential Geometry Glossary
3. From Forms To Equations (And Back Again)
4. Some Useful (2D) Tools
5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



Select References

La Rochelle Seminar

Lewis Napper

- Napper, Roulstone, Rubtsov, and Wolf, *Monge–Ampère Geometry and Vortices*, Nonlinearity 37, (2024).
- Kushner, Lychagin, and Rubtsov, *Contact Geometry and Non-Linear Differential Equations*, (2007).
- Lychagin, Rubtsov, and Chekalov, *A Classification of Monge–Ampère Equations*, Ann. Sci. Ec. Norm. Sup. 26 (1993).
- Bryant, *On the Geometry of Almost Complex 6-Manifolds*, Asian J. Math. 10, (2006).
- Cantrijn, Ibort, and De León, *On the Geometry of Multisymplectic Manifolds*, J. Austr. Math. Soc. 66, (1999).

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



1. Monge–Ampère Equations

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



What is a Monge–Ampère Equation?

La Rochelle Seminar

Lewis Napper

- $x^1, x^2 \dots x^n$ and ψ are the independent and dependent variables.
- Non-linear, second-order PDEs, given by quasi-linear combinations of the determinant of the Hessian of ψ and its minors.

$$\text{Hess}(\psi) = \begin{pmatrix} \psi_{x^1 x^1} & \psi_{x^1 x^2} & \cdots & \psi_{x^1 x^n} \\ \psi_{x^2 x^1} & \psi_{x^2 x^2} & \cdots & \psi_{x^2 x^n} \\ \vdots & \vdots & \cdots & \vdots \\ \psi_{x^n x^1} & \psi_{x^n, x^2} & \cdots & \psi_{x^n x^n} \end{pmatrix}$$

- Minors of a matrix M are $k \times k$ sub-matrices with entries given by the intersections of k rows and k columns of M .

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



General Form in Two Dimensions

La Rochelle Seminar

Lewis Napper

- Set $x^1 = x$ and $x^2 = y$.
- In two dimensions, MAEs take the form

$$A\psi_{xx} + 2B\psi_{xy} + C\psi_{yy} + D(\psi_{xx}\psi_{yy} - \psi_{xy}^2) + E = 0.$$

where $A, B, \dots E$ can depend on $x, y, \psi, \psi_x, \psi_y$ non-linearly.

- If $A, B, \dots E$ do not depend on ψ , we have a symplectic Monge–Ampère equation, e.g. 2D Poisson.

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



Some Examples You May Know

La Rochelle Seminar

Lewis Napper

- 2D Reaction-Diffusion: $\psi^\alpha \psi_{xx} + [\alpha \psi^{\alpha-1} \psi_x - \psi_t + F(\psi)] = 0.$
- 3D Chynoweth–Sewell: $[\psi_{xx} \psi_{yy} - (\psi_{xy})^2] + \psi_{zz} = 0.$
- 4D Khokhlov–Zabolotskaya: $\psi_{tt} + \psi_{yy} + \psi_{zz} - \psi_{xt} + (\psi_t)^2 = 0.$
- Laplace: $\Delta \psi := \psi_{x^1 x^1} + \psi_{x^2 x^2} + \cdots + \psi_{x^n x^n} = 0.$
- Wave: $\square \psi := \psi_{tt} - \psi_{x^1 x^1} - \psi_{x^2 x^2} - \cdots - \psi_{x^n x^n} = 0 .$
- 2D Poisson: $\psi_{xx} \psi_{yy} - (\psi_{xy})^2 = F(x, y).$

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



1. Monge–Ampère Equations
 2. Differential Geometry Glossary
 3. From Forms To Equations (And Back Again)
 4. Some Useful (2D) Tools
 5. An Example: The Poisson Equation For Incompressible Fluids
- Part 2: Curved Backgrounds and High-Dimensional Systems

2. Differential Geometry Glossary



Terminology: Configuration and Phase Space

La Rochelle Seminar

Lewis Napper

- Configuration Space/Background – \mathbb{R}^n with coordinates x^1, x^2, \dots, x^n representing independent variables and points $x \in \mathbb{R}^n$ representing positions in space.
- A particle at point $x \in \mathbb{R}^n$ has an m -dimensional space of possible momenta (one in each direction x^i) with coordinates q_1, q_2, \dots, q_m . These will be related to the dependent variable.
- Phase space T^*M – $2m$ -dimensional space (manifold), with coordinates $x^1 \dots x^m, q_1 \dots q_m$, representing all possible combinations of positions and momenta.

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



Terminology: Tangent Vectors and Forms

La Rochelle Seminar

Lewis Napper

Let M be a space (manifold) with coordinates y^1, y^2, \dots, y^n .

- At each $p \in M$, there is a vector space of Tangent Vectors to M . Space has a basis $\partial_{y^1}, \dots, \partial_{y^n}$.
- At each $p \in M$ there is a dual vector space of 1-Forms on M . Space has a basis dy^1, \dots, dy^n , satisfying $dy^i(\partial_{y^j}) = \delta_j^i$.
- Coefficients are in $\mathcal{C}^\infty(M)$, so a 1-form on $T^*\mathbb{R}^2$ looks like

$$\alpha = \alpha_i(x^1, x^2, q_1, q_2) dx^i + \tilde{\alpha}^j(x^1, x^2, q_1, q_2) dq_j$$

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



Terminology: k -Forms and Operators

La Rochelle Seminar

Lewis Napper

- Wedge Product \wedge – A skew-symmetric bilinear operator on forms. For 1-forms α, β , we have

$$\alpha \wedge \beta = -\beta \wedge \alpha,$$

$$(\alpha + \tilde{\alpha}) \wedge \beta = \alpha \wedge \beta + \tilde{\alpha} \wedge \beta.$$

- k -Forms – totally skew-symmetric, k -linear operator on tangent vectors $\gamma = \gamma_{i_1, \dots, i_k} dy^{i_1} \wedge dy^{i_2} \wedge \dots \wedge dy^{i_k}$.

- The wedge product of a k -form α with an ℓ -form β is a $(k + \ell)$ -form satisfying $\alpha \wedge \beta = (-1)^{k\ell} \beta \wedge \alpha$:

$$\alpha \wedge \beta = (\alpha_i dy^i) \wedge (\beta_j dy^j) = (\alpha_i \beta_j) dy^i \wedge dy^j.$$

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



- Exterior Derivative – Operator d taking k -forms to $(k + 1)$ -forms.
- Exterior derivative of a function (think total derivative):
 $d(f) = (\partial_{y^i} f)dy^i = (f_{y^i})dy^i.$
- Exterior derivative of a 1-form:

$$\begin{aligned}d(\alpha_i dy^i) &= (d\alpha_i) \wedge dy^i + \alpha_i d(dy^i) \\ &= (\partial_{y^j} \alpha_i) dy^j \wedge dy^i.\end{aligned}$$

- Note: $d^2 = 0$

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



1. Monge–Ampère
Equations

2. Differential Geometry
Glossary

3. From Forms To
Equations (And Back
Again)

4. Some Useful (2D)
Tools

5. An Example: The
Poisson Equation For
Incompressible Fluids

Part 2: Curved
Backgrounds and
High-Dimensional
Systems

3. From Forms To Equations (And Back Again)



Example: Equation from Form

Consider a 2-form on $T^*\mathbb{R}^2$, where x^1, x^2 are our independent variables,

$$\alpha = dq_1 \wedge dx^2 - dq_2 \wedge dx^1.$$

Define $L := \{(x^1, x^2, \psi_{x^1}, \psi_{x^2})\} \subset T^*\mathbb{R}^2$ (fix q_1 and q_2 at each x).

$$\begin{aligned}\alpha|_L &= d(\psi_{x^1}) \wedge dx^2 - d(\psi_{x^2}) \wedge dx^1 \\ &= (\psi_{x^1 x^1} + \psi_{x^2 x^2}) dx^1 \wedge dx^2\end{aligned}$$

So $\alpha|_L = 0$ if and only if $\Delta\psi = 0$, that is, ψ solves $\Delta\psi = 0$ when L solves $\alpha|_L = 0$.

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



Symplectic Forms and Equivalent Equations

La Rochelle Seminar

Lewis Napper

A symplectic form ω on $T^*\mathbb{R}^n$ is

- a 2-form: skew-symmetric and bilinear,
- closed: $d\omega \equiv 0$,
- non-degenerate: $\omega(X, \cdot) \equiv 0$ if and only if $X \equiv 0$.

The canonical choice is

$$\omega = dq_i \wedge dx^i = \begin{pmatrix} 0_n & -I_n \\ I_n & 0_n \end{pmatrix}$$

Then $\omega|_L = 0$ is trivial, so $\alpha|_L = 0$ and $(\alpha + \omega)|_L = 0$ are the same equation! Which one do we pick?

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



Effective Forms in Two Dimensions

La Rochelle Seminar

Lewis Napper

- An n -form α on $T^*\mathbb{R}^n$ is called ω -effective if it is skew-orthogonal to ω , i.e. $\alpha \wedge \omega = 0$.
- (Hodge–Lepage–Lychagin) For any 2-form β and symplectic form ω ,

$$\beta = \alpha + F\omega,$$

for unique ω -effective 2-form α and $F \in \mathcal{C}^\infty(T^*\mathbb{R}^2)$. ($n = 2$)

- If $\omega|_L = 0$, all β corresponding to a given α correspond to the same equation. So we only need to consider the ω -effective forms. More formally...

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



- A Monge–Ampère Structure on $T^*\mathbb{R}^n$ is a pair (ω, α) , where ω is a symplectic form and α is an ω -effective n -form [Banos 2002].
- Fixing canonical ω , then $\omega|_L = 0$ trivially and $\alpha|_L = 0$ is a Monge–Ampère equation.
- Classical Solutions $\psi \in \mathcal{C}^\infty(\mathbb{R}^n)$ of the Monge–Ampère equations correspond to choices of $L = \{(x^1, \dots, x^n, \psi_{x^1}, \dots, \psi_{x^n})\}$.
- Generalised Solutions are n -dimensional spaces (submanifolds) $L \subset T^*\mathbb{R}^n$ such that $\omega|_L = 0$ and $\alpha|_L = 0$.

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems





Classical and Generalised Solutions

Classical $L = \{(x^1, \dots, x^n, \psi_{x^1}, \dots, \psi_{x^n})\}$:

- $\pi : L \rightarrow \mathbb{R}^n$ is bijective (diffeomorphic).

Generalised L :

- When π is not surjective, ψ is not defined on the whole domain.
- When π is not injective, ψ is a multivalued solution. [Vinogradov 1973]
- When π is not immersive, we have Arnold's singularities.



1. Monge–Ampère Equations
2. Differential Geometry Glossary
3. From Forms To Equations (And Back Again)
4. Some Useful (2D) Tools
5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



Monge–Ampère Forms in Two Dimensions

La Rochelle Seminar

Lewis Napper

For canonical ω on $T^*\mathbb{R}^2$, the ω -effective 2-forms are

$$\begin{aligned}\alpha = & A \, dq_1 \wedge dx^2 + B \, (dx^1 \wedge dq_1 + dq_2 \wedge dx^2) \\ & + C \, dx^1 \wedge dq_2 + D \, dq_1 \wedge dq_2 + E \, dx^1 \wedge dx^2\end{aligned}$$

For classical solutions $L = \{(x, y, \psi_x, \psi_y)\}$, the constraint $\alpha|_{d\psi} = 0$ gives the Monge–Ampère equation

$$A\psi_{xx} + 2B\psi_{xy} + C\psi_{yy} + D(\psi_{xx}\psi_{yy} - \psi_{xy}^2) + E = 0.$$

This is a bijection between Monge–Ampère equations and ω -effective forms.

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



1. Monge–Ampère Equations
2. Differential Geometry Glossary
3. From Forms To Equations (And Back Again)
4. Some Useful (2D) Tools
5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved
Backgrounds and
High-Dimensional
Systems

4. Some Useful (2D) Tools



The Pfaffian in Two Dimensions

La Rochelle Seminar

Lewis Napper

- The Pfaffian is defined by $\alpha \wedge \alpha =: f_\alpha \omega \wedge \omega$ where $f_\alpha = AC - B^2 - DE$.
- Note that f_α on L is the determinant of the linearisation matrix of the equation $\alpha|_L = 0$ (Rellich invariant).
- Hence, the Monge–Ampère equation $\alpha|_L = 0$ is
 - elliptic* $\Leftrightarrow f_\alpha > 0$.
 - hyperbolic* $\Leftrightarrow f_\alpha < 0$.
 - parabolic* $\Leftrightarrow f_\alpha = 0$.

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



Almost (Para-)Complex Structure

La Rochelle Seminar

Lewis Napper

- Set $\tilde{\alpha} = \frac{1}{\sqrt{|f_\alpha|}}\alpha$, so that $f_{\tilde{\alpha}} = \text{sign}(f_\alpha)$.

Multiples of α have the same $\tilde{\alpha}$ (removes choice).

- Define endomorphism of vector fields $J : \mathfrak{X}(T^*\mathbb{R}^2) \rightarrow \mathfrak{X}(T^*\mathbb{R}^2)$ by

$$\tilde{\alpha}(\cdot, \cdot) =: \omega(J\cdot, \cdot) \quad (J = \omega^{-1}\tilde{\alpha} \text{ as matrices}) ,$$

- $f_\alpha \leq 0 \Leftrightarrow J^2 = \pm I_4$. [Lychagin et al. 1993]

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



The Lychagin–Rubtsov Theorem and Equivalence

La Rochelle Seminar

Lewis Napper

Two Monge–Ampère equations are locally equivalent if there exists a bijection (diffeomorphism) $F : (T^*\overline{\mathbb{R}^2}, \omega, \alpha_1) \rightarrow (T^*\overline{\mathbb{R}^2}, \omega, \alpha_2)$

$$\omega(F_*\cdot, F_*\cdot) = \omega(\cdot, \cdot) \text{ and } \alpha_2(F_*\cdot, F_*\cdot) = \alpha_1(\cdot, \cdot)$$

[Lychagin–Rubtsov] The following conditions are equivalent:

- $\alpha|_L = 0$ is locally equivalent to $\square\psi = 0$ or $\Delta\psi = 0$.
- $d(\tilde{\alpha}) = 0$ ($f_\alpha \leq 0$).
- J is integrable ($J^2 = \pm I_2$).

These criteria do not always hold.

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



The Lychagin–Rubtsov Metric

La Rochelle Seminar

Lewis Napper

- Picking a non-degenerate, ω -effective, and α -effective 2-form K , we can define a symmetric, bilinear form

$$\hat{g}(\cdot, \cdot) := -K(J\cdot, \cdot)$$

called a Lychagin–Rubtsov metric. [Roulstone et al. 2001]

- There exists a choice of K s.t. the metric in (x^i, q_i) coordinates is

$$\hat{g} = \begin{pmatrix} f_\alpha I_2 & 0 \\ 0 & I_2 \end{pmatrix}$$

with signature dictated by the sign of f_α .

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



1. Monge–Ampère Equations
 2. Differential Geometry Glossary
 3. From Forms To Equations (And Back Again)
 4. Some Useful (2D) Tools
 5. An Example: The Poisson Equation For Incompressible Fluids
- Part 2: Curved Backgrounds and High-Dimensional Systems

5. An Example: The Poisson Equation For Incompressible Fluids



Pressure, Vorticity, and Strain

La Rochelle Seminar

Lewis Napper

➤ Homogeneous, Incompressible Navier–Stokes on \mathbb{R}^m

$$\partial_t v^j = -v^i \nabla_i v^j - \nabla_j p + \nu \Delta v^j \quad (-c_j).$$

$$\nabla_i v^i = 0$$

Here $\nabla_i := \partial_{x^i}$, time t is a parameter, ν is viscosity and v^i is the x^i -component of velocity.

1. Monge–Ampère Equations
 2. Differential Geometry Glossary
 3. From Forms To Equations (And Back Again)
 4. Some Useful (2D) Tools
 5. An Example: The Poisson Equation For Incompressible Fluids
- Part 2: Curved Backgrounds and High-Dimensional Systems



Pressure, Vorticity, and Strain

La Rochelle Seminar

Lewis Napper

- Homogeneous, Incompressible Navier–Stokes on \mathbb{R}^m

$$\partial_t v^j = -v^i \nabla_i v^j - \nabla_j p + \nu \Delta v^j \quad (-c_j).$$

- Taking the divergence and applying $\nabla_i v^i = 0$ one finds

$$\zeta_{ij} \zeta^{ij} - S_{ij} S^{ij} = \Delta p \quad (+\nabla_i c^i).$$

where $\zeta_{ij} = \frac{1}{2}(\nabla_j v_i - \nabla_i v_j)$ is the vorticity form
and $S_{ij} = \frac{1}{2}(\nabla_j v_i + \nabla_i v_j)$ is the strain-rate tensor.

1. Monge–Ampère
Equations

2. Differential Geometry
Glossary

3. From Forms To
Equations (And Back
Again)

4. Some Useful (2D)
Tools

5. An Example: The
Poisson Equation For
Incompressible Fluids


Part 2: Curved
Backgrounds and
High-Dimensional
Systems



Vorticity, Divergence, and Strain

La Rochelle Seminar

Lewis Napper



Based on Figure from Clough et al. 2014

$$(\zeta_{ij})_{2D} = \frac{1}{2} \begin{pmatrix} 0 & \zeta \\ -\zeta & 0 \end{pmatrix} \quad (\zeta_{ij})_{3D} = \frac{1}{2} \begin{pmatrix} 0 & \zeta_3 & -\zeta_2 \\ -\zeta_3 & 0 & \zeta_1 \\ \zeta_2 & -\zeta_1 & 0 \end{pmatrix}$$

1. Monge–Ampère Equations
2. Differential Geometry Glossary
3. From Forms To Equations (And Back Again)
4. Some Useful (2D) Tools
5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



Pressure Equation in Two Dimensions

La Rochelle Seminar

Lewis Napper

- In 2D, there exists a stream function $\psi \in \mathcal{C}^\infty(\mathbb{R}^2)$ such that $v^1 = -\psi_y$ and $v^2 = \psi_x$.
- The constraint $\nabla_i v^i = 0$ is trivially satisfied and the pressure equation is a Monge–Ampère equation for ψ

$$\Delta p = 2 \left(\psi_{xx} \psi_{yy} - (\psi_{xy})^2 \right) .$$

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



Pressure Equation in Two Dimensions

La Rochelle Seminar

Lewis Napper

- In 2D, there exists a stream function $\psi \in \mathcal{C}^\infty(\mathbb{R}^2)$ such that $v^1 = -\psi_y$ and $v^2 = \psi_x$.
- The constraint $\nabla_i v^i = 0$ is trivially satisfied and the pressure equation is a Monge–Ampère equation for ψ

$$\zeta_{ij}\zeta^{ij} - S_{ij}S^{ij} = \Delta p = 2(\psi_{xx}\psi_{yy} - (\psi_{xy})^2) .$$

- *Vorticity dominates* $\Leftrightarrow \Delta p > 0 \Leftrightarrow$ *Elliptic equation.*
Strain dominates $\Leftrightarrow \Delta p < 0 \Leftrightarrow$ *Hyperbolic equation.*
No dominance $\Leftrightarrow \Delta p = 0 \Leftrightarrow$ *Parabolic equation.*
[Weiss 1991, Larchevêque 1993]

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



What Can the Geometry Tell Us?

La Rochelle Seminar

Lewis Napper

- Which differential form α corresponds to the equation?
- Does the geometry see vorticity/strain dominating regions?
- Can we apply the LR theorem to get (local) solutions of the Poisson equation from the Laplace or Wave equation?
- Does the Lychagin–Rubtsov metric pick up on any interesting properties of solutions?
- What does this tell us about solutions to Euler/Navier–Stokes?

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



Question 1: What Differential Form?

La Rochelle Seminar

Lewis Napper

One can recover the pressure equation

$$\frac{\Delta p}{2} = (\psi_{xx}\psi_{yy} - \psi_{xy}^2)$$

by choosing the Monge–Ampère form [Roulstone et al. 2009]

$$\alpha = dq_1 \wedge dq_2 - f_\alpha dx^1 \wedge dx^2,$$

1. Monge–Ampère
Equations

2. Differential Geometry
Glossary

3. From Forms To
Equations (And Back
Again)

4. Some Useful (2D)
Tools

5. An Example: The
Poisson Equation For
Incompressible Fluids

Part 2: Curved
Backgrounds and
High-Dimensional
Systems



Question 2: Does Geometry See Existing Results?

La Rochelle Seminar

Lewis Napper

- The Pfaffian of α is given by

$$f_\alpha = \frac{\Delta p(x, y)}{2}.$$

Recall that the sign of f_α tells us if a Monge–Ampère equation is elliptic/hyperbolic/parabolic.

- Hence, we have a geometric justification for the Poisson equation being:

$$\begin{aligned}\text{elliptic} &\Leftrightarrow \Delta p > 0, \\ \text{hyperbolic} &\Leftrightarrow \Delta p < 0, \\ \text{parabolic} &\Leftrightarrow \Delta p = 0.\end{aligned}$$

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



Question 3: Do We Have Equivalence?

- For $\tilde{\alpha} = \frac{1}{\sqrt{|f_\alpha|}}\alpha$, we find $d\tilde{\alpha} = 0$ if and only if Δp is constant.
- Hence, by the Lychagin–Rubtsov Theorem,

$$\frac{\Delta p}{2} = (\psi_{xx}\psi_{yy} - \psi_{xy}^2)$$

is locally equivalent to $\Delta\psi = 0$ or $\square\psi = 0$ if and only if Δp is constant.

- So this equivalence cannot be applied to some physical problems.

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



Question 4: What Does the Metric on $T^*\mathbb{R}^2$ Say?

La Rochelle Seminar

Lewis Napper

The Lychagin–Rubtsov metric on $T^*\mathbb{R}^2$ given by

$$\hat{g} = \begin{pmatrix} \frac{\Delta p}{2} I & 0 \\ 0 & I \end{pmatrix}$$

is

Riemannian $\Leftrightarrow \Delta p > 0$.

Kleinian $\Leftrightarrow \Delta p < 0$.

Degenerate $\Leftrightarrow \Delta p = 0$.

These degeneracies are where the scalar curvature of \hat{g} blows up (curvature singularities) — they persist under coordinate changes.

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



Question 4: What Does the Metric on L Say?

- If we restrict \hat{g} to the classical solution $L = \{(x, y, \psi_x, \psi_y)\}$, we get

$$\hat{g}|_L = \zeta \begin{pmatrix} \psi_{xx} & \psi_{xy} \\ \psi_{xy} & \psi_{yy} \end{pmatrix}$$

where $\zeta = \Delta\psi$ is the vorticity.

- Degenerate when $\zeta = 0$ or $\Delta p = 0$.
Riemannian when $\Delta p > 0$.
Kleinian when $\Delta p < 0$.
- Degeneracy when $\zeta = 0$ not always curvature singularity.

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



Summary of Relationship

La Rochelle Seminar

Lewis Napper

Δp	> 0	< 0	$= 0$
Dominance	Vorticity	Strain	None
$\alpha _L = 0$	Elliptic	Hyperbolic	Parabolic
f_α	> 0	< 0	$= 0$
J^2	$-I_2$	I_2	Singular
\hat{g}	Riemannian (4, 0)	Kleinian (2, 2)	Degenerate
$\hat{g} _L$	Riemannian (2, 0)	Kleinian (1, 1)*	Degenerate

*Except when $\zeta = 0$, in which case it is degenerate.

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids









Part 2: Curved Backgrounds and High-Dimensional Systems



2D ABC Flow: $\psi(x, y) = \frac{3}{2} \cos(y) + \sin(x) = -\zeta$

La Rochelle Seminar

Lewis Napper



1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)

4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



Summary and Outlook

La Rochelle Seminar

Lewis Napper

This time we:

- Introduced MA equations and their geometry as a tool for studying solutions and discussed the LR theorem and metric.
- Applied this to the Poisson equation in 2D to recover existing results and pick up interesting features of solutions.

Next time we will:

- show how our fluid dynamical results can be extended to curved backgrounds, such as the sphere, and what they tell us about the topology of vortices.
- discuss what happens when we take a non-canonical symplectic form and how to consider the Poisson equation in three dimensions, when it is not Monge–Ampère.

1. Monge–Ampère Equations

2. Differential Geometry Glossary

3. From Forms To Equations (And Back Again)


4. Some Useful (2D) Tools

5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



Thank you!



Any questions?

(Image Credit [Kushner, Lychagin, Rubtsov. 2007])

1. Monge–Ampère Equations
2. Differential Geometry Glossary
3. From Forms To Equations (And Back Again)
4. Some Useful (2D) Tools
5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems



Part 2: Curved Backgrounds and High-Dimensional Systems

1. Monge–Ampère Equations
2. Differential Geometry Glossary
3. From Forms To Equations (And Back Again)
4. Some Useful (2D) Tools
5. An Example: The Poisson Equation For Incompressible Fluids

Part 2: Curved Backgrounds and High-Dimensional Systems

