BFGS can converge to non-Clarke-stationary points with positive probability

Lexiao Lai*

Abstract

We construct a coercive, locally Lipschitz, semialgebraic function on $\mathbb{R}^2$ for which BFGS with Armijo–Wolfe line search converges to a non-Clarke-stationary point from a nonempty open set of initial points and inverse-Hessian estimates. This disproves the almost-sure stationarity assertion in the 2013 challenge of Lewis and Overton [LO13, Challenge 7.1(2)]. The construction and proof combine analytical arguments with computer-assisted verification of explicit inequalities.[1]

Keywords: Nonsmooth optimization, BFGS, Clarke stationarity, semialgebraic functions, computer-assisted proof.

1 Introduction and main result

We consider the unconstrained minimization problem $\min_{x\in\mathbb{R}^n} f(x)$, where $f : \mathbb{R}^n \to \mathbb{R}$ is locally Lipschitz and may be nonsmooth and nonconvex. The Broyden–Fletcher–Goldfarb–Shanno (BFGS) method approximates curvature using gradient differences without evaluating Hessians [Bro70, Fle70, Gol70, Sha70]. Its numerical effectiveness and convergence properties in smooth optimization have been studied extensively [DM77]. For twice continuously differentiable convex objectives with bounded initial sublevel sets, BFGS function values converge to the minimum under an Armijo–Wolfe line search [Pow76]. Explicit global convergence rates are available for twice differentiable, strongly convex objectives with Lipschitz gradients when the Armijo parameter is below $1/2$ [JJM24, Theorem 4.1]. For convex objectives with Lipschitz gradients, rates have also been established for the smallest gradient norm reached when the objective is bounded below, and for the function-value gap when the initial sublevel set is bounded [DYZ26, Theorem 3.1].

BFGS has also shown empirical success on nonsmooth objectives [LO13, CMO17]. This empirical behavior is supported by convergence results for the Euclidean norm in two dimensions with exact line search, and for $|x|$ with a particular inexact line search using a zero Armijo parameter [LO13, Theorems 3.2 and 5.2]. A doubling/bisection Armijo–Wolfe line search also terminates finitely for bounded-below, locally Lipschitz, semialgebraic objectives along directions with negative directional derivative [LO13, Theorem 4.7]. These results and the numerical evidence motivate the broader question posed in [LO13, Challenge 7.1(2)]: for locally Lipschitz, semialgebraic functions with bounded sublevel sets, is every cluster point generated by BFGS with this line search Clarke stationary almost surely under Gaussian initialization of the point and Wishart initialization of the inverse-Hessian estimate?

*`lai.lexiao@hku.hk`, Department of Mathematics, The University of Hong Kong, Hong Kong.

[1]Verification code and Lean formalization: `https://github.com/LexiaoLai/Nonsmooth-BFGS-Nonstationary`.

Further results extend the analysis to broader nonsmooth classes. For convex objectives satisfying additional structural assumptions, BFGS function values converge to the minimum [GL18]; in particular, convergence holds for the Euclidean norm in every dimension [GL18, Corollary 4.2]. For a class of unbounded-below convex functions with two affine pieces, BFGS detects a direction of unbounded descent after finitely many iterations, provided that the iterates remain differentiability points [XW17, Theorem IV.1 and Remark IV.1]. For a class of piecewise-differentiable objectives, convergent sequences initialized sufficiently near a minimizer have Clarke-stationary limits under additional structural and asymptotic assumptions on the objective, iterates, and quasi-Newton matrices [Geb26, Theorem 3.1]. These results do not settle the general nonsmooth convex case; convergence for $x_1^2 + |x_2|$ is identified as unresolved in [Geb26, Section 1].

Existing counterexamples show that BFGS and related methods can fail to converge or converge to nonminimizers. For smooth nonconvex objectives, examples exhibit cycling with Wolfe line searches [Dai02] or failure with exact line searches, including one with bounded sublevel sets [Mas04, Mas14]. A polynomial example with Armijo–Wolfe line search has bounded iterates with nonstationary cluster points [Dai13, Sections 3.1–3.4]; it can easily be modified to have bounded sublevel sets without changing the iterates. For nonsmooth objectives, BFGS with an Armijo–Wolfe line search can converge to a Clarke-stationary point admitting linear descent [LZ15, Proposition 5.3]. Scaled memoryless BFGS can converge to non-Clarke-stationary points on an unbounded-below convex piecewise-affine objective [AO21, Corollary 3.12]. These examples do not establish failure of Clarke stationarity on an open set of initializations for BFGS with the specified Armijo–Wolfe line search on locally Lipschitz, semialgebraic objectives with bounded sublevel sets.

In this paper, we disprove the almost-sure stationarity assertion in [LO13, Challenge 7.1(2)]. We construct a coercive, locally Lipschitz, semialgebraic function on $\mathbb{R}^2$ for which, with the Armijo–Wolfe line search, the entire BFGS sequence converges to the same non-Clarke-stationary point from every initialization in a fixed nonempty open set of points and positive definite inverse-Hessian estimates. Consequently, failure has positive probability under the Gaussian–Wishart initialization specified in the challenge.

1.1 Algorithm and main result

We consider BFGS with the unit-trial doubling/bisection line search and strict Armijo–Wolfe tests specified in [LO13, Eqs. (4.2)–(4.3), Algorithm 4.6]; See Algorithm 1 below. At iteration k, x^k is the current point and $H^k \succ 0$ is the positive definite inverse-Hessian estimate. The constants c_1 and c_2 are the Armijo and Wolfe parameters, respectively, and I denotes the identity matrix. Whenever f is differentiable at the accepted point x^{k+1}, the Wolfe condition (1.2) gives $(s^k)^T y^k > \alpha^k(c_2 - 1)(g^k)^T p^k > 0$. The update therefore preserves positive definiteness of inverse-Hessian estimates, so each subsequent nonzero gradient generates a descent direction.

Algorithm 1 BFGS with Armijo–Wolfe line search

Input: f, $x^0 \in \mathbb{R}^n$, $H^0 \succ 0$, and $0 < c_1 < c_2 < 1$.

$g^0 \leftarrow \nabla f(x^0)$.
for $k = 0, 1, 2, \ldots$ **do**
  Stop if $g^k = 0$; set $p^k = -H^k g^k$.
  Set $\varphi^k(\alpha) = f(x^k + \alpha p^k)$ and $(\ell, u, \alpha) = (0, +\infty, 1)$.
  Accept a trial when both strict tests hold:

$$\varphi^k(\alpha) < \varphi^k(0) + c_1 \alpha (g^k)^T p^k, \qquad \text{(Armijo)}, \tag{1.1}$$

$$(\varphi^k)'(\alpha) > c_2 (g^k)^T p^k, \qquad \text{(Wolfe)}. \tag{1.2}$$

  The Wolfe test fails if $(\varphi^k)'(\alpha)$ does not exist.
  while either (1.1) or (1.2) fails **do**
    If (1.1) fails, set $u = \alpha$; otherwise set $\ell = \alpha$.
    Set $\alpha = (\ell + u)/2$ if $u < +\infty$, and $\alpha = 2\ell$ otherwise.
  end while
  $\alpha^k \leftarrow \alpha$, $s^k \leftarrow \alpha^k p^k$, $x^{k+1} \leftarrow x^k + s^k$.
  $g^{k+1} \leftarrow \nabla f(x^{k+1})$, $y^k \leftarrow g^{k+1} - g^k$.
  $H^{k+1} \leftarrow \mathcal{B}(H^k, s^k, y^k)$, where

$$\mathcal{B}(H, s, y) = \left(I - \frac{sy^T}{s^T y}\right) H \left(I - \frac{ys^T}{s^T y}\right) + \frac{ss^T}{s^T y}. \tag{1.3}$$

end for

To state the main result, recall that for locally Lipschitz f, the Clarke subdifferential $\partial_C f(x)$ is the closed convex hull of limiting gradients at differentiability points approaching x [Cla75, Cla90]. A point x is Clarke stationary when $0 \in \partial_C f(x)$. A function is semialgebraic if its graph is a finite Boolean combination of sets defined by polynomial equalities and inequalities [BCR98, Sections 2.1–2.2]. We use $\|\cdot\|$ for the Euclidean norm and call f coercive if $f(x) \to +\infty$ as $\|x\| \to \infty$.

Let $\mathbb{S}^2_{++} \subset \mathbb{R}^{2\times 2}$ be the set of symmetric positive definite 2×2 matrices. The following is the main result of this paper.

Theorem 1.1. *There exist a coercive, locally Lipschitz, semialgebraic function $f : \mathbb{R}^2 \to \mathbb{R}$ with $0 \notin \partial_C f(0)$ and a nonempty open set $\mathcal{U} \subset \mathbb{R}^2 \times \mathbb{S}^2_{++}$ such that, for every $(x^0, H^0) \in \mathcal{U}$ and every parameter pair (c_1, c_2) satisfying*

$$0 < c_1 \le 0.03 < 0.6 \le c_2 < 1, \tag{1.4}$$

Algorithm 1 generates a well-defined infinite sequence with $x^k \to 0$.

Figure 1 shows the objective in Theorem 1.1 and BFGS iterates converging to the non-Clarke-stationary origin. Figure 2 plots the function values and iterate norms for 100 BFGS runs initialized by randomly perturbing both the initial point and inverse-Hessian estimate used in Figure 1.

Remark 1.2 (Positive probability of failure). If x^0 is a nondegenerate Gaussian vector with a positive definite covariance matrix and, independently, $H^0 = X^T X$ with independent standard Gaussian entries in the 2×2 matrix X, every nonempty open subset of $\mathbb{R}^2 \times \mathbb{S}^2_{++}$ has positive probability. The same event $\mathcal{U}$ gives failure for every parameter pair in (1.4), disproving the almost-sure stationarity assertion in [LO13, Challenge 7.1(2)].

Theorem 1.1 is proved in Section 2. The proof constructs a positively homogeneous function with an exact shrinking four-step BFGS cycle. A stability argument extends the failure to an open




Figure 1: BFGS convergence to a non-Clarke-stationary origin. Left: the objective landscape near the cycle. Right: the first 120 BFGS iterations with $(c_1, c_2) = (0.01, 0.9)$.




Figure 2: BFGS from 100 random initial states for the objective in Figure 1, with $(c_1, c_2) = (0.01, 0.9)$ and 1000 iterations per run. The initial states are sampled uniformly from the five-dimensional Euclidean ball of radius $\varepsilon = 10^{-9}$ centered at the constructed state (x^0, H^0), in coordinates $(x_1, x_2, H_{11}, H_{12}, H_{22})$. The left panel shows the initial position displacements $(\Delta x_1, \Delta x_2)$, while the middle and right panels show the objective values and iterate norms, respectively. Runs converging to the origin are shown in blue, and the remaining runs in orange.

set of initial states. In Section 2.1, a scalar polynomial equation and a linear involution determine a four-step pattern. After four updates, both the iterate and the inverse-Hessian estimate are scaled by 64/81. Once realized by a positively homogeneous function, this pattern repeats at successively

smaller scales, so the iterates converge to the origin. Section 2.2 constructs this function by interpolation. It matches the prescribed values and gradients while keeping its Clarke subgradients uniformly separated from zero. The interpolation also allows us to prescribe compatible Hessians for the subsequent stability argument. Section 2.3 verifies that, with the chosen Hessians and after dividing out the geometric decay, the four-step return map contracts directions transverse to the ray of scaled cycle states, and establishes an invariant neighborhood. Finally, Section 2.4, makes the objective coercive while preserving local Lipschitz continuity, then proves convergence from an open set of initial states.

The proof relies on computer-assisted certification of the finite inequalities in Appendix A, using rigorous interval arithmetic [MKC09, Rum10]. We also use symbolic computation in Python to confirm the cycle identities proved analytically in Section 2.1. The verification details, executed Jupyter notebook, and Lean 4 formalization of the main result are available in the accompanying GitHub repository.

2 Construction and proof

Recall that a function $\phi : \mathbb{R}^n \to \mathbb{R}$ is positively homogeneous of degree one if $\phi(tx) = t\phi(x)$ for every $x \in \mathbb{R}^n$ and $t \geq 0$. We use "homogeneous" in this sense throughout. We first record the consequences of homogeneity used in the construction and in the analysis of BFGS.

Proposition 2.1 (Homogeneity identities)**.** *Let $\phi : \mathbb{R}^n \to \mathbb{R}$ be positively homogeneous and C^2 near $x \in \mathbb{R}^n$. Then, for every $\tau > 0$, ϕ is C^2 near τx, and*

$$x^T \nabla\phi(x) = \phi(x), \qquad \nabla\phi(\tau x) = \nabla\phi(x), \qquad \nabla^2\phi(\tau x) = \tau^{-1}\nabla^2\phi(x). \tag{2.1}$$

Proof. Let U be an open neighborhood of x on which ϕ is C^2. For fixed $\tau > 0$, homogeneity gives $\phi(z) = \tau\phi(z/\tau)$ on τU. The right-hand side is C^2 there, so ϕ is C^2 in a neighborhood of τx. Differentiating $\phi(tx) = t\phi(x)$ with respect to t at $t = 1$ gives the first identity in (2.1). Differentiating $\phi(\tau z) = \tau\phi(z)$ once and twice with respect to z at x gives the derivative scaling identities. □

We next record how BFGS updates transform under scaling and invertible linear changes of coordinates. For homogeneous objectives, scaling the point and inverse-Hessian estimate together also preserves the line-search decisions.

Lemma 2.2 (BFGS scaling and covariance)**.** *For $H \in \mathbb{R}^{n\times n}$, invertible $A \in \mathbb{R}^{n\times n}$, $s, y \in \mathbb{R}^n$ with $s^T y \neq 0$, and $\tau > 0$, the update in* (1.3) *satisfies*

$$\mathcal{B}(\tau AHA^T, \tau As, A^{-T}y) = \tau A\mathcal{B}(H, s, y)A^T. \tag{2.2}$$

Let ϕ and x satisfy the hypotheses of Proposition 2.1, and let $H \succ 0$, $\nabla\phi(x) \neq 0$ and $\tau > 0$. Applied to ϕ with the same line-search parameters, Algorithm 1 tests the same step sizes and makes the same acceptance decisions from (x, H) and $(\tau x, \tau H)$. If the first run accepts a step and produces (x^1, H^1), and ϕ is C^2 near x^1, then the second accepts the same step size and produces $(\tau x^1, \tau H^1)$.

Proof. The transformed scalar product is $(\tau As)^T(A^{-T}y) = \tau s^T y$, and

$$I - \frac{(\tau As)(A^{-T}y)^T}{\tau s^T y} = A\left(I - \frac{sy^T}{s^T y}\right)A^{-1},$$
$$\frac{(\tau As)(\tau As)^T}{\tau s^T y} = \tau A\frac{ss^T}{s^T y}A^T.$$

Substitution into (1.3), together with the transpose of the first identity, proves (2.2).

Set $p = -H\nabla\phi(x)$. By the gradient scaling identity in Proposition 2.1, the search direction at $(\tau x, \tau H)$ is τp. For the line functions $\varphi(\alpha) = \phi(x + \alpha p)$ and $\widehat{\varphi}(\alpha) = \phi(\tau x + \alpha\tau p)$, homogeneity gives $\widehat{\varphi} = \tau\varphi$. Thus a line derivative exists at a trial in one search exactly when it exists in the other, and then $\widehat{\varphi}' = \tau\varphi'$. Both sides of the Armijo and Wolfe tests (1.1)–(1.2) are multiplied by $\tau > 0$, so the decisions and subsequent trial step sizes coincide. If ϕ is C^2 near the accepted iterate $x^1 = x + \alpha p$, Proposition 2.1 applies at both x and x^1. Thus the displacement is multiplied by τ and the gradient difference is unchanged. The Wolfe condition gives positive curvature, so the update is defined; (2.2) with $A = I$ proves the assertion about the resulting states. □

2.1 An exact four-step cycle

We construct the cycle from a scalar parameter and a linear involution. The involution determines the last two updates from the first two, and makes the common contraction factor explicit. Let

$$P(b) = 777924b^4 - 576632b^3 - 587665b^2 + 352256b + 65536. \tag{2.3}$$

Lemma 2.3 (Admissible root)**.** *The polynomial P has a unique root in* $(0.9178, 0.9179)$.

Proof. Set $a = 0.9178$ and $c = 0.9179$. Exact rational evaluation gives $P(a) < -2$ and $P(c) > 19$. For $z \in [a, c]$,

$$P'(z) \geq 3111696a^3 - 1729896c^2 - 1175330c + 352256 > 200000.$$

The intermediate value theorem gives a root in (a, c), and the positive derivative makes it unique. □

For the rest of the construction, fix the root b supplied by Lemma 2.3, and put $\rho = 8/9$ and $r = \rho^2 = 64/81$. Let $e_1 = (1, 0)^T$ and $e_2 = (0, 1)^T$ be the standard basis vectors. Define

$$\begin{gathered} p = -\frac{9(882b^2 - 463b - 256)}{640(2b - 1)}, \qquad t = -\frac{3087b^2 - 2233b - 256}{320(2b - 1)}, \\ q = \frac{15876b^3 - 4316b^2 - 6815b - 2304}{1280b(2b - 1)}, \qquad J = \begin{pmatrix} p & t \\ q & -p \end{pmatrix}. \end{gathered} \tag{2.4}$$

All denominators are nonzero because $b > 0.9178$. The rational identity

$$p^2 + qt - 1 = \frac{9P(b)}{409600b(2b - 1)} \tag{2.5}$$

gives $J^2 = I$ and $\det J = -1$. Set

$$(g^0, g^1, g^2, g^3, g^4) = (e_1, e_2, Je_1, Je_2, e_1) \tag{2.6}$$

and

$$\begin{gathered} H^0 = \begin{pmatrix} 1 & 1/2 \\ 1/2 & b/4 + 1/8 \end{pmatrix}, \qquad H^1 = \begin{pmatrix} b + 3/2 & b + 1/2 \\ b + 1/2 & b \end{pmatrix}, \\ H^2 = \rho J^T H^0 J, \qquad H^3 = \rho J^T H^1 J, \qquad H^4 = rH^0. \end{gathered} \tag{2.7}$$

For $v_0 = (b + 3/2, b + 1/2)^T$, define the positions by

$$\begin{gathered} x^0 = \frac{(I + \rho J^T)v_0}{1 - r}, \qquad x^1 = x^0 - (1, 1/2)^T, \\ x^2 = \rho J^T x^0, \qquad x^3 = \rho J^T x^1, \qquad x^4 = rx^0. \end{gathered} \tag{2.8}$$

Set $f^i = (g^i)^T x^i$ for $0 \le i \le 4$, and define

$$s^i = -H^i g^i, \qquad y^i = g^{i+1} - g^i \quad (0 \le i < 4). \tag{2.9}$$

The next lemma verifies a proposed two-dimensional BFGS update by checking its secant equation and determinant.

Lemma 2.4 (A two-dimensional update test)**.** *Let $H \in \mathbb{S}^2_{++}$ and $s, y \in \mathbb{R}^2$ satisfy $s^T y > 0$. A symmetric matrix $K \in \mathbb{R}^{2\times 2}$ equals $\mathcal{B}(H, s, y)$ if and only if*

$$Ky = s, \qquad \det K = \det H \, \frac{s^T H^{-1} s}{s^T y}. \tag{2.10}$$

Proof. The update (1.3) is symmetric and satisfies $\mathcal{B}(H, s, y)y = s$. Choose an orthonormal basis with $s = (\sigma, 0)^T$, where $\sigma = \|s\| > 0$, and write $y = (u, v)^T$ and $H = \left(\begin{smallmatrix} a & h \\ h & d \end{smallmatrix}\right)$. Since $\sigma u = s^T y > 0$, we have $u > 0$, and substitution into (1.3) gives

$$\mathcal{B}(H, s, y) = \begin{pmatrix} dv^2/u^2 + \sigma/u & -dv/u \\ -dv/u & d \end{pmatrix}.$$

Its determinant is $d\sigma/u$. Since $s^T H^{-1} s = d\sigma^2/\det H$, this is the determinant in (2.10), proving necessity.

Conversely, in an orthonormal basis whose first vector is $y/\|y\|$, the equation $Ky = s$ fixes the first column and, by symmetry, the first row of K. Its first diagonal entry is $s^T y/\|y\|^2 > 0$, so its determinant fixes the remaining entry. Hence the two conditions in (2.10) determine K uniquely, and the matrix is $\mathcal{B}(H, s, y)$. □

The following lemma supplies the exact cycle and the strict positivity needed for all its updates, together with bounds on the cycle positions.

Lemma 2.5 (Exact cycle)**.** *For the data just constructed, $H^i \succ 0$ and $\|x^i\|^2 < 8$ for $0 \le i \le 4$, and*

$$\begin{gathered} x^{i+1} = x^i + s^i, \qquad H^{i+1} = \mathcal{B}(H^i, s^i, y^i) \quad (0 \le i < 4), \\ H^4 = rH^0, \qquad x^4 = rx^0. \end{gathered} \tag{2.11}$$

The curvature products are positive and satisfy

$$\left((s^i)^T y^i\right)_{i=0}^3 = \left(\frac{1}{2}, \frac{81b}{32}, \frac{4}{9}, \frac{9b}{4}\right). \tag{2.12}$$

Proof. *Step 1: Positive definiteness and position bounds.* By Lemma 2.3, $b \in (0.9178, 0.9179)$ and $P(b) = 0$. Thus (2.5) gives $J^2 = I$. For H^0 and H^1 in (2.7), the determinants are $(2b-1)/8$ and $(2b-1)/4$, respectively. Since $b > 1/2$, both determinants and both leading diagonal entries are positive. The remaining matrices in (2.7) are positive multiples of congruences of these matrices, so all five are positive definite.

Substituting the bounds on b into (2.4) and (2.8) gives $|x^i_j| < 2$ for $j = 1, 2$ and $0 \le i < 4$. The definition $x^4 = rx^0$, with $0 < r < 1$, gives the same coordinate bounds for x^4. Hence $\|x^i\|^2 < 8$ for $0 \le i \le 4$.

Step 2: The first update. By (2.9), $(s^i)^T (H^i)^{-1} s^i = (g^i)^T H^i g^i$ for $0 \le i < 4$. For the first update, (2.6) gives $g^0 = e_1$ and $g^1 = e_2$, so $s^0 = -H^0 e_1 = (-1, -1/2)^T$ and $y^0 = e_2 - e_1 = (-1, 1)^T$. Hence

$$H^1 y^0 = \begin{pmatrix} b + 3/2 & b + 1/2 \\ b + 1/2 & b \end{pmatrix} \begin{pmatrix} -1 \\ 1 \end{pmatrix} = \begin{pmatrix} -(b + 3/2) + (b + 1/2) \\ -(b + 1/2) + b \end{pmatrix} = \begin{pmatrix} -1 \\ -1/2 \end{pmatrix} = s^0.$$

Moreover, $(s^0)^T y^0 = (-1)(-1)+(-1/2)(1) = 1/2$ and $(g^0)^T H^0 g^0 = e_1^T H^0 e_1 = (H^0)_{11} = 1$. Together with $\det H^1 / \det H^0 = 2$, these identities verify (2.10), so Lemma 2.4 proves the first update.

Step 3: The second update. Equation (2.6) gives $g^1 = e_2$ and $g^2 = Je_1 = (p, q)^T$. The definitions in (2.7) and (2.9) give $s^1 = (-(b + 1/2), -b)^T$ and $y^1 = (p, q - 1)^T$. Since $\rho = 8/9$, expanding the matrix products gives

$$\rho H^0 J e_2 - J^T H^1 e_2 - \rho H^0 e_1 = \begin{pmatrix} \frac{8}{9}(t - \frac{1}{2}p) - [(b + \frac{1}{2})p + bq] - \frac{8}{9} \\ \frac{1}{9}[(7b - 1)p - (9b + \frac{1}{2})t - 4] \end{pmatrix}. \tag{2.13}$$

For the first entry, substituting (2.4) and collecting terms gives

$$t - \tfrac{1}{2}p = \frac{-4410b^2 + 4765b - 1280}{1280(2b - 1)} = 1 - \frac{441b}{256},$$
$$(b + \tfrac{1}{2})p + bq = \frac{-3920b^2 + 1960b}{1280(2b - 1)} = -\frac{49b}{32}.$$

Thus the first entry is $\frac{8}{9}(1 - 441b/256) + 49b/32 - \frac{8}{9} = 0$. For the second entry, the same substitution gives

$$(7b - 1)p - (9b + \tfrac{1}{2})t = \frac{5120b - 2560}{640(2b - 1)} = 4,$$

so this entry is $(4 - 4)/9 = 0$. Hence the vector in (2.13) vanishes. Substituting the definitions of H^2 and y^1 from (2.7) and (2.9), respectively, then using $J^2 = I$ and the vanishing identity (2.13) multiplied by J^T, gives

$$H^2 y^1 = \rho J^T H^0 J(Je_1 - e_2) = \rho J^T H^0 (e_1 - Je_2) = -H^1 e_2 = s^1.$$

The curvature product is

$$(s^1)^T y^1 = -(b + \tfrac{1}{2})p - b(q - 1) = \frac{49b}{32} + b = \frac{81b}{32} > 0,$$

and $(g^1)^T H^1 g^1 = e_2^T H^1 e_2 = b$. Finally, $\det J = -1$ and the matrices in (2.7) give

$$\frac{\det H^2}{\det H^1} = \rho^2 (\det J)^2 \frac{\det H^0}{\det H^1} = \left(\frac{8}{9}\right)^2 \frac{1}{2} = \frac{32}{81} = \frac{(g^1)^T H^1 g^1}{(s^1)^T y^1}.$$

These identities verify (2.10), so Lemma 2.4 proves the second update.

Step 4: The third and fourth updates. We express their data in terms of those of the first two updates. By (2.6) and $J^2 = I$, the gradients satisfy $g^{i+2} = Jg^i$ for $i = 0, 1, 2$. The definitions in (2.7) give $H^{i+2} = \rho J^T H^i J$ for $i = 0, 1$; the same identity holds for $i = 2$ because

$$\rho J^T H^2 J = \rho^2 (J^T)^2 H^0 J^2 = rH^0 = H^4.$$

Thus, for $i = 0, 1$, the definitions in (2.9) give

$$\begin{aligned} s^{i+2} &= -H^{i+2} g^{i+2} = -\rho J^T H^i J^2 g^i = \rho J^T s^i, \\ y^{i+2} &= g^{i+3} - g^{i+2} = J(g^{i+1} - g^i) = Jy^i. \end{aligned} \tag{2.14}$$

In particular,

$$(s^{i+2})^T y^{i+2} = \rho (s^i)^T J^2 y^i = \rho (s^i)^T y^i > 0 \qquad (i = 0, 1).$$

Apply Lemma 2.2 with $A = J^T$ and $\tau = \rho$, so that $A^{-T} = J$. Using the first two updates proved in Steps 2–3, (2.2) gives

$$\mathcal{B}(H^{i+2}, s^{i+2}, y^{i+2}) = \rho J^T \mathcal{B}(H^i, s^i, y^i) J = \rho J^T H^{i+1} J = H^{i+3} \qquad (i = 0, 1).$$

Taking $i = 0$ and $i = 1$ proves the third and fourth updates, respectively. The curvature identity also gives $(s^2)^T y^2 = \rho/2 = 4/9$ and $(s^3)^T y^3 = \rho(81b/32) = 9b/4$.

Step 5: Position identities. Since $s^0 + s^1 = -v_0$ and $(I - \rho J^T)(I + \rho J^T) = (1 - r)I$, the definition of x^0 in (2.8) gives $x^0 + s^0 + s^1 = \rho J^T x^0 = x^2$. The other position identities follow from (2.8) and the step transformations in (2.14). □

2.2 Realizing the objective and the line search

We now realize the cycle of Lemma 2.5 by a single objective. The prescribed function values enforce the line-search decisions, while the gradients generate the cycle. The Hessians will make convergence persist under perturbations of the initial point and matrix. The construction reduces to interpolation of two univariate functions. The following consequence of shape-preserving Hermite interpolation [Man01, Sections 4.1–4.2] matches the prescribed values and first two derivatives while keeping the derivative positive.

Lemma 2.6 (Monotone interpolation)**.** *Let $t_1 < \cdots < t_N$ be interpolation points, with $N \geq 1$. For $1 \leq j \leq N$, let $\eta_j, p_j, P_j \in \mathbb{R}$ be prescribed values of the function and its first two derivatives at t_j. Suppose $0 < m < 1$ and*

$$p_j > m \quad (1 \leq j \leq N), \qquad \frac{\eta_{j+1} - \eta_j}{t_{j+1} - t_j} > m \quad (1 \leq j < N).$$

Then there is a C^2 semialgebraic function $h : \mathbb{R} \to \mathbb{R}$ with $h'(t) \geq m$ for every $t \in \mathbb{R}$ and

$$h(t_j) = \eta_j, \qquad h'(t_j) = p_j, \qquad h''(t_j) = P_j \quad (1 \leq j \leq N).$$

Moreover, the affine tails can be chosen as

$$h(t) = t + \eta_1 - t_1 \quad (t \leq t_1 - 1), \qquad h(t) = t + \eta_N - t_N \quad (t \geq t_N + 1).$$

Proof. Adjoin the points $t_0 = t_1 - 1$ and $t_{N+1} = t_N + 1$, with

$$\eta_0 = \eta_1 - 1, \qquad \eta_{N+1} = \eta_N + 1, \qquad p_0 = p_{N+1} = 1, \qquad P_0 = P_{N+1} = 0.$$

The two new secant slopes are one, so the extended data satisfy the same strict inequalities. Consider the shifted Hermite data

$$(t_j, \eta_j - m t_j, p_j - m, P_j) \qquad (0 \leq j \leq N + 1).$$

Their consecutive value increments and prescribed first derivatives are positive. They therefore satisfy the monotonicity conditions in [Man01, Definition 2.1]: the strict positivity of $p_j - m$ makes the endpoint conditions hold for sufficiently small positive parameters, regardless of P_j. The parametric construction [Man01, Section 4.1 and Section 4.2, equations (4.7)–(4.9)] then gives a C^2 interpolant g on $[t_0, t_{N+1}]$ with $g' \geq 0$. Its graph consists of finitely many polynomially parametrized arcs, so it is semialgebraic by the Tarski–Seidenberg theorem [BCR98, Theorem 2.2.1].

Set $h(t) = g(t) + mt$ on $[t_0, t_{N+1}]$ and extend it by the two affine formulas in the statement. The function g matches the shifted data, so h matches the original data and satisfies $h' \geq m$ on this interval. The affine extensions match its endpoint values, first derivatives one and second derivatives zero. Hence h is C^2 and semialgebraic on $\mathbb{R}$, with $h' \geq m$ everywhere. □

Recall from Section 2.1 the cycle positions x^i, matrices H^i, gradients g^i, values f^i, steps s^i, gradient differences y^i, and contraction factor r. To realize these data, consider the change of coordinates

$$M = \begin{pmatrix} 1 & 0.05 \\ 0 & 1 \end{pmatrix}, \qquad v = Me_2 = (0.05, 1)^T, \qquad z^i = M^{-1}x^i \quad (0 \le i < 4). \tag{2.15}$$

We seek two functions h_-, h_+ from Lemma 2.6 and use them to define

$$F(z) = \begin{cases} z_1 h_+(z_2/z_1), & z_1 > 0, \\ -z_1 h_-(z_2/(-z_1)), & z_1 < 0, \\ z_2, & z_1 = 0, \end{cases} \qquad \widetilde{f}(x) = F(M^{-1}x). \tag{2.16}$$

For $z_1 \neq 0$, the first formula is $F(z) = |z_1| h_{\mathrm{sign}(z_1)}(z_2/|z_1|)$. Positive scaling leaves the ratio unchanged and scales the prefactor, so F and $\widetilde{f}$ are positively homogeneous. Moreover, $\partial_2 F(z) = h'_{\mathrm{sign}(z_1)}(z_2/|z_1|)$ for $z_1 \neq 0$. A common positive lower bound on h'_+ and h'_- will therefore separate the Clarke subgradients from zero.

The next proposition determines which cycle data are assigned to h_+ and h_- and orders their interpolation points $z_2^i/|z_1^i|$.

Proposition 2.7 (Signs and slope order)**.** *The points in* (2.15) *satisfy* $z_1^0, z_1^1 > 0$ *and* $z_1^2, z_1^3 < 0$. *Writing* $t^i = z_2^i/|z_1^i|$ *for* $0 \le i < 4$, *we have* $-1 < t^0 < t^1 < 6$ *and* $-1 < t^2 < t^3 < 6$.

Proof. Lemma 2.3 gives $0.9178 < b < 0.9179$. Substitution into (2.4), (2.8) and (2.15) bounds z_1^i away from zero. Dividing z_2^i by $|z_1^i|$ then gives the following enclosures:

i	z_1^i	t^i
0	$(1.1, 1.3)$	$(1.3, 1.5)$
1	$(0.2, 0.3)$	$(4, 5.8)$
2	$(-1.2, -1)$	$(0.2, 0.4)$
3	$(-0.3, -0.2)$	$(2.6, 4.4)$

These bounds give the asserted signs and both strict orderings. □

To match the cycle positions, prescribe the following data

$$\eta^i = \frac{f^i}{|z_1^i|}, \qquad d^i = (g^i)^T v \quad (0 \le i < 4). \tag{2.17}$$

We assign $(h_+(t^i), h'_+(t^i)) = (\eta^i, d^i)$ for $i = 0, 1$, and the corresponding data for h_- when $i = 2, 3$. These data guarantee that the gradients of $\widetilde{f}$ at x^i are g^i; see the proof of Lemma 2.10. Adding the identity values at -1 and 6 gives the two ordered lists of interpolation points and function values:

$$\begin{aligned} h_+ :&\quad (-1,-1),\ (t^0,\eta^0),\ (t^1,\eta^1),\ (6,6), \\ h_- :&\quad (-1,-1),\ (t^2,\eta^2),\ (t^3,\eta^3),\ (6,6). \end{aligned} \tag{2.18}$$

The next proposition verifies the first-derivative and secant-slope conditions of Lemma 2.6. Its proof uses the enclosures in Tables 1 and 2 of Appendix A.1, obtained from the root bounds in Lemma 2.3.

Proposition 2.8 (Interpolation certificate)**.** *The first derivatives prescribed in* (2.17) *satisfy* $0.02 < d^i \le 1$ *for* $0 \le i < 4$*. In either list in* (2.18)*, every two consecutive entries* $(a, A), (c, B)$ *satisfy*

$$c - a > 1, \qquad 0.5 < \frac{B - A}{c - a} < 2.$$

Proof. Equations (2.6), (2.15) and (2.17) give $d^0 = 0.05$, $d^1 = 1$, $d^2 = 0.05p + q$ and $d^3 = 0.05t - p$. The enclosures in Table 1 place d^2 and d^3 strictly between 0.02 and 1. For the six consecutive gaps in (2.18), Table 2 gives lower endpoints greater than one for $c - a$, and encloses each secant slope $(B - A)/(c - a)$ strictly between 0.5 and 2. □

We next consider the line-search requirements and define

$$\mathsf{A}^i = f^i + 0.03(g^i)^T s^i - f^{i+1}, \qquad \mathsf{W}^i = (g^{i+1})^T s^i - 0.6(g^i)^T s^i \quad (0 \le i < 4). \tag{2.19}$$

Lemma 2.5 gives $H^i \succ 0$ and $(s^i)^T y^i > 0$. Thus $s^i \neq 0$ and $g^i \neq 0$ by (2.9), so $(g^i)^T s^i = -(g^i)^T H^i g^i < 0$. Consequently A^i bounds the unit-step Armijo margin from below when $c_1 \le 0.03$, and W^i bounds the Wolfe margin from below when $c_2 \ge 0.6$. The next proposition verifies that both margins are positive.

Proposition 2.9 (Line-search margins)**.** *The margins in* (2.19) *satisfy* $\mathsf{A}^i > 0$ *and* $\mathsf{W}^i > 0$ *for* $0 \le i < 4$*.*

Proof. The definitions (2.6) and (2.8), together with $J^2 = I$, give $f^{i+2} = \rho f^i$ for $i = 0, 1, 2$. Using (2.6) and (2.14) in (2.19) therefore yields

$$\mathsf{A}^{i+2} = \rho \mathsf{A}^i, \qquad \mathsf{W}^{i+2} = \rho \mathsf{W}^i \qquad (i = 0, 1).$$

Since $\rho > 0$, it suffices to check the first two phases.

For the Armijo margins, substitute $f^i = (g^i)^T x^i$ and the data in (2.4)–(2.9) into (2.19). Simplifying with $P(b) = 0$ from Lemma 2.3 gives

$$\mathsf{A}^0 = -\frac{158760b^3 + 163620b^2 - 172579b - 103680}{27200b},$$
$$\mathsf{A}^1 = \frac{79380b^3 + 58957b^2 - 70380b - 46080}{13600b}.$$

Rational interval evaluation on $[0.9178, 0.9179]$, which contains b by Lemma 2.3, gives

$$0.0575 < \mathsf{A}^0 < 0.0611, \qquad 0.0281 < \mathsf{A}^1 < 0.0312. \tag{2.20}$$

For the Wolfe margins, (2.6)–(2.7) and (2.9) give $(g^0)^T s^0 = -1$ and $(g^1)^T s^1 = -b$. Combining these identities with (2.12) gives

$$\mathsf{W}^0 = \frac{1}{2} - 0.4 = 0.1 > 0, \qquad \mathsf{W}^1 = \frac{81b}{32} - 0.4b = \frac{341b}{160} > 0.$$

Thus all four Armijo and Wolfe margins are positive. □

Lemma 2.6 allows arbitrary second derivatives. The next lemma uses this freedom to realize the exact cycle with adjustable Hessians. Their coefficients will be chosen in the next subsection so that convergence persists under perturbations of the initialization.

Lemma 2.10 (Realization and accepted steps)**.** *For arbitrary $\beta^0, \dots, \beta^3 \in \mathbb{R}$, set*

$$q^i = (-x_2^i, x_1^i)^T, \qquad K^i = \frac{\beta^i}{\|x^i\|^2} q^i (q^i)^T \quad (0 \le i < 4). \tag{2.21}$$

Then there is a globally Lipschitz, positively homogeneous, semialgebraic function $\widetilde{f} : \mathbb{R}^2 \to \mathbb{R}$ with $0 \notin \partial_{\mathrm{C}} \widetilde{f}(0)$. It is C^2 on $\mathbb{R}^2 \setminus \{0\}$ and satisfies

$$\widetilde{f}(x^i) = f^i, \qquad \nabla \widetilde{f}(x^i) = g^i, \qquad \nabla^2 \widetilde{f}(x^i) = K^i \quad (0 \le i < 4).$$

For every parameter pair in (1.4)*, Algorithm 1 applied to $\widetilde{f}$ from (x^0, H^0) accepts the initial trial $\alpha = 1$ at every iteration, and*

$$x^{4j+i} = r^j x^i, \qquad H^{4j+i} = r^j H^i \quad (j \ge 0,\ 0 \le i < 4).$$

Proof. Step 1: Construction and interpolation. The vectors q^i in (2.21) are perpendicular to x^i, so $K^i x^i = 0$. Complete the interpolation data by prescribing

$$h''_{\mathrm{sign}(z_1^i)}(t^i) = k^i, \qquad k^i = |z_1^i| v^T K^i v \quad (0 \le i < 4). \tag{2.22}$$

Apply Lemma 2.6 separately to the two lists in (2.18), using the bounds in Proposition 2.8, with $m = 0.01$, first derivative one and second derivative zero at the anchors $-1, 6$, and the data (2.17) and (2.22) at the other points. The second derivatives are unrestricted, so this construction applies to every choice of $\beta^0, \dots, \beta^3$. Choose any interpolants supplied by the lemma. The prescribed values at the extreme anchors are -1 and 6, so its affine-tail formulas give $h_\pm(t) = t$ for $t \notin (-2, 7)$. Define F and $\widetilde{f}$ by (2.16). These functions are positively homogeneous and semialgebraic. For $z_1 \ne 0$, put $a = |z_1|$, $\sigma = \mathrm{sign}(z_1)$ and $t = z_2/a$. Differentiation gives

$$\nabla F(z) = \begin{pmatrix} \sigma(h_\sigma(t) - t h'_\sigma(t)) \\ h'_\sigma(t) \end{pmatrix}, \qquad \nabla^2 F(z) = \frac{h''_\sigma(t)}{a} \begin{pmatrix} t^2 & -\sigma t \\ -\sigma t & 1 \end{pmatrix}. \tag{2.23}$$

The identity tails just established give $F(z) = z_2$ in a neighborhood of every nonzero point of the axis $z_1 = 0$. Thus F and $\widetilde{f}$ are C^2 away from zero.

For $0 \le i < 4$, the prescriptions in (2.17) give $\widetilde{f}(x^i) = F(z^i) = f^i$ and $\partial_2 F(z^i) = d^i$. Proposition 2.1 gives $(z^i)^T \nabla F(z^i) = f^i$. By (2.15) and (2.17), $x^i = z_1^i M e_1 + z_2^i M e_2$ and $d^i = (g^i)^T M e_2$. Since $f^i = (g^i)^T x^i$ and $z_1^i \ne 0$ by Proposition 2.7, the first component is

$$\partial_1 F(z^i) = \frac{f^i - z_2^i d^i}{z_1^i} = \frac{(g^i)^T (x^i - z_2^i M e_2)}{z_1^i} = (g^i)^T M e_1.$$

Together with $\partial_2 F(z^i) = d^i = (g^i)^T M e_2$, this gives $\nabla F(z^i) = M^T g^i$. The chain rule applied to (2.16) then yields $\nabla \widetilde{f}(x^i) = M^{-T} \nabla F(z^i) = g^i$.

For the Hessian, the symmetric matrix $M^T K^i M$ satisfies $(M^T K^i M) z^i = 0$, because $M z^i = x^i$ and $K^i x^i = 0$ by (2.21). At $z = z^i$, the notation in (2.23) gives $z^i = a(\sigma, t)^T$ and $\sigma^2 = 1$, so multiplying the displayed Hessian by z^i gives $\nabla^2 F(z^i) z^i = 0$. Since $z_1^i \ne 0$ by Proposition 2.7, the two rows of $B z^i = 0$ for a symmetric matrix B give

$$B_{12} = -\frac{z_2^i}{z_1^i} B_{22}, \qquad B_{11} = \left(\frac{z_2^i}{z_1^i}\right)^2 B_{22}.$$

Thus B is determined by B_{22}. The $(2,2)$ entries of M^TK^iM and $\nabla^2F(z^i)$ both equal $v^TK^iv = k^i/a$ by (2.15), (2.22) and (2.23). Hence $\nabla^2F(z^i) = M^TK^iM$. The chain rule for $\widetilde{f}(x) = F(M^{-1}x)$ therefore gives $\nabla^2\widetilde{f}(x^i) = K^i$.

Step 2: Lipschitz continuity and nonstationarity. Continuity on the unit circle and homogeneity give $\widetilde{f}(x) = \|x\|\widetilde{f}(x/\|x\|) \to 0 = \widetilde{f}(0)$ as $x \to 0$. Since $\widetilde{f}$ is C^2 away from zero, the gradient scaling identity in (2.1) gives

$$\|\nabla\widetilde{f}(x)\| \le L := \max_{\|u\|=1}\|\nabla\widetilde{f}(u)\| < \infty \qquad (x \neq 0).$$

Integrating along any segment avoiding zero gives $|\widetilde{f}(x)-\widetilde{f}(y)| \le L\|x-y\|$. If the segment contains zero, split it there and pass to the limit using continuity. The two lengths add to $\|x-y\|$, so the same inequality holds, proving global Lipschitz continuity.

By Lemma 2.6 and (2.23), the second component of ∇F is at least 0.01 off zero, including on the nonzero axis, where $F(z) = z_2$ locally. With $v = Me_2$ from (2.15), the chain rule gives $\nabla\widetilde{f}(x)^Tv = \partial_2F(M^{-1}x) \ge 0.01$ at every nonzero point. All limiting gradients at zero, and their closed convex hull, satisfy the same inequality. Thus $0 \notin \partial_{\mathrm{C}}\widetilde{f}(0)$.

Step 3: Line-search decisions. By Proposition 2.1, $x^4 = rx^0$ from Lemma 2.5, and $g^4 = g^0$ from (2.6), we have $\widetilde{f}(x^4) = f^4$ and $\nabla\widetilde{f}(x^4) = g^4 = g^0$. Proposition 2.9 gives $\mathsf{A}^i, \mathsf{W}^i > 0$ for the margins in (2.19). As noted after that definition, these are lower bounds for the unit-step Armijo and Wolfe margins throughout (1.4). Since $p^i = s^i$ and $x^i + s^i = x^{i+1}$, every cycle state accepts its initial unit trial. For $\tau > 0$, Lemma 2.2 shows that a step from $(\tau x^i, \tau H^i)$ makes the same line-search decisions and ends at $(\tau x^{i+1}, \tau H^{i+1})$. Since $(x^4, H^4) = (rx^0, rH^0)$ by Lemma 2.5, induction over successive four-step blocks gives the asserted formulas. □

2.3 Stability of the shrinking cycle

Lemma 2.10 proves convergence to the origin for the exact cycle and its positive rescalings. To obtain an open set of initial states, we must show that this behavior persists under perturbations of both x and H. We divide the state after $4j$ steps by r^j, the decay factor of the exact cycle, and show that these normalized states remain bounded. A related use of homogeneity and attracting cycles in normalized coordinates appears in a construction for stochastic subgradient methods [Li26, Propositions 3.1 and 4.1].

We first express one accepted BFGS step as a map of the current state. Along the exact cycle, Lemma 2.10 gives an accepted step size of one. We study the map with this fixed step size; the final proof verifies that the line search accepts its steps throughout an invariant neighborhood. Represent a state $(x, H) \in \mathbb{R}^2 \times \mathbb{S}^2_{++}$ by $\omega = (x_1, x_2, H_{11}, H_{12}, H_{22}) \in \mathbb{R}^5$ and write $\omega_* = (x_1^0, x_2^0, 1, 1/2, b/4+1/8)$ for the base state from (2.7)–(2.8). For any fixed choice of $\beta^0, \dots, \beta^3$ and the corresponding function $\widetilde{f}$ constructed in Lemma 2.10, define the unit-step BFGS map by

$$\begin{aligned} s = -H\nabla\widetilde{f}(x), \qquad y = \nabla\widetilde{f}(x+s) - \nabla\widetilde{f}(x), \qquad \widehat{H} = \mathcal{B}(H,s,y), \\ \mathcal{T}(\omega) = (x_1 + s_1, x_2 + s_2, \widehat{H}_{11}, \widehat{H}_{12}, \widehat{H}_{22}), \end{aligned} \tag{2.24}$$

where $\mathcal{B}$ is given by (1.3). Its domain is the open set

$$\mathcal{D} = \{\omega \in \mathbb{R}^5 : H \succ 0,\ x \neq 0,\ x + s \neq 0,\ s^Ty > 0\}. \tag{2.25}$$

Write $\mathcal{T}^j$ for the j-fold composition wherever defined, with $\mathcal{T}^0 = \mathrm{id}$. The next lemma records the smoothness and scaling properties needed to study the shrinking cycle.

Lemma 2.11 (Local BFGS dynamics)**.** *The map $\mathcal{T}$ is C^1 on $\mathcal{D}$. For every $\omega \in \mathcal{D}$ and every $\tau > 0$,*

$$\mathcal{T}(\tau\omega) = \tau\mathcal{T}(\omega), \qquad \mathcal{T}^4(\omega_*) = r\omega_*.$$

Proof. On $\mathcal{D}$ in (2.25), $\widetilde{f}$ is C^2 near x and $x + s$, and $s^T y > 0$ keeps the denominators in (1.3) nonzero. Hence (2.24) defines a C^1 map there. By the gradient scaling in Proposition 2.1, replacing ω by $\tau\omega$ multiplies s by τ and leaves y unchanged. This preserves $H \succ 0$, the conditions $x \neq 0$ and $x + s \neq 0$, and the inequality $s^T y > 0$. Hence $\tau\omega \in \mathcal{D}$ for every $\tau > 0$. The update scaling in Lemma 2.2 and (2.24) then give $\mathcal{T}(\tau\omega) = \tau\mathcal{T}(\omega)$. Applying $\mathcal{T}$ along the exact cycle in Lemma 2.10 gives $\mathcal{T}^4(\omega_*) = r\omega_*$. □

For a sequence $\omega^{k+1} = \mathcal{T}(\omega^k)$, normalize each four-step block by setting $\xi^j = r^{-j}\omega^{4j}$. Write $\mathcal{D}_4 = \mathrm{dom}(\mathcal{T}^4)$, an open set invariant under positive scaling by Lemma 2.11. On this set, define the normalized return map

$$\mathcal{F}(\omega) = r^{-1}\mathcal{T}^4(\omega). \tag{2.26}$$

The same lemma gives, whenever the updates are defined,

$$\xi^{j+1} = \mathcal{F}(\xi^j), \qquad \omega^{4j+i} = r^j\mathcal{T}^i(\xi^j) \quad (j \geq 0,\ 0 \leq i < 4). \tag{2.27}$$

Thus, if the normalized states ξ^j form a bounded sequence, then $\omega^{4j} = r^j\xi^j \to 0$, since $r = 64/81 \in (0, 1)$ by the construction in Section 2.1. The map $\mathcal{F}$ is C^1 and positively homogeneous on $\mathcal{D}_4$, and $\mathcal{F}(a\omega_*) = a\omega_*$ for every $a > 0$. We will show that, for ξ^0 sufficiently close to ω_*, the sequence (ξ^j) converges to a point of the fixed ray $\{a\omega_* : a > 0\}$ and is therefore bounded.

The Hessians prescribed in (2.21) determine the derivative of this return map along the cycle. To obtain attraction, choose

$$(\beta^0, \beta^1, \beta^2, \beta^3) = (1.9, 1.6, -4.1, 0.2). \tag{2.28}$$

For the rest of the proof, fix the corresponding objective $\widetilde{f}$ and maps $\mathcal{T}, \mathcal{F}$. The following lemma keeps nearby normalized states close to the exact cycle.

Lemma 2.12 (Attraction to the fixed ray)**.** *There are arbitrarily small open neighborhoods $U \subset \mathcal{D}_4$ of ω_* such that $\mathcal{F}(U) \subset U$ and, for every $\xi^0 \in U$, the iterates $\xi^{j+1} = \mathcal{F}(\xi^j)$ converge to a positive multiple of ω_*.*

Proof. *Step 1: Integral identities.* Since $(\omega_*)_3 = 1$, write each state uniquely as $\xi = \xi_3\omega_* + u$, where $u \in W := \{u \in \mathbb{R}^5 : u_3 = 0\}$. For $\xi' = \mathcal{F}(\xi)$, set $u' = \xi' - \xi'_3\omega_*$. Suppose $\xi_3 > 0$ and the segment $\xi(t) = \xi_3\omega_* + tu$, $0 \leq t \leq 1$, lies in $\mathcal{D}_4$. Since $\mathcal{F}$ is C^1 and $\mathcal{F}(\xi_3\omega_*) = \xi_3\omega_*$, the fundamental theorem of calculus applied to its third component gives

$$\xi'_3 - \xi_3 = (\mathcal{F}(\xi))_3 - (\mathcal{F}(\xi_3\omega_*))_3 = \int_0^1 (D\mathcal{F}(\xi(t))u)_3\, dt.$$

Using this identity and applying the same theorem to each component of $\mathcal{F}$ gives

$$\begin{aligned} u' &= \xi' - \xi'_3\omega_* \\ &= \big(\mathcal{F}(\xi) - \mathcal{F}(\xi_3\omega_*)\big) - (\xi'_3 - \xi_3)\omega_* \\ &= \int_0^1 D\mathcal{F}(\xi(t))u\, dt - \left(\int_0^1 (D\mathcal{F}(\xi(t))u)_3\, dt\right)\omega_* \\ &= \int_0^1 \big(D\mathcal{F}(\xi(t))u - (D\mathcal{F}(\xi(t))u)_3\omega_*\big)\, dt. \end{aligned}$$

The vector in the last integrand belongs to W, since its third coordinate is zero. For any norm $\|\cdot\|_\perp$ on W, the triangle inequalities for these integrals yield

$$\begin{aligned} |\xi_3' - \xi_3| &\le \int_0^1 |(D\mathcal{F}(\xi(t))u)_3|\,dt, \\ \|u'\|_\perp &\le \int_0^1 \|D\mathcal{F}(\xi(t))u - (D\mathcal{F}(\xi(t))u)_3\omega_*\|_\perp\,dt. \end{aligned} \tag{2.29}$$

Step 2: Operator-norm bounds. For each fixed $z \in \mathcal{D}_4$, define the linear maps $L_z : W \to W$ and $\ell_z : W \to \mathbb{R}$ by

$$L_z(v) = D\mathcal{F}(z)v - (D\mathcal{F}(z)v)_3\omega_*, \qquad \ell_z(v) = (D\mathcal{F}(z)v)_3.$$

The two integrands in (2.29) are therefore $|\ell_{\xi(t)}(u)|$ and $\|L_{\xi(t)}(u)\|_\perp$. We choose a norm on W in which L_{ω_*} is a contraction, then use continuity to bound both operators uniformly near ω_*. Define $R : \mathbb{R}^5 \to \mathbb{R}^4$ by $R(v) = (v_1, v_2, v_4, v_5)$. Since $R|_W : W \to \mathbb{R}^4$ is an isomorphism, there is a unique matrix $A \in \mathbb{R}^{4\times 4}$ satisfying

$$R\big(L_{\omega_*}(v)\big) = AR(v) \qquad (v \in W). \tag{2.30}$$

For a vector y, write $\|y\|_\infty = \max_i |y_i|$. For a matrix $B = (b_{ij})$, the induced matrix norm is

$$\|B\|_\infty = \sup_{y\neq 0} \frac{\|By\|_\infty}{\|y\|_\infty} = \max_i \sum_j |b_{ij}|.$$

Proposition A.1, proved in Section A.2, supplies an invertible matrix S with $\|SAS^{-1}\|_\infty \le 0.78$. Equip W with the norm $\|v\|_\perp = \|SR(v)\|_\infty$. We use operator norms induced by this norm on W and the absolute value on $\mathbb{R}$. By (2.30) and the isomorphism $SR : W \to \mathbb{R}^4$,

$$\|L_{\omega_*}\|_{\mathrm{op}} = \sup_{v\in W\setminus\{0\}} \frac{\|SAR(v)\|_\infty}{\|SR(v)\|_\infty} = \|SAS^{-1}\|_\infty \le 0.78 < 1.$$

Choose $\gamma \in (0.78, 1)$ and $C > \|\ell_{\omega_*}\|_{\mathrm{op}}$. Because $D\mathcal{F}$ is continuous, both L_z and ℓ_z depend continuously on z in these operator norms. Choose $\varepsilon \in (0,1)$ and $\delta > 0$ small enough that every state z satisfying $|z_3 - 1| \le \varepsilon$ and $\|z - z_3\omega_*\|_\perp \le \delta$ belongs to $\mathcal{D}_4$ and obeys

$$\|L_z\|_{\mathrm{op}} \le \gamma, \qquad \|\ell_z\|_{\mathrm{op}} \le C.$$

If $|\xi_3 - 1| \le \varepsilon$ and $\|u\|_\perp \le \delta$, then $(\xi(t))_3 = \xi_3$ and $\|\xi(t) - \xi_3\omega_*\|_\perp = t\|u\|_\perp \le \delta$. Thus every point of the segment satisfies these operator bounds, and

$$\begin{aligned} |\ell_{\xi(t)}(u)| &\le \|\ell_{\xi(t)}\|_{\mathrm{op}}\|u\|_\perp \le C\|u\|_\perp, \\ \|L_{\xi(t)}(u)\|_\perp &\le \|L_{\xi(t)}\|_{\mathrm{op}}\|u\|_\perp \le \gamma\|u\|_\perp \qquad (0 \le t \le 1). \end{aligned}$$

Substituting into (2.29) gives

$$\|u'\|_\perp \le \gamma\|u\|_\perp, \qquad |\xi_3' - \xi_3| \le C\|u\|_\perp \tag{2.31}$$

throughout this range of ξ_3 and u.

Step 3: An invariant neighborhood. Choose $M \ge \max\{C/(1-\gamma), \varepsilon/\delta\}$ and set

$$U = \{\xi \in \mathbb{R}^5 : |\xi_3 - 1| + M\|\xi - \xi_3\omega_*\|_\perp < \varepsilon\},$$

which is an open neighborhood of ω_* in $\mathbb{R}^5$, and $M \geq \varepsilon/\delta$ gives $U \subset \mathcal{D}_4$. Decreasing ε and δ makes U arbitrarily small. By (2.31),

$$|\xi_3' - 1| + M\|u'\|_\perp \leq |\xi_3 - 1| + (C + M\gamma)\|u\|_\perp \leq |\xi_3 - 1| + M\|u\|_\perp < \varepsilon,$$

so $\mathcal{F}(U) \subset U$. For $\xi^0 \in U$, write $\xi^j = \xi_3^j \omega_* + u^j$. Iterating (2.31) gives $\|u^j\|_\perp \leq \gamma^j \|u^0\|_\perp$ and

$$\sum_{j=0}^{\infty} |\xi_3^{j+1} - \xi_3^j| \leq \frac{C\|u^0\|_\perp}{1-\gamma} < \infty.$$

Hence $u^j \to 0$, and ξ_3^j converges. Since $\xi^j \in U$ for every j, the definition of U gives $\xi_3^j > 1 - \varepsilon$. Its limit is therefore at least $1 - \varepsilon > 0$. □

2.4 Completion of the main proof

We choose an invariant neighborhood on which every four-step block satisfies the line-search tests and stays inside the ball of radius ten. We then define the coercive objective and prove by induction that its BFGS iterates follow the scaled blocks.

Proof of Theorem 1.1. *Step 1: The initial set.* Use the objective $\widetilde{f}$ fixed in Section 2.3 with the coefficients (2.28). For $\xi \in \mathcal{D}_4$, let $(\widehat{x}^i, \widehat{H}^i)$ be the state represented by $\mathcal{T}^i(\xi)$ and set $\widehat{g}^i = \nabla \widetilde{f}(\widehat{x}^i)$ for $0 \leq i \leq 4$. For $0 \leq i < 4$, write $\widehat{s}^i = \widehat{x}^{i+1} - \widehat{x}^i = -\widehat{H}^i \widehat{g}^i$. These quantities depend on ξ and are continuous on $\mathcal{D}_4$ by Lemma 2.11. The domain (2.25) ensures smooth evaluation points, positive definite matrices and $(\widehat{s}^i)^T(\widehat{g}^{i+1} - \widehat{g}^i) > 0$. In particular, $\widehat{g}^i \neq 0$ and $(\widehat{g}^i)^T \widehat{s}^i < 0$ for $0 \leq i < 4$.

At $\xi = \omega_*$, Lemmas 2.5 and 2.10 give $\|\widehat{x}^i\| < 10$ for $0 \leq i \leq 4$, and Proposition 2.9 gives strict Armijo and Wolfe inequalities at $(c_1, c_2) = (0.03, 0.6)$. These finitely many strict inequalities persist on an open neighborhood of ω_* by continuity. Since Lemma 2.12 supplies arbitrarily small invariant neighborhoods, choose one such $U \subset \mathcal{D}_4$ with $\mathcal{F}(U) \subset U$ so that, for every $\xi \in U$,

$$\begin{aligned} \|\widehat{x}^i\| &< 10 && (0 \leq i \leq 4), \\ \widetilde{f}(\widehat{x}^{i+1}) &< \widetilde{f}(\widehat{x}^i) + 0.03(\widehat{g}^i)^T \widehat{s}^i && (0 \leq i < 4), \\ (\widehat{g}^{i+1})^T \widehat{s}^i &> 0.6(\widehat{g}^i)^T \widehat{s}^i && (0 \leq i < 4). \end{aligned} \tag{2.32}$$

Because $(\widehat{g}^i)^T \widehat{s}^i < 0$, these tests also hold for every parameter pair in (1.4). Define the initial set in the theorem by

$$\mathcal{U} = \{(x, H) \in \mathbb{R}^2 \times \mathbb{S}^2_{++} : (x_1, x_2, H_{11}, H_{12}, H_{22}) \in U\}. \tag{2.33}$$

The state-coordinate map is a homeomorphism onto its image, so $\mathcal{U}$ is open and contains the cycle state (x^0, H^0) constructed in Section 2.1.

Step 2: The coercive objective. Define

$$f(x) = \widetilde{f}(x) + \max\{0, \|x\|^2 - 100\}. \tag{2.34}$$

By Lemma 2.10, $\widetilde{f}$ has a global Lipschitz constant L. Since $\widetilde{f}(0) = 0$, we have $\widetilde{f}(x) \geq -L\|x\|$, and hence

$$f(x) \geq \|x\|^2 - L\|x\| - 100 \longrightarrow +\infty \qquad \text{as } \|x\| \to \infty.$$

Thus f is coercive. It is locally Lipschitz and semialgebraic, and agrees with $\widetilde{f}$ on the ball of radius ten. In particular, $\partial_{\mathrm{C}} f(0) = \partial_{\mathrm{C}} \widetilde{f}(0)$, so Lemma 2.10 gives $0 \notin \partial_{\mathrm{C}} f(0)$.

Step 3: Induction and convergence. Fix a parameter pair in (1.4) and an initial state in $\mathcal{U}$ from (2.33), represented by $\omega^0 \in U$. Define $\xi^0 = \omega^0$ and $\xi^{j+1} = \mathcal{F}(\xi^j)$. Here $\mathcal{F}$ is given by (2.26). Since $\mathcal{F}(U) \subset U$, this sequence is well defined, with $\xi^j \in U$ for every $j \geq 0$. We prove by induction that the actual algorithm applied to f is defined at every iteration, accepts every unit trial, and has states

$$\omega^{4j+i} = r^j \mathcal{T}^i(\xi^j) \qquad (j \geq 0,\ 0 \leq i \leq 4). \tag{2.35}$$

The initial identity is $\omega^0 = \xi^0$. Suppose the algorithm has reached $\omega^{4j} = r^j \xi^j$ with all preceding unit trials accepted. Keep j fixed and, for $0 \leq i \leq 4$, let $(\widehat{x}^i, \widehat{H}^i)$ be the state represented by $\mathcal{T}^i(\xi^j)$. We show that the state identity propagates from phase i to phase $i+1$. Suppose the state at $k = 4j + i$, $0 \leq i < 4$, is $\omega^k = r^j \mathcal{T}^i(\xi^j)$, so $(x^k, H^k) = (r^j \widehat{x}^i, r^j \widehat{H}^i)$. At $(\widehat{x}^i, \widehat{H}^i)$, the tests in (2.32) accept the unit trial for $\widetilde{f}$, and (2.24) gives the next state $(\widehat{x}^{i+1}, \widehat{H}^{i+1})$. Lemma 2.2, with $\tau = r^j$, therefore gives an accepted unit step from $(r^j \widehat{x}^i, r^j \widehat{H}^i)$ to $(r^j \widehat{x}^{i+1}, r^j \widehat{H}^{i+1})$.

Both evaluation points are nonzero by (2.25), and their norms are below ten by (2.32) and $0 < r^j \leq 1$. Thus (2.34) makes f and $\widetilde{f}$ agree locally at the current point and the unit trial. The objective values and gradients used by Algorithm 1 coincide, so it accepts the same trial and produces $\omega^{k+1} = r^j \mathcal{T}^{i+1}(\xi^j)$. By (2.24)–(2.25), each unscaled phase has positive curvature and hence a nonzero gradient. These properties persist under positive scaling, excluding termination or breakdown.

Applying this implication successively for $i = 0, 1, 2, 3$ shows that the iterations $4j, \ldots, 4j+3$ are well defined and accept their unit trials, and that (2.35) holds throughout this block. By (2.26), the state after these four iterations is

$$\omega^{4(j+1)} = r^j \mathcal{T}^4(\xi^j) = r^{j+1} \mathcal{F}(\xi^j) = r^{j+1} \xi^{j+1},$$

which closes the induction on j. Finally, combining (2.35) with the position bounds in (2.32) yields $\|x^{4j+i}\| < 10 r^j$ for $j \geq 0$ and $0 \leq i < 4$. Since $0 < r < 1$, the iterates converge to the origin. □

3 Concluding remarks

We give a coercive, locally Lipschitz, semialgebraic objective on $\mathbb{R}^2$ for which BFGS converges to a non-Clarke-stationary point from a nonempty open set of initial states. This disproves the almost-sure stationarity assertion of Lewis and Overton [LO13, Challenge 7.1(2)].

An important remaining question is whether Algorithm 1 has only Clarke-stationary cluster points almost surely for convex, locally Lipschitz, semialgebraic objectives with bounded sublevel sets. For nonconvex objectives with the same regularity and boundedness assumptions, which alternative line-search rules or safeguards on the inverse-Hessian estimates can recover this guarantee?

AI use statement

The author used Codex to assist with the mathematical construction, numerical experiments, verification code, and manuscript preparation. GPT-6 Pro was consulted on developing and simplifying the example and proofs. AI-assisted material was refined through repeated discussion, mathematical and computational checks, and editing. The author independently verified all mathematical arguments and takes full responsibility for the content of the paper.

A Rational certificates for the finite inequalities

This appendix certifies the interpolation bounds in Proposition 2.8 and the transverse contraction estimate used in Lemma 2.12. Verification details and reproducible computations are provided in the accompanying GitHub repository linked in the introduction.

Certified arithmetic. We use standard interval arithmetic with outward rounding; see [MKC09, Sections 2.3–2.4 and 3.2]. Every terminating decimal denotes an exact rational number. An interval $[a, b]$ represents all real numbers between its rational endpoints. Addition, multiplication and negation are evaluated by

$$[a, b] + [c, d] = [a + c, b + d], \qquad [a, b][c, d] = [\min(ac, ad, bc, bd), \max(ac, ad, bc, bd)],$$

with $-[a, b] = [-b, -a]$. If $0 \notin [a, b]$, its reciprocal is $[a, b]^{-1} = [1/b, 1/a]$; division by an interval containing zero is rejected. With $Q = 2^{256}$, round every input and computed interval outward by

$$[a, b] \longmapsto \left[\frac{\lfloor Qa \rfloor}{Q}, \frac{\lceil Qb \rceil}{Q}\right]. \tag{A.1}$$

Printed decimal endpoints are again rounded outward. For interval matrices, membership and arithmetic are understood entrywise, with matrix products formed by interval sums and products.

A.1 Interpolation bounds

Consider the data constructed in Section 2.1. Substituting (2.4) into (2.8) and simplifying gives the expression used in these calculations:

$$x^0 = \left(-\frac{9(8978b^2 - 6527b - 2304)}{5440b}, \frac{9(1764b^2 - 2671b + 1488)}{2720}\right)^T, \tag{A.2}$$

where $b \in [0.9178, 0.9179]$ by Lemma 2.3. Use the positions from (A.2) and (2.8) and the gradients from (2.6). With the signs of z_1^i established in Proposition 2.7, evaluate $t^i = z_2^i/|z_1^i|$ and the data in (2.17). Their enclosing intervals are given in Table 1.

Table 1: Certified interpolation data at the four cycle positions.

i	t^i	η^i	d^i
0	[1.4003, 1.4070]	[1.0665, 1.0739]	[0.0499, 0.0501]
1	[4.7424, 4.8319]	[4.7424, 4.8319]	[1.0000, 1.0000]
2	[0.2714, 0.2900]	[1.0334, 1.0730]	[0.0271, 0.0330]
3	[3.2562, 3.4625]	[4.2658, 4.5914]	[0.9874, 0.9914]

For each list in (2.18), number the three consecutive gaps $j = 0, 1, 2$ from left to right. For adjacent entries $(a, A), (c, B)$, evaluate $\ell = c - a$ and $(B - A)/\ell$ by the interval rules above. The resulting enclosures, used in the proof of Proposition 2.8, are given in Table 2.

Table 2: Certified distances and secant slopes between consecutive interpolation points.

Sign	j	ℓ	$(B-A)/\ell$
+	0	[2.4003, 2.4070]	[0.8585, 0.8640]
+	1	[3.3355, 3.4316]	[1.0690, 1.1289]
+	2	[1.1681, 1.2576]	[0.9288, 1.0766]
-	0	[1.2714, 1.2900]	[1.5763, 1.6304]
-	1	[2.9662, 3.1910]	[1.0006, 1.1995]
-	2	[2.5375, 2.7438]	[0.5133, 0.6834]

A.2 Contraction toward the fixed ray

Lemma 2.3 gives a unique root b of P in $(0.9178, 0.9179)$. For the transverse contraction estimate needed in the proof of Lemma 2.12, we refine its enclosure to the rational interval

$$b \in Y = [0.91781039164, 0.91781039165]. \tag{A.3}$$

Using exact fractions, P takes a negative value at the left endpoint of Y and a positive value at the right endpoint. The intermediate value theorem gives a root in the interior of Y; since $Y \subset (0.9178, 0.9179)$, uniqueness in Lemma 2.3 identifies this root with b.

Set

$$S = \begin{pmatrix} 464 & 1077 & 4501 & -1708 \\ 1366 & 2793 & 10000 & 552 \\ -448 & -1058 & -4343 & 1648 \\ 16 & 293 & 1787 & -2075 \end{pmatrix}. \tag{A.4}$$

The following estimate supplies the contraction used in Lemma 2.12.

Proposition A.1. *For the matrix A in* (2.30) *and the matrix S in* (A.4)*, S is invertible and $\|SAS^{-1}\|_\infty \le 0.78$.*

Proof. We first compute A by differentiating at the exact cycle, then enclose the resulting rational expression by interval arithmetic over Y.

Step 1: Differentiating one BFGS step. Recall the cycle data g^i, H^i, x^i from (2.6)–(2.8), the steps s^i and gradient differences y^i from (2.9), and the Hessians K^i from (2.21), with the coefficients (2.28). Set $K^4 = K^0/r$. Lemma 2.10 and the Hessian scaling in Proposition 2.1 give $K^i = \nabla^2 \widetilde{f}(x^i)$ for $0 \le i \le 4$.

In the state coordinates of (2.24), write $\omega^i = (x_1^i, x_2^i, H_{11}^i, H_{12}^i, H_{22}^i)$ for $0 \le i \le 4$. Thus $\omega^0 = \omega_*$ and $\mathcal{T}(\omega^i) = \omega^{i+1}$ for $0 \le i < 4$ by Lemmas 2.5 and 2.10. For a state variation $\delta\omega \in \mathbb{R}^5$, set

$$\delta x = \big((\delta\omega)_1, (\delta\omega)_2\big)^T, \qquad \delta H = \begin{pmatrix} (\delta\omega)_3 & (\delta\omega)_4 \\ (\delta\omega)_4 & (\delta\omega)_5 \end{pmatrix}.$$

Fix $0 \le i < 4$. The symbol δ denotes differentiation at $\varepsilon = 0$ along $\omega = \omega^i + \varepsilon\delta\omega$. Write $\widehat{\omega} = \mathcal{T}(\omega)$ for the updated state, whose point and matrix are $\widehat{x} = x + s$ and $\widehat{H} = \mathcal{B}(H, s, y)$, and set $g = \nabla \widetilde{f}(x)$. Since $x^i + s^i = x^{i+1}$, differentiation gives

$$\begin{aligned} &\delta g = K^i \delta x, \qquad \delta s = -((\delta H) g^i + H^i \delta g), \\ &\delta\widehat{x} = \delta x + \delta s, \quad \delta y = K^{i+1}\delta\widehat{x} - \delta g. \end{aligned} \tag{A.5}$$

All quantities in the following formulas are evaluated at this cycle state, so $H = H^i$, $s = s^i$ and $y = y^i$. Introduce the scalars c, d, the vector t and the matrices N, V by

$$c = s^T y, \qquad t = Hy, \qquad d = y^T H y, \qquad N = ts^T + st^T, \qquad V = ss^T.$$

Expansion of (1.3) gives

$$\widehat{H} = H - \frac{N}{c} + \frac{c+d}{c^2} V. \tag{A.6}$$

The product and quotient rules give

$$\begin{aligned}
\delta c &= (\delta s)^T y + s^T \delta y, \qquad \delta t = (\delta H) y + H \delta y, \\
\delta d &= 2(\delta y)^T t + y^T (\delta H) y, \qquad \delta V = (\delta s) s^T + s (\delta s)^T, \\
\delta N &= (\delta t) s^T + t (\delta s)^T + (\delta s) t^T + s (\delta t)^T, \\
\delta \widehat{H} &= \delta H - \frac{\delta N}{c} + \frac{N \delta c}{c^2} + \frac{c+d}{c^2} \delta V + \left(\frac{\delta c + \delta d}{c^2} - \frac{2(c+d)\delta c}{c^3} \right) V.
\end{aligned} \tag{A.7}$$

Step 2: The four-step derivative. Set $D^i = D\mathcal{T}(\omega^i) \in \mathbb{R}^{5 \times 5}$. Equations (A.5) and (A.7) give

$$\delta \widehat{\omega} = D^i \delta \omega = \big((\delta \widehat{x})_1, (\delta \widehat{x})_2, (\delta \widehat{H})_{11}, (\delta \widehat{H})_{12}, (\delta \widehat{H})_{22} \big)^T. \tag{A.8}$$

Its jth column is obtained by taking $\delta\omega$ to be the jth standard basis vector of $\mathbb{R}^5$. Since $\omega^0 = \omega_*$ and $\mathcal{F} = r^{-1}\mathcal{T}^4$ by (2.26), the chain rule gives

$$D_{\mathrm{tot}} := D^3 D^2 D^1 D^0 = D\mathcal{T}^4(\omega_*), \qquad D\mathcal{F}(\omega_*) = r^{-1} D_{\mathrm{tot}}. \tag{A.9}$$

Recall $W = \{v \in \mathbb{R}^5 : v_3 = 0\}$, $R(v) = (v_1, v_2, v_4, v_5)$ and the map L_{ω_*} used in (2.30). By (A.9),

$$L_{\omega_*}(v) = r^{-1} \big(D_{\mathrm{tot}} v - (D_{\mathrm{tot}} v)_3 \omega_* \big) \qquad (v \in W).$$

The subtraction makes the third coordinate zero because $(\omega_*)_3 = 1$. Equation (2.30) states that $R(L_{\omega_*}(v)) = AR(v)$ for $v \in W$. Writing $(\kappa_1, \kappa_2, \kappa_3, \kappa_4) = (1, 2, 4, 5)$ for the coordinates retained by R, we obtain

$$A_{ij} = \frac{(D_{\mathrm{tot}})_{\kappa_i \kappa_j} - (\omega_*)_{\kappa_i} (D_{\mathrm{tot}})_{3 \kappa_j}}{r} \qquad (1 \le i, j \le 4). \tag{A.10}$$

Step 3: Entry bounds and row sums. The formulas above express every entry of A rationally in b. To enclose them, replace b by an independent variable $\lambda \in Y$ in the formulas defining $g^i, H^i, x^i, K^i, s^i, y^i$, using (A.2) for x^0. An argument λ denotes this substitution, for example $x^i(\lambda)$ and $s^i(\lambda)$. Evaluate (A.5), (A.7) and (A.8) with these data to obtain $D^i(\lambda)$. Form $D_{\mathrm{tot}}(\lambda) = D^3(\lambda) D^2(\lambda) D^1(\lambda) D^0(\lambda)$ and set

$$\omega^0(\lambda) = \big(x^0_1(\lambda), x^0_2(\lambda), 1, 1/2, \lambda/4 + 1/8 \big).$$

In the right-hand side of (A.10), replace D_{tot} and ω_* by $D_{\mathrm{tot}}(\lambda)$ and $\omega^0(\lambda)$ to define $\widehat{A}(\lambda)$. Since $D_{\mathrm{tot}}(b) = D_{\mathrm{tot}}$ and $\omega^0(b) = \omega_*$, we have $\widehat{A}(b) = A$. The derivative interpretation established above applies at $\lambda = b$.

Apart from fixed nonzero constants, the denominator factors in (2.4), (2.8) and (2.21) are λ, $2\lambda - 1$ and $\|x^i(\lambda)\|^2$. Formula (A.7) also divides by powers of $s^i(\lambda)^T y^i(\lambda)$. For every $\lambda \in Y$, we have $\lambda > 0.9$ and $2\lambda - 1 > 0.8$, and interval evaluation gives

$$\|x^i(\lambda)\|^2 > 0.7, \qquad s^i(\lambda)^T y^i(\lambda) > 0.4 \qquad (0 \le i < 4).$$

The remaining factors $r = 64/81$ and $1 - r = 17/81$ are fixed and positive.

Since $\det S = -22\,230\,591\,050 \neq 0$, S is invertible. Evaluate $S\widehat{A}(\lambda)S^{-1}$ over Y by interval arithmetic and denote the resulting 4×4 interval matrix by $\mathbf{M}$. Every denominator interval excludes zero, so the inclusion principle [Rum10, equation (5.17)] ensures that each interval $\mathbf{M}_{ij}$ contains $(S\widehat{A}(\lambda)S^{-1})_{ij}$ for every $\lambda \in Y$.

For an interval $[a, c]$, set $\operatorname{mag}([a, c]) = \max\{|a|, |c|\}$. Rounding $\operatorname{mag}(\mathbf{M}_{ij})$ upward to three decimal places gives the nonnegative matrix Γ:

$$|(S\widehat{A}(\lambda)S^{-1})_{ij}| \leq \operatorname{mag}(\mathbf{M}_{ij}) \leq \Gamma_{ij} \qquad (\lambda \in Y),$$
$$\Gamma = \begin{pmatrix} 0.133 & 0.164 & 0.049 & 0.011 \\ 0.240 & 0.091 & 0.094 & 0.015 \\ 0.042 & 0.006 & 0.521 & 0.207 \\ 0.024 & 0.002 & 0.211 & 0.478 \end{pmatrix}. \tag{A.11}$$

Summing each row of this displayed matrix gives

$$\left(\sum_{j=1}^{4} |(S\widehat{A}(\lambda)S^{-1})_{ij}| \right)_{i=1}^{4} \leq (0.357,\, 0.440,\, 0.776,\, 0.715) \qquad (\lambda \in Y), \tag{A.12}$$

where the inequality is componentwise. Since $\widehat{A}(b) = A$, the maximum absolute row-sum formula for the induced norm gives $\|SAS^{-1}\|_\infty \leq 0.776 < 0.78$. □

References

[AO21] A. Asl and M. L. Overton. Analysis of limited-memory BFGS on a class of nonsmooth convex functions. *IMA Journal of Numerical Analysis*, 41(1):1–27, 2021.

[BCR98] J. Bochnak, M. Coste, and M.-F. Roy. *Real Algebraic Geometry*, volume 36 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge*. Springer-Verlag, Berlin, 1998.

[Bro70] C. G. Broyden. The convergence of a class of double-rank minimization algorithms. 2. The new algorithm. *Journal of the Institute of Mathematics and its Applications*, 6(3):222–231, 1970.

[Cla75] F. H. Clarke. Generalized gradients and applications. *Transactions of the American Mathematical Society*, 205:247–262, 1975.

[Cla90] F. H. Clarke. *Optimization and Nonsmooth Analysis*, volume 5 of *Classics in Applied Mathematics*. Society for Industrial and Applied Mathematics, Philadelphia, PA, 1990.

[CMO17] F. E. Curtis, T. Mitchell, and M. L. Overton. A BFGS-SQP method for nonsmooth, nonconvex, constrained optimization and its evaluation using relative minimization profiles. *Optimization Methods and Software*, 32(1):148–181, 2017.

[Dai02] Y.-H. Dai. Convergence properties of the BFGS algoritm. *SIAM Journal on Optimization*, 13(3):693–701, 2002.

[Dai13] Y.-H. Dai. A perfect example for the BFGS method. *Mathematical Programming*, 138(1–2):501–530, 2013.

[DM77] J. E. Dennis, Jr. and J. J. Moré. Quasi-Newton methods, motivation and theory. *SIAM Review*, 19(1):46–89, 1977.

[DYZ26] L. Ding, J. Yang, and B. Zhou. On the complexity of BFGS method for smooth convex optimization. Preprint, 2026. arXiv:2608.16009.

[Fle70] R. Fletcher. A new approach to variable metric algorithms. *The Computer Journal*, 13(3):317–322, 1970.

[Geb26] B. Gebken. Technical results on the convergence of quasi-Newton methods for nonsmooth optimization. *Journal of Optimization Theory and Applications*, 209, 2026. Article 57.

[GL18] J. Guo and A. S. Lewis. Nonsmooth variants of Powell's BFGS convergence theorem. *SIAM Journal on Optimization*, 28(2):1301–1311, 2018.

[Gol70] D. Goldfarb. A family of variable-metric methods derived by variational means. *Mathematics of Computation*, 24(109):23–26, 1970.

[JJM24] Q. Jin, R. Jiang, and A. Mokhtari. Non-asymptotic global convergence analysis of BFGS with the Armijo–Wolfe line search. In *Advances in Neural Information Processing Systems*, volume 37, pages 16810–16851. Curran Associates, Inc., 2024.

[Li26] J. Li. Nonminimizing attractors of stochastic subgradient methods on semialgebraic functions. Preprint, 2026. arXiv:2610.11298.

[LO13] A. S. Lewis and M. L. Overton. Nonsmooth optimization via quasi-Newton methods. *Mathematical Programming*, 141(1–2):135–163, 2013.

[LZ15] A. S. Lewis and S. Zhang. Nonsmoothness and a variable metric method. *Journal of Optimization Theory and Applications*, 165(1):151–171, 2015.

[Man01] C. Manni. On shape preserving C^2 Hermite interpolation. *BIT Numerical Mathematics*, 41(1):127–148, 2001.

[Mas04] W. F. Mascarenhas. The BFGS method with exact line searches fails for non-convex objective functions. *Mathematical Programming*, 99(1):49–61, 2004.

[Mas14] W. F. Mascarenhas. The divergence of the BFGS and Gauss Newton methods. *Mathematical Programming*, 147(1–2):253–276, 2014.

[MKC09] R. E. Moore, R. B. Kearfott, and M. J. Cloud. *Introduction to Interval Analysis*. Society for Industrial and Applied Mathematics, Philadelphia, PA, 2009.

[Pow76] M. J. D. Powell. Some global convergence properties of a variable metric algorithm for minimization without exact line searches. In R. W. Cottle and C. E. Lemke, editors, *Nonlinear Programming*, volume 9 of *SIAM–AMS Proceedings*, pages 53–72. American Mathematical Society, Providence, RI, 1976.

[Rum10] S. M. Rump. Verification methods: Rigorous results using floating-point arithmetic. *Acta Numerica*, 19:287–449, 2010.

[Sha70] D. F. Shanno. Conditioning of quasi-Newton methods for function minimization. *Mathematics of Computation*, 24(111):647–656, 1970.

[XW17] Y. Xie and A. Wächter. On the convergence of BFGS on a class of piecewise linear non-smooth functions. Preprint, 2017. arXiv:1712.08571.