



Eye Injury / Complaint

Assessment

Pediatric Pearls:

- Use pediatric dosing of medications or electrical therapy for a pediatric patient < 37 kg and as defined by HandTevy.
- Pediatric hypotension is defined as SBP < 70 + (age in years x 2) mmHg

Signs & Symptoms:

- Pain, swelling, blood
- Deformity, contusion
- Visual deficit / loss of vision
- Leaky aqueous / vitreous humor
- Upwardly fixed eye
- Shooting or streaking light
- Visual contaminants
- Rust ring
- Lacrimation

Differential:

- Trauma
- Infection – Viral URI or Sinusitis
- Allergic rhinitis
- Lesions – Polyps, Ulcers
- Hypertension

Clinical Management Options

- | | | | | | | |
|---|---|---|---|---|---|---|
| P | P | P | P | P | P | <ul style="list-style-type: none"> • Evaluate pupils • Complete neurological exam • Screen for unrecognized chemical and/or agent exposure • Cover both eyes • If out of socket, then cover with sterile water or isotonic crystalloid soaked gauze • If impaled object, then stabilize the object then cover both eyes • If chemical exposure or burn, then irrigate with copious amounts of sterile water or isotonic crystalloid <ul style="list-style-type: none"> ◦ Perform an initial and repeat respiratory assessments. • Vascular access • May use Lidocaine in 1 L Isotonic Crystalloid for irrigation |
| L | L | L | L | L | L | |
| 1 | 2 | 3 | 4 | 5 | 6 | |
| | | | | | | |
| | | | | | | |

Consult Online Medical Control As Needed

Pearls:

- Refer to drug formulary charts for all medication dosing for both adults and pediatric patients.
- Normal visual acuity can present even with severe injury.
- Remove contact lens when possible. If adherent to globe, then do not force. Irrigation may assist with removal.
- Any chemical or thermal burns to the face/eyes should raise concern for respiratory insult.
- Orbital fracture raises concern for globe or nerve injury, or compartment syndrome. This requires repeat assessments.
- Always cover both eyes to prevent further insult.
- Use shield not pads for physical trauma to the eye. Pads are acceptable for the uninjured eye.
- DO NOT remove impaled objects
- Suspected globe rupture or compartment syndrome requires emergent evaluation.