



Hypotension Non-Traumatic

Assessment

Pediatric Pearls:

- Use pediatric dosing of medications or electrical therapy for a pediatric patient < 37 kg and as defined by HandTevy.
- Pediatric hypotension is defined as SBP < 70 + (age in years x 2) mmHg

Signs & Symptoms:

- Restlessness, confusion, weakness
- Syncope
- Tachycardia
- Diaphoresis
- Pale, cool, clammy skin
- Delayed capillary refill
- Coffee-ground emesis
- Tarry stools

Differential:

- Infection/Sepsis
- Dehydration
- Vomiting
- Diarrhea
- Congenital heart disease
- Medication or Toxin
- Anaphylaxis
- Cardiac Failure (myocarditis)

Clinical Management Options

P	P	P	P	P	P	• Oxygen
L	L	L	L	L	L	• Supine position, keep patient warm
1	2	3	4	5	6	• Oral rehydration, if tolerated
						• Vascular access • These fluid boluses are for volume depletion – NOT for active bleeding. • Pediatric: Isotonic Crystalloid bolus 20 ml/kg may repeat 10ml/kg bolus x 2 PRN • Adult non-cardiac: Isotonic Crystalloid 500-1000 ml bolus, may repeat up to 2 liters • Adult Cardiac: Isotonic Crystalloid 250-500 ml bolus, may repeat up to 1 liter • Tranexamic acid (TXA) for hypotension due to hemorrhage
						• Adult: Norepinephrine (Levophed) infusion, titrated to MAP $\geq$ 65 following fluid resuscitation or Epinephrine • Push dose Epinephrine or Norepinephrine • Pediatric: Epinephrine infusion titrated to effect • Blood product administration procedure, including Calcium Chloride

Consult Online Medical Control As Needed

Pearls:

- Refer to drug formulary charts for all medication dosing for both adults and pediatric patients.
- Adult hypotension can be defined as a MAP < 65 and signs or symptoms of hypoperfusion – altered mental status, increased respirations, tachycardia, poor pulses, skin changes.
- Consider all possible causes of shock and treat per appropriate COG.
- Patients should always have adequate intravascular fluid load before using vasopressors.
- Isotonic Crystalloid should be avoided in patients in whom hemorrhage is suspected.

$$MAP \text{ Calculation} = \frac{(2 \times \text{Diastolic}) + \text{Systolic}}{3}$$