



# Nausea / Vomiting

## Assessment

### Pediatric Pearls:

- Use pediatric dosing of medications or electrical therapy for a pediatric patient < 37 kg and as defined by HandTevy.
- Pediatric hypotension is defined as SBP < 70 + (age in years x 2) mmHg

### Signs & Symptoms:

- Fever
- Pain
- Constipation
- Diarrhea
- Anorexia
- Hematemesis

### Differential:

- CNS (Increased ICP, headache, stroke, CNS lesions, Trauma or hemorrhage)
- Vestibular
- AMI
- Drugs (NSAIDS, antibiotics, narcotics, chemotherapy)
- GI or GU disorders
- Uremia
- Gynecologic disease (Ovarian Cyst / PID)
- Infections (pneumonia, influenza)
- Electrolyte abnormalities
- Food or Toxin induced
- Pregnancy

## Clinical Management Options

|   |   |   |   |   |   |  |                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|---|---|---|---|---|---|--|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| P | P | P | P | P | P |  | <ul style="list-style-type: none"> <li>• <a href="#">Oxygen</a></li> <li>• <a href="#">BGL</a> assessment</li> </ul>                                                                                                                                                                                                                                                                                                                          |
| L | L | L | L | L | L |  |                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| 1 | 2 | 3 | 4 | 5 | 6 |  | <ul style="list-style-type: none"> <li>• Inhaled isopropyl alcohol wipe to help control nausea</li> <li>• Adult &amp; Pediatric: <a href="#">Ondansetron</a></li> </ul>                                                                                                                                                                                                                                                                       |
|   |   |   |   |   |   |  | <ul style="list-style-type: none"> <li>• <a href="#">Vascular access</a></li> <li>• Adult: <a href="#">Diphenhydramine</a> if patient allergic to Ondansetron <b>and</b> Droperidol</li> <li>• IV fluid with <a href="#">Isotonic Crystalloid</a> as needed for dehydration</li> <li>• Adult: <a href="#">Tranexamic acid (TXA)</a> with confirmed upper or lower GI bleeding AND hypotension if &lt; 3 hours since bleeding onset</li> </ul> |
|   |   |   |   |   |   |  | <ul style="list-style-type: none"> <li>• ECG monitoring and interpretation</li> <li>• Adult: <a href="#">Droperidol</a></li> </ul>                                                                                                                                                                                                                                                                                                            |

## Consult Online Medical Control As Needed

### Pearls:

- Refer to drug formulary charts for all medication dosing for both adults and pediatric patients.
- If patient is conscious, place alcohol wipe under patient's nose and allow them to breathe it in to help control nausea.
- Assess number of times of emesis
- Appearance of emesis: bloody, coffee ground, bilious – green bile – solids and liquid or just liquid
- Heart rate: One of the first clinical signs of dehydration, almost always increased heart rate, tachycardia increases as dehydration becomes more severe, very unlikely to be significantly dehydrated if heart rate is close to normal.
- Droperidol may be a good option for cyclic vomiting or Cannabinoid Hyperemesis Syndrome.
- Due precaution of long QT syndrome for both Ondansetron and Droperidol, providers should choose one therapy only, **Contact OLMC if necessary**