



Overdose

Assessment

Pediatric Pearls:

- Use pediatric dosing of medications or electrical therapy for a pediatric patient < 37 kg and as defined by HandTevy.

Signs & Symptoms:

- Salivation
- Lacrimation
- Urination
- Defecation
- GI distress
- Emesis
- Bronchospasm
- Bronchorrhea
- Bradycardia
- Seizure

Differential:

- Stroke
- MI
- Asthma / COPD
- Other chemical agent / weapon
- Biologic agent / weapon
- Overdose
- Food borne illness
- Airborne irritant (hydrogen sulfide, chlorine, etc.)

Clinical Management Options

P	P	P	P	P	P	• Scene safety, PPE
L	L	L	L	L	L	• Basic Airway Maneuvers
1	2	3	4	5	6	• Narcotic OD: Naloxone IN, only if respirations are depressed • Provide patient and/or acquaintance a Naloxone Rescue Kit if available
						• ECG Application • EtCO₂ application and monitoring
						• Vascular access • Narcotic OD: Naloxone if respirations are depressed
						• Monitoring and interpretation of ECG & EtCO₂ • Advance airway maneuvers • Adult: Midazolam • Calcium Channel / Beta Blocker OD: Calcium Chloride, Epinephrine Infusion & Glucagon (Beta blocker OD only) • Tricyclic or Phenobarbital OD: Sodium Bicarbonate • Transcutaneous Cardiac Pacing PRN

Consult Online Medical Control As Needed

Pearls:

- Contact CHP early in cases of opiate use or withdrawal.
- Do not rely on patient history of ingestion, especially in suicide attempts.
- Tricyclic: 4 major areas of toxicity: seizures, dysrhythmias, hypotension, decreased mental status or coma, rapid progression from AMS to death.
- Depressants: decreased HR, BP, body temperature, and respirations; non-specific pupils.
- Stimulants: increased HR, BP, body temperature; dilated pupils, seizures.
- Anticholinergic: increased HR, body temperature; dilated pupils, mental status changes.
- Cardiac meds: dysrhythmias and mental status changes.
- Solvents: nausea, vomiting, and mental status changes.
- Cholinergics / Insecticides: increased or decreased HR, increased HR, nausea / vomiting, diarrhea, pinpoint pupils
- Consider contacting Poison Control for Guidance on a recorded line – 1-800-222-1222
- Decon or Haz-Mat patients should be performed by trained personnel prior to initial patient contact or transport.
- Novel opioids, such as fentanyl and carfentanyl, may require very high doses of naloxone (10-20mg)