

Problem

Consider a matrix problem (s.p.d.) of the form

$$A\mathbf{u} = \mathbf{f}, \quad A \in \mathbb{R}^{n \times n}$$

Suppose we have a multilevel iteration process \mathcal{M}_{MG}

$$I - \mathcal{M}_{\text{MG}}A = (I - M^\top A)^{\nu_{\text{pre}}} \left(I - P(P^\top AP)^{-1}P^\top A \right) (I - MA)^{\nu_{\text{post}}}$$

so that

$$\mathbf{e} \leftarrow (I - \mathcal{M}_{\text{MG}}A)\mathbf{e}$$

The iteration converges for any \mathbf{f} and \mathbf{u}_0 iff $\rho(I - \mathcal{M}_{\text{MG}}A) < 1$.

we are seeking a method that yields **a bound on the error reduction in each iteration that is independent of n** .

Notation

Fine grid $\Omega = \{1, \dots, n\} = C \cup F$ and coarse grid $\Omega_c = C$. Interpolation / restriction:

$$P : \Omega_c \rightarrow \Omega \quad \text{and} \quad R : \Omega \rightarrow \Omega_c$$

A is s.p.d., $D = \text{diag}(A)$ — defining an inner product:

$$\begin{aligned} \langle \mathbf{u}, \mathbf{v} \rangle_A &= \langle A\mathbf{u}, \mathbf{v} \rangle \\ \langle \mathbf{u}, \mathbf{v} \rangle_D &= \langle D\mathbf{u}, \mathbf{v} \rangle \\ \langle \mathbf{u}, \mathbf{v} \rangle_{AD^{-1}A} &= \langle D^{-1}A\mathbf{u}, A\mathbf{v} \rangle \end{aligned}$$

Define S as post-relaxation

(and \mathcal{S} as the affine version $\mathbf{u} \leftarrow \mathcal{S}(\mathbf{u}, \mathbf{f})$):

$$\mathbf{u} \leftarrow S\mathbf{u} + (I - S)A^{-1}\mathbf{f} \quad \text{or} \quad \mathbf{e} \leftarrow S\mathbf{e}$$

Algorithm

$$\begin{aligned} \mathbf{u} &\leftarrow \hat{S}(\mathbf{u}, \mathbf{f}) \\ \mathbf{r}_c &\leftarrow R\mathbf{r} \\ \mathbf{e}_c &\leftarrow A_c^{-1}\mathbf{r}_c \\ \hat{\mathbf{e}} &\leftarrow P\mathbf{e}_c \\ \mathbf{u} &\leftarrow \mathbf{u} + \hat{\mathbf{e}} \\ \mathbf{u} &\leftarrow \mathcal{S}(\mathbf{u}, \mathbf{f}) \end{aligned}$$

In general, consider relaxation as

$$S = I - MA$$

Assumptions:

- M is norm convergent (in A): $\|S\|_A < 1$
- P is full rank


- A is s.p.d.

Let the coarse grid correction step be

$$T = I - P(P^\top AP)^{-1}P^\top A$$

T is an A -orthogonal projection onto the range of P : After coarse grid correction, the error is minimized in the energy norm over $\text{range}(P)$

$$\begin{aligned} &\text{if } w \quad \text{s.t.} \quad Aw \in \mathcal{N}(P^\top A) \\ &\left(I - P(P^\top AP)^{-1}P^\top A \right) w = w \\ &\| \cdot \| \geq 1 \end{aligned}$$



Focus on $V(0, 1)$

The A -adjoint of ST is

$$TS^+ \quad S^+ = I - M^\top A$$

The symmetric $V(1, 1)$ cycle is

$$\begin{aligned} (I - MA) \left(I - P(P^\top AP)^{-1}P^\top A \right) (I - M^\top A) &= STS^+ \\ &= STTS^+ \end{aligned}$$

Since $\|ST\|_A = \|TS^+\|_A$ (A -adjoints) we have

$$\|STS^+\|_A = \|ST\|_A^2$$

Ok, so we can focus focus on the $V(0, 1)$ cycle, the other cycles follow.

Upper Bound

We will measure convergence or reduction in $\|e\|_A$. Note:

$$\|e\|_A^2 = \|(I - T)e\|_A^2 + \|Te\|_A^2$$

For a $V(0, 1)$ cycle, the reduction in e is

$$\|STe\|_A^2 \leq (1 - \delta^*)\|e\|_A^2$$

we seek a sharp bound

$$\|ST\|_A^2 := \sup_{e \neq 0} \frac{\|STe\|_A^2}{\|e\|_A^2} = 1 - \delta^*$$

Sufficient conditions

What should we assume on relaxation and interpolation?

- relaxation is effective on the range of coarse grid correction

There exists $\delta > 0$ such that

$$\|STe\|_A^2 \leq (1 - \delta)\|Te\|_A^2 \text{ for all } e.$$

Then, since T is an A -orthogonal projector

$$\|STe\|_A^2 \leq (1 - \delta)\|e\|_A^2 \text{ for all } e$$

- No side effects on the range of interpolation

$$\|Sv\|_A^2 \leq \|v\|_A^2 \quad \text{for all } v \perp \text{range}(T)$$

Combine these into an assumption

$$\|Sv\|_A^2 \leq \|STv\|_A^2 + \|S(I - T)v\|_A^2 \leq (1 - \delta)\|Tv\|_A^2 + \|(I - T)v\|_A^2 = \|v\|_A^2 - \delta\|Tv\|_A^2$$

Theorem

If there exists $\delta > 0$ so that

$$\|Se\|_A^2 \leq \|e\|_A^2 - \delta\|Te\|_A^2 \quad \text{for all } e,$$

then

$$\|ST\|_A^2 \leq 1 - \delta$$

To be sharp, the largest δ , say $\hat{\delta}$

$$\hat{\delta} = \inf_{e: Te \neq 0} \frac{\|e\|_A^2 - \|Se\|_A^2}{\|Te\|_A^2},$$

should be δ^* .

Proof

Since $Te = \mathbf{0}$ gives $\|STe\|_A = 0$:

$$\|ST\|_A^2 = \sup_{e: Te \neq 0} \frac{\|STe\|_A^2}{\|e\|_A^2} = \sup_{e: Te \neq 0} \frac{\|STe\|_A^2}{\|Te\|_A^2 + \|(I - T)e\|_A^2}$$

Let \hat{e} be the arg sup

Then $T\hat{e}$ is also at the supremum.

Thus we have an error at the supremum with $(I - T)\hat{e} = 0$

$$\|ST\|_A^2 = \sup_{e: Te \neq 0} \frac{\|STe\|_A^2}{\|e\|_A^2} = \sup_{e: Te \neq 0} \frac{\|S(Te + (I - T)e)\|_A^2}{\|Te\|_A^2} = \sup_{e: Te \neq 0} \frac{\|Se\|_A^2}{\|Te\|_A^2},$$

And

$$1 - \|ST\|_A^2 = \inf_{e:Te \neq 0} \frac{\|Te\|_A^2 - \|Se\|_A^2}{\|Te\|_A^2} = \inf_{e:Te \neq 0} \frac{\|e\|_A^2 - \|Se\|_A^2}{\|Te\|_A^2} = \hat{\delta}$$

The worst δ is sharp

$$\hat{\delta} = \inf_{e:Te \neq 0} \frac{\|e\|_A^2 - \|Se\|_A^2}{\|Te\|_A^2},$$

The early theory split this in two part

For some $g(e)$ define δ , α_g , and β_g as in

$$\delta(e) = \underbrace{\frac{\|e\|_A^2 - \|Se\|_A^2}{g(e)}}_{\alpha_g(e)} \underbrace{\frac{g(e)}{\|Te\|_A^2}}_{1/\beta_g(e)}$$

Consider the smallest α_g and the largest β_g :

$$\hat{\alpha}_g = \inf_{e:g(e) \neq 0} \alpha_g(e) \quad \hat{\beta}_g = \sup_{e:g(e) \neq 0} \beta_g(e)$$

For e such that $g(Te) \neq 0$,

$$\begin{aligned} \|STe\|_A^2 &\leq \|Te\|_A^2 - \hat{\alpha}_g g(Te) \leq \|Te\|_A^2 - \frac{\hat{\alpha}_g}{\hat{\beta}_g} \|Te\|_A^2 = \left(1 - \frac{\hat{\alpha}_g}{\hat{\beta}_g}\right) \|Te\|_A^2 \\ &\leq \left(1 - \frac{\hat{\alpha}_g}{\hat{\beta}_g}\right) \|e\|_A^2 \end{aligned}$$

Ok, so this is generally worse than the sharp bound

$$\|ST\|_A = \sqrt{1 - \hat{\delta}} \leq \sqrt{1 - \frac{\hat{\alpha}_g}{\hat{\beta}_g}}$$

(α_g and β_g are not generally simultaneously satisfied)

Early works, e.g. Ruge-Stüben 1987, use $g(e) = \|e\|_{AD^{-1}A}^2$ Or the weaker form $g(e) = \|Te\|_{AD^{-1}A}^2$

Smoothing and Approximation

if there exists $\bar{\alpha}_g > 0$ such that

$$\|Se\|_A^2 \leq \|e\|_A^2 - \bar{\alpha}_g g(e) \quad \text{for all } e \quad (\text{smoothing})$$

and there exists $\bar{\beta}_g > 0$ such that

$$\|Te\|_A^2 \leq \bar{\beta}_g g(Te) \quad \text{for all } e \quad (\text{approximation})$$

$$\text{then } \|ST\|_A \leq \sqrt{1 - \bar{\alpha}_g / \bar{\beta}_g}$$

Select $g(e) = \|e\|_{AD^{-1}A}^2$

Since T is an A -orthogonal projection we have

$$\|Te\|_A = \inf_{e_c} \|e - Pe_c\|_A$$

(strong approximation) Assume there is a $\bar{\beta}_s$ such that

$$\inf_{e_c} \|e - Pe_c\|_A^2 \leq \bar{\beta}_s \|e\|_{AD^{-1}A}^2 \quad \text{for all } e.$$

The weaker version looks like (for some $\hat{\beta}$)

$$\|Te\|_A^2 \leq \bar{\beta} \|Te\|_{AD^{-1}A}^2 \quad \text{for all } e.$$

Weaker, means weaker norm. And we can make this a bit more practical. The range of T is A -orthogonal to the range of P , so

$$\begin{aligned} \|Te\|_A^2 &= \langle ATe, Te \rangle = \langle ATe, Te - Pe_c \rangle \\ &\leq \|Te\|_{AD^{-1}A} \|Te - Pe_c\|_D. \end{aligned}$$

(weak approximation) Assume that

$$\inf_{e_c} \|e - Pe_c\|_D^2 \leq \bar{\beta}_w \|e\|_A^2 \quad \text{for all } e,$$