Diagonal Difference *

11.2 more points to get your next star!

Rank: 499282 | Points: 463.8/475



X

You have successfully solved Diagonal Difference

Share

Post

You are now 11.2 points away from the 4th star for your problem solving badge.

Try the next challenge | Try a Random Challenge

Problem

Submissions

Leaderboard

Editorial

RATE THIS CHALLENGE



Given a square matrix, calculate the absolute difference between the sums of its diagonals.

For example, the square matrix arr is shown below:

- 1 2 3
- 4 5 6
- 9 8 9
- The left-to-right diagonal = 1 + 5 + 9 = 15.
- The right-to-left diagonal = 3 + 5 + 9 = 17.

Their absolute difference is |15 - 17| = 2.

Function description

Complete the ${\it diagonal Difference}$ function with the following parameter:

• int arr[n][m]: a 2-D array of integers

Return

• int: the absolute difference in sums along the diagonals

Input Format

The first line contains a single integer, n, the number of rows and columns in the square matrix arr.

Each of the next n lines describes a row, arr[i], and consists of n space-separated integers arr[i][j].

Constraints

• $-100 \le arr[i][j] \le 100$

Sample Input

```
STDIN Function
-----
3 arr[][] sizes n = 3, m = 3
11 2 4 arr = [[11, 2, 4], [4, 5, 6], [10, 8, -12]]
4 5 6
10 8 -12
```

Sample Output

15

Explanation

The primary diagonal is:

```
11
5
-12
```

Sum across the primary diagonal: 11 + 5 - 12 = 4.

The secondary diagonal is:

```
5
10
```

Sum across the secondary diagonal: 4 + 5 + 10 = 19

Difference: |4-19|=15

Note: |x| is the absolute value of x.

```
Change Theme Language Python 3
                                                                                                          10 23 ...
          #!/bin/python3
      2
      3
          import math
          import os
      5
          import random
          import re
      6
          import sys
      7
      8
      9
          # Complete the 'diagonalDifference' function below.
     10
     11
     12
          # The function is expected to return an INTEGER.
          # The function accepts 2D_INTEGER_ARRAY arr as parameter.
     13
     14
     15
          def diagonalDifference(arr):
     16
     17
              # Write your code here
     18
              n = len(arr)
     19
              primary_sum = 0
     20
              secondary_sum = 0
     21
              for i in range(n):
     22
     23
                  primary_sum += arr[i][i]
                  secondary_sum += arr[i][n - 1 - i]
     24
     25
     26
              return abs(primary_sum - secondary_sum)
     27
          if __name__ == '__main__':
              fptr = open(os.environ['OUTPUT_PATH'], 'w')
     28
     29
     30
              n = int(input().strip())
     31
     32
              arr = []
     33
              for in range(n):
     34
                                                                                                       Line: 42 Col: 1
EMACS
```

Run Code Submit Code

Expected Output

Test against custom input Congratulations Next Challenge You solved this challenge. Would you like to challenge your friends? Compiler Message Success Download Input (stdin) 11 2 4 Test case 3 △ Test case 4 △ 10 8 -12

Blog | Scoring | Environment | FAQ | About Us | Helpdesk | Careers | Terms Of Service | Privacy Policy

Download