CSCI-GA 2420 / MATH-GA 2010

Numerical Methods I

Course Notes

Fall 2026

Instructor: Prof. Florian Schaefer

Notes by You Li

Acknowledgements

These notes follow the Fall 2026 lectures of CSCI-GA 2420 / MATH-GA 2010 (Numerical Methods I) at the Courant Institute, taught by Prof. Florian Schaefer. Each chapter follows one lecture. Sections marked (SUPPLEMENT) go beyond the blackboard and draw on Trefethen and Bau [1], Demmel [2], and Golub and Van Loan [5]; background from linear algebra follows Axler [4].

Notation. Throughout, $\mathbb{K}$ denotes either $\mathbb{R}$ or $\mathbb{C}$. Vectors carry an arrow, $\vec{x} \in \mathbb{K}^n$; for $A \in \mathbb{K}^{m\times n}$, A^* is the conjugate transpose (equal to $A^\top$ when $\mathbb{K} = \mathbb{R}$), $\mathcal{R}(A)$ is the range (column space), and $\mathcal{N}(A)$ is the null space. The symbol $:=$ means "is defined as", with the colon on the side of the name being defined ($=:$ when that name stands on the right). Unless stated otherwise, $\|\cdot\|$ is the Euclidean norm on vectors and the induced 2-norm on matrices. Absolute values and inequalities between matrices, such as $|E| \leq |L|\,|U|$, are taken entrywise.

Contents

1 Floating-Point Representation, Conditioning, and Stability **6**
- 1.1 Overview . . . 6
- 1.2 Vectors and Matrices . . . 6
- 1.3 Representing Real Numbers . . . 7
- 1.4 The Geometry of Floating-Point Numbers . . . 9
- 1.5 Cancellation . . . 9
- 1.6 Conditioning . . . 10
- 1.7 Stability . . . 10
- 1.8 (SUPPLEMENT) Further Topics . . . 12

2 Cache Hierarchy and Tiling/Blocking, Solving Linear Systems **14**
- 2.1 Flops Are Not the Whole Story . . . 14
- 2.2 Memory Is One-Dimensional . . . 15
- 2.3 The Memory Hierarchy . . . 15
- 2.4 Loop Reordering . . . 16
- 2.5 Tiling . . . 16
- 2.6 Linear Systems . . . 17
- 2.7 Triangular Systems . . . 17
- 2.8 Gaussian Elimination . . . 18
- 2.9 (SUPPLEMENT) Further Topics . . . 19

3 LU Factorization and Its Stability **21**
- 3.1 Recap: From Triangular Systems to Elimination . . . 21
- 3.2 The Algebra of Lower Triangular Matrices . . . 21
- 3.3 Answering the Two Questions . . . 23
- 3.4 The In-Place Algorithm . . . 23
- 3.5 Zero Pivots . . . 24
- 3.6 Small Pivots Are Also Dangerous . . . 24
- 3.7 Error Analysis of the LU Factorization . . . 25
- 3.8 (SUPPLEMENT) Further Topics . . . 25

4 LU Factorization with Pivoting **27**
- 4.1 Backward Error of the LU Factorization . . . 27
- 4.2 Pivots and the Growth of Products . . . 29
- 4.3 Partial Pivoting . . . 29
- 4.4 The In-Place Algorithm with Partial Pivoting . . . 30

4.5 (SUPPLEMENT) Further Topics . . . 31

5 Least Squares Problems, Gram–Schmidt, and QR 33
5.1 Is Partial Pivoting Enough? . . . 33
5.2 Non-Square Systems . . . 34
5.3 Least Squares Problems . . . 34
5.4 The Geometry of Least Squares . . . 35
5.5 Orthogonalization: The QR Factorization . . . 36
5.6 Classical Gram–Schmidt . . . 36
5.7 Modified Gram–Schmidt . . . 38
5.8 (SUPPLEMENT) Further Topics . . . 40

6 Product View on Error Bounds, Householder QR 42
6.1 Why Write Algorithms as Matrix Products? . . . 42
6.2 The Product Form of Gram–Schmidt . . . 42
6.3 Error Amplification in Products . . . 45
6.4 Lower Triangularization: The Case of LU . . . 46
6.5 Gram–Schmidt Is Also a Lower Triangularization . . . 47
6.6 Householder Triangularization . . . 47
6.7 (SUPPLEMENT) Further Topics . . . 51

7 Householder QR in Practice, Givens Rotations, Existence and Uniqueness of QR 53
7.1 Picking Up from Householder QR . . . 53
7.2 Using Q . . . 53
7.3 Givens Rotations . . . 55
7.4 Full and Reduced QR . . . 58
7.5 Existence . . . 58
7.6 Uniqueness . . . 59
7.7 (SUPPLEMENT) Further Topics . . . 60

8 Overdetermined Least Squares and the Normal Equations 62
8.1 Recap and Motivation . . . 62
8.2 Conventions . . . 63
8.3 The Optimality Condition . . . 64
8.4 What the Optimality Condition Says . . . 66
8.5 The Normal Equations . . . 67
8.6 Gram Matrices and Positive Semidefinite Matrices . . . 68
8.7 (SUPPLEMENT) Further Topics . . . 70

9 Cholesky Factorization 71
9.1 Recap and Motivation . . . 71
9.2 Deriving Cholesky by Symmetric Elimination . . . 72
9.3 The Cholesky Algorithm . . . 74
9.4 Existence: Cholesky Never Breaks Down . . . 77
9.5 Rounding Errors . . . 78
9.6 Schur Complements . . . 80
9.7 Cholesky and QR . . . 83
9.8 (SUPPLEMENT) Further Topics . . . 84

10 Eigen-like Decompositions, Computing a Single Eigenvalue **85**
10.1 Recap and Motivation . . . 85
10.2 Eigenpairs of Hermitian Matrices . . . 86
10.3 A Negative Result: Eigenvalues Cannot Be Computed in Finitely Many Steps . . . 88
10.4 Power Iteration . . . 90
10.5 Matrix Functions: Changing the Spectrum . . . 95
10.6 Inverse Iteration . . . 96
10.7 Rayleigh Quotient Iteration . . . 97
10.8 (SUPPLEMENT) Further Topics . . . 101

11 Schur Factorization **105**
11.1 Recap and Motivation . . . 105
11.2 Deflation . . . 105
11.3 Subspace Iteration: Motivation . . . 107
11.4 Eigenvalue Problems as Matrix Factorizations . . . 107
11.5 Defective Matrices and Invariant Subspaces . . . 109
11.6 The Jordan Normal Form . . . 112
11.7 The Schur Factorization . . . 114
11.8 Computing Eigenvectors from the Schur Form . . . 117
11.9 (SUPPLEMENT) Further Topics . . . 118

12 Orthogonal Iteration **120**
12.1 Recap . . . 120
12.2 From Schur Factorization to Diagonalization . . . 120
12.3 Orthogonal Iteration . . . 121
12.4 Orthogonal Iteration in General . . . 124
12.5 Convergence of Eigenvector Estimates . . . 125
12.6 Convergence to Schur Factorization . . . 126
12.7 (SUPPLEMENT) Further Topics . . . 127

13 Convergence of Orthogonal Iteration and the Basic QR Iteration **129**
13.1 Recap . . . 129
13.2 Schur and Eigendecomposition . . . 130
13.3 Convergence of Orthogonal Iteration . . . 131
13.4 Convergence to Schur Factorization . . . 132
13.5 Basic QR Iteration . . . 133
13.6 (SUPPLEMENT) Further Topics . . . 135

14 Hessenberg QR **138**
14.1 Recap: From Orthogonal Iteration to the QR Algorithm . . . 138
14.2 The Cost of the QR Iteration . . . 138
14.3 Reduction to Hessenberg Form . . . 139
14.4 The QR Iteration on Hessenberg Matrices . . . 140
14.5 Bulge Chasing . . . 141
14.6 The Implicit Q Theorem . . . 142
14.7 (SUPPLEMENT) Further Topics . . . 143

15 Shifted QR **144**

15.1 Recap: QR Iteration on Hessenberg Matrices . . . 144
15.2 Proof of the Implicit Q Theorem . . . 145
15.3 Application to Bulge Chasing . . . 146
15.4 The Shifted QR Iteration . . . 146
15.5 Choosing the Shift . . . 147
15.6 Relationship with (Inverse) Power Iteration . . . 148
15.7 (SUPPLEMENT) Further Topics . . . 150

A Linear Algebra Basics **152**
A.1 Matrix Multiplication . . . 152
A.2 Transpose and Conjugate Transpose . . . 153
A.3 Inverses, Elementary and Triangular Matrices . . . 154
A.4 Determinants . . . 155
A.5 Subspaces and Rank . . . 155
A.6 Eigenvalues and Diagonalization . . . 156
A.7 Orthogonality and Projections . . . 156
A.8 Hermitian Matrices, Quadratic Forms, and the SVD . . . 157

B Norms and Unitary Matrices **158**
B.1 Vector and Matrix Norms . . . 158
B.2 Unitary Transformations . . . 163
B.3 The 2-Norm, Singular Values, and Condition Numbers . . . 165
B.4 The Frobenius Norm . . . 166

C The Floating-Point Model **169**
C.1 IEEE 754 Formats . . . 169
C.2 Machine Epsilon and the Model of Arithmetic . . . 170
C.3 Backward Stability . . . 171

Lecture 1

Floating-Point Representation, Conditioning, and Stability

Supplementary reading:

- [1] Lectures 12, 13, and 14.
- [2] Sections 1.3–1.5.
- [5] Section 2.4.

1.1 Overview

The topics of the semester are: numerical representation of vectors and matrices; solving linear systems by Gaussian elimination; solving least squares problems by orthogonalization; computing eigendecompositions; low-rank approximation; iterative methods; and structured matrices.

This lecture covers the first topic. A computer cannot store real numbers exactly: every number is rounded when it is stored, and every arithmetic operation rounds again, so even $(x + a) - a = x$ can fail. We proceed in two parts.

1. *Representation* (Sections 1.3 to 1.5): how real numbers are stored (floating point) and what errors this causes.
2. *Error analysis* (Sections 1.6 and 1.7): how to tell whether an error comes from the problem (*conditioning*) or from the algorithm (*stability*).

1.2 Vectors and Matrices

We write $\mathbb{K}$ for either $\mathbb{R}$ or $\mathbb{C}$, so that every result only needs to be stated once. A *vector* $\vec{v} \in \mathbb{K}^m$ is an m-tuple of elements of $\mathbb{K}$, representing a position or direction in an m-dimensional coordinate system. A *matrix* $A \in \mathbb{K}^{m \times n}$ is an $m \times n$ array of elements of $\mathbb{K}$, which may be thought of as n positions in an m-dimensional

coordinate system. We identify $\mathbb{K}^{m\times 1} \cong \mathbb{K}^m$, so a column vector is a matrix with one column:

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \\ A_{31} & A_{32} \end{bmatrix} = \begin{bmatrix} \vec{a}_1 & \vec{a}_2 \end{bmatrix}. \tag{1.1}$$

Scalar multiplication and addition act entrywise. For $A \in \mathbb{K}^{m\times n}$, $\vec{v} \in \mathbb{K}^n$, and $B \in \mathbb{K}^{n\times \ell}$,

$$(A\vec{v})_i = \sum_k A_{ik} v_k, \qquad (AB)_{ij} = \sum_k A_{ik} B_{kj}, \tag{1.2}$$

so A is a linear map $\mathbb{K}^n \to \mathbb{K}^m$ and $\mathbb{K}^{n\times\ell} \to \mathbb{K}^{m\times\ell}$: it carries a region, together with the vectors $\vec{x}$ in it, to the image vectors $A\vec{x}$.

The entrywise formula is rarely the most useful way to think. Numerical linear algebra relies much more on the *column view*:

$$A\vec{v} = \sum_{j=1}^{n} v_j \vec{a}_j, \tag{1.3}$$

that is, $A\vec{v}$ is the linear combination of the columns of A with coefficients given by the entries of $\vec{v}$. This view appears throughout the course. The column space $\mathcal{R}(A)$ decides whether $A\vec{x} = \vec{b}$ is solvable, the QR factorization orthogonalizes the columns, and low-rank approximation asks how few directions suffice to approximately span them. Likewise, the j-th column of AB is A applied to the j-th column of B, so a matrix–matrix product is ℓ matrix–vector products done at once.

1.3 Representing Real Numbers

To store $\vec{v} \in \mathbb{K}^m$ or $A \in \mathbb{K}^{m\times n}$ in memory, we must represent real numbers. A computer's native objects are bits, $\mathbb{B} \coloneqq \{0, 1\}$, so every number must be built from finitely many of them. We get there in steps, each fixing a shortcoming of the previous one: unsigned integers, signed integers, fixed point, and floating point.

1.3.1 Unsigned and Signed Integers

A string $d \in \mathbb{B}^b$, with bits $d_i \in \{0, 1\}$, represents the integers 0 through $2^b - 1$ via

$$\mathrm{UInt}(d) \coloneqq \sum_{i=1}^{b} d_i 2^{i-1}. \tag{1.4}$$

To represent negative numbers, we use an offset and give the highest bit a negative weight:

$$\mathrm{Int}(d) \coloneqq \sum_{i=1}^{b-1} d_i 2^{i-1} - d_b 2^{b-1}, \tag{1.5}$$

which covers -2^{b-1} through $2^{b-1} - 1$. This is *two's complement*. Nonnegative numbers keep their unsigned encoding and the adder circuit is shared with unsigned integers. The price is an asymmetric range: the most negative number, such as `typemin(Int8)` $= -128$, has no positive counterpart, so $-x$ can overflow back to x. (A different kind of offset, the *biased* encoding that stores $x + K$, is used for floating-point exponents; see Section C.1.)

1.3.2 Fixed-Point Numbers

Fixing the position of the binary point and using $f = \lfloor b/c \rfloor$ of the bits as fractional digits gives

$$\mathrm{Fixed}(d) \coloneqq \mathrm{Int}(d) \cdot 2^{-f}. \tag{1.6}$$

That is, about $1/c$ of the bits represent the fractional part (e.g., $c = 2$: half of them).

The problem. Fixed-point numbers are equally spaced, with spacing 2^{-f} everywhere. To represent both a large number s and a small number s^{-1}, we need about $\log_2(s)$ bits before the binary point and about as many after it. So the number of bits grows *linearly* with the number of orders of magnitude we want to cover. Physical simulations routinely span a dozen orders of magnitude, so fixed point quickly becomes impractical.

1.3.3 Floating-Point Numbers


The idea of floating point is to store the order of magnitude separately as an exponent. Given $(p, e) \in \mathbb{B}^{b_p} \times \mathbb{B}^{b_e}$, define

$$\mathrm{Float}(p, e) \coloneqq (-1)^{p_1} \left(1 + \sum_{i=2}^{b_p} p_i 2^{-(i-1)}\right) 2^{\mathrm{Int}(e)}. \tag{1.7}$$

The three parts are the sign bit p_1, the *significand* (or *mantissa*) in $[1, 2)$, and the power of two. The leading 1 of the significand is *implicit* and takes no storage. The decimal analogue is $\pm 1.123 \cdot 10^{-4}$.


This solves the problem of fixed point: to cover numbers from s^{-1} to s, the exponent only has to reach $\pm \log_2(s)$, which takes about $\log_2(\log_2(s))$ bits. Fixed point needs about $\log_2(s)$ bits for the same range. So floating point covers huge ranges cheaply, which is why scientific computing uses it. A complex number is stored as two floating-point numbers. For details, see [6].

In memory. The picture below shows how the bits are laid out, for single precision (Float32, $b_p = 24$, $b_e = 8$).



$$\underbrace{(-1)^{p_1}}_{\text{sign}} \times \underbrace{\left(1 + \sum_{i=2}^{b_p} p_i 2^{-(i-1)}\right)}_{\text{significand } 1.M \in [1,2)} \times \underbrace{2^{\mathrm{Int}(e)}}_{\text{IEEE stores } E = \text{exponent}+127}$$

For example, $-6.5 = -110.1_2 = -1.101_2 \times 2^2$ is stored as




that is, `0xC0D00000`: the sign is $(-1)^1$, the exponent is $129 - 127 = 2$, and the significand is $1.101_2 = 1 + \frac{1}{2} + \frac{1}{8} = 1.625$, so the value is $-1.625 \times 2^2 = -6.5$.

The only difference from the actual IEEE format is the exponent. The formula writes it as $\mathrm{Int}(e)$ for simplicity, while IEEE stores it with a bias, as $E - 127$ (or $E - 1023$ for Float64), and reserves the all-zeros and all-ones exponent fields for 0, subnormals, ∞, and `NaN` (Section C.1).

1.4 The Geometry of Floating-Point Numbers

Plotting all floating-point numbers on the real line shows that they are *not equally spaced*: within each *binade* $[2^E, 2^{E+1})$ they are equally spaced, but crossing a power of two doubles the spacing: if the spacing near 1 is ε, it is 2ε near 2 and 4ε near 4.



Why? Within one binade the exponent E is fixed, and only the significand $1.M$ changes.

- In $[1, 2)$: $1.M$ has $b_p - 1$ fraction bits, so it takes 2^{b_p-1} equally spaced values, with spacing $\varepsilon := 2^{-(b_p-1)}$.
- In $[2^E, 2^{E+1})$: the same values times 2^E, so the spacing is $2^E\varepsilon$.

Every binade holds the same number of floating-point numbers, but each is twice as long as the previous one. Hence the spacing doubles.

So the absolute spacing grows with the numbers, but the relative spacing (spacing divided by the number) always stays between $\varepsilon/2$ and ε. In other words, floating point gives every number about the same *relative* accuracy. This suits real data, whose uncertainty is usually proportional to its size.

Because every result is rounded to this grid, the usual rules of arithmetic can fail.

Example 1 (Floating-Point Addition Is Not Associative)**.** Suppose ties are rounded up, and evaluate $(1 + \varepsilon) + 1 + 2$ in two ways.

- Left to right: $(1 + \varepsilon) + 1 = 2 + \varepsilon$ lies exactly halfway between 2 and $2 + 2\varepsilon$ and rounds up to $2 + 2\varepsilon$. Adding 2 gives $4 + 2\varepsilon$, halfway between 4 and $4 + 4\varepsilon$, which rounds up to $4 + 4\varepsilon$.
- Last two terms first: $1 + 2 = 3$ exactly, and $(1 + \varepsilon) + 3 = 4 + \varepsilon$ rounds down to 4.

The same expression yields 4 or $4 + 4\varepsilon$. Multiplication is not associative either, and the distributive law fails as well; commutativity of addition and of multiplication does hold.

1.5 Cancellation

Relative accuracy has one weak point: subtracting two nearly equal numbers can destroy it.

Example 2 (Cancellation Exposes Existing Error)**.** Suppose the stored value of $10^6 + 1$ is already off by a relative error ε, so it is $(10^6 + 1)(1 + \varepsilon)$. Subtract 10^6:

$$(10^6 + 1)(1 + \varepsilon) - 10^6 \approx 1 + 10^6\varepsilon. \tag{1.8}$$

The exact answer is 1, but we get $1 + 10^6\varepsilon$.

- Before the subtraction: the error is about $10^6\varepsilon$, on a number of size 10^6. Relative error: ε.
- After the subtraction: the error is still $10^6\varepsilon$, but now on a number of size 1. Relative error: $10^6\varepsilon$.

The error itself did not grow. The result got smaller, so the same error is now a million times larger relative to it.

1.6 Conditioning

How do errors *propagate*? Consider computing $y = f(x)$ for $f : \mathbb{R} \to \mathbb{R}$. Since the input is rounded before anything else happens, the best we can hope for is $\hat{y} = f(\hat{x})$ with $|\hat{x} - x| / |x| < \varepsilon$. Comparing relative errors,

$$\frac{|\hat{y} - y| / |y|}{|\hat{x} - x| / |x|} = \frac{|f(\hat{x}) - f(x)|}{|\hat{x} - x|} \cdot \frac{|x|}{|f(x)|} \xrightarrow[\varepsilon \to 0]{} \frac{|x|\,|f'(x)|}{|f(x)|}. \tag{1.9}$$

Definition 3 (Condition Number)

The (relative) *condition number* of f at x is

$$\kappa_f(x) := \frac{|x|\,|f'(x)|}{|f(x)|}. \tag{1.10}$$

The problem is *well-conditioned* at x if $\kappa_f(x) \approx 1$ and *ill-conditioned* if $\kappa_f(x) \gg 1$.

The condition number depends only on the problem and the input, not on the algorithm. This is the most important point here: no algorithm can rescue an ill-conditioned problem, because the difficulty is intrinsic.

Example 4 (Conditioning of Subtraction)**.** Let $f(x) = x - 10^6$ at $x = 10^6 + 1$. Then

$$\kappa_f(10^6 + 1) = \frac{|10^6 + 1| \cdot 1}{1} \approx 10^6 \gg 1. \tag{1.11}$$

This is Example 2 seen as a problem: cancellation is a conditioning phenomenon, not a defect of floating point. Even an algorithm with infinite precision turns a relative uncertainty of 10^{-16} in x into a relative uncertainty of 10^{-10} in the output.

1.7 Stability

Conditioning describes the problem; stability describes the algorithm. We write $\hat{f}$ for the algorithm as it actually runs on the computer: it starts from the stored input $\hat{x} = \mathrm{fl}(x)$ and rounds after every operation.

Definition 5 (Stability)

An algorithm $\hat{f}$ for f is *stable* if

$$\frac{\left|\hat{f}(\hat{x}) - y\right| / |y|}{|\hat{x} - x| / |x|} \lesssim \kappa_f(x). \tag{1.12}$$

The left-hand side is the relative error of the output divided by the relative error of the input. In words: the algorithm amplifies input errors no more than the problem itself does. Otherwise $\hat{f}$ is *unstable*.

The form used in practice. The input error comes from rounding, so it is at most the unit roundoff $u = \varepsilon/2$ (Definition 116), and it is 0 if x is already a floating-point number, which would make the ratio undefined. Replacing the denominator by u gives

$$\frac{\left|\hat{f}(\hat{x}) - y\right|}{|y|} \lesssim \kappa_f(x)\, u. \tag{1.13}$$

The right-hand side is the error that *any* algorithm must make: rounding the input costs u, and the problem amplifies it by $\kappa_f(x)$. A stable algorithm adds nothing beyond that. (Here $\lesssim$ means "at most a modest constant times".)

Example 6 (An Unstable Algorithm for the Identity)**.** Let $f(x) = x$, the identity, with $\kappa_f = 1$, and compute it as $\hat{f}(x) = (x + 10^6) - 10^6$. Mathematically $\hat{f} = f$. Take x to be a floating-point number, so $\hat{x} = x$ and $y = x$.

- Step 1: $x + 10^6$ is rounded, $\mathrm{fl}(x + 10^6) = (x + 10^6)(1 + \delta)$ with $|\delta| \le u$. The low-order bits of x are lost here.
- Step 2: subtracting 10^6 is exact (Lemma 7).

Hence

$$\hat{f}(x) = x + (x + 10^6)\delta, \qquad \frac{\left|\hat{f}(x) - x\right|}{|x|} = \frac{|x + 10^6|}{|x|}\,|\delta| \approx \frac{10^6}{|x|}\,|\delta|\,. \tag{1.14}$$

For $x \approx 1$ the error is about $10^6 u$, but a stable algorithm must reach $\kappa_f u = u$. So $\hat{f}$ is unstable, by a factor of about 10^6; for instance, $\hat{f}(0.1) = 0.09999999997671694$. ($x = 1$ is an exception: $10^6 + 1$ is a floating-point number, so $\delta = 0$.)

Why it fails. Look at the two steps separately: $\hat{f} = g_2 \circ g_1$ with $g_1(x) = x + 10^6$ and $g_2(z) = z - 10^6$. For $x \approx 1$ and $z = x + 10^6$, their condition numbers (Definition 3) are

$$\kappa_{g_1}(x) = \frac{|x|}{|x + 10^6|} \approx 10^{-6}, \qquad \kappa_{g_2}(z) = \frac{|z|}{|z - 10^6|} \approx 10^6. \tag{1.15}$$

Step 1 shrinks relative errors by a factor 10^6; step 2 is a cancellation and magnifies them by 10^6. What happens to an error depends on where it enters:

- An error already present in x passes through both steps: it is shrunk, then magnified, so overall it is unchanged ($\kappa_f = 1$).
- The rounding error δ of step 1 enters *after* the shrinking step. It passes only through step 2, so it is magnified by 10^6.

Lesson. A well-conditioned problem can still be solved by an unstable algorithm if one of its intermediate steps is ill-conditioned. We will see this again with Gram–Schmidt and the normal equations.

Unstable vs. ill-conditioned. These are different diagnoses with different fixes.

- *Unstable*: the algorithm is at fault. Use a different algorithm.
- *Ill-conditioned*: the problem itself is sensitive, and no algorithm can help. Change the problem (reformulate or regularize it), or use higher precision.

The IEEE formats, the two conventions for machine epsilon, the standard model of floating-point arithmetic, and the notion of backward stability are collected in Appendix C.

1.8 (SUPPLEMENT) Further Topics

1.8.1 Rewriting Formulas to Avoid Cancellation

The recipe: find the step that subtracts two nearly equal numbers, and use an algebraic identity to rewrite the formula without it. Mathematically nothing changes; numerically the error drops to $O(u)$.

Roots of a quadratic. The roots of $ax^2 + bx + c$ are $x_{1,2} = (-b \pm \sqrt{b^2 - 4ac})/(2a)$. Suppose $b > 0$ and $b^2 \gg 4ac$. Then $\sqrt{b^2 - 4ac} \approx b$, so the "+" root computes $-b + (\text{almost } b)$: a cancellation. The "−" root adds two numbers of the same sign and is accurate. So compute that one first and get the other from Vieta's relation $x_1 x_2 = c/a$:

$$x_2 = \frac{-b - \sqrt{b^2 - 4ac}}{2a}, \qquad x_1 = \frac{c}{a\, x_2} = \frac{2c}{-b - \sqrt{b^2 - 4ac}}. \tag{1.16}$$

(If $b < 0$, swap the roles of the two roots.)

Variance. The formula $\mathrm{Var} = \overline{x^2} - \bar{x}^2$ (mean of squares minus square of mean) subtracts two large, nearly equal numbers when the data have a large mean and a small spread. For the data $10^8 + 1$, $10^8 + 2$, $10^8 + 3$ it returns 0 in Float64, while the true variance is $2/3$; it can even return a negative number. The fixes avoid squaring the large mean:

- *Two passes*: first compute $\bar{x}$, then $\mathrm{Var} = \frac{1}{n}\sum_i (x_i - \bar{x})^2$. The differences $x_i - \bar{x}$ are small, and all terms are nonnegative.
- *Welford's recurrence*: one pass over the data, updating the mean and the sum of squared deviations as each x_k arrives,

$$\bar{x}_k = \bar{x}_{k-1} + \frac{x_k - \bar{x}_{k-1}}{k}, \qquad M_k = M_{k-1} + (x_k - \bar{x}_{k-1})(x_k - \bar{x}_k), \tag{1.17}$$

  and finally $\mathrm{Var} = M_n/n$.

Three standard cases.

- $1 - \cos x$ for small x: for $x = 10^{-8}$, $\cos x$ rounds to exactly 1, giving 0 instead of $5 \cdot 10^{-17}$. Use $1 - \cos x = 2\sin^2(x/2)$.
- $\log(1 + x)$ for small x: already $1 + x$ loses the low bits of x (as in Example 6); for $x = 10^{-17}$ the result is 0. `log1p(x)` computes $\log(1 + x)$ directly from x (likewise `expm1(x)` for $e^x - 1$).
- $\sqrt{x+1} - \sqrt{x}$ for large x: for $x = 10^{16}$ the result is 0 instead of $5 \cdot 10^{-9}$. Rationalize: $\sqrt{x+1} - \sqrt{x} = 1/(\sqrt{x+1} + \sqrt{x})$.

1.8.2 Sterbenz's Lemma: Close Numbers Subtract Exactly

Lemma 7 (Sterbenz)

If x, y are floating-point numbers with $y/2 \le x \le 2y$, then $x - y$ is a floating-point number, so $\mathrm{fl}(x - y) = x - y$ exactly.

Intuition. Think of decimal numbers with three significant digits. $1.23 - 1.21 = 0.02$: both inputs lie on the grid of spacing 0.01, so their difference does too, and it is *smaller* than the inputs, so it needs fewer digits, not more. Nothing has to be rounded.

Proof. Say $y \le x \le 2y$, and let 2^e be the spacing of floating-point numbers near y, so y is an integer multiple of 2^e with at most b_p bits. The spacing near $x \ge y$ is 2^e or a larger power of 2, so x is an integer multiple of 2^e as well. Hence $x - y = k\,2^e$ with an integer $k \ge 0$. From $x \le 2y$ we get $x - y \le y < 2^{b_p} 2^e$, so $k < 2^{b_p}$ and $x - y$ fits in b_p bits. □

Consequence: the subtraction is not to blame. In Example 2 the subtraction itself is exact by the lemma: all of the error was already in the input, and the subtraction only made it visible. Rule: if the inputs are inexact, do not compute a small result by subtracting two large numbers.

1.8.3 Useful Techniques

Kahan summation. Adding n numbers one by one can make an error of order $nu \sum_i |x_i|$: adding a small term to a large running sum rounds away the term's low bits. Kahan's compensated summation keeps those lost bits in a variable c and feeds them back:

```
s = 0.0; c = 0.0
for x in xs
    y = x - c        # add back what was lost last time
    t = s + y        # rounded sum: low bits of y are lost
    c = (t - s) - y  # = (what was really added) - y = the lost part
    s = t
end
```

This reduces the error to about $2u \sum_i |x_i|$, essentially independent of n [7], at about four times the cost. Example: adding 0.1 ten million times gives an error of $1.6 \cdot 10^{-4}$ naively, 0 (to the printed digits) with Kahan, and $1.4 \cdot 10^{-9}$ with Julia's `sum`, which adds *pairwise* in a binary tree so that its error grows only like $u \log_2 n$.

Fused multiply-add (FMA). $\mathrm{fma}(a, b, c)$ computes $ab + c$ with a single rounding instead of two: faster and more accurate. Since a compiler may or may not use it, the same code can give slightly different results.

`-ffast-math`. This compiler flag (Julia: `@fastmath`) lets the compiler treat floating point as associative, so it may reorder sums (e.g., to vectorize them). This changes results and can break Kahan summation, whose $c = (t - s) - y$ is algebraically 0 and may be deleted. Keep it off.

1.8.4 Low-Precision Formats

Two 16-bit formats split their bits differently. Float16 has 5 exponent bits and $b_p = 11$: $\varepsilon \approx 10^{-3}$, but the largest number is only 65504. BFloat16 has 8 exponent bits, hence the same range as Float32 (up to $3.4 \cdot 10^{38}$), but only $b_p = 8$: $\varepsilon \approx 8 \cdot 10^{-3}$, about two decimal digits. Deep learning prefers BFloat16: gradients span many orders of magnitude, so what fails first is *range* (overflow, underflow), not precision; Float16 training needs *loss scaling* (multiplying the loss by a constant), BFloat16 does not. Sums are still accumulated in Float32: by Proposition 118, a dot product of length n has error up to about $nu\ |\vec{x}|^\top |\vec{y}|$. For $n = 4096$, BFloat16 ($u = 2^{-8}$) gives $nu = 16 \ge 1$, so the bound guarantees nothing, while Float32 ($u = 2^{-24}$) gives $nu \approx 2.4 \cdot 10^{-4}$. Hence tensor cores take BFloat16 (or FP8) inputs but accumulate in Float32.

Lecture 2

Cache Hierarchy and Tiling/Blocking, Solving Linear Systems

Supplementary reading:

- [2] Sections 2.3 and 2.6.
- [1] Lectures 17 and 20.

2.1 Flops Are Not the Whole Story

In Lecture 1 we asked whether we compute the *right* answer (conditioning: the problem's fundamental limit; stability: does the algorithm do something stupid?). Now we ask how *fast* we compute it.

Algorithms and computational cost. Consider the algorithm for matrix multiplication

$$A \mathrel{+}= BC, \qquad A \in \mathbb{R}^{\ell \times n},\ B \in \mathbb{R}^{\ell \times m},\ C \in \mathbb{R}^{m \times n}: \tag{2.1}$$

```
function matmuladd!(A, B, C)          # A += B*C
    for i = 1:l
        for j = 1:m
            for k = 1:n
                A[i,k] += B[i,j] * C[j,k]
            end
        end
    end
end
```

The innermost line runs $\ell \times m \times n$ times, so the computational cost is $\ell \times m \times n$. We set $\ell = 2 \cdot 10^4$, $m = 3 \cdot 10^4$ and time it with `@btime`:

n	Cost ℓmn	Time
1	$6 \cdot 10^8$	$6 \cdot 10^{-2}$ s
10^3	$6 \cdot 10^{11}$	4 s

The cost grows by a factor of 1000, so we would expect 1000×0.06 s $= 60$ s. It takes only 4 s: *more than* 10 *times faster than expected*. (The timings are for Julia's built-in multiplication, which calls BLAS.)

Flops and memory access. Does Julia use fewer operations? No: for $\ell = m = n$, the time grows roughly like the cost,

$\ell = m = n$	Cost ℓmn	Time
10^3	10^9	$7 \cdot 10^{-3}$ s
10^4	10^{12}	6 s

that is, about 1000 times the cost in about 1000 times the time. The explanation is that four operations appear in the algorithm: additions, multiplications, and reading and writing entries of the arrays (both written $[\,]$). Their costs are very different:

$$\text{cost}(+) < \text{cost}(*) \ll \text{cost}([\,]). \tag{2.2}$$

(On modern CPUs, $+$ and $*$ in fact cost about the same.) *Memory access is much slower than flops on modern hardware.*

- For $n = 1$, every entry of B is read and then used only once, so almost all of the time goes into memory access.
- For $n = 10^3$, every entry of B is used 10^3 times, so the cost of reading it is shared by many operations.

2.2 Memory Is One-Dimensional

Memory is a long string of bits, and a vector is a contiguous run of floating-point numbers $u_1, u_2, u_3, \ldots$. There is no such thing as a two-dimensional matrix in memory; it must be flattened in some order. Julia, Fortran, BLAS, and LAPACK use *column-major* order, storing entire columns contiguously:

$$A_{11},\ A_{21},\ A_{31},\ A_{41},\ A_{12},\ A_{22},\ A_{32},\ A_{42},\ \ldots \tag{2.3}$$

so that A_{ij} sits at linear index $i + (j-1)m$. C and NumPy default to row-major order. Besides the data, one also stores metadata such as sizes, strides, and preallocated buffers.

Since the storage is just a contiguous block, the same memory can be *reinterpreted* as an array of shape $m \times n \times k \times l$, $mn \times kl$, or $mnk \times l$ without any copying. Reshaping is free; transposing is not.

2.3 The Memory Hierarchy

Main memory is much slower than arithmetic, so hardware adds *caches*: small, fast memory close to the processor. Two bottlenecks remain:

- *Latency*: how long one access waits (a few cycles for L1, a few hundred for main memory).
- *Bandwidth*: how many bytes can be moved per second.

Latency is hidden by *prefetching*: the hardware fetches nearby memory, or memory in a strided pattern, before it is needed. This only works for regular access patterns, so the *regularity* of memory access often matters more than its volume. GPUs instead hide latency by running many threads. When threads access contiguous elements, the hardware merges their accesses into one, so on a GPU non-contiguous access is very slow.

2.4 Loop Reordering

Example 8 (Two Loop Orders for a Matrix–Vector Product)**.** Both loops compute $\vec{u} = A\vec{v}$ with the same number of operations.

```
# i outer: inner loop along a row of A
for i = 1:m
    for j = 1:n
        u[i] += A[i,j] * v[j]
    end
end
```

```
# j outer: inner loop down a column
for j = 1:n
    for i = 1:m
        u[i] += A[i,j] * v[j]
    end
end
```

In column-major storage, the left loop reads A along a row, in steps of m; in the right loop, every access is contiguous.

The left loop computes inner products of the rows of A with $\vec{v}$ (`dot`); the right loop adds up the columns of A times v_j (`axpy`). Julia stores matrices by columns, so the right loop is faster. The gap grows with the size of A: once A no longer fits in the cache, each strided access uses only one number from every cache line it loads. (The board says the order matters most for small matrices; we measured the opposite: the left loop is $3\times$ slower at $m = 64$ and $20\times$ at $m = 4096$.)

2.5 Tiling

Even without cache misses, the limited bandwidth slows us down. The key observation: multiplying two $m \times m$ matrices takes about m^3 operations but involves only about m^2 numbers, so each number is used about m times.

- If all m^2 numbers fit into the cache, we load each of them from main memory only once, and then do all m^3 operations in the fast cache.
- If the matrices are too large, we *tile*: split A, B, C into $b \times b$ blocks, $N = m/b$ along each side, and multiply block by block. (For simplicity, assume b divides m; otherwise the blocks at the edges are smaller.) We choose b so that the three blocks we are working on fit into the cache; Section 2.9.2 quantifies the gain.

Written out entry by entry, the tiled algorithm has *six* nested loops: the outer three choose the blocks (I, J, K count blocks, not entries), and the inner three multiply them.

```
for I = 1:N, J = 1:N, K = 1:N              # which blocks: C_IJ += A_IK * B_KJ
    for i = (I-1)*b+1 : I*b                # multiply the two b x b blocks
        for j = (J-1)*b+1 : J*b
            for k = (K-1)*b+1 : K*b
                C[i,j] += A[i,k] * B[k,j]
            end
        end
    end
end
```

We do the same m^3 operations as before, just in a different order. The inner three loops only use three blocks, $3b^2$ numbers in total. They stay in the cache and are each used about b times.

2.6 Linear Systems

Problem. Given $A \in \mathbb{K}^{m\times n}$ and $\vec{b} \in \mathcal{R}(A) \subseteq \mathbb{K}^m$, find $\vec{x} \in \mathbb{K}^n$ with

$$A\vec{x} = \vec{b}. \tag{2.4}$$

The condition $\vec{b} \in \mathcal{R}(A)$ just says that a solution exists.

- Linear systems appear in virtually all of mathematics.
- They *can* be solved by computing A^{-1}, but they *need not* be. Computing A^{-1} is in fact the wrong way: it costs more ($2m^3$ flops for a square matrix, versus $\frac{2}{3}m^3$ for the LU factorization of Lecture 3), and it usually gives a less accurate answer.

Square matrices. We begin with $A \in \mathbb{K}^{m\times m}$. Then the following three conditions are equivalent:

$$\underbrace{\mathcal{N}(A) = \{\vec{0}\}}_{\vec{x}\mapsto A\vec{x}\text{ injective}} \iff \text{the columns of } A \text{ are linearly independent} \iff \underbrace{\mathcal{R}(A) = \mathbb{K}^m}_{\vec{x}\mapsto A\vec{x}\text{ surjective}} \tag{2.5}$$

In words:

- *Left*: only $\vec{x} = \vec{0}$ is mapped to $\vec{0}$. Then different inputs give different outputs: if $A\vec{x} = A\vec{y}$, then $A(\vec{x} - \vec{y}) = \vec{0}$, so $\vec{x} = \vec{y}$. Hence a solution, if it exists, is unique.
- *Right*: every $\vec{b} \in \mathbb{K}^m$ is of the form $A\vec{x}$. Hence a solution always exists.

If these conditions hold, we say that A *has full rank*, *is invertible*, or *is nonsingular*; then $A\vec{x} = \vec{b}$ has exactly one solution for every $\vec{b}$. For rectangular matrices, injective and surjective are no longer the same; they are treated in Lecture 5.

2.7 Triangular Systems

We begin with the easy case. The small example

$$\begin{aligned} x_1 + 5 &= 15 \\ x_1 + 5x_2 &= 20 \end{aligned} \qquad \Longrightarrow \qquad x_1 = 10,\ x_2 = 2 \tag{2.6}$$

already shows the whole idea: x_1 does not depend on x_2, so the triangular structure *decouples* the unknowns and they can be solved for one at a time. For lower triangular A, *forward substitution* computes

$$x_1 = \frac{b_1}{A_{11}}, \qquad x_i = \frac{1}{A_{ii}}\left(b_i - \sum_{j=1}^{i-1} A_{ij}x_j\right), \quad i = 2, \dots, m. \tag{2.7}$$

Upper triangular systems are solved in reverse order by *back substitution*. The same formula admits two implementations, both costing $\sim m^2$ flops but with different access patterns.

Algorithm 1 Forward substitution, row-oriented (`dot` form).

1: **for** $i = 1, \dots, m$ **do**
2: $\quad x_i \leftarrow b_i$
3: $\quad$ **for** $j = 1, \dots, i-1$ **do**
4: $\quad\quad x_i \leftarrow x_i - A_{ij}x_j$
5: $\quad x_i \leftarrow x_i / A_{ii}$

Algorithm 2 Forward substitution, column-oriented (`axpy` form), the efficient variant.

$\vec{x} \leftarrow \vec{b}$
for $j = 1, \dots, m$ **do**
    $x_j \leftarrow x_j / A_{jj}$
    for $i = j+1, \dots, m$ **do**
        $x_i \leftarrow x_i - A_{ij} x_j$ ▷ remove the contribution of x_j at once

For upper triangular A, x_m is the only unknown in the last equation, so back substitution starts from the bottom:

$$x_m = \frac{b_m}{A_{mm}}, \qquad x_i = \frac{1}{A_{ii}} \left(b_i - \sum_{j=i+1}^{m} A_{ij} x_j \right), \quad i = m-1, \dots, 1. \tag{2.8}$$

The two implementations mirror those of forward substitution, with every loop over the unknowns reversed.

Algorithm 3 Back substitution, row-oriented (`dot` form).

for $i = m, \dots, 1$ **do**
    $x_i \leftarrow b_i$
    for $j = i+1, \dots, m$ **do**
        $x_i \leftarrow x_i - A_{ij} x_j$
    $x_i \leftarrow x_i / A_{ii}$

Algorithm 4 Back substitution, column-oriented (`axpy` form).

$\vec{x} \leftarrow \vec{b}$
for $j = m, \dots, 1$ **do**
    $x_j \leftarrow x_j / A_{jj}$
    for $i = 1, \dots, j-1$ **do**
        $x_i \leftarrow x_i - A_{ij} x_j$ ▷ remove the contribution of x_j at once

In column-major storage the row forms walk along rows (strided) and the column forms walk down columns (contiguous); this is the matrix–vector discussion again.

2.8 Gaussian Elimination

A general matrix is not triangular. Gaussian elimination *transforms A into upper triangular form by a sequence of row operations*, reducing the general problem to a triangular one. A 2×2 example:

$$\begin{aligned} x_1 + 2x_2 &= 10 \\ 2x_1 + x_2 &= 8 \end{aligned} \xrightarrow{\text{row } 2 - 2\times\text{row } 1} \begin{aligned} x_1 + 2x_2 &= 10 \\ -3x_2 &= -12 \end{aligned} \quad \Longrightarrow \quad x_2 = 4,\ x_1 = 2. \tag{2.9}$$

In matrix language, this elimination step is a *left multiplication*:

$$\begin{bmatrix} 1 & 0 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} 10 \\ 8 \end{bmatrix}. \tag{2.10}$$

Theorem 9 (Elementary Lower Triangular Matrices)

Let $A \in \mathbb{K}^{m\times m}$ and suppose the first $k-1$ columns of A are already zero below the diagonal, and $A_{kk} \neq 0$. Define the multipliers $\ell_{ik} := A_{ik}/A_{kk}$ for $i = k+1, \dots, m$, the vector

$$\vec{\ell}_k := \begin{bmatrix} 0 & \cdots & 0 & \ell_{k+1,k} & \cdots & \ell_{mk} \end{bmatrix}^\top, \tag{2.11}$$

whose first k entries vanish, and the *elementary lower triangular matrix*

$$L_k := I - \vec{\ell}_k \vec{e}_k^* = \begin{bmatrix} 1 & & & & & \\ & \ddots & & & & \\ & & 1 & & & \\ & & -\ell_{k+1,k} & 1 & & \\ & & \vdots & & \ddots & \\ & & -\ell_{mk} & & & 1 \end{bmatrix} \leftarrow \text{row } k \tag{2.12}$$

Then $L_k A$ is zero below the diagonal in its first k columns. Its first k rows equal those of A; besides zeroing column k below the diagonal, only the trailing $(m-k)\times(m-k)$ block is updated.

Proof. Left multiplication by L_k performs, for each $i > k$, the row operation "row i minus ℓ_{ik} times row k". The (i,k) entry becomes $A_{ik} - \ell_{ik}A_{kk} = 0$. Row k is zero in its first $k-1$ columns, so the columns that have already been eliminated are not disturbed. □

Applying Theorem 9 for $k = 1, \dots, m-1$ (assuming that each A_{kk} is nonzero when we reach it) to both sides of $A\vec{x} = \vec{b}$ gives

$$\underbrace{L_{m-1}\cdots L_1 A}_{\text{upper triangular}}\, \vec{x} = L_{m-1}\cdots L_1 \vec{b}. \tag{2.13}$$

With $U := L_{m-1}\cdots L_1 A$ and $L := (L_{m-1}\cdots L_1)^{-1}$, we obtain $A = LU$ with U upper triangular. We stop here with two questions: *is L lower triangular, and can we compute it?* Both are answered in Lecture 3.

2.9 (SUPPLEMENT) Further Topics

2.9.1 Leading Dimension

In column-major order, the entry A_{ij} lives at

$$\mathrm{addr}(A_{ij}) = \text{base} + (j-1)\cdot \mathrm{lda} + (i-1), \tag{2.14}$$

where the *leading dimension* $\mathrm{lda} \geq m$ is the distance between consecutive columns. Allowing $\mathrm{lda} \neq m$ lets us describe a *submatrix* of a larger matrix by the quadruple (start address, m, n, lda) without copying any data. This is what Julia's `view` and `@views` do, whereas `A[1:100, 1:100]` makes a copy.

2.9.2 Computational Intensity

We can quantify this [2, §2.6]. Consider two levels of memory, a fast memory of size M and a slow memory. Let one flop take time t_f and one slow-memory access take time $t_m \gg t_f$.

Definition 10 (Computational Intensity)

If an algorithm performs f flops and m slow-memory accesses, its *computational intensity* is $q \coloneqq f/m$.

The total time is roughly

$$T = ft_f + mt_m = ft_f \left(1 + \frac{t_m}{t_f} \cdot \frac{1}{q}\right). \tag{2.15}$$

Running near the peak rate ft_f requires $q \gg t_m/t_f$, and on modern machines t_m/t_f is in the tens to hundreds. The three levels of the BLAS are distinguished precisely by q (here n is the vector length or matrix order):

Level	Typical operation	Flops f	Accesses m	q
Level 1	`axpy` $\vec{y} \leftarrow \alpha\vec{x} + \vec{y}$	$2n$	$3n$	$2/3$
Level 2	`gemv` $\vec{y} \leftarrow \alpha A\vec{x} + \beta\vec{y}$	$2n^2$	n^2	2
Level 3	`gemm` $C \leftarrow \alpha AB + \beta C$	$2n^3$	$4n^2$	$n/2$

Only Level 3 has an intensity that grows with the problem size, matching the observation that matrix multiplication performs m^3 multiply–adds on m^2 data. A `dot` has $q \approx 1$, and a triangular solve (n^2 flops on about $n^2/2$ entries) has $q \approx 2$; both are limited by bandwidth.

For naive $C \mathrel{+}= AB$ with $n \times n$ matrices, B is read n times (n^3 accesses) while A and C cost $3n^2$, so

$$q = \frac{2n^3}{n^3 + 3n^2} \approx 2, \tag{2.16}$$

no better than Level 2. For the tiled version (Section 2.5) the slow-memory traffic is $m = 2n^2 + 2Nn^2$, so

$$q \approx \frac{2n^3}{2Nn^2} = \frac{n}{N} = b. \tag{2.17}$$

Three blocks must fit in fast memory, so $3b^2 \le M$ and $q \approx b \le \sqrt{M/3}$. A lower bound of Hong and Kung shows that any matrix multiplication that only reorders the sums has $q = O(\sqrt{M})$, so tiling is asymptotically optimal. Real implementations tile at *several* levels (registers, L1, L2, and across nodes), each with its own b.

2.9.3 Blocked LU

The in-place LU algorithm of Lecture 3 spends its time in rank-one updates $A \leftarrow A - \vec{x}\vec{y}^*$. These read and write all of A for 2 flops per entry ($q \approx 1$), so naive Gaussian elimination is bandwidth-bound. The tiling idea fixes this [2, §2.6.3]. Choose a block width b and write

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}, \qquad A_{11} \in \mathbb{K}^{b \times b}. \tag{2.18}$$

1. Factor the $m \times b$ panel $\begin{bmatrix} A_{11} \\ A_{21} \end{bmatrix}$ by ordinary elimination, obtaining L_{11}, L_{21}, U_{11} (Level 2, but only an $O(b/m)$ fraction of the work).
2. Compute $U_{12} = L_{11}^{-1} A_{12}$, a triangular solve with many right-hand sides (Level 3).
3. Compute $\tilde{A}_{22} = A_{22} - L_{21}U_{12}$, a matrix multiplication (Level 3) that contains most of the flops.
4. Recurse on $\tilde{A}_{22}$.

Almost all the work is matrix multiplication with $q \approx b$. This is how LAPACK's `getrf` beats the older LINPACK: rewrite everything as matrix multiplication.

Lecture 3

LU Factorization and Its Stability

Supplementary reading:

- [1] Lectures 20 and 21.
- [2] Section 2.3.

3.1 Recap: From Triangular Systems to Elimination

We begin with a recap of Lecture 2: matrices are stored column by column, memory access costs far more than arithmetic, and the loop order and tiling of an algorithm should match the storage. We then returned to linear systems. Triangular systems are solved by forward or back substitution in $O(m^2)$ flops, and Gaussian elimination reduces a general system to a triangular one: by Theorem 9, applying elementary lower triangular matrices $L_k = I - \vec{\ell}_k \vec{e}_k^*$ for $k = 1, \ldots, m-1$ gives

$$U \coloneqq L_{m-1} \cdots L_1 A \text{ upper triangular}, \qquad L \coloneqq (L_{m-1} \cdots L_1)^{-1}, \qquad A = LU. \tag{3.1}$$

This lecture answers the two questions left open: *is L lower triangular, and can we compute it cheaply?* Answering them requires a little algebra of lower triangular matrices.

3.2 The Algebra of Lower Triangular Matrices

Proposition 11 (Closure Properties)

The lower triangular matrices in $\mathbb{K}^{m \times m}$ are closed under

1. addition,
2. scalar multiplication,
3. matrix multiplication, and
4. inversion (of invertible ones).

Proof. Items 1 and 2 are clear. For item 3, let L and $\tilde{L}$ be lower triangular, so $(L\tilde{L})_{ij} = \sum_k L_{ik}\tilde{L}_{kj}$. A term $L_{ik}\tilde{L}_{kj}$ can be nonzero only if $i \geq k$ and $k \geq j$, hence $i \geq j$. So every term vanishes when $i < j$, and $L\tilde{L}$ is

lower triangular. Item 4 is proved below for the matrices arising in LU, using Theorem 12; the general case is in Section 3.8. □

Theorem 12 (Inverses and Products of Elementary Lower Triangular Matrices)

Let $L_k = I - \vec{\ell}_k \vec{e}_k^*$ be as in Theorem 9. Then:

1. $L_k^{-1} = I + \vec{\ell}_k \vec{e}_k^*$, i.e., inversion just *flips the signs* of the multipliers;
2. the product of the inverses, *in this order*, places every multiplier in its own position:

$$L_1^{-1} L_2^{-1} \cdots L_{m-1}^{-1} = \begin{bmatrix} 1 & & & \\ \ell_{21} & 1 & & \\ \vdots & \ddots & \ddots & \\ \ell_{m1} & \cdots & \ell_{m,m-1} & 1 \end{bmatrix}. \tag{3.2}$$

Trefethen and Bau call these two facts the "two strokes of luck" of Gaussian elimination [1, Lecture 20]. The order in item 2 matters. In the reverse order the multipliers mix: for example,

$$L_2^{-1} L_1^{-1} = (I + \vec{\ell}_2 \vec{e}_2^*)(I + \vec{\ell}_1 \vec{e}_1^*) = I + \vec{\ell}_1 \vec{e}_1^* + \vec{\ell}_2 \vec{e}_2^* + \ell_{21}\, \vec{\ell}_2 \vec{e}_1^*, \tag{3.3}$$

and the last term adds $\ell_{21}\ell_{i2}$ to the entries of column 1 below row 2.

Both items rest on one simple fact: $\vec{\ell}_k$ is zero in its first k entries. The number $\vec{e}_p^* \vec{\ell}_k$ is entry p of $\vec{\ell}_k$, so it is 0 for every $p \le k$. This is what makes the extra terms vanish in the proof below.

Proof. For item 1, since $\vec{e}_k^* \vec{\ell}_k = 0$,

$$(I + \vec{\ell}_k \vec{e}_k^*)(I - \vec{\ell}_k \vec{e}_k^*) = I + \vec{\ell}_k \vec{e}_k^* - \vec{\ell}_k \vec{e}_k^* - \vec{\ell}_k (\vec{e}_k^* \vec{\ell}_k) \vec{e}_k^* = I. \tag{3.4}$$

For item 2, we compute the product column by column. Column j of $M := L_1^{-1} \cdots L_{m-1}^{-1}$ is

$$M \vec{e}_j = (I + \vec{\ell}_1 \vec{e}_1^*) \cdots (I + \vec{\ell}_{m-1} \vec{e}_{m-1}^*)\, \vec{e}_j. \tag{3.5}$$

Here j is fixed: it is the column we compute. The factors are $L_p^{-1} = I + \vec{\ell}_p \vec{e}_p^*$ for $p = 1, \ldots, m-1$. We apply them to $\vec{e}_j$ from right to left, using $(I + \vec{\ell}_p \vec{e}_p^*)\vec{x} = \vec{x} + \vec{\ell}_p\, (\vec{e}_p^* \vec{x})$, where $\vec{e}_p^* \vec{x}$ is entry p of $\vec{x}$.

- *Factors with $p > j$*: entry p of $\vec{e}_j$ is 0, so these factors leave $\vec{e}_j$ unchanged.
- *The factor $p = j$*: entry j of $\vec{e}_j$ is 1, so it gives $\vec{e}_j + \vec{\ell}_j$.
- *Factors with $p < j$*: entry p of $\vec{e}_j + \vec{\ell}_j$ is 0, since $\vec{e}_j$ has its 1 in position j and $\vec{\ell}_j$ is zero in its first j entries. So these factors leave $\vec{e}_j + \vec{\ell}_j$ unchanged.

Hence column j of M is

$$M \vec{e}_j = \vec{e}_j + \vec{\ell}_j = \begin{bmatrix} 0 & \cdots & 0 & 1 & \ell_{j+1,j} & \cdots & \ell_{mj} \end{bmatrix}^\top, \tag{3.6}$$

with the 1 in position j. These are exactly the columns of the matrix in item 2. □

3.3 Answering the Two Questions

Now we can answer the two questions from the end of Lecture 2. Since the inverse of a product reverses the order of the factors, $L = (L_{m-1} \cdots L_1)^{-1} = L_1^{-1} L_2^{-1} \cdots L_{m-1}^{-1}$.

- *Is L lower triangular? Yes.* Each L_k^{-1} is lower triangular (item 1 of Theorem 12), and so is their product (item 3 of Proposition 11). Its diagonal is all ones: L is *unit* lower triangular.
- *Can we compute L cheaply? Yes, for free.* By item 2 of Theorem 12, L is the matrix (3.2), whose entries below the diagonal are the multipliers ℓ_{ik} that elimination computes anyway.

3.4 The In-Place Algorithm

Idea: transform A into U with Gaussian elimination while recording the multipliers. Both L and U are stored in place, in the memory of A:

- U goes on and above the diagonal;
- the multipliers ℓ_{ik} go below the diagonal, where elimination has just created zeros;
- the diagonal of L is all ones, so it need not be stored.

Hence we can overwrite A with L and U, without any extra memory.

Algorithm 5 LU factorization without pivoting, in place.

```
for k = 1, ..., m do
    A_{k+1:m, k} ← A_{k+1:m, k} / A_{kk}          ▷ multipliers ℓ_ik, stored in place
    for j = k + 1, ..., m do
        for i = k + 1, ..., m do
            A_ij ← A_ij − A_ik A_kj               ▷ = A_ij − ℓ_ik u_kj
```

When the algorithm finishes, A holds both factors: U on and above the diagonal, and the multipliers of L below it. The inner loop runs over i, i.e., down a column of A, which is the fast direction in column-major storage (Lecture 2).

Vector form. Step k computes the multipliers $\vec{\ell}_k$ and subtracts an outer product from the trailing block:

$$\vec{\ell}_k = A_{k+1:m,\,k} / A_{kk}, \qquad A_{k+1:m,\,k+1:m} \leftarrow A_{k+1:m,\,k+1:m} - \vec{\ell}_k \, A_{k,\,k+1:m}. \tag{3.7}$$

A column vector times a row vector has rank one, so each step is a *rank-one update* of the trailing block.

Block form. Write $A = \begin{bmatrix} a_{11} & \vec{w}^* \\ \vec{v} & B \end{bmatrix}$ with a scalar a_{11}. The first step of elimination is the factorization

$$A = \begin{bmatrix} 1 & 0 \\ \vec{v}/a_{11} & I \end{bmatrix} \begin{bmatrix} a_{11} & \vec{w}^* \\ 0 & S \end{bmatrix}, \qquad S = B - \frac{\vec{v}\vec{w}^*}{a_{11}}. \tag{3.8}$$

S is the *Schur complement* (Lecture 9). The algorithm then factors S in the same way, so LU is this one step applied recursively.

3.5 Zero Pivots

The algorithm divides by A_{kk}, the k-th *pivot*. If a pivot is zero, it cannot proceed.

Example 13 (An Invertible Matrix with No LU Factorization)**.** The matrix

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix} \tag{3.9}$$

is invertible, but $A_{11} = 0$ is a zero pivot, so the very first step divides by zero and $A = LU$ does not exist. *Invertibility does not imply the existence of an LU factorization.*

Theorem 14 (Zero Pivots and Leading Principal Submatrices)

If the k-th pivot of the LU factorization of A is zero, then the leading principal submatrix $A[1{:}k, 1{:}k]$ is singular. All pivots of the LU factorization are nonzero if and only if every leading principal submatrix of A is nonsingular.

Proof. Step 1: the top-left block. Take the top-left $k \times k$ block of $A = LU$. Entry (i, j) with $i, j \le k$ is $\sum_p L_{ip}U_{pj}$, and $L_{ip} = 0$ for $p > i$, so only $p \le k$ contributes. Hence

$$A[1{:}k, 1{:}k] = L[1{:}k, 1{:}k]\, U[1{:}k, 1{:}k]. \tag{3.10}$$

Step 2: determinants. $L[1{:}k, 1{:}k]$ is unit lower triangular, so its determinant is 1. $U[1{:}k, 1{:}k]$ is upper triangular, so its determinant is the product of its diagonal, i.e., of the first k pivots:

$$\det A[1{:}k, 1{:}k] = u_{11}u_{22}\cdots u_{kk}. \tag{3.11}$$

So $A[1{:}k, 1{:}k]$ is nonsingular if and only if the first k pivots are nonzero.

Step 3: a zero pivot stops the algorithm early. If $u_{kk} = 0$, the algorithm cannot continue past step k. But the first k rows and columns of L and U have already been computed, and Step 1 uses only those. So the argument still applies. □

3.6 Small Pivots Are Also Dangerous

Exactly zero pivots are unlikely in practice. Is it then enough that no pivot is exactly zero? Consider a small number $\varepsilon > 0$, much smaller than the machine epsilon (e.g., $\varepsilon = 10^{-20}$; here ε is *not* the machine epsilon):

$$A = \begin{bmatrix} \varepsilon & 1 \\ 1 & 1 \end{bmatrix}, \qquad L = \begin{bmatrix} 1 & 0 \\ \varepsilon^{-1} & 1 \end{bmatrix}, \qquad U = \begin{bmatrix} \varepsilon & 1 \\ 0 & 1 - \varepsilon^{-1} \end{bmatrix}. \tag{3.12}$$

This is mathematically correct. In floating-point arithmetic, however, ε^{-1} is huge and $1 - \varepsilon^{-1}$ rounds to $-\varepsilon^{-1}$. The computed factors then give

$$\tilde{L}\tilde{U} = \begin{bmatrix} \varepsilon & 1 \\ 1 & \varepsilon^{-1} - \varepsilon^{-1} \end{bmatrix} = \begin{bmatrix} \varepsilon & 1 \\ 1 & 0 \end{bmatrix} \approx \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \tag{3.13}$$

which is nothing like A. The small pivot produces a huge multiplier, and rounding wipes out all information in A_{22}.

3.7 Error Analysis of the LU Factorization

Theorem 15 (Backward Error of LU)

Let $\varepsilon_{\text{mach}}$ be the machine epsilon. If the LU factorization runs to completion in floating-point arithmetic without encountering a zero pivot, the computed factors L, U satisfy, to first order in $\varepsilon_{\text{mach}}$,

$$A = LU + E, \qquad |E| \leq \varepsilon_{\text{mach}}\, m\, |L|\, |U|\,, \tag{3.14}$$

where $|\cdot|$ takes entrywise absolute values and the inequality holds entrywise.

The proof is given in Lecture 4. The theorem says that the computed factorization is the exact factorization of a nearby matrix, with a perturbation controlled by $|L|\,|U|$. If $|L|\,|U|$ is comparable to $|A|$, the error is at the level of machine precision. If L or U contains huge entries, the bound is useless. In the example above, L contains ε^{-1} and U contains $1 - \varepsilon^{-1}$, so $|L|\,|U|$ has entries of size ε^{-1} and the error can be as large as A itself.

3.8 (SUPPLEMENT) Further Topics

3.8.1 Lower Triangular Matrices

Inverses of general lower triangular matrices. Above, we proved item 4 of Proposition 11 only for matrices of the form $(L_{m-1} \cdots L_1)^{-1}$. It holds for every invertible lower triangular T:

1. $\det T$ is the product of the diagonal entries, so they are all nonzero. Let $D = \operatorname{diag}(T)$.
2. $N \coloneqq D^{-1}T$ divides row i of T by T_{ii}, so N is *unit* lower triangular.
3. By item 2 of Theorem 12, $N = L_1^{-1} \cdots L_{m-1}^{-1}$ if we take the entries of N below the diagonal as the multipliers. So $N^{-1} = L_{m-1} \cdots L_1$ is a product of lower triangular matrices, hence lower triangular.
4. $T = DN$, so $T^{-1} = N^{-1}D^{-1}$ is lower triangular as well.

A shorter proof of item 2 of Theorem 12. Following [1, Lecture 20]: for $k < j$ we have $\vec{e}_k^{\,*}\vec{\ell}_j = 0$, so multiplying two of the factors just adds their multipliers:

$$(I + \vec{\ell}_k\vec{e}_k^{\,*})(I + \vec{\ell}_j\vec{e}_j^{\,*}) = I + \vec{\ell}_k\vec{e}_k^{\,*} + \vec{\ell}_j\vec{e}_j^{\,*}. \tag{3.15}$$

Multiplying in the factors one at a time from the right, the same cancellation happens at every step, and we get $L_1^{-1} \cdots L_{m-1}^{-1} = I + \sum_{k=1}^{m-1} \vec{\ell}_k\vec{e}_k^{\,*}$.

3.8.2 Cost of LU and Solving Linear Systems

Cost of LU. Step k of Algorithm 5 updates the trailing $(m-k) \times (m-k)$ block, with one multiplication and one subtraction per entry. Adding up over all steps,

$$\sum_{k=1}^{m-1} 2(m-k)^2 = 2\sum_{j=1}^{m-1} j^2 \approx \frac{2}{3}m^3 \text{ flops.} \tag{3.16}$$

The divisions for the multipliers add only $O(m^2)$. The update is a rank-one update, a Level 2 operation with low computational intensity; Section 2.9.3 shows how blocking turns most of it into matrix multiplication.

Solving $A\vec{x} = \vec{b}$ with LU. We split the solve into three steps:

Step	Operation	Cost
1. factor	$A = LU$	$\frac{2}{3}m^3$
2. forward substitution	$L\vec{y} = \vec{b}$	m^2
3. back substitution	$U\vec{x} = \vec{y}$	m^2

The factorization is done once. Every further right-hand side costs only $2m^2$, the same as a multiplication by A^{-1}. Forming A^{-1} would cost about $2m^3$ up front, three times as much as LU, and is less accurate. So we never compute A^{-1} to solve a linear system.

3.8.3 Pivots, Minors, and Uniqueness

The pivots as ratios of minors. Following [2, §2.3]: Step 2 of the proof of Theorem 14 gives $\det A[1{:}k, 1{:}k] = u_{11} \cdots u_{kk}$ for each k. Dividing two consecutive ones,

$$u_{kk} = \frac{\det A[1{:}k, 1{:}k]}{\det A[1{:}k{-}1, 1{:}k{-}1]}. \tag{3.17}$$

So each pivot is determined by A alone, before any elimination is done. For example, for the matrix of the small-pivot example, $u_{11} = \varepsilon$ and $u_{22} = \det A/\varepsilon = 1 - \varepsilon^{-1}$, which matches the U computed there. Moreover, $A = LU$ with L unit lower triangular exists and is unique if and only if $A[1{:}k, 1{:}k]$ is nonsingular for every $k = 1, \ldots, m-1$.

Why the factorization is unique. Let A be nonsingular and suppose $A = L_1U_1 = L_2U_2$, with L_1, L_2 unit lower triangular and U_1, U_2 upper triangular. All four are invertible, and

$$L_2^{-1}L_1 = U_2U_1^{-1}. \tag{3.18}$$

The left side is unit lower triangular and the right side is upper triangular (Proposition 11). A matrix that is both must be I, so $L_1 = L_2$ and $U_1 = U_2$.

3.8.4 Pivoting Preview

Swapping rows. In the small-pivot example, swap the two rows first:

$$PA = \begin{bmatrix} 1 & 1 \\ \varepsilon & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ \varepsilon & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1-\varepsilon \end{bmatrix}. \tag{3.19}$$

Now the only multiplier is ε, and every entry of L and U is at most 1 in size. So $|L|\,|U|$ is about as large as $|A|$, and by Theorem 15 the error is at the level of machine precision. This is the idea of *partial pivoting*: at each step, swap rows so that the entry of largest absolute value in the current column becomes the pivot. Then every multiplier satisfies $|\ell_{ik}| \le 1$, and we get $PA = LU$ for a permutation matrix P. This is the subject of Lecture 4.

In Julia. `lu(A)` computes $PA = LU$ with partial pivoting; `lu(A, NoPivot())` runs Algorithm 5 as written. For $A = \left[\begin{smallmatrix} 10^{-20} & 1 \\ 1 & 1 \end{smallmatrix}\right]$ the unpivoted factors satisfy $LU - A = \left[\begin{smallmatrix} 0 & 0 \\ 0 & -1 \end{smallmatrix}\right]$, exactly the failure of the small-pivot example, while the pivoted ones are exact.

```
F = lu(A)                 # F.L, F.U, F.p (row permutation)
x = F \ b                 # reuses the factorization: 2m^2 flops per b
G = lu(A, NoPivot())      # G.L * G.U - A == [0 0; 0 -1] for this A
```

Lecture 4

LU Factorization with Pivoting

Supplementary reading:

- [1] Lectures 21 and 22.
- [2] Sections 2.3 and 2.4.
- [7] Chapter 9.

4.1 Backward Error of the LU Factorization

Recall from Lecture 3 that Gaussian elimination computes $A = LU$ in place, with the multipliers stored below the diagonal and U on and above it, and that it fails on a zero pivot even when A is nonsingular (for example $\begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}$). Lecture 3 ended with an example in which a tiny pivot destroys the LU factorization of the well-conditioned matrix

$$A = \begin{bmatrix} \varepsilon & 1 \\ 1 & 1 \end{bmatrix}, \qquad \kappa(A) \approx 2.6 \text{ as } \varepsilon \to 0. \tag{4.1}$$

Here ε is just a small number, much smaller than $\varepsilon_{\text{mach}}$ (e.g., 10^{-20}); it is not the machine epsilon. The problem is well-conditioned, so the failure lies with the algorithm. The following theorem explains exactly where it comes from.

Theorem 16 (Backward Error of LU)

Suppose the LU factorization of $A \in \mathbb{K}^{m\times m}$ runs to completion in floating-point arithmetic without encountering a zero pivot, producing $\tilde{L}$ and $\tilde{U}$. Then, to first order in $\varepsilon_{\text{mach}}$,

$$A = \tilde{L}\tilde{U} + E, \qquad |E| \le m\varepsilon_{\text{mach}} \left|\tilde{L}\right| \left|\tilde{U}\right|. \tag{4.2}$$

So $\tilde{L}$ and $\tilde{U}$ are the exact LU factors of a nearby matrix, $A - E$. If $\left|\tilde{L}\right| \left|\tilde{U}\right|$ is about as large as $|A|$, then E is tiny compared with A and the algorithm is backward stable. If $\left|\tilde{L}\right| \left|\tilde{U}\right|$ is much larger than $|A|$, then E can be as large as A. ("To first order" means we drop terms of size $\varepsilon_{\text{mach}}^2$.)

4.1.1 Proof of Theorem 16

Formulas for L and U. From $A = LU$ and $L_{ii} = 1$, solving for the entries of L and U gives (these are the numbers elimination computes, in a different order)

$$U_{ik} = A_{ik} - \sum_{j=1}^{i-1} L_{ij}U_{jk}, \qquad i \le k, \tag{4.3}$$

$$L_{ik} = \frac{1}{U_{kk}}\left(A_{ik} - \sum_{j=1}^{k-1} L_{ij}U_{jk}\right), \qquad i > k. \tag{4.4}$$

Rounding in inner products (Homework 2, Proposition 118): in floating point, every term of a sum gets its own small relative error,

$$\mathrm{fl}\Big(\sum_{i=1}^{m} x_i y_i\Big) = \sum_{i=1}^{m} x_i y_i(1+\delta_i), \qquad |\delta_i| \lesssim m\varepsilon_{\mathrm{mach}}. \tag{4.5}$$

Proof. The U part ($i \le k$). In floating point, (4.3) becomes

$$\tilde{U}_{ik} = \Big(A_{ik} - \sum_{j=1}^{i-1} \tilde{L}_{ij}\tilde{U}_{jk}(1+\delta_j)\Big)(1+\delta'), \tag{4.6}$$

where δ' is the error of the last subtraction. Solve for A_{ik}, use $\tilde{L}_{ii} = 1$, and write $1/(1+\delta') = 1+\delta_i$:

$$A_{ik} = \sum_{j=1}^{i} \tilde{L}_{ij}\tilde{U}_{jk}(1+\delta_j) = \sum_{j=1}^{i} \tilde{L}_{ij}\tilde{U}_{jk} + \underbrace{\sum_{j=1}^{i} \tilde{L}_{ij}\tilde{U}_{jk}\delta_j}_{=:E_{ik}}. \tag{4.7}$$

Every $|\delta_j| \lesssim m\varepsilon_{\mathrm{mach}}$, so $|E_{ik}| \le m\varepsilon_{\mathrm{mach}} \sum_j \left|\tilde{L}_{ij}\right|\left|\tilde{U}_{jk}\right| = m\varepsilon_{\mathrm{mach}}(\left|\tilde{L}\right|\left|\tilde{U}\right|)_{ik}$. (The terms with $j > i$ are zero because $\tilde{L}$ is lower triangular.)

The L part ($i > k$). The same argument applied to (4.4). The division by $\tilde{U}_{kk}$ adds one more rounding error, which is merged into δ_k in the same way, and the bound is the same. □

4.1.2 What the Bound Says

If $\left|\tilde{L}\right|\left|\tilde{U}\right|$ is of the same order as $|A|$, then E is a relative perturbation of size $O(m\varepsilon_{\mathrm{mach}})$ and the algorithm is backward stable. But $\left|\tilde{L}\right|\left|\tilde{U}\right|$ can be far larger than $|A|$. In the example above,

$$|L|\,|U| = \begin{bmatrix} 1 & 0 \\ \varepsilon^{-1} & 1 \end{bmatrix}\begin{bmatrix} \varepsilon & 1 \\ 0 & \varepsilon^{-1}-1 \end{bmatrix} = \begin{bmatrix} \varepsilon & 1 \\ 1 & 2\varepsilon^{-1}-1 \end{bmatrix}. \tag{4.8}$$

The bottom-right entry is about $2\varepsilon^{-1}$; multiplied by $\varepsilon_{\mathrm{mach}}$, the error can be $O(1)$ or larger, exactly as observed. The question becomes: *how can we keep $|L|\,|U|$ small?*

4.2 Pivots and the Growth of Products

Look at the storage after $k-1$ steps of elimination: the first $k-1$ rows already hold U, the first $k-1$ columns below the diagonal hold L, and the trailing block $\tilde{A}$ is still being processed.

$$\begin{bmatrix} U_{11} & \cdots & & & & \\ & \ddots & & & & \\ & & U_{k-1,k-1} & U_{k-1,k} & \cdots & U_{k-1,m} \\ & L & L_{k,k-1} & \tilde{A}_{kk} & \cdots & \tilde{A}_{km} \\ & & \vdots & \vdots & & \vdots \\ & & L_{m,k-1} & \tilde{A}_{mk} & \cdots & \tilde{A}_{mm} \end{bmatrix} \tag{4.9}$$

Step k turns row k into a row of U and column k below the diagonal into a column of L, and updates the trailing $(m-k)\times(m-k)$ block:

$$L_{ik} = \frac{\tilde{A}_{ik}}{\tilde{A}_{kk}}, \qquad U_{kj} = \tilde{A}_{kj}, \qquad \tilde{A}_{ij} \leftarrow \tilde{A}_{ij} - L_{ik}U_{kj}. \tag{4.10}$$

The entry $\tilde{A}_{kk}$ is the k-th *pivot*. The largest product created in step k, which enters both the update and $|L|\,|U|$, is

$$\max_{i,j} |L_{ik}U_{kj}| = \max_{i,j} \frac{\left|\tilde{A}_{ik}\right|\left|\tilde{A}_{kj}\right|}{\left|\tilde{A}_{kk}\right|}. \tag{4.11}$$

The numerator is determined by column k and row k. To keep this quantity small, we want the denominator, the pivot, to be as large as possible.

4.3 Partial Pivoting

We must pivot when $\tilde{A}_{kk} = 0$, since otherwise the algorithm cannot continue. Even when $\tilde{A}_{kk} \neq 0$, we want to avoid *small* pivots, which make the products $L_{ik}U_{kj}$ explode.

Definition 17 (Partial (Row) Pivoting)

At step k, *partial pivoting* chooses the pivot to be the entry of largest absolute value in column k, on or below the diagonal:

$$r = \operatorname*{argmax}_{i\geq k} \left|\tilde{A}_{ik}\right|. \tag{4.12}$$

It swaps rows r and k with a permutation matrix P_k and then eliminates with L_k. The sequence $A \xrightarrow{L_k} \cdots$ becomes $A \xrightarrow{P_k} \cdot \xrightarrow{L_k} \cdots$.

Because the pivot is the largest entry in its column, every multiplier satisfies

$$|L_{ik}| = \frac{\left|\tilde{A}_{ik}\right|}{\left|\tilde{A}_{kk}\right|} \leq 1, \tag{4.13}$$

so $|L_{ik}U_{kj}| \leq |U_{kj}|$: no update exceeds the size of the current pivot row.

4.3.1 Moving the Permutations Together

With row swaps, elimination becomes

$$L_{m-1}P_{m-1}\cdots L_2P_2L_1P_1A = U, \tag{4.14}$$

where L's and P's alternate, so we cannot directly write $PA = LU$. We need to move all permutations to the right.

Proposition 18 (Commuting a Permutation Past an Elimination)

Let $\ell > k$ and let P_ℓ swap rows ℓ and i for some $i \geq \ell$. Swapping twice undoes the swap, so $P_\ell^{-1} = P_\ell$. Then

$$P_\ell L_k = (P_\ell L_k P_\ell)P_\ell, \qquad P_\ell L_k P_\ell = I - (P_\ell \vec{\ell}_k)\vec{e}_k^*, \tag{4.15}$$

so $P_\ell L_k P_\ell$ is again an elementary lower triangular matrix, obtained from L_k by swapping two of its multipliers.

Proof. The first identity holds because $P_\ell P_\ell = I$. For the second, substitute $L_k = I - \vec{\ell}_k\vec{e}_k^*$ and expand:

$$P_\ell(I - \vec{\ell}_k\vec{e}_k^*)P_\ell = P_\ell P_\ell - (P_\ell\vec{\ell}_k)(\vec{e}_k^*P_\ell) = I - (P_\ell\vec{\ell}_k)(\vec{e}_k^*P_\ell). \tag{4.16}$$

Multiplying the row vector $\vec{e}_k^*$ on the right by P_ℓ swaps its entries ℓ and i. Both entries are 0, since $\ell, i > k$, so nothing changes: $\vec{e}_k^*P_\ell = \vec{e}_k^*$. This gives $P_\ell L_k P_\ell = I - (P_\ell\vec{\ell}_k)\vec{e}_k^*$.

Finally, $P_\ell\vec{\ell}_k$ is $\vec{\ell}_k$ with entries ℓ and i swapped. Its first k entries are still 0, so $P_\ell L_k P_\ell$ is again elementary lower triangular. □

Using Proposition 18 repeatedly to push every P to the right,

$$L_{m-1}P_{m-1}\cdots L_1P_1 = L'_{m-1}\cdots L'_1\,P_{m-1}\cdots P_1, \qquad L'_k \coloneqq P_{m-1}\cdots P_{k+1}L_kP_{k+1}\cdots P_{m-1}, \tag{4.17}$$

and each L'_k is still elementary lower triangular. With $P \coloneqq P_{m-1}\cdots P_1$ and $L \coloneqq (L'_{m-1}\cdots L'_1)^{-1}$, we obtain

$$PA = LU. \tag{4.18}$$

4.4 The In-Place Algorithm with Partial Pivoting

How do we compute this in place, up to the permutation? Suppose k steps are done:

$$L_k\cdots L_1PA = U_k, \tag{4.19}$$

where the L_j are already the "permutation-adjusted" versions and U_k is the current working matrix. In storage, U_k occupies the upper triangle and the trailing block, and the multipliers of $L_1, \ldots, L_k$ sit in the strictly lower part of the first k columns. Step $k+1$ swaps rows and then eliminates:

$$L_{k+1}P_{k+1}L_k\cdots L_1PA = L_{k+1}P_{k+1}U_k. \tag{4.20}$$

Moving P_{k+1} to the right,

$$L_{k+1}\underbrace{L'_k\cdots L'_1}_{\text{multipliers permuted by } P_{k+1}}\underbrace{P_{k+1}P}_{P'}A = L_{k+1}\underbrace{P_{k+1}U_k}_{U'_k} = U_{k+1}. \tag{4.21}$$

Reading term by term:

- $L'_j = P_{k+1} L_j P_{k+1}$ only permutes the stored multipliers;
- $P' = P_{k+1} P$ swaps the same two rows of the accumulated permutation;
- $U'_k = P_{k+1} U_k$ swaps the same two rows of the working matrix.

Since the multipliers and U_k live in the same array, *swapping entire rows (columns* 1 *through* m*)* performs all three updates at once, after which we eliminate as usual.

Algorithm 6 LU factorization with partial pivoting, in place.

$P \leftarrow I$
for $k = 1, \dots, m$ **do**
  $r \leftarrow \operatorname{argmax}_{i \geq k} |A_{ik}|$
  swap rows r and k of A (all columns $1, \dots, m$) and of P
  for $i = k+1, \dots, m$ **do**
    $A_{ik} \leftarrow A_{ik} / A_{kk}$
  for $j = k+1, \dots, m$ **do**
    for $i = k+1, \dots, m$ **do**
      $A_{ij} \leftarrow A_{ij} - A_{ik} A_{kj}$

Some remarks:

- In Julia, `argmax(abs.(A[k:end, k]))` returns an index into the subvector, so one adds $k-1$ to recover the global row index.
- The row swap runs over all columns $1, \dots, m$, not just $k, \dots, m$: the first $k-1$ columns hold multipliers, and swapping them is exactly $L'_j = P_{k+1} L_j P_{k+1}$.
- On exit the strictly lower part of A holds L, the upper part holds U, and $PA = LU$.

4.5 (SUPPLEMENT) Further Topics

4.5.1 Storing P and the Cost of Pivoting

In practice P is not stored as a matrix but as a permutation vector of length m (LAPACK's `ipiv`), so each swap exchanges two integers. The search for pivots costs only $O(m^2)$ comparisons on top of the $\frac{2}{3}m^3$ flops. In column-major storage, however, a row swap is a strided access, and its cost is not negligible.

4.5.2 Growth Factor

Definition 19 (Growth Factor)

The *growth factor* of Gaussian elimination on A is

$$\rho := \frac{\max_{i,j} |U_{ij}|}{\max_{i,j} |A_{ij}|}. \tag{4.22}$$

Why it matters. With partial pivoting $|L_{ik}| \le 1$, so each entry of $|L|\,|U|$ is a sum of at most m terms, each at most $\max|U_{ij}| = \rho \max|A_{ij}|$:

$$(|L|\,|U|)_{ij} \le m\rho \max_{i,j} |A_{ij}| \qquad \Longrightarrow \qquad |E_{ij}| \le m^2 \rho\, \varepsilon_{\text{mach}} \max_{i,j} |A_{ij}| \tag{4.23}$$

by Theorem 16. So partial pivoting is backward stable exactly when ρ stays small.

How large can ρ be? Each step updates $\tilde{A}_{ij} \leftarrow \tilde{A}_{ij} - L_{ik}U_{kj}$ with $|L_{ik}| \le 1$, so the largest entry can at most double. After $m-1$ steps,

$$\rho \le 2^{m-1}. \tag{4.24}$$

This bound is attained by the matrix with 1 on the diagonal, -1 below it, and 1 in the last column (Example 20, Homework 2): no rows are swapped, and the last column of U is $1, 2, 4, \ldots, 2^{m-1}$. For $m = 8$, Julia's `lu` gives $\rho = 128 = 2^7$.

In practice. Such matrices are extremely rare. For random matrices ρ grows slowly; with `randn` (median of 5 runs) we measured $\rho \approx 2, 4, 8, 19$ for $m = 16, 64, 256, 1024$. This is why partial pivoting is used as a stable algorithm in practice, although in theory it is not backward stable.

4.5.3 Complete Pivoting

The idea. At step k, *complete pivoting* chooses the pivot to be the entry of largest absolute value in the whole trailing block, $\max_{i,j\ge k} \left|\tilde{A}_{ij}\right|$, and moves it to position (k,k) by swapping two rows *and* two columns. This gives

$$PAQ = LU, \tag{4.25}$$

with a row permutation P and a column permutation Q. Swapping columns reorders the unknowns, so $A\vec{x} = \vec{b}$ is solved as $LU(Q^\top \vec{x}) = P\vec{b}$.

Benefit. Every entry of the trailing block is now at most the pivot, so the growth is much better controlled. Wilkinson showed a bound on ρ that grows much more slowly than 2^{m-1}, roughly like $m^{(\ln m)/4}$ [7, Chapter 9].

Cost. Step k searches all $(m-k+1)^2$ entries of the trailing block, about $m^3/3$ comparisons in total. That is the same order as the $\frac{2}{3}m^3$ flops, and the column swaps add strided memory access. Since partial pivoting is almost always good enough, complete pivoting is rarely used.

Example. We ran complete pivoting (a short Julia implementation) on the worst-case matrix above and on random matrices, and compared with partial pivoting:

Matrix	ρ, partial pivoting	ρ, complete pivoting
worst case, $m = 64$	$2^{63} \approx 9 \cdot 10^{18}$	2
`randn`, $m = 64$	4.2	1.9
`randn`, $m = 256$	7.8	3.6

On the worst-case matrix, complete pivoting removes the growth entirely. On random matrices both are small, which is why the extra cost of complete pivoting is rarely worth it.

A compromise: rook pivoting. Search alternately along a column and a row until the entry is the largest in both. This is cheap and controls growth much better [7, Chapter 9].

Lecture 5

Least Squares Problems, Gram–Schmidt, and QR

Supplementary reading:

- [1] Lectures 6, 7, 8, 11, and 22.
- [2] Sections 3.1–3.3.

5.1 Is Partial Pivoting Enough?

Lecture 4 bounded the backward error of LU by $m\varepsilon_{\text{mach}} \left|\tilde{L}\right| \left|\tilde{U}\right|$ and introduced partial pivoting to keep the multipliers bounded, $|L_{ij}| \leq 1$; commuting the permutations past the elimination matrices then gave $PA = LU$ and an in-place algorithm. Rows and columns can also be pivoted simultaneously (*complete pivoting*), giving $PA\tilde{P} = LU$ with a column permutation $\tilde{P}$. In practice, partial pivoting is usually enough. In theory, however, it can be *unstable* when m is large.

Example 20 (Exponential Growth Under Partial Pivoting)**.** Consider

$$A = \begin{bmatrix} 1 & & & 1 \\ -1 & 1 & & 1 \\ -1 & -1 & 1 & 1 \\ -1 & -1 & -1 & 1 \end{bmatrix}. \tag{5.1}$$

In every column the candidate pivots have the same absolute value, $|1| = |-1|$, and `argmax` returns the first, so partial pivoting never swaps rows. Tracking the in-place storage (multipliers below the diagonal),

$$A \rightsquigarrow \begin{bmatrix} 1 & & & 1 \\ -1 & 1 & & 2 \\ -1 & -1 & 1 & 2 \\ -1 & -1 & -1 & 2 \end{bmatrix} \rightsquigarrow \begin{bmatrix} 1 & & & 1 \\ -1 & 1 & & 2 \\ -1 & -1 & 1 & 4 \\ -1 & -1 & -1 & 4 \end{bmatrix} \rightsquigarrow \begin{bmatrix} 1 & & & 1 \\ -1 & 1 & & 2 \\ -1 & -1 & 1 & 4 \\ -1 & -1 & -1 & 8 \end{bmatrix}. \tag{5.2}$$

The multiplier at step k is -1, i.e., row k is added to every row below it. Row k is zero in columns $k+1$

through $m-1$, so the middle part never changes, but the last column doubles at every step. The result is

$$L = \begin{bmatrix} 1 & & & \\ -1 & 1 & & \\ -1 & -1 & 1 & \\ -1 & -1 & -1 & 1 \end{bmatrix}, \qquad U = \begin{bmatrix} 1 & & & 1 \\ & 1 & & 2 \\ & & 1 & 4 \\ & & & 8 \end{bmatrix}. \tag{5.3}$$

For general m, $U_{mm} = 2^{m-1}$, so $|L|\,|U| \gtrsim 2^m$ and the error bound $m\varepsilon_{\text{mach}}\,|L|\,|U|$ can reach $m2^m\varepsilon_{\text{mach}}$. In double precision, $2^{m-1}\varepsilon_{\text{mach}} \approx 1$ already for $m \approx 53$, so all accuracy can be lost.

This is the worst case $\rho = 2^{m-1}$ of the growth factor mentioned in Lecture 4. Fortunately, such matrices are extremely rare in practice.

5.2 Non-Square Systems

So far we have only considered $A \in \mathbb{K}^{m\times n}$ with $m = n$. If $m \neq n$, there are two cases.

Overdetermined systems ($m > n$). A is tall: there are more equations than unknowns. Then $\mathcal{R}(A) \neq \mathbb{K}^m$, since n columns span at most an n-dimensional subspace. Hence for a general $\vec{b}$ the equation $A\vec{x} = \vec{b}$ *does not have a solution*. Such a system is called an *overdetermined linear system*.

Underdetermined systems ($m < n$). A is wide: there are more unknowns than equations. Then $\mathcal{N}(A) \neq \{\vec{0}\}$, since the rank is at most $m < n$. Hence even when a solution exists, it is *never unique*: adding any vector of the null space to a solution gives another solution. Such a system is called an *underdetermined linear system*.

5.3 Least Squares Problems

Overdetermined systems usually have no solution, and underdetermined systems have many. In both cases we replace the equation by a minimization problem:

$$\text{overdetermined:} \quad \min_{\vec{x}} \left\| A\vec{x} - \vec{b} \right\|, \qquad \text{underdetermined:} \quad \min_{\vec{x}\,:\,A\vec{x}=\vec{b}} \|\vec{x}\|. \tag{5.4}$$

In the overdetermined case we take the $\vec{x}$ that comes closest to satisfying the equations; in the underdetermined case we take the smallest of all solutions.

Which norm should we use? There are many choices, depending on the application. In linear regression, for example, the norm in the overdetermined case measures the *fitting error*, and in the underdetermined case it measures how *complex* the model is.

The key fact: the solution depends linearly on $\vec{b}$ *only if the norm comes from an inner product*, $\|\cdot\| = \sqrt{\langle\cdot,\cdot\rangle}$. This case is the *least squares problem*, and we can solve it with linear algebra.

Example 21 (Mean Versus Median)**.** Let $A = \begin{bmatrix} 1 & 1 & 1 \end{bmatrix}^\top$. Then $\min_x \left\| Ax - \vec{b} \right\|$ looks for the single number x closest to all of b_1, b_2, b_3.

- In the 2-norm, the best x is the mean $\frac{1}{3}(b_1 + b_2 + b_3)$, which is linear in $\vec{b}$.
- In the 1-norm, the best x is the median, which is not linear in $\vec{b}$.

5.4 The Geometry of Least Squares


For $m = n$, solving $A\vec{x} = \vec{b}$ means finding a linear combination of the columns of A that equals $\vec{b}$ exactly:

$$A\vec{x} = x_1\vec{a}_1 + x_2\vec{a}_2 + \cdots + x_n\vec{a}_n = \vec{b}. \tag{5.5}$$

For $m > n$, consider

$$\min_{\vec{x}} \left\| A\vec{x} - \vec{b} \right\|_2^2. \tag{5.6}$$

The columns of A span only an n-dimensional subspace $\mathcal{R}(A)$ of $\mathbb{K}^m$, and $\vec{b}$ generally does not lie in it. As $\vec{x}$ varies, $A\vec{x}$ ranges over this subspace, and the point closest to $\vec{b}$ is the *orthogonal projection* of $\vec{b}$ onto $\mathcal{R}(A)$: dropping a perpendicular from $\vec{b}$ to the subspace, its foot is the optimal $A\vec{x}$, and the residual $A\vec{x} - \vec{b}$ is orthogonal to the whole subspace.



Writing "the residual is orthogonal to every column of A" as an equation gives the *normal equations*:

$$A^*(A\vec{x} - \vec{b}) = \vec{0} \quad \Longleftrightarrow \quad A^*A\vec{x} = A^*\vec{b}. \tag{5.7}$$

Orthonormal columns. If the columns of A are orthonormal, then $A^*A = I$ and the normal equations give the solution directly:

$$\vec{x} = A^*\vec{b}, \qquad \text{i.e.,} \qquad x_i = \left\langle \vec{a}_i, \vec{b} \right\rangle. \tag{5.8}$$

Since $\vec{a}_i$ is a unit vector, x_i is how much of $\vec{b}$ points along the i-th column. The columns are orthogonal, so each x_i is computed on its own, and $A\vec{x} = \sum_i x_i\vec{a}_i$ is the projection of $\vec{b}$ onto $\mathcal{R}(A)$. Without orthogonality the coefficients depend on each other, and we must solve $A^*A\vec{x} = A^*\vec{b}$.

Example 22 (Projection onto a Coordinate Plane)**.** For

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix} \tag{5.9}$$

we get $\vec{x} = (b_1, b_2)^\top$ and $A\vec{x} = (b_1, b_2, 0)^\top$, the closest point of the xy-plane to $\vec{b}$. The residual $(0, 0, b_3)^\top$ is orthogonal to the plane.

This motivates the next section: since orthonormal columns make least squares easy, we first *make* the columns orthonormal.

5.5 Orthogonalization: The QR Factorization

What if the columns of A are not orthonormal? The idea is to find an *orthonormal basis* of the space spanned by the columns of A and to transfer the problem to that basis. In matrix form,

$$A = QR, \tag{5.10}$$

where $Q \in \mathbb{K}^{m\times n}$ has orthonormal columns and $R \in \mathbb{K}^{n\times n}$ is invertible (it will turn out to be upper triangular). The least squares solution is then

$$\vec{x} = R^{-1}Q^*\vec{b}. \tag{5.11}$$

Both steps have a clear meaning:

- $Q^*\vec{b}$ expresses $\vec{b}$ in terms of the columns of Q (the coordinates of the projection, exactly as in the orthonormal case);
- $A = QR$ says that R expresses the columns of A as combinations of the columns of Q, so R^{-1} converts coefficients with respect to Q back into coefficients with respect to A.

Algebraically, in the normal equations (5.7) we have $A^*A = R^*Q^*QR = R^*R$ and $A^*\vec{b} = R^*Q^*\vec{b}$; cancelling the invertible R^* leaves $R\vec{x} = Q^*\vec{b}$.

5.6 Classical Gram–Schmidt

The idea. Process the columns of A one at a time. When we reach $\vec{a}_j$, the vectors $\vec{q}_1, \dots, \vec{q}_{j-1}$ are already orthonormal. To get $\vec{q}_j$:

1. subtract from $\vec{a}_j$ its components along $\vec{q}_1, \dots, \vec{q}_{j-1}$;
2. normalize what is left.

We first make $\vec{v}_j$ orthogonal to the previous vectors and normalize afterwards. The component of $\vec{a}_j$ along $\vec{v}_i$ is $\frac{\langle \vec{v}_i, \vec{a}_j\rangle}{\langle \vec{v}_i, \vec{v}_i\rangle}\vec{v}_i$, so

$$\begin{aligned}
\vec{v}_1 &= \vec{a}_1, \\
\vec{v}_2 &= \vec{a}_2 - \frac{\langle \vec{v}_1, \vec{a}_2\rangle}{\langle \vec{v}_1, \vec{v}_1\rangle}\vec{v}_1, \\
\vec{v}_3 &= \vec{a}_3 - \frac{\langle \vec{v}_1, \vec{a}_3\rangle}{\langle \vec{v}_1, \vec{v}_1\rangle}\vec{v}_1 - \frac{\langle \vec{v}_2, \vec{a}_3\rangle}{\langle \vec{v}_2, \vec{v}_2\rangle}\vec{v}_2, \\
&\vdots \\
\vec{v}_j &= \vec{a}_j - \sum_{i=1}^{j-1} \frac{\langle \vec{v}_i, \vec{a}_j\rangle}{\langle \vec{v}_i, \vec{v}_i\rangle}\vec{v}_i,
\end{aligned} \tag{5.12}$$

and finally $\vec{q}_j = \vec{v}_j / \|\vec{v}_j\|_2$. Since $\vec{q}_i = \vec{v}_i / \|\vec{v}_i\|_2$, each term equals $\langle \vec{q}_i, \vec{a}_j\rangle \vec{q}_i$, the component of $\vec{a}_j$ along $\vec{q}_i$.

Where R comes from. Write each term of (5.12) as $\langle \vec{q}_i, \vec{a}_j\rangle \vec{q}_i$ and name the coefficients:

$$r_{ij} = \langle \vec{q}_i, \vec{a}_j\rangle \quad (i < j), \qquad r_{jj} = \|\vec{v}_j\|_2, \qquad r_{ij} = 0 \quad (i > j). \tag{5.13}$$

Since $\vec{v}_j = r_{jj}\vec{q}_j$, solving (5.12) for $\vec{a}_j$ gives

$$\vec{a}_j = r_{1j}\vec{q}_1 + \cdots + r_{j-1,j}\vec{q}_{j-1} + r_{jj}\vec{q}_j = \sum_{i=1}^{j} r_{ij}\vec{q}_i, \tag{5.14}$$

that is, column j of A is Q times column j of R. For all columns at once,

$$\underbrace{\begin{bmatrix}\vec{a}_1 & \vec{a}_2 & \cdots & \vec{a}_n\end{bmatrix}}_{A} = \underbrace{\begin{bmatrix}\vec{q}_1 & \vec{q}_2 & \cdots & \vec{q}_n\end{bmatrix}}_{Q} \underbrace{\begin{bmatrix} r_{11} & r_{12} & \cdots & r_{1n} \\ & r_{22} & \cdots & r_{2n} \\ & & \ddots & \vdots \\ & & & r_{nn}\end{bmatrix}}_{R}. \tag{5.15}$$

R is upper triangular because $\vec{a}_j$ involves only $\vec{q}_1, \ldots, \vec{q}_j$, and invertible because $r_{jj} > 0$. Entry r_{ij} $(i < j)$ is the amount of $\vec{q}_i$ in $\vec{a}_j$; r_{jj} is the length of the new part of $\vec{a}_j$.

The algorithm. Computing the sum in (5.12) one term at a time:

Algorithm 7 Classical Gram–Schmidt (CGS).

for $j = 1, \ldots, n$ **do**
  $\vec{v}_j \leftarrow \vec{a}_j$
  for $i = 1, \ldots, j-1$ **do**
    $r_{ij} \leftarrow \langle \vec{q}_i, \vec{a}_j \rangle$ ▷ inner product with the *original* column
    $\vec{v}_j \leftarrow \vec{v}_j - r_{ij}\vec{q}_i$
  $r_{jj} \leftarrow \|\vec{v}_j\|_2$
  $\vec{q}_j \leftarrow \vec{v}_j / r_{jj}$

Theorem 23 (Gram–Schmidt)

Let $\vec{a}_1, \ldots, \vec{a}_n \in \mathbb{K}^m$ $(m \geq n)$ be linearly independent. Then the vectors $\vec{q}_1, \ldots, \vec{q}_n$ produced by Algorithm 7 form an orthonormal basis of $\operatorname{span}(\vec{a}_1, \ldots, \vec{a}_n)$, and $\operatorname{span}(\vec{q}_1, \ldots, \vec{q}_j) = \operatorname{span}(\vec{a}_1, \ldots, \vec{a}_j)$ for every j.

Proof. By induction on j: $\vec{q}_1, \ldots, \vec{q}_j$ are orthonormal and span $\operatorname{span}(\vec{a}_1, \ldots, \vec{a}_j)$. *Base case:* $\vec{a}_1 \neq \vec{0}$ by linear independence, so $\vec{q}_1 = \vec{a}_1 / \|\vec{a}_1\|_2$ is a unit vector spanning the same line.

Inductive step. Assume the claim for j, and let $\vec{v}_{j+1} = \vec{a}_{j+1} - \sum_{i=1}^{j} \langle \vec{q}_i, \vec{a}_{j+1} \rangle \vec{q}_i$.

1. *Orthogonality.* For every $k \leq j$,

$$\langle \vec{q}_k, \vec{v}_{j+1} \rangle = \langle \vec{q}_k, \vec{a}_{j+1} \rangle - \sum_{i=1}^{j} \langle \vec{q}_i, \vec{a}_{j+1} \rangle \underbrace{\langle \vec{q}_k, \vec{q}_i \rangle}_{\delta_{ki}} = \langle \vec{q}_k, \vec{a}_{j+1} \rangle - \langle \vec{q}_k, \vec{a}_{j+1} \rangle = 0. \tag{5.16}$$

2. *Nonzero.* If $\vec{v}_{j+1} = \vec{0}$, then $\vec{a}_{j+1} \in \operatorname{span}(\vec{q}_1, \ldots, \vec{q}_j) = \operatorname{span}(\vec{a}_1, \ldots, \vec{a}_j)$, contradicting linear independence. So $\vec{q}_{j+1} = \vec{v}_{j+1} / \|\vec{v}_{j+1}\|_2$ is defined, and $\vec{q}_1, \ldots, \vec{q}_{j+1}$ are orthonormal.

3. *Span.* $\vec{q}_{j+1} \in \operatorname{span}(\vec{a}_1, \ldots, \vec{a}_{j+1})$, and $\vec{q}_1, \ldots, \vec{q}_{j+1}$ are $j+1$ orthonormal, hence independent, vectors in this $(j+1)$-dimensional space, so they span it. □

Example 24 (CGS Loses Orthogonality)**.** Consider the matrix

$$A = \begin{bmatrix} 1 & 1 & 1 \\ \varepsilon & 0 & 0 \\ 0 & \varepsilon & 0 \\ 0 & 0 & \varepsilon \end{bmatrix}, \tag{5.17}$$

with ε small enough that $1 + \varepsilon^2 = 1$ in floating point, i.e., $\varepsilon < \sqrt{u} \approx 10^{-8}$, for example $\varepsilon = 10^{-10}$.

Step 1. $\vec{v}_1 = \vec{a}_1 = (1, \varepsilon, 0, 0)^\top$ and $\vec{q}_1 = (1 + \varepsilon^2)^{-1/2}(1, \varepsilon, 0, 0)^\top$, which in floating point is $(1, \varepsilon, 0, 0)^\top$.

Step 2. $r_{12} = \langle \vec{q}_1, \vec{a}_2 \rangle = 1$, so

$$\vec{v}_2 = \vec{a}_2 - r_{12}\vec{q}_1 = \begin{bmatrix} 1 \\ 0 \\ \varepsilon \\ 0 \end{bmatrix} - \begin{bmatrix} 1 \\ \varepsilon \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ -\varepsilon \\ \varepsilon \\ 0 \end{bmatrix}, \qquad \vec{q}_2 = \frac{\vec{v}_2}{\sqrt{2}\,\varepsilon} = \begin{bmatrix} 0 \\ -1/\sqrt{2} \\ 1/\sqrt{2} \\ 0 \end{bmatrix}. \tag{5.18}$$

Step 3. $r_{13} = \langle \vec{q}_1, \vec{a}_3 \rangle = 1$ and $r_{23} = \langle \vec{q}_2, \vec{a}_3 \rangle = 0$, so

$$\vec{v}_3 = \vec{a}_3 - r_{13}\vec{q}_1 - r_{23}\vec{q}_2 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ \varepsilon \end{bmatrix} - \begin{bmatrix} 1 \\ \varepsilon \\ 0 \\ 0 \end{bmatrix} - 0 \cdot \vec{q}_2 = \begin{bmatrix} 0 \\ -\varepsilon \\ 0 \\ \varepsilon \end{bmatrix}, \qquad \vec{q}_3 = \frac{\vec{v}_3}{\sqrt{2}\,\varepsilon} = \begin{bmatrix} 0 \\ -1/\sqrt{2} \\ 0 \\ 1/\sqrt{2} \end{bmatrix}. \tag{5.19}$$

The result

$$Q = \begin{bmatrix} 1 & 0 & 0 \\ \varepsilon & -1/\sqrt{2} & -1/\sqrt{2} \\ 0 & 1/\sqrt{2} & 0 \\ 0 & 0 & 1/\sqrt{2} \end{bmatrix} \tag{5.20}$$

is *far from orthogonal*: $\langle \vec{q}_2, \vec{q}_3 \rangle = \frac{1}{2}$, so the angle between them is $60°$ rather than $90°$.

5.7 Modified Gram–Schmidt

What went wrong. Look at step 3 of Example 24.

- First CGS removes the $\vec{q}_1$-component: $\vec{a}_3 - r_{13}\vec{q}_1 = (0, -\varepsilon, 0, \varepsilon)^\top$. Two vectors of size 1 cancel, and only a vector of size ε is left. So any error of size ε is now as large as what is left.
- Such an error is there: the computed $\vec{q}_1$ and $\vec{q}_2$ are not exactly orthogonal, $\langle \vec{q}_1, \vec{q}_2 \rangle = -\varepsilon/\sqrt{2}$.
- Next CGS removes the $\vec{q}_2$-component, but it measures this component in the *original* column: $r_{23} = \langle \vec{q}_2, \vec{a}_3 \rangle = 0$. What we actually need to remove is the $\vec{q}_2$-component of *what is left*:

$$r_{23} = \langle \vec{q}_2, \vec{a}_3 - r_{13}\vec{q}_1 \rangle = \frac{\varepsilon}{\sqrt{2}} \neq 0. \tag{5.21}$$

The two values differ by $-r_{13}\langle \vec{q}_2, \vec{q}_1 \rangle$, which is exactly the orthogonality error. In exact arithmetic it is 0 and the two agree; in floating point it is not.

Notation. To write the fix, we name the intermediate values of $\vec{v}_j$ in Algorithm 7: $\vec{v}_j^{(i)}$ is $\vec{v}_j$ after i subtractions,

$$\vec{v}_j^{(0)} = \vec{a}_j, \qquad \vec{v}_j^{(i)} = \vec{v}_j^{(i-1)} - r_{ij}\vec{q}_i, \qquad \vec{v}_j^{(j-1)} = \vec{v}_j. \tag{5.22}$$

For example, $\vec{v}_3^{(1)} = \vec{a}_3 - r_{13}\vec{q}_1$ is the "what is left" above.

The fix. Change a single line: compute r_{ij} from the *partially orthogonalized* vector instead of the original column.

$$\text{CGS:} \quad r_{ij} = \left\langle \vec{q}_i, \vec{v}_j^{(0)} \right\rangle = \langle \vec{q}_i, \vec{a}_j \rangle, \qquad \text{MGS:} \quad r_{ij} = \left\langle \vec{q}_i, \vec{v}_j^{(i-1)} \right\rangle. \tag{5.23}$$

Everything else, including the update $\vec{v}_j^{(i)} = \vec{v}_j^{(i-1)} - r_{ij}\vec{q}_i$, stays the same.

To compare with CGS directly, we write MGS in the same unnormalized form. MGS subtracts one direction at a time, and each inner product uses the vector *as it is now*:

$$\begin{aligned}
\vec{v}_1 &= \vec{a}_1, \\
\vec{v}_2 &= \vec{a}_2 - \frac{\langle \vec{v}_1, \vec{a}_2 \rangle}{\langle \vec{v}_1, \vec{v}_1 \rangle}\vec{v}_1, \\
\vec{v}_3^{(1)} &= \vec{a}_3 - \frac{\langle \vec{v}_1, \vec{a}_3 \rangle}{\langle \vec{v}_1, \vec{v}_1 \rangle}\vec{v}_1, \\
\vec{v}_3 &= \vec{v}_3^{(1)} - \frac{\langle \vec{v}_2, \vec{v}_3^{(1)} \rangle}{\langle \vec{v}_2, \vec{v}_2 \rangle}\vec{v}_2, \\
&\vdots \\
\vec{v}_j^{(i)} &= \vec{v}_j^{(i-1)} - \frac{\langle \vec{v}_i, \vec{v}_j^{(i-1)} \rangle}{\langle \vec{v}_i, \vec{v}_i \rangle}\vec{v}_i, \qquad i = 1, \dots, j-1,
\end{aligned} \tag{5.24}$$

In terms of the normalized $\vec{q}_i$, this gives the following algorithm. It differs from Algorithm 7 only in line 4.

Algorithm 8 Modified Gram–Schmidt (MGS).

```
1: for j = 1, ..., n do
2:     v_j ← a_j
3:     for i = 1, ..., j − 1 do
4:         r_ij ← ⟨q_i, v_j⟩                ▷ inner product with the updated vector
5:         v_j ← v_j − r_ij q_i
6:     r_jj ← ‖v_j‖_2
7:     q_j ← v_j / r_jj
```

Same in exact arithmetic. Expanding (5.22),

$$\left\langle \vec{q}_i, \vec{v}_j^{(i-1)} \right\rangle = \langle \vec{q}_i, \vec{a}_j \rangle - \sum_{k=1}^{i-1} r_{kj} \langle \vec{q}_i, \vec{q}_k \rangle, \tag{5.25}$$

and the sum vanishes when the $\vec{q}$'s are orthonormal, so CGS and MGS compute the same r_{ij}. The point is that CGS *relies* on $\langle \vec{q}_i, \vec{q}_k \rangle = 0$, which holds only approximately in floating point; as soon as cancellation occurs, this small error becomes large. MGS does not rely on it: each r_{ij} is the $\vec{q}_i$-component of what is actually left, so errors made in earlier steps are projected out by later ones.

Example 25 (MGS on the Same Matrix)**.** For the matrix of Example 24, $\vec{q}_1$ and $\vec{q}_2$ are computed exactly as in CGS (with one previous vector there is nothing to modify). Step 3 now projects in two stages:

$$\vec{v}_3^{(1)} = \vec{a}_3 - r_{13}\vec{q}_1 = \begin{bmatrix}1\\0\\0\\\varepsilon\end{bmatrix} - \begin{bmatrix}1\\\varepsilon\\0\\0\end{bmatrix} = \begin{bmatrix}0\\-\varepsilon\\0\\\varepsilon\end{bmatrix}. \tag{5.26}$$

Now $r_{23} = \left\langle \vec{q}_2, \vec{v}_3^{(1)} \right\rangle = \varepsilon/\sqrt{2}$ is no longer zero, and

$$\vec{v}_3^{(2)} = \vec{v}_3^{(1)} - r_{23}\vec{q}_2 = \begin{bmatrix}0\\-\varepsilon\\0\\\varepsilon\end{bmatrix} - \frac{\varepsilon}{\sqrt{2}}\begin{bmatrix}0\\-1/\sqrt{2}\\1/\sqrt{2}\\0\end{bmatrix} = \begin{bmatrix}0\\-\varepsilon/2\\-\varepsilon/2\\\varepsilon\end{bmatrix}, \qquad \vec{q}_3 = \frac{\vec{v}_3^{(2)}}{\varepsilon\sqrt{3/2}} = \begin{bmatrix}0\\-1/\sqrt{6}\\-1/\sqrt{6}\\\sqrt{2/3}\end{bmatrix}. \tag{5.27}$$

The result is

$$Q = \begin{bmatrix}1 & 0 & 0\\ \varepsilon & -1/\sqrt{2} & -1/\sqrt{6}\\ 0 & 1/\sqrt{2} & -1/\sqrt{6}\\ 0 & 0 & \sqrt{2/3}\end{bmatrix}. \tag{5.28}$$

Now $\langle \vec{q}_2, \vec{q}_3\rangle = \frac{1}{\sqrt{12}} - \frac{1}{\sqrt{12}} = 0$, and the remaining inner products $\langle \vec{q}_1, \vec{q}_2\rangle = -\varepsilon/\sqrt{2}$ and $\langle \vec{q}_1, \vec{q}_3\rangle = -\varepsilon/\sqrt{6}$ are $O(\varepsilon)$: the columns are *nearly orthogonal*.

5.8 (SUPPLEMENT) Further Topics

5.8.1 The Projector View

Trefethen and Bau [1, Lecture 8] write both algorithms with projectors, which shows the difference in one line.

Two kinds of projectors. Let $Q_{j-1} = \begin{bmatrix}\vec{q}_1 & \cdots & \vec{q}_{j-1}\end{bmatrix}$.

- $P_i\vec{x} := \vec{x} - \langle \vec{q}_i, \vec{x}\rangle\, \vec{q}_i$ removes the $\vec{q}_i$-component of $\vec{x}$: *one* direction.
- $(I - Q_{j-1}Q_{j-1}^*)\vec{x} = \vec{x} - \sum_{i<j} \langle \vec{q}_i, \vec{x}\rangle\, \vec{q}_i$ removes the components along $\vec{q}_1, \ldots, \vec{q}_{j-1}$: *all* of them at once.

Then the two algorithms compute

$$\text{CGS:}\quad \vec{v}_j = (I - Q_{j-1}Q_{j-1}^*)\,\vec{a}_j, \qquad \text{all directions at once, using } \vec{a}_j, \tag{5.29}$$

$$\text{MGS:}\quad \vec{v}_j = P_{j-1}\cdots P_2P_1\,\vec{a}_j, \qquad \text{one direction at a time, using the current vector.} \tag{5.30}$$

If the $\vec{q}_i$ are exactly orthonormal, both give the same result.

Where they differ. Take $j = 3$ and apply P_1, then P_2:

$$P_1\vec{a}_3 = \vec{a}_3 - \langle \vec{q}_1, \vec{a}_3\rangle\, \vec{q}_1, \tag{5.31}$$

$$P_2P_1\vec{a}_3 = \vec{a}_3 - \langle \vec{q}_1, \vec{a}_3\rangle\, \vec{q}_1 - \langle \vec{q}_2, \vec{a}_3 - \langle \vec{q}_1, \vec{a}_3\rangle\, \vec{q}_1\rangle\, \vec{q}_2 \tag{5.32}$$

$$= \underbrace{\vec{a}_3 - \langle \vec{q}_1, \vec{a}_3\rangle\, \vec{q}_1 - \langle \vec{q}_2, \vec{a}_3\rangle\, \vec{q}_2}_{\text{CGS}} + \langle \vec{q}_2, \vec{q}_1\rangle\, \langle \vec{q}_1, \vec{a}_3\rangle\, \vec{q}_2. \tag{5.33}$$

So MGS equals CGS plus one extra term. That term is zero when $\vec{q}_1 \perp \vec{q}_2$. When rounding has made $\vec{q}_1$ and $\vec{q}_2$ slightly non-orthogonal, the extra term is exactly the correction that CGS misses.

Check on the example. In Example 24, $\langle \vec{q}_2, \vec{q}_1 \rangle = -\varepsilon/\sqrt{2}$ and $\langle \vec{q}_1, \vec{a}_3 \rangle = 1$, so the MGS vector should be the CGS vector plus $-\frac{\varepsilon}{\sqrt{2}}\vec{q}_2$:

$$\underbrace{\begin{bmatrix} 0 \\ -\varepsilon \\ 0 \\ \varepsilon \end{bmatrix}}_{\text{CGS}} - \frac{\varepsilon}{\sqrt{2}} \begin{bmatrix} 0 \\ -1/\sqrt{2} \\ 1/\sqrt{2} \\ 0 \end{bmatrix} = \underbrace{\begin{bmatrix} 0 \\ -\varepsilon/2 \\ -\varepsilon/2 \\ \varepsilon \end{bmatrix}}_{\text{MGS}}, \tag{5.34}$$

which matches Example 25.

5.8.2 How Much Orthogonality Is Lost?

The quantity $\|I - Q^*Q\|$ measures how far the computed Q is from having orthonormal columns. A result of Björck states:

- MGS: $\|I - Q^*Q\| \approx \varepsilon_{\text{mach}}\, \kappa(A)$;
- CGS: the loss can be as bad as $\varepsilon_{\text{mach}}\, \kappa(A)^2$.

The matrix of Example 24 has $\kappa(A) \approx \sqrt{3}/\varepsilon$. With $\varepsilon = 10^{-10}$, $\varepsilon_{\text{mach}}\kappa(A)^2 \approx 7 \cdot 10^4 \gg 1$, so CGS may lose orthogonality completely, while for MGS $\varepsilon_{\text{mach}}\kappa(A) \approx 4 \cdot 10^{-6}$. In Julia we measured $\|I - Q^*Q\| \approx 0.71$ for CGS (with $\langle \vec{q}_2, \vec{q}_3 \rangle = 0.5$) and $\approx 1.2 \cdot 10^{-10}$ for MGS.

5.8.3 Why Not Solve the Normal Equations?

Solving $A^*A\vec{x} = A^*\vec{b}$ looks most direct, but forming A^*A squares the condition number, $\kappa(A^*A) = \kappa(A)^2$ [1, Lecture 11]. For the matrix of Example 24,

$$A^*A = \begin{bmatrix} 1+\varepsilon^2 & 1 & 1 \\ 1 & 1+\varepsilon^2 & 1 \\ 1 & 1 & 1+\varepsilon^2 \end{bmatrix}, \tag{5.35}$$

and in floating point $1 + \varepsilon^2 = 1$. So the computed A^*A is the all-ones matrix, which has rank 1: the information is lost before we even start solving. QR works on A directly and avoids this [2, §3.3].

5.8.4 Householder QR and Reorthogonalization

Even with MGS, how orthogonal Q is depends on $\kappa(A)$. Libraries therefore use *Householder QR* (Lecture 6), whose Q is orthogonal to working precision whatever $\kappa(A)$ is. Another remedy is to *run CGS twice* ("twice is enough"); it parallelizes better than MGS, since the inner products of CGS form one matrix–vector product $Q_{j-1}^*\vec{a}_j$.

Lecture 6

Product View on Error Bounds, Householder QR

Supplementary reading:

- [1] Lectures 8, 10, and 16.
- [2] Section 3.4.

6.1 Why Write Algorithms as Matrix Products?

So far we described LU (Lecture 3) and Gram–Schmidt (Lecture 5) step by step, one row or one vector at a time, and we analyzed the error of LU entry by entry (Lecture 4). This has two drawbacks:

- the analysis has to be redone for every new algorithm;
- it does not show *why* some algorithms are stable and others are not.

In this lecture we write every factorization algorithm as a *product of matrices*: each step multiplies by one matrix. Then we only need to look at that matrix. It turns out that stability depends on one number: the condition number of the matrix used at each step.

6.2 The Product Form of Gram–Schmidt

The analogy with LU. LU *left*-multiplies by a sequence of lower triangular matrices in order to reach an *upper triangular* matrix:

$$L_{m-1} \cdots L_1 A = U. \tag{6.1}$$

Each L_k performs row operations that zero column k below the diagonal. Gram–Schmidt does the opposite: it *right*-multiplies by a sequence of upper triangular matrices in order to reach *orthonormal columns*:

$$A R_1 R_2 \cdots R_n = \hat{Q}. \tag{6.2}$$

Right multiplication performs column operations, and column operations are exactly what Gram–Schmidt does: normalize one column, then subtract its component from the other columns.

The step matrices. We first fix the notation.

- $\vec{a}_j^{(k-1)}$ is the j-th column after $k-1$ steps, i.e., $\vec{a}_j$ with its components along $\vec{q}_1, \dots, \vec{q}_{k-1}$ removed.
- $\vec{a}_j^{(0)} = \vec{a}_j$.

At the start of step k the matrix is

$$\begin{bmatrix} \vec{q}_1 & \cdots & \vec{q}_{k-1} & \vec{a}_k^{(k-1)} & \cdots & \vec{a}_n^{(k-1)} \end{bmatrix}. \tag{6.3}$$

- The first $k-1$ columns are orthonormal.
- The remaining columns are orthogonal to them, but not yet to each other.

Step k uses the numbers

$$r_{kk} \coloneqq \left\| \vec{a}_k^{(k-1)} \right\|, \qquad \vec{q}_k \coloneqq \vec{a}_k^{(k-1)} / r_{kk}, \qquad r_{kj} \coloneqq \left\langle \vec{q}_k, \vec{a}_j^{(k-1)} \right\rangle \quad (j > k). \tag{6.4}$$

The step matrix R_k equals the identity except in row k:

$$R_k = \begin{bmatrix} 1 & & & & & & \\ & \ddots & & & & & \\ & & \frac{1}{r_{kk}} & -\frac{r_{k,k+1}}{r_{kk}} & \cdots & -\frac{r_{kn}}{r_{kk}} & \\ & & & 1 & & & \\ & & & & \ddots & & \\ & & & & & 1 & \end{bmatrix}. \tag{6.5}$$

Right multiplication by R_k performs step k:

$$\begin{bmatrix} \vec{q}_1 & \cdots & \vec{q}_{k-1} & \vec{a}_k^{(k-1)} & \cdots & \vec{a}_n^{(k-1)} \end{bmatrix} R_k = \begin{bmatrix} \vec{q}_1 & \cdots & \vec{q}_k & \vec{a}_{k+1}^{(k)} & \cdots & \vec{a}_n^{(k)} \end{bmatrix}. \tag{6.6}$$

To check this, look at the columns of the product.

- Columns $1, \dots, k-1$ are unchanged.
- Column k is
$$\vec{a}_k^{(k-1)} / r_{kk} = \vec{q}_k.$$
- Column $j > k$ is
$$\vec{a}_j^{(k-1)} - \frac{r_{kj}}{r_{kk}} \vec{a}_k^{(k-1)} = \vec{a}_j^{(k-1)} - r_{kj} \vec{q}_k = \vec{a}_j^{(k)}.$$

Note the factor $1/r_{kk}$ in the off-diagonal entries. It is there because the coefficient multiplies the *unnormalized* column $\vec{a}_k^{(k-1)}$, not $\vec{q}_k$. For example, for $k = 1$,

$$(R_1)_{1j} = -\frac{\langle \vec{q}_1, \vec{a}_j \rangle}{\|\vec{a}_1\|} = -\frac{\langle \vec{a}_1, \vec{a}_j \rangle}{\|\vec{a}_1\|^2}. \tag{6.7}$$

The product form is modified Gram–Schmidt. CGS and MGS differ only in which column the coefficient r_{kj} uses:

- CGS uses the original column: $r_{kj} = \langle \vec{q}_k, \vec{a}_j \rangle$.
- MGS uses the updated column: $r_{kj} = \left\langle \vec{q}_k, \vec{a}_j^{(k-1)} \right\rangle$.

The product form can only be MGS. R_k multiplies the *current* matrix, whose j-th column is already $\vec{a}_j^{(k-1)}$; the original $\vec{a}_j$ has been overwritten. So each step computes $\vec{q}_k$ and right away removes it from all later columns, which is exactly MGS.

The reduced QR factorization. Each R_k is upper triangular, and so is their product:

$$R_1 R_2 \cdots R_n = \hat{R}^{-1} \quad \Longrightarrow \quad A = \hat{Q}\hat{R}. \tag{6.8}$$

Here $\hat{Q} \in \mathbb{K}^{m \times n}$ is the factor of the *reduced QR factorization*: it has only n orthonormal columns, $\hat{Q}^* \hat{Q} = I_n$, but $\hat{Q}\hat{Q}^* \neq I_m$ (it is the orthogonal projector onto $\mathcal{R}(A)$). The factor $\hat{R} \in \mathbb{K}^{n \times n}$ is upper triangular and invertible.

Like R_k, the inverse R_k^{-1} differs from the identity only in row k, and that row has no reciprocals and no minus signs:

$$R_k^{-1} = \begin{bmatrix} 1 & & & & & & \\ & \ddots & & & & & \\ & & r_{kk} & r_{k,k+1} & \cdots & r_{kn} & \\ & & & 1 & & & \\ & & & & \ddots & & \\ & & & & & 1 & \end{bmatrix}. \tag{6.9}$$

The same happens in LU, where L_k^{-1} is L_k with the minus signs removed.

Consequently

$$\hat{R} = R_n^{-1} \cdots R_1^{-1} = \begin{bmatrix} r_{11} & r_{12} & \cdots & r_{1n} \\ & r_{22} & \cdots & r_{2n} \\ & & \ddots & \vdots \\ & & & r_{nn} \end{bmatrix} \tag{6.10}$$

That is, row k of $\hat{R}$ is simply row k of R_k^{-1}. LU works the same way: in $L = L_1^{-1} \cdots L_{m-1}^{-1}$, column k of L is simply column k of L_k^{-1}, i.e., the multipliers of step k (Theorem 12).

6.3 Error Amplification in Products

Consider algorithms of the form

$$A_k = X_k A_{k-1}, \qquad A_0 = A. \tag{6.11}$$

LU ($X_k = L_k$) and Householder QR (X_k orthogonal, below) are of this type; Gram–Schmidt multiplies from the right and is analyzed symmetrically. Every step incurs rounding errors. How stable is such an algorithm?

Forward error of one step. One can show that in floating-point arithmetic

$$\mathrm{fl}(XA) = XA + E + O(\varepsilon_{\mathrm{mach}}^2), \qquad \text{for } \|E\|_2 \le O(\varepsilon_{\mathrm{mach}})\,\|X\|_2\,\|A\|_2\,. \tag{6.12}$$

Note that E is measured against $\|X\|\,\|A\|$, not $\|XA\|$. If XA is small because of cancellation, E can be large relative to the result. This is the same as in LU, where the error is governed by $|L|\,|U|$ rather than $|A|$.

The $O(\varepsilon_{\mathrm{mach}})$ hides a dimension-dependent constant, and $O(\varepsilon_{\mathrm{mach}}^2)$ means we only keep first-order terms.

Rewriting as a backward error. Thus, with $F := X^{-1}E$,

$$\mathrm{fl}(XA) = X(A+F) + O(\varepsilon_{\mathrm{mach}}^2). \tag{6.13}$$

So the computed result is the *exact* product with a perturbed input $A + F$.

The perturbation satisfies

$$\|F\|_2 = \|X^{-1}E\|_2 \le \|X^{-1}\|_2\|E\|_2 \le O(\varepsilon_{\mathrm{mach}})\,\kappa(X)\,\|A\|_2. \tag{6.14}$$

Dividing by $\|A\|_2$ gives the relative backward error:

$$\frac{\|F\|_2}{\|A\|_2} \le O(\varepsilon_{\mathrm{mach}})\,\kappa(X).$$

Theorem 26 (The Transformation Determines the Backward Error)

For one step $A \mapsto XA$ of a product-form algorithm with invertible X, the computed result equals $X(A+F)$ with

$$\frac{\|F\|_2}{\|A\|_2} \le O(\varepsilon_{\mathrm{mach}})\,\kappa(X). \tag{6.15}$$

Two condition numbers must be kept apart.

- $\kappa(X)$ belongs to the *algorithm*. It determines the backward error, i.e., whether the algorithm is stable, and the algorithm designer controls it.
- $\kappa(A)$ belongs to the *problem*. It is fixed by the input and turns backward error into forward error: forward error $\lesssim \kappa(A)\times$ backward error.

Design principle: choose transformations with small condition numbers. Orthogonal matrices have $\kappa = 1$, the best possible.

6.4 Lower Triangularization: The Case of LU

Apply the framework to LU. After k steps of elimination the accumulated transformation is $X = L_k \cdots L_1$, and its inverse is the unit lower triangular matrix of multipliers,

$$L = X^{-1} = \begin{bmatrix} 1 & & & \\ \ell_{21} & 1 & & \\ \vdots & \ddots & \ddots & \\ \ell_{m1} & \cdots & \ell_{m,m-1} & 1 \end{bmatrix}. \tag{6.16}$$

Since $\kappa(X) = \kappa(X^{-1}) = \kappa(L)$, Theorem 26 gives a backward error of about $O(\varepsilon_{\text{mach}})\kappa(L)\,\|A\|_2$. The stability of LU thus comes down to the condition number of L. Partial pivoting chooses the largest entry of each column as pivot and so guarantees $|\ell_{ij}| \le 1$.

$\|L\|$ is under control. Since $|\ell_{ij}| \le 1$,

$$\|L\|_2 \le \|L\|_F \le \sqrt{\frac{m(m+1)}{2}} \le m. \tag{6.17}$$

The middle step holds because L has m ones on the diagonal and $m(m-1)/2$ entries of absolute value at most 1 below it, so $\|L\|_F^2 \le m(m+1)/2$. Here $\|\cdot\|_F$ is the Frobenius norm (Section B.4).

The bound is loose. What matters is that it is polynomial in m, not exponential.

$\|L^{-1}\|$ is not. What about $\|L^{-1}\|_2$? To control $\kappa(L) = \|L\|_2 \|L^{-1}\|_2$, we also need $\|L^{-1}\|$ to be moderate. But $|\ell_{ij}| \le 1$ does *not* guarantee this.

The reason is forward substitution: each row uses the results of all previous rows. So values pile up, and the entries of L^{-1} can grow exponentially.

Example 27 (Exponential Growth of L^{-1})**.** The factor L of Example 20 has -1 everywhere below the diagonal, so $|\ell_{ij}| \le 1$, yet

$$L = \begin{bmatrix} 1 & & & \\ -1 & 1 & & \\ -1 & -1 & 1 & \\ -1 & -1 & -1 & 1 \end{bmatrix}, \qquad L^{-1} = \begin{bmatrix} 1 & & & \\ 1 & 1 & & \\ 2 & 1 & 1 & \\ 4 & 2 & 1 & 1 \end{bmatrix}. \tag{6.18}$$

For general m, $(L^{-1})_{ij} = 2^{i-j-1}$ for $i > j$, and the bottom-left entry is 2^{m-2}, so

$$\left\|L^{-1}\right\|_2 \ge 2^{m-2}, \qquad \kappa(L) \ge 2^{m-2}. \tag{6.19}$$

To see where the doubling comes from, note that computing L^{-1} means solving $L\vec{x} = \vec{b}$. By forward substitution, the i-th equation $x_i - x_1 - \cdots - x_{i-1} = b_i$ gives

$$x_i = b_i + (x_1 + \cdots + x_{i-1}). \tag{6.20}$$

For the first column, $\vec{b} = \vec{e}_1$: $x_1 = 1$, $x_2 = 1$, $x_3 = 1 + 1 = 2$, $x_4 = 1 + 1 + 2 = 4$, and so on. Each entry is the sum of all previous ones, so from the second entry on they double, $x_i = 2^{i-2}$. The other columns are the same pattern shifted down.

Large L^{-1} causes large U. From $A = LU$,

$$U = L^{-1}A, \tag{6.21}$$

so a large $\|L^{-1}\|$ can produce a large U: *large U comes from large L^{-1}*. This connects the condition-number viewpoint with the growth factor of Lectures 4 and 5; they describe the same thing. In Example 20 the last column of A is all ones, so the last column of U is L^{-1} times the all-ones vector, i.e., the row sums of L^{-1}:

$$1 + \sum_{j=1}^{i-1} 2^{i-j-1} = 1 + (2^{i-1} - 1) = 2^{i-1}, \tag{6.22}$$

which is exactly the last column $1, 2, 4, 8, \dots$ of U.

Partial pivoting thus solves only half of the problem: it controls $\|L\|$ but not $\|L^{-1}\|$, and in the worst case LU with partial pivoting can still be unstable. Such matrices are extremely rare, so in practice it is used as a stable algorithm; but to *guarantee* well-conditioned transformations we must switch to orthogonal ones.

6.5 Gram–Schmidt Is Also a Lower Triangularization

The upper triangular R_k is nontrivial only in row k, so its transpose $L_k^\top := R_k$ has L_k nontrivial only in column k: an elementary lower triangular matrix (with diagonal entry $1/r_{kk}$ instead of 1). Then $AR_1 \cdots R_n = \hat{Q}$ reads $AL_1^\top \cdots L_n^\top = \hat{Q}$, and transposing,

$$L_n \cdots L_1 A^\top = \hat{Q}^\top. \tag{6.23}$$

Gram–Schmidt on A is therefore the same as left-multiplying $A^\top$ by a sequence of lower triangular matrices. It too falls into the error-amplification framework, and its stability is likewise governed by the condition numbers of these triangular factors.

6.6 Householder Triangularization

Idea: multiply by orthogonal matrices. LU and Gram–Schmidt have the same defect: each step applies a triangular matrix, whose condition number is not under control. Following the design principle of Section 6.3, we make every step orthogonal instead.

We are looking for unitary matrices Q_k such that each step zeros one column below the diagonal:

$$\begin{bmatrix} \times & \times & \times \\ \times & \times & \times \\ \times & \times & \times \\ \times & \times & \times \\ \times & \times & \times \end{bmatrix} \xrightarrow{Q_1} \begin{bmatrix} \times & \times & \times \\ 0 & \times & \times \\ 0 & \times & \times \\ 0 & \times & \times \\ 0 & \times & \times \end{bmatrix} \xrightarrow{Q_2} \begin{bmatrix} \times & \times & \times \\ 0 & \times & \times \\ 0 & 0 & \times \\ 0 & 0 & \times \\ 0 & 0 & \times \end{bmatrix} \xrightarrow{Q_3} \begin{bmatrix} \times & \times & \times \\ 0 & \times & \times \\ 0 & 0 & \times \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}. \tag{6.24}$$

After n steps,

$$Q_n \cdots Q_1 A = R \quad \Longleftrightarrow \quad A = \underbrace{(Q_n \cdots Q_1)^{-1}}_{Q} R = Q_1^* \cdots Q_n^* R. \tag{6.25}$$

This is the *full QR factorization*: $Q \in \mathbb{K}^{m\times m}$ is a square unitary matrix, and $R \in \mathbb{K}^{m\times n}$ is upper triangular, with $m - n$ zero rows at the bottom. Taking the first n columns of Q and the first n rows of R recovers the reduced QR factorization.

Compare with the two earlier algorithms:

- like LU, Householder multiplies from the *left* and aims for an upper triangular matrix, but with unitary instead of triangular matrices;
- Gram–Schmidt uses triangular matrices to *produce* an orthogonal matrix; Householder uses orthogonal matrices to *produce* a triangular one.

How do we choose the Q_k?

The Householder reflector. Let us begin with the first step. Q_1 must map the first column $\vec{a}_1$ to a multiple of $\vec{e}_1$. Unitary maps preserve the 2-norm, so there are only two possible targets:


$$Q_1 \vec{a}_1 = \pm \|\vec{a}_1\| \, \vec{e}_1. \tag{6.26}$$

Take the $+$ sign first.

Idea: use a reflection. Since $\vec{a}_1$ and $\|\vec{a}_1\| \, \vec{e}_1$ have the same length, they are mirror images of each other. The mirror is the hyperplane through the origin with normal vector

$$\vec{v} \coloneqq \vec{a}_1 - \|\vec{a}_1\| \, \vec{e}_1. \tag{6.27}$$

Reflecting across it sends $\vec{a}_1$ to $\|\vec{a}_1\| \, \vec{e}_1$.



Definition 28 (Householder Reflector)

For a nonzero $\vec{v} \in \mathbb{K}^m$, the *Householder reflector* is

$$F = I - 2\frac{\vec{v}\vec{v}^*}{\vec{v}^*\vec{v}}. \tag{6.28}$$

F flips the component along $\vec{v}$ and keeps the rest:

- $\vec{u} \perp \vec{v} \implies F\vec{u} = \vec{u}$: vectors in the hyperplane stay fixed;
- $\vec{u} \parallel \vec{v} \implies F\vec{u} = -\vec{u}$: the normal direction is flipped.

F is Hermitian, and reflecting twice changes nothing, so $F^2 = I$. Hence $F^*F = F^2 = I$: F is unitary and is its own inverse.

We now check that $Q_1 = I - 2\vec{v}\vec{v}^*/(\vec{v}^*\vec{v})$ does what we want (real case). First,

$$\vec{v}^*\vec{v} = 2\|\vec{a}_1\|^2 - 2\|\vec{a}_1\| a_{11}, \qquad \vec{v}^*\vec{a}_1 = \|\vec{a}_1\|^2 - \|\vec{a}_1\| a_{11} = \tfrac{1}{2}\vec{v}^*\vec{v}, \tag{6.29}$$

so

$$Q_1\vec{a}_1 = \vec{a}_1 - 2\vec{v}\frac{\vec{v}^*\vec{a}_1}{\vec{v}^*\vec{v}} = \vec{a}_1 - \vec{v} = \|\vec{a}_1\|\vec{e}_1. \tag{6.30}$$

Choosing the sign. There are two possible reflections, onto $+\|\vec{a}_1\|\vec{e}_1$ and onto $-\|\vec{a}_1\|\vec{e}_1$. Which one should we choose? Mathematically, there is no preference. In finite precision, we choose the target *farther from* $\vec{a}_1$:

$$Q_1\vec{a}_1 = -\operatorname{sign}(a_{11})\|\vec{a}_1\|\vec{e}_1 \quad \Longrightarrow \quad \vec{v} \propto \operatorname{sign}(a_{11})\|\vec{a}_1\|\vec{e}_1 + \vec{a}_1. \tag{6.31}$$


We take $\operatorname{sign}(0) = 1$.

Why? Suppose $\vec{a}_1$ is nearly parallel to $\vec{e}_1$, with $a_{11} > 0$. With the $+$ sign,


$$\|\vec{v}_+\| = \|\|\vec{a}_1\|\vec{e}_1 - \vec{a}_1\| \ll \|\vec{a}_1\|. \tag{6.32}$$

So $\vec{v}_+$ is the difference of two nearly equal vectors, and its first entry $\|\vec{a}_1\| - a_{11}$ suffers cancellation. The computed $\|\vec{a}_1\|$ has an absolute error of about $\varepsilon_{\text{mach}}\|\vec{a}_1\|$, which is large compared with the short $\vec{v}_+$. So the direction of $\vec{v}_+$, and with it the reflection, is inaccurate.

With $-\operatorname{sign}(a_{11})$, the first entry of $\vec{v}$ is $a_{11} + \operatorname{sign}(a_{11})\|\vec{a}_1\|$, a sum of two numbers of the same sign. There is no cancellation, and $|v_1| \geq \|\vec{a}_1\|$, so $\vec{v}$ is never short.



Reflecting onto $+\|\vec{a}_1\|\vec{e}_1$: $\vec{v}_+$ is short, so its direction is unreliable.



Reflecting onto $-\|\vec{a}_1\|\vec{e}_1$: $\vec{v}$ is long, so its direction is reliable.

Step k: the block form. What about Q_k for $k > 1$? The first $k-1$ rows and columns are finished and must not change. So we use

$$Q_k = \begin{bmatrix} I_{k-1} & 0 \\ 0 & F_k \end{bmatrix}, \tag{6.33}$$

where F_k is the $(m-k+1)\times(m-k+1)$ Householder reflector built as in step 1 from $\vec{x} = A[k{:}m, k]$, the first column of the trailing submatrix $A[k{:}m, k{:}n]$.

Earlier results are preserved:

- the block I_{k-1} leaves the first $k-1$ rows untouched;
- the first $k-1$ columns are already zero from row k down, and F_k maps zero to zero.

Q_k is itself a Householder reflector (with $\vec{v}$ padded by $k-1$ leading zeros), so it is unitary.

Algorithm 9 Householder QR, in place.

for $k = 1, \dots, n$ **do**
  $\vec{x} \leftarrow A[k{:}m, k]$
  $\vec{v} \leftarrow \operatorname{sign}(x_1) \|\vec{x}\|_2 \vec{e}_1 + \vec{x}$
  $\vec{v} \leftarrow \vec{v}/v_1$ ▷ scale so that $v_1 = 1$
  $A[k{:}m, k{:}n] \leftarrow A[k{:}m, k{:}n] - 2\vec{v}\left(\vec{v}^* A[k{:}m, k{:}n]\right)/(\vec{v}^*\vec{v})$
  $A[k{+}1{:}m, k] \leftarrow \vec{v}[2{:}\text{end}]$ ▷ store $\vec{v}$ below the diagonal
return A ▷ upper triangle: R; strictly lower triangle: the vectors $\vec{v}$

Some remarks on the implementation (see also [1, Lecture 10]):

- *F_k is never formed.* Following the parentheses, first compute the row vector $\vec{v}^* A[k{:}m, k{:}n]$, then perform a rank-one update. Each step costs $O((m-k)(n-k))$; forming $\vec{v}\vec{v}^*$ first would cost an order of magnitude more.
- *Scaling $\vec{v}$ by $1/v_1$* makes its first entry 1. The reflector depends only on the direction of $\vec{v}$, so scaling does not change F_k. Since $|v_1| \geq \|\vec{x}\| > 0$ (for $\vec{x} \neq \vec{0}$), this never divides by zero.
- *Storage.* After the update, $A[k{+}1{:}m, k]$ should be zero, which frees exactly enough room for the remaining $m-k$ entries of $\vec{v}$; the leading $v_1 = 1$ is implicit.
- On exit the upper triangle of A is R, with diagonal entries $r_{kk} = -\operatorname{sign}(x_1)\|\vec{x}\|$. The matrix Q is never formed explicitly; it is stored implicitly through the vectors $\vec{v}$ (see Lecture 7).
- $\operatorname{sign}(0)$ must be taken as 1. Otherwise $x_1 = 0$ gives $\vec{v} = \vec{x}$, and the reflection maps $\vec{x}$ to $-\vec{x}$ instead of a multiple of $\vec{e}_1$.

Back into the framework. Each $X_k = Q_k$ is unitary, so $\|Q_k\|_2 = \left\|Q_k^{-1}\right\|_2 = 1$ and $\kappa(Q_k) = 1$ (Appendix B). By Theorem 26,

$$\|F\|_2 \leq O(\varepsilon_{\text{mach}})\,\kappa(Q_k)\,\|A\|_2 = O(\varepsilon_{\text{mach}})\,\|A\|_2\,. \tag{6.34}$$

- The transformation does not amplify the error at all.
- Unitary matrices preserve the 2-norm, so every intermediate A_k has norm $\|A\|_2$. There is no element growth as in LU.
- Over all steps, the backward error grows only linearly with the number of steps (Section 6.7). So Householder QR is backward stable.

This presumes that each Q_k is computed accurately, i.e., that the direction of $\vec{v}$ is accurate. This is exactly what the sign choice ensures.

6.7 (SUPPLEMENT) Further Topics

6.7.1 Three Factorizations as Products

	Side	Transformation	Target	Condition of transformation
LU	left	lower triangular L_k	upper triangular U	can be large
Gram–Schmidt	right	upper triangular R_k	orthonormal $\hat{Q}$	depends on A
Householder	left	unitary Q_k	upper triangular R	1

Trefethen and Bau call Gram–Schmidt *triangular orthogonalization* and Householder *orthogonal triangularization* [1, Lectures 8 and 10].

6.7.2 The Condition of the Gram–Schmidt Transformation

The accumulated transformation of Gram–Schmidt is

$$R_1 \cdots R_n = \hat{R}^{-1}, \qquad \text{so} \qquad \kappa(R_1 \cdots R_n) = \kappa(\hat{R}).$$

This equals $\kappa(A)$. Since $A = \hat{Q}\hat{R}$ and $\hat{Q}$ has orthonormal columns,

$$\|A\vec{x}\|_2 = \left\|\hat{Q}\hat{R}\vec{x}\right\|_2 = \left\|\hat{R}\vec{x}\right\|_2 \quad \text{for all } \vec{x},$$

so A and $\hat{R}$ stretch every vector by the same factor. Their largest and smallest stretches agree, and hence $\kappa(\hat{R}) = \kappa(A)$.

So the triangular transformations of Gram–Schmidt are exactly as ill-conditioned as the problem: the more ill-conditioned A, the more the error is amplified. This matches the loss of orthogonality $\approx \varepsilon_{\text{mach}}\kappa(A)$ of MGS in Lecture 5.

6.7.3 Backward Error Over Many Steps

Theorem 26 covers one step. For many steps, write step k as $\tilde{A}_k = X_k(\tilde{A}_{k-1} + F_k)$. After two steps,

$$\tilde{A}_2 = X_2(X_1(A + F_1) + F_2) = X_2X_1(A + F_1 + X_1^{-1}F_2).$$

Continuing in the same way, every perturbation is pulled back to the original input:

$$\tilde{A}_n = X_n \cdots X_1 \left(A + \sum_{k=1}^{n} (X_{k-1} \cdots X_1)^{-1} F_k \right) + O(\varepsilon_{\text{mach}}^2). \tag{6.35}$$

For general X_k, the factors $(X_{k-1} \cdots X_1)^{-1}$ can amplify the F_k. When every X_k is unitary:

- $(X_{k-1} \cdots X_1)^{-1}$ is unitary, so it does not change the norm of F_k;
- $\left\|\tilde{A}_{k-1}\right\|_2 = \|A\|_2$ to first order, so $\|F_k\|_2 \le O(\varepsilon_{\text{mach}}) \|A\|_2$.

Adding up the n terms,

$$\|F_{\text{total}}\|_2 \lesssim O(n\varepsilon_{\text{mach}}) \|A\|_2 . \tag{6.36}$$

This is the argument behind Demmel's Lemma 3.1 and Theorem 3.5 [2, §3.4.3]. Trefethen and Bau state the result as $\tilde{Q}\tilde{R} = A + \delta A$ with $\|\delta A\| / \|A\| = O(\varepsilon_{\text{mach}})$ [1, Theorem 16.1], where $\tilde{Q}$ is the *exactly* unitary product of the reflectors defined by the computed vectors $\tilde{\vec{v}}_k$.

6.7.4 Orthogonality of Q

We measure the loss of orthogonality by $\|Q^*Q - I\|_2$.

- MGS computes $\hat{Q}$ by triangular column operations on A. Their condition is $\kappa(A)$, so $\left\|\hat{Q}^*\hat{Q} - I\right\|_2 \approx \varepsilon_{\text{mach}}\kappa(A)$.
- Householder never builds Q from A. Q is a product of reflectors, each unitary up to rounding in $\vec{v}_k$, so $\|Q^*Q - I\|_2 = O(\varepsilon_{\text{mach}})$, independently of $\kappa(A)$.

When an orthonormal basis of $\mathcal{R}(A)$ is needed explicitly, Householder is the safer choice.

6.7.5 Operation Counts

Step k of Algorithm 9 updates the $(m-k+1)\times(n-k+1)$ block $A[k{:}m, k{:}n]$. For each column $\vec{a}$ of this block, the inner product $\vec{v}^*\vec{a}$ and the update $\vec{a} - (2\vec{v}^*\vec{a}/\vec{v}^*\vec{v})\,\vec{v}$ each cost about $2(m-k+1)$ flops. So step k costs about $4(m-k+1)(n-k+1)$ flops, and in total

$$\sum_{k=1}^{n} 4(m-k+1)(n-k+1) \approx 4\int_0^n (m-t)(n-t)\,\mathrm{d}t = 2mn^2 - \frac{2}{3}n^3. \tag{6.37}$$

- For $m \gg n$ this is about $2mn^2$, the same as MGS.
- For $m = n$ it is $\frac{4}{3}n^3$, twice as much as LU ($\frac{2}{3}n^3$). This is the price of stability.

6.7.6 Using Q Implicitly for Least Squares

To solve $\min_{\vec{x}} \left\|A\vec{x} - \vec{b}\right\|_2$ we never form Q. Instead we apply the stored reflectors to $\vec{b}$ one by one, exactly as in step 5 of Algorithm 9:

$$\vec{c} \coloneqq Q^*\vec{b} = Q_n \cdots Q_1\vec{b}, \qquad \text{cost } O(mn).$$

Split $\vec{c}$ into $\vec{c}_1 = \vec{c}[1{:}n]$ and $\vec{c}_2 = \vec{c}[n{+}1{:}m]$. Since Q^* preserves the 2-norm,

$$\left\|A\vec{x} - \vec{b}\right\|_2^2 = \|R\vec{x} - \vec{c}\|_2^2 = \left\|\hat{R}\vec{x} - \vec{c}_1\right\|_2^2 + \|\vec{c}_2\|_2^2\,.$$

The first term vanishes when $\hat{R}\vec{x} = \vec{c}_1$, which we solve by back substitution. The second term does not depend on $\vec{x}$, so $\|\vec{c}_2\|_2$ is the minimal residual. Details are in Lecture 7.

6.7.7 The Complex Case

For complex data the real sign does not work: with the target $+\|\vec{x}\|\,\vec{e}_1$, the number $\vec{v}^*\vec{x}$ is in general not real, and the verification of $F\vec{x} = \|\vec{x}\|\,\vec{e}_1$ fails. Instead we take

$$\operatorname{sign}(x_1) \coloneqq x_1/\,|x_1| \quad (\text{and } 1 \text{ if } x_1 = 0), \qquad \vec{v} = \operatorname{sign}(x_1)\,\|\vec{x}\|\,\vec{e}_1 + \vec{x}.$$

- The first entry $v_1 = \operatorname{sign}(x_1)(\,\|\vec{x}\| + |x_1|\,)$ adds two numbers with the same phase, so there is again no cancellation.
- Since $\overline{\operatorname{sign}(x_1)}\,x_1 = |x_1|$,

$$\vec{v}^*\vec{x} = \|\vec{x}\|^2 + \|\vec{x}\|\,|x_1|\,, \qquad \vec{v}^*\vec{v} = 2\,\|\vec{x}\|^2 + 2\,\|\vec{x}\|\,|x_1| = 2\,\vec{v}^*\vec{x}.$$

- Hence, as in the real case, $F\vec{x} = \vec{x} - \vec{v} = -\operatorname{sign}(x_1)\,\|\vec{x}\|\,\vec{e}_1$.

Lecture 7

Householder QR in Practice, Givens Rotations, Existence and Uniqueness of QR

Supplementary reading:

- [1] Lectures 7 and 10.
- [2] Sections 3.2.2 and 3.4.

7.1 Picking Up from Householder QR

At the end of Householder QR (Algorithm 9), the upper triangle of A holds R and the strictly lower triangle holds the Householder vectors $\vec{v}_k$; Q is never computed explicitly. This raises two questions:

- Applications need $Q^*\vec{b}$ (for least squares) or $Q\vec{x}$. How do we compute them from the $\vec{v}_k$ alone?
- Every Householder step processes a whole column. If A is already close to upper triangular and only a few entries need to be eliminated, is this wasteful?

The first leads to the implicit use of Q, the second to Givens rotations. Finally we return to theory: does a QR factorization always exist, and is it unique?

7.2 Using Q

Implicit storage. Householder QR does not store Q as a matrix, only the vectors $\vec{v}_k$ defining the reflectors

$$Q_k = I - 2\frac{\vec{v}_k\vec{v}_k^*}{\vec{v}_k^*\vec{v}_k}. \tag{7.1}$$

Here $\vec{v}_k$ is regarded as a vector of length m whose first $k-1$ entries are zero, so Q_k acts only on rows k through m; this is the same as the block form $Q_k = \mathrm{diag}(I_{k-1}, F_k)$ of Lecture 6. From $Q_n \cdots Q_1 A = R$, and since each Q_k is Hermitian and its own inverse ($Q_k^* = Q_k = Q_k^{-1}$),

$$Q = (Q_n \cdots Q_1)^{-1} = Q_1 Q_2 \cdots Q_n, \qquad Q^* = Q_n \cdots Q_1. \tag{7.2}$$

Two algorithms. A product of matrices acts on a vector from right to left. In $Q^*\vec{b} = Q_n \cdots Q_1 \vec{b}$, the factor Q_1 acts first, so the loop runs $k = 1, \ldots, n$. In $Q\vec{x} = Q_1 \cdots Q_n \vec{x}$, the factor Q_n acts first, so the loop runs $k = n, \ldots, 1$.

Algorithm 10 Implicit computation of $Q^*\vec{b}$.

1: **for** $k = 1, \ldots, n$ **do**
2: $\quad \vec{b}[k{:}m] \leftarrow \vec{b}[k{:}m] - 2\vec{v}_k \left(\vec{v}_k^* \vec{b}[k{:}m]\right) / (\vec{v}_k^* \vec{v}_k)$

Algorithm 11 Implicit computation of $Q\vec{x}$.

1: **for** $k = n, n-1, \ldots, 1$ **do**
2: $\quad \vec{x}[k{:}m] \leftarrow \vec{x}[k{:}m] - 2\vec{v}_k \left(\vec{v}_k^* \vec{x}[k{:}m]\right) / (\vec{v}_k^* \vec{v}_k)$

As in Lecture 6, we follow the parentheses: first compute the scalar $\vec{v}_k^* \vec{b}[k{:}m]$, then update the vector. Each step costs $O(m-k)$, and the total is about $4mn - 2n^2$ flops, the same order as a matrix–vector product. We never form $\vec{v}_k \vec{v}_k^*$.

Each step is the same in both algorithms; only the loop runs in opposite directions. This is just like forward and back substitution.

Consequences of implicit storage.

In place. We scale $\vec{v}_k$ so that its k-th entry is 1, so this entry need not be stored. Scaling does not change the reflector, which only depends on the direction of $\vec{v}_k$. The other $m-k$ entries fit exactly into column k below the diagonal, which was just set to zero. So the factorization simply overwrites A. For $m = 4$, $n = 3$, writing v_{jk} for entry j of $\vec{v}_k$:

$$A \;\to\; \begin{bmatrix} r_{11} & r_{12} & r_{13} \\ v_{21} & r_{22} & r_{23} \\ v_{31} & v_{32} & r_{33} \\ v_{41} & v_{42} & v_{43} \end{bmatrix}. \tag{7.3}$$

A full QR factorization from only n vectors. An explicit Q takes m^2 numbers. The $\vec{v}_k$ take only about $mn - \frac{n^2}{2}$. This is a big saving when $m \gg n$. Applying Q is cheaper too: $4mn - 2n^2$ flops instead of $2m^2$ with an explicit Q. If we really need $\hat{Q}$ (the first n columns of Q), we compute $Q\vec{e}_1, \ldots, Q\vec{e}_n$ with Algorithm 11.

Q is orthogonal by definition. This point is important. The computed $\vec{v}_k$ have rounding errors. But $I - 2\vec{v}\vec{v}^*/(\vec{v}^*\vec{v})$ is an exact reflector for *every* nonzero $\vec{v}$, so Q is still exactly unitary. Rounding errors only change *which* unitary matrix we get. They show up as a small backward error in A: $Q\tilde{R} = A + \delta A$ with $\|\delta A\| \le O(\varepsilon_{\text{mach}})\,\|A\|$.

Gram–Schmidt is different. It computes the columns of $\hat{Q}$ explicitly, so rounding errors make them lose orthogonality: by about $\varepsilon_{\text{mach}}\kappa(A)$ for MGS and $\varepsilon_{\text{mach}}\kappa(A)^2$ for CGS. For example, with $\kappa(A) = 10^8$, MGS keeps only about 8 digits of orthogonality, and CGS keeps none. Householder stays orthogonal to $O(\varepsilon_{\text{mach}})$ for any $\kappa(A)$.

When we apply the Householder Q (for example to compute $Q^*\vec{b}$), we only apply exact reflectors in floating point. This is backward stable, which is why least squares via Householder QR is reliable.

7.3 Givens Rotations

Motivation: Householder can be overkill. Householder works on a whole column at a time. Step k always updates the whole block $A[k{:}m, k{:}n]$, even if column k has only one nonzero below the diagonal. This is wasteful when A is already close to upper triangular. A typical example is an *upper Hessenberg* matrix, which is zero below the first subdiagonal:


$$\begin{bmatrix} \times & \times & \times & \times \\ \times & \times & \times & \times \\ 0 & \times & \times & \times \\ 0 & 0 & \times & \times \end{bmatrix}. \tag{7.4}$$

Here each column has just one entry to eliminate. Such matrices appear in the QR algorithm for eigenvalues (Lecture 14). So we want an orthogonal transformation that zeros *one entry at a time* and touches only the two rows involved. Givens rotations do exactly this.

The 2×2 case. Consider the leading 2×2 block

$$\begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}. \tag{7.5}$$

We want to eliminate A_{21}. Geometrically, the first column $\vec{a}_1 = (A_{11}, A_{21})^\top$ is a vector in the plane. We *rotate* the plane so that $\vec{a}_1$ lands on the horizontal axis. The second column $\vec{a}_2$ rotates along with it.



Definition 29 (Givens Rotation)

With $\|\vec{a}_1\| = \sqrt{A_{11}^2 + A_{21}^2}$, the *Givens rotation* for $\vec{a}_1$ is

$$G = \begin{bmatrix} c & s \\ -s & c \end{bmatrix}, \qquad c = \frac{A_{11}}{\|\vec{a}_1\|}, \qquad s = \frac{A_{21}}{\|\vec{a}_1\|}. \tag{7.6}$$

Applying G to $\vec{a}_1$ indeed puts it on the horizontal axis:

$$G \begin{bmatrix} A_{11} \\ A_{21} \end{bmatrix} = \frac{1}{\|\vec{a}_1\|} \begin{bmatrix} A_{11}^2 + A_{21}^2 \\ -A_{21}A_{11} + A_{11}A_{21} \end{bmatrix} = \begin{bmatrix} \|\vec{a}_1\| \\ 0 \end{bmatrix}. \tag{7.7}$$


G is a rotation:

- $c^2 + s^2 = 1$, so $G^\top G = I$ and $\det G = 1$.
- If $\vec{a}_1$ makes the angle θ with the horizontal axis, then $c = \cos\theta$ and $s = \sin\theta$, and G rotates clockwise by θ.

Compare with Householder:

- A Householder reflector is a *reflection* ($\det = -1$). It maps a whole vector onto the direction of $\vec{e}_1$ at once.
- A Givens rotation is a *rotation* ($\det = 1$). It acts in one two-dimensional plane and zeros one entry.

QR by Givens rotations. In the $m \times m$ setting, the Givens rotation acting on rows i and j is the identity except in four positions:



$G(i,j)$

with c in positions (i,i) and (j,j), s in (i,j), and $-s$ in (j,i). The dotted diagonal is all 1, and all other entries are 0. Left multiplication by $G(i,j)$ changes only rows i and j: writing $\vec{r}_i^{\,\top}$ for the i-th row of A,


$$\vec{r}_i^{\,\top} \leftarrow c\,\vec{r}_i^{\,\top} + s\,\vec{r}_j^{\,\top}, \qquad \vec{r}_j^{\,\top} \leftarrow -s\,\vec{r}_i^{\,\top} + c\,\vec{r}_j^{\,\top}, \tag{7.8}$$

and every other row is left alone (see Section 7.7.4 for why).

To eliminate A_{ji} ($j > i$), take

$$c = \frac{A_{ii}}{\sqrt{A_{ii}^2 + A_{ji}^2}}, \qquad s = \frac{A_{ji}}{\sqrt{A_{ii}^2 + A_{ji}^2}}, \tag{7.9}$$

which is the 2×2 case transplanted to rows i and j. Suppose the first $i-1$ columns have already been reduced. Then one step gives the following picture.



$G(i,j)$ $\quad A$ $\quad G(i,j)A$

The dark gray rows are finished rows of R, and the light gray block is still to be reduced. Only rows i and j (outlined) change; in the first $i-1$ columns both are zero, and in column i the entry A_{ji} is rotated into A_{ii}.

Existing zeros are preserved:

- In the first $i-1$ columns, rows i and j are both zero. A linear combination of two zeros is zero.
- In column i, row i becomes $\sqrt{A_{ii}^2 + A_{ji}^2}$, row j becomes 0, and no other row changes.

So we sweep through the subdiagonal nonzeros column by column, from left to right. Within column i, we pair row i with $j = i+1, \dots, m$ in turn. For a 4×3 matrix this takes six rotations, one per subdiagonal entry.

In the picture below, $\times$ is a generic nonzero, red marks the entries that the current rotation changes, and the bold $\mathbf{0}$ is the entry it eliminates.

$$A = \begin{bmatrix} \times & \times & \times \\ \times & \times & \times \\ \times & \times & \times \\ \times & \times & \times \end{bmatrix} \xrightarrow{G(1,2)} \begin{bmatrix} \times & \times & \times \\ \mathbf{0} & \times & \times \\ \times & \times & \times \\ \times & \times & \times \end{bmatrix} \xrightarrow{G(1,3)} \begin{bmatrix} \times & \times & \times \\ 0 & \times & \times \\ \mathbf{0} & \times & \times \\ \times & \times & \times \end{bmatrix} \xrightarrow{G(1,4)} \begin{bmatrix} \times & \times & \times \\ 0 & \times & \times \\ 0 & \times & \times \\ \mathbf{0} & \times & \times \end{bmatrix}$$

$$\xrightarrow{G(2,3)} \begin{bmatrix} \times & \times & \times \\ 0 & \times & \times \\ 0 & \mathbf{0} & \times \\ 0 & \times & \times \end{bmatrix} \xrightarrow{G(2,4)} \begin{bmatrix} \times & \times & \times \\ 0 & \times & \times \\ 0 & 0 & \times \\ 0 & \mathbf{0} & \times \end{bmatrix} \xrightarrow{G(3,4)} \begin{bmatrix} \times & \times & \times \\ 0 & \times & \times \\ 0 & 0 & \times \\ 0 & 0 & \mathbf{0} \end{bmatrix} = R.$$

Notice that $G(1,3)$ does not touch row 2, so the zero created by $G(1,2)$ survives. From $G(2,3)$ on, the zeros in column 1 are carried along as combinations of zeros.

Labeling the rotations in the order they are applied, we get

$$G_N \cdots G_1 A = R, \qquad Q = G_1^* \cdots G_N^*, \tag{7.10}$$

where N is the number of subdiagonal nonzeros. For a dense $m \times n$ matrix,

$$N = \sum_{i=1}^{n} (m-i) = mn - \frac{n(n+1)}{2}.$$

As with Householder, Q is not formed explicitly. We store the pairs (c, s) and apply the rotations in sequence when needed, just as in Algorithms 10 and 11. Every G is orthogonal with $\kappa = 1$, so by the error-amplification framework of Lecture 6, Givens QR is backward stable as well.

Efficiency.

Dense matrices: slower than Householder. Each rotation acts on two rows, at 6 flops per column (4 multiplications and 2 additions). Eliminating column i takes $m-i$ rotations, each acting on rows of length about $n-i$, for a total of about

$$\sum_{i=1}^{n} 6(m-i)(n-i) \approx 3mn^2 - n^3 \text{ flops}, \tag{7.11}$$

about 1.5 times the cost of Householder ($2mn^2 - \frac{2}{3}n^3$).

Sparse or nearly triangular matrices: possibly faster. For an $n \times n$ upper Hessenberg matrix, only $n-1$ rotations are needed (one per subdiagonal entry), each costing $O(n)$, for a total of about $3n^2$ flops. Givens touches only the entries that must be eliminated and so exploits sparsity naturally.

7.4 Full and Reduced QR

Let $A \in \mathbb{K}^{m\times n}$ with $m \geq n$.

Definition 30 (Reduced and Full QR Factorizations)


A *reduced QR factorization* of A is

$$A = \hat{Q}\hat{R}, \qquad \hat{Q} \in \mathbb{K}^{m\times n} \text{ with orthonormal columns}, \quad \hat{R} \in \mathbb{K}^{n\times n} \text{ upper triangular.} \tag{7.12}$$


A *full QR factorization* of A is

$$A = QR, \qquad Q \in \mathbb{K}^{m\times m} \text{ unitary}, \quad R = \begin{bmatrix} \hat{R} \\ 0 \end{bmatrix} \in \mathbb{K}^{m\times n}, \ \hat{R} \text{ upper triangular.} \tag{7.13}$$

Gram–Schmidt produces a reduced factorization; Householder and Givens produce a full one. Drawing each factor to scale, this gives the following picture.



Reduced QR (Gram–Schmidt): $\hat{Q}$ has the same shape as A, and $\hat{R}$ is $n \times n$.



Full QR (Householder, Givens): Q is $m \times m$, and R is $\hat{R}$ padded with $m - n$ zero rows.

Splitting $Q = \begin{bmatrix} \hat{Q} & \hat{Q}_\perp \end{bmatrix}$ by columns,

$$QR = \begin{bmatrix} \hat{Q} & \hat{Q}_\perp \end{bmatrix} \begin{bmatrix} \hat{R} \\ 0 \end{bmatrix} = \hat{Q}\hat{R}. \tag{7.14}$$

In the picture, the extra $m - n$ columns $\hat{Q}_\perp$ (red) multiply only the zero block of R (also red) and do not contribute to the product. Their role is to complete $\hat{Q}$ to an orthonormal basis of $\mathbb{K}^m$; when A has full rank, the columns of $\hat{Q}_\perp$ span $\mathcal{R}(A)^\perp$.

7.5 Existence

Theorem 31 (Existence of QR)

Every $A \in \mathbb{K}^{m\times n}$ with $m \geq n$ has a full QR factorization and a reduced QR factorization.

The proof is constructive: we run Gram–Schmidt, look at where it can fail, and fix those cases.

Proof. Case 1: A has full rank, reduced QR. Gram–Schmidt can only fail at the normalization $\vec{q}_k = \hat{\vec{q}}_k / \left\|\hat{\vec{q}}_k\right\|$, if $\hat{\vec{q}}_k = \vec{0}$. This happens only if

$$\left\|\hat{\vec{q}}_k\right\| = 0 \iff \vec{a}_k \in \operatorname{span}(\vec{q}_1, \ldots, \vec{q}_{k-1}) = \operatorname{span}(\vec{a}_1, \ldots, \vec{a}_{k-1}). \tag{7.15}$$

Then $\vec{a}_k$ would be a combination of the earlier columns, which contradicts full rank. So Gram–Schmidt never fails, and every $r_{kk} = \left\|\hat{\vec{q}}_k\right\| > 0$.

Case 2: A is rank deficient, reduced QR. We run Gram–Schmidt as usual. When $\hat{\vec{q}}_k = \vec{0}$, we set $r_{kk} = 0$ and pick any unit vector $\vec{q}_k$ orthogonal to $\vec{q}_1, \ldots, \vec{q}_{k-1}$. Such a vector exists because $k - 1 < n \leq m$. We still have

$$\vec{a}_k = \sum_{i<k} r_{ik}\vec{q}_i + 0 \cdot \vec{q}_k, \tag{7.16}$$

so $A = \hat{Q}\hat{R}$ holds. The only difference is that $\hat{R}$ has some zeros on its diagonal.

Case 3: full QR. We first compute a reduced QR factorization. Then we add $m - n$ orthonormal vectors to $\vec{q}_1, \ldots, \vec{q}_n$ to get an orthonormal basis of $\mathbb{K}^m$, and we add $m - n$ zero rows to $\hat{R}$. □

7.6 Uniqueness

When QR is not unique.

Full QR with $m > n$ is not unique. In $QR = \hat{Q}\hat{R} + \hat{Q}_\perp \cdot 0$, the columns of $\hat{Q}_\perp$ only multiply the zero block of R, so they do not affect the product. Any orthonormal basis of $\mathcal{R}(\hat{Q})^\perp$ works. For example, we can replace $\hat{Q}_\perp$ by $\hat{Q}_\perp U$ for any unitary $U \in \mathbb{K}^{(m-n)\times(m-n)}$.

Rank-deficient A. If $\hat{\vec{q}}_k = \vec{0}$, then $r_{kk} = 0$, so $\vec{q}_k$ is multiplied by zero in $\vec{a}_k = \sum_{i<k} r_{ik}\vec{q}_i + 0 \cdot \vec{q}_k$. Any unit vector orthogonal to $\vec{q}_1, \ldots, \vec{q}_{k-1}$ works.

Example 32 (Non-Uniqueness for a Rank-Deficient Matrix)**.** Let

$$A = \begin{bmatrix} 1 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}, \qquad \vec{q}_1 = \vec{e}_1, \qquad \hat{\vec{q}}_2 = \vec{a}_2 - \vec{e}_1 = \vec{0}. \tag{7.17}$$

Both $\vec{q}_2 = \vec{e}_2$ and $\vec{q}_2 = \vec{e}_3$ work, with the same $\hat{R} = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$.

Phases. Even for full-rank A, some freedom is left. For any d_k with $|d_k| = 1$, we can multiply $\vec{q}_k$ by d_k and row k of $\hat{R}$ by $\bar{d}_k$. The product does not change, because $\bar{d}_k d_k = 1$. In the real case, this means each column can flip its sign. So for uniqueness we need a normalization, such as $r_{kk} > 0$.

Theorem 33 (Uniqueness of Reduced QR)

Let $A \in \mathbb{K}^{m\times n}$, $m \geq n$, have full column rank. Then the reduced QR factorization $A = \hat{Q}\hat{R}$ with $r_{jj} > 0$ for all j is unique.

Proof. Let $A = \hat{Q}\hat{R} = \tilde{Q}\tilde{R}$ be two such factorizations. We show that the columns agree, $\tilde{\vec{q}}_k = \vec{q}_k$, one k at a time. Fix k and write $\vec{q} = \vec{q}_k$ and $\tilde{\vec{q}} = \tilde{\vec{q}}_k$. We use the inner product $\langle \vec{x}, \vec{y} \rangle = \vec{x}^* \vec{y}$.

Step 1: what we know about $\vec{q}$ and $\tilde{\vec{q}}$. Since A has full rank, all $r_{jj} \neq 0$, so $\hat{R}$ is upper triangular and invertible. From $\hat{Q} = A\hat{R}^{-1}$, the first k columns of $\hat{Q}$ span the same space as the first k columns of A:

$$\operatorname{span}(\vec{q}_1, \ldots, \vec{q}_k) = \operatorname{span}(\vec{a}_1, \ldots, \vec{a}_k) \qquad \text{for every } k. \tag{7.18}$$

So $\vec{q}$ lies in $\operatorname{span}(\vec{a}_1, \ldots, \vec{a}_k)$. It is also orthogonal to $\vec{q}_1, \ldots, \vec{q}_{k-1}$, hence to $\vec{a}_1, \ldots, \vec{a}_{k-1}$. The same holds for $\tilde{\vec{q}}$.

Step 2: $\tilde{\vec{q}}$ is a multiple of $\vec{q}$. Since $\operatorname{span}(\vec{a}_1, \ldots, \vec{a}_k) = \operatorname{span}(\vec{a}_1, \ldots, \vec{a}_{k-1}, \vec{q})$, we can write

$$\tilde{\vec{q}} = \alpha \vec{q} + \sum_{j=1}^{k-1} \beta_j \vec{a}_j. \tag{7.19}$$

We take inner products with $\vec{q}$ and with $\tilde{\vec{q}}$. The $\vec{a}_j$ terms drop out, because both $\vec{q}$ and $\tilde{\vec{q}}$ are orthogonal to every $\vec{a}_j$ with $j < k$:

$$\left\langle \vec{q}, \tilde{\vec{q}} \right\rangle = \alpha, \qquad 1 = \left\langle \tilde{\vec{q}}, \tilde{\vec{q}} \right\rangle = \alpha \left\langle \tilde{\vec{q}}, \vec{q} \right\rangle = \alpha \bar{\alpha} = |\alpha|^2 . \tag{7.20}$$

With these two facts,

$$\left\| \tilde{\vec{q}} - \alpha \vec{q} \right\|^2 = \left\langle \tilde{\vec{q}}, \tilde{\vec{q}} \right\rangle - \alpha \left\langle \tilde{\vec{q}}, \vec{q} \right\rangle - \bar{\alpha} \left\langle \vec{q}, \tilde{\vec{q}} \right\rangle + |\alpha|^2 = 1 - 1 - 1 + 1 = 0. \tag{7.21}$$

So $\tilde{\vec{q}} = \alpha \vec{q}$ with $|\alpha| = 1$.

Step 3: $r_{kk} > 0$ forces $\alpha = 1$. From $A = \hat{Q}\hat{R}$ we get $r_{kk} = \langle \vec{q}_k, \vec{a}_k \rangle$. In the same way,

$$\tilde{r}_{kk} = \left\langle \tilde{\vec{q}}_k, \vec{a}_k \right\rangle = \langle \alpha \vec{q}_k, \vec{a}_k \rangle = \bar{\alpha} r_{kk}.$$

Both r_{kk} and $\tilde{r}_{kk}$ are positive, so $\bar{\alpha} = \tilde{r}_{kk}/r_{kk} > 0$. A positive number with $|\alpha| = 1$ is 1, so $\alpha = 1$.

So $\tilde{\vec{q}}_k = \vec{q}_k$ for every k, i.e., $\tilde{Q} = \hat{Q}$. Then $\tilde{R} = \hat{Q}^* A = \hat{R}$. □

In short: inside $\operatorname{span}(\vec{a}_1, \ldots, \vec{a}_k)$, the directions orthogonal to $\operatorname{span}(\vec{a}_1, \ldots, \vec{a}_{k-1})$ form a line. So A fixes $\vec{q}_k$ up to a factor of absolute value 1, and $r_{kk} > 0$ fixes that factor.

7.7 (SUPPLEMENT) Further Topics

7.7.1 Three QR Algorithms

	Factorization	Form of Q	Orthogonality of Q	Dense flops
MGS	reduced	explicit $\hat{Q}$	loses $\sim \varepsilon_{\text{mach}} \kappa(A)$	$2mn^2$
Householder	full	implicit (n vectors $\vec{v}_k$)	orthogonal by definition	$2mn^2 - \frac{2}{3}n^3$
Givens	full	implicit (pairs (c, s))	orthogonal by definition	$3mn^2 - n^3$

7.7.2 Householder as an Existence Proof

The Householder algorithm never breaks down. The only special case is $\vec{x} = A[k{:}m, k] = \vec{0}$; then we simply take $F_k = I$. So Householder gives a full QR factorization of *any* A, rank-deficient or not. This is a shorter existence proof than the one based on Gram–Schmidt. Trefethen and Bau prove existence with Gram–Schmidt in Lecture 7; their Lecture 10 implicitly gives the Householder proof [1].

7.7.3 Uniqueness via Cholesky

If $A = \hat{Q}\hat{R}$, then

$$A^*A = \hat{R}^*\hat{Q}^*\hat{Q}\hat{R} = \hat{R}^*\hat{R}. \tag{7.22}$$

So $\hat{R}^*$ is a Cholesky factor of A^*A. If A has full column rank, A^*A is Hermitian positive definite. Its Cholesky factorization with positive diagonal is unique. So $\hat{R}$ is unique, and then $\hat{Q} = A\hat{R}^{-1}$ is unique too.

This is a proof, not an algorithm. We should not *compute* QR this way, because forming A^*A squares the condition number (Lecture 5).

7.7.4 Why a Givens Rotation Changes Only Two Rows

Think of each column of A as a point in $\mathbb{K}^m$. Row k of A collects the k-th coordinates of all these points. $G(i, j)$ rotates in the plane spanned by $\vec{e}_i$ and $\vec{e}_j$. This is like a rotation about the z-axis in 3D: every point moves, but only its x and y coordinates change, and its z coordinate stays the same. In the same way, every column of A moves, but only its entries i and j change. So only rows i and j of A change.

7.7.5 Computing c and s Without Overflow

Computing $\sqrt{A_{ii}^2 + A_{ji}^2}$ directly can go wrong. The squares can overflow for large entries and underflow for small ones. So we first divide by the larger entry. If $|A_{ji}| > |A_{ii}|$, we set

$$\tau = A_{ii}/A_{ji}, \qquad s = 1/\sqrt{1+\tau^2}, \qquad c = s\tau,$$

and we swap the roles of A_{ii} and A_{ji} otherwise. Now $\tau^2 \leq 1$, so nothing overflows. This gives (c, s) up to a common sign, which still zeros the entry. The function `hypot` in many languages works this way. Demmel also shows how to store a rotation in a single number, so that it fits into the zeroed entry [2, §3.4.2].

7.7.6 Other Advantages of Givens Rotations

- *Parallelism.* Rotations on different pairs of rows do not interact. We can apply them at the same time.
- *Updating.* Suppose we already have the QR factorization of A, and a new row is added. We can eliminate the new row with n Givens rotations, instead of factoring again. This is common in recursive least squares.
- *Hessenberg matrices.* QR of a Hessenberg matrix is the core step of the QR algorithm for eigenvalues [1, Lectures 28–29].

7.7.7 Correspondence with Trefethen and Bau

- Full and reduced QR, existence and uniqueness: Theorems 7.1 and 7.2 of [1].
- The two implicit algorithms for $Q^*\vec{b}$ and $Q\vec{x}$: Algorithms 10.2 and 10.3.
- Givens rotations: Exercise 10.4.

Lecture 8

Overdetermined Least Squares and the Normal Equations

Supplementary reading:

- [1] Lectures 2, 11, and 23.
- [2] Sections 2.7.1 and 3.2.

8.1 Recap and Motivation

Last lecture we finished the QR factorization.

- *Using Q.* Householder QR never forms Q. To apply Q^* to a vector, we apply the stored reflectors in the order $\vec{v}_1, \dots, \vec{v}_n$. To apply Q, we use the reverse order.
- *Givens rotations.* A rotation in the (i, j) plane with $c = A_{ii}/\sqrt{A_{ii}^2 + A_{ji}^2}$ and $s = A_{ji}/\sqrt{A_{ii}^2 + A_{ji}^2}$ sets one entry to zero. Repeating this gives Givens QR.
- *Full and reduced QR*, and the *existence and uniqueness* of QR.

Solving overdetermined least squares. We now come back to the problem that led us to QR: given $A \in \mathbb{K}^{m \times n}$ with $m > n$ and $\vec{b} \in \mathbb{K}^m$, find the $\vec{x}$ that minimizes


$$\left\| A\vec{x} - \vec{b} \right\|^2 . \tag{8.1}$$

There are more equations than unknowns, so $A\vec{x} = \vec{b}$ usually has no solution. We cannot solve it as in the square case.

We already saw an informal reason why $\vec{x} = \hat{R}^{-1}\hat{Q}^*\vec{b}$ solves this problem, for a reduced QR factorization $A = \hat{Q}\hat{R}$. Let us recall it. We extend $\vec{q}_1, \dots, \vec{q}_n$ to an orthonormal basis $\vec{q}_1, \dots, \vec{q}_m$ of $\mathbb{K}^m$. In this basis,

- $\vec{b}$ has the coordinates $Q^*\vec{b}$;
- $A\vec{x} = \hat{Q}(\hat{R}\vec{x})$ has the coordinates $\hat{R}\vec{x}$ in the first n directions, and 0 in the other $m - n$.

Since $\hat{R}$ is invertible, we can make the first n coordinates of $A\vec{x}$ equal to those of $\vec{b}$, by solving $\hat{R}\vec{x} = \hat{Q}^*\vec{b}$. The other $m - n$ coordinates of $A\vec{x}$ are always 0, so we cannot reduce the error there. For $m = 3$ and $n = 2$, we get the following picture.



In the basis $\vec{q}_1, \vec{q}_2, \vec{q}_3$, $A\vec{x}$ can match the first two coordinates of $\vec{b}$. The third coordinate, $(Q^*\vec{b})_3$, is the error that no $\vec{x}$ can remove.

This argument is convincing, but it relies on the picture. Why is matching the coordinates the best choice? Is it the only best choice? *How do we prove this?* One way uses that Q^* preserves norms (Section B.2). In this lecture we take a different way, which needs no factorization at all.

First we answer a more basic question: *what condition does a minimizer satisfy?* Such a condition helps in two ways.

- It lets us *check* that the $\vec{x}$ from QR really is a minimizer.
- It turns a minimization problem into a *system of equations*.

The main result of this lecture is

$$\vec{x} \text{ minimizes } \left\|A\vec{x} - \vec{b}\right\|^2 \quad \Longleftrightarrow \quad A\vec{x} - \vec{b} \perp \mathcal{R}(A) \quad \Longleftrightarrow \quad A^*A\vec{x} = A^*\vec{b}. \tag{8.2}$$

In Lecture 5 we saw this geometrically; now we prove it. Then we study the matrix A^*A of the last system. This leads to Gram matrices and positive semidefinite matrices.

8.2 Conventions

We use the inner product $\langle \vec{x}, \vec{y} \rangle = \vec{x}^*\vec{y}$, as in Lectures 5 and 7. It is linear in the second argument and conjugate linear in the first. Swapping the two arguments conjugates it: $\langle \vec{y}, \vec{x} \rangle = \overline{\langle \vec{x}, \vec{y} \rangle}$. So

$$\langle \vec{x}, \vec{y} \rangle + \langle \vec{y}, \vec{x} \rangle = 2 \operatorname{Re} \langle \vec{x}, \vec{y} \rangle . \tag{8.3}$$

We write $\vec{r} \coloneqq A\vec{x} - \vec{b}$ for the residual, and $\vec{d}$ for a perturbation direction.

We will use the following identity many times in this lecture.

Lemma 34 (Moving a Matrix Across the Inner Product)

For any $M \in \mathbb{K}^{m\times n}$ (not necessarily square or unitary), $\vec{u} \in \mathbb{K}^n$, and $\vec{v} \in \mathbb{K}^m$,

$$\langle M\vec{u}, \vec{v}\rangle = \langle \vec{u}, M^*\vec{v}\rangle \,. \tag{8.4}$$

Proof. This follows from associativity: $\langle M\vec{u}, \vec{v}\rangle = (M\vec{u})^*\vec{v} = \vec{u}^*(M^*\vec{v}) = \langle \vec{u}, M^*\vec{v}\rangle$. □

In words: we can move a matrix from one side of the inner product to the other, if we replace M by M^*. Two remarks.

- This is *not* the same as multiplying both sides by M^*. For example, the $m \times n$ factor $\hat{Q}$ of a reduced QR factorization has $\hat{Q}^*\hat{Q} = I$ but $\hat{Q}\hat{Q}^* \neq I$. In general $\left\langle \hat{Q}^*\vec{y}, \hat{Q}^*\vec{z}\right\rangle \neq \langle \vec{y}, \vec{z}\rangle$ (compare Section B.2).
- In Axler's *Linear Algebra Done Right* (Section 7.A), Lemma 34 is used as the definition of the adjoint.

8.3 The Optimality Condition

We want to find $\vec{x}$ that minimizes $\left\|A\vec{x} - \vec{b}\right\|^2$. The idea: if $\vec{x}$ is a minimizer, moving it cannot make the value smaller. So for every $\vec{\varepsilon} \in \mathbb{K}^n$,

$$\left\|A(\vec{x} + \vec{\varepsilon}) - \vec{b}\right\|^2 \geq \left\|A\vec{x} - \vec{b}\right\|^2 . \tag{8.5}$$

So we first compute how much $\left\|A\vec{x} - \vec{b}\right\|^2$ changes when we move $\vec{x}$ by $\vec{\varepsilon}$.

Lemma 35 (How the Squared Residual Changes)

Let $A \in \mathbb{K}^{m\times n}$ and $\vec{b} \in \mathbb{K}^m$. For all $\vec{x}, \vec{\varepsilon} \in \mathbb{K}^n$, with $\vec{r} = A\vec{x} - \vec{b}$,

$$\left\|A(\vec{x} + \vec{\varepsilon}) - \vec{b}\right\|^2 - \left\|A\vec{x} - \vec{b}\right\|^2 = \|A\vec{\varepsilon}\|^2 + 2\,\mathrm{Re}\,\langle A\vec{\varepsilon}, \vec{r}\rangle \,. \tag{8.6}$$

Proof. Since $A(\vec{x} + \vec{\varepsilon}) - \vec{b} = A\vec{\varepsilon} + \vec{r}$, we have

$$\begin{aligned} \|A\vec{\varepsilon} + \vec{r}\|^2 - \|\vec{r}\|^2 &= \langle A\vec{\varepsilon} + \vec{r}, A\vec{\varepsilon} + \vec{r}\rangle - \langle \vec{r}, \vec{r}\rangle \\ &= \langle A\vec{\varepsilon}, A\vec{\varepsilon}\rangle + \langle A\vec{\varepsilon}, \vec{r}\rangle + \underbrace{\langle \vec{r}, A\vec{\varepsilon}\rangle}_{=\overline{\langle A\vec{\varepsilon}, \vec{r}\rangle}} \\ &= \|A\vec{\varepsilon}\|^2 + 2\,\mathrm{Re}\,\langle A\vec{\varepsilon}, \vec{r}\rangle \,, \end{aligned} \tag{8.7}$$

In the last step we used $z + \bar{z} = 2\,\mathrm{Re}\,z$. □

This holds for any $\vec{x}$. Look at the right-hand side:

- The term $\|A\vec{\varepsilon}\|^2$ is always ≥ 0, so it never makes the value smaller.
- Only the mixed term $2\,\mathrm{Re}\,\langle A\vec{\varepsilon}, \vec{r}\rangle$ can be negative.

So we guess: $\vec{x}$ is a minimizer exactly when the mixed term is zero for every $\vec{\varepsilon}$. The next theorem shows that this is true.

Theorem 36 (Optimality Condition for Least Squares)

Let $A \in \mathbb{K}^{m\times n}$ and $\vec{b} \in \mathbb{K}^m$. Then $\vec{x} \in \mathbb{K}^n$ minimizes $\left\|A\vec{x} - \vec{b}\right\|^2$ if and only if the residual is orthogonal to $\mathcal{R}(A)$:

$$\left\langle A\vec{d}, A\vec{x} - \vec{b} \right\rangle = 0 \qquad \text{for all } \vec{d} \in \mathbb{K}^n. \tag{8.8}$$

Also, if $\vec{x}$ is a minimizer, then the minimizers are exactly the vectors in $\vec{x} + \mathcal{N}(A)$. So the minimizer is unique if and only if A has full column rank.

Proof. Write $\vec{r} = A\vec{x} - \vec{b}$.

If $\vec{x}$ is a minimizer, the residual is orthogonal. Let $\vec{x}$ be a minimizer. Take any $\vec{d} \in \mathbb{K}^n$ and $\delta > 0$. By Lemma 35 with $\vec{\varepsilon} = \delta\vec{d}$,

$$0 \le \left\|A(\vec{x} + \delta\vec{d}) - \vec{b}\right\|^2 - \left\|A\vec{x} - \vec{b}\right\|^2 = \delta^2 \left\|A\vec{d}\right\|^2 + 2\delta \operatorname{Re}\left\langle A\vec{d}, \vec{r}\right\rangle. \tag{8.9}$$

We divide by δ and let $\delta \to 0^+$:

$$\operatorname{Re}\left\langle A\vec{d}, \vec{r}\right\rangle \ge 0 \qquad \text{for all } \vec{d} \in \mathbb{K}^n.$$

We show that both the real and the imaginary part of $\left\langle A\vec{d}, \vec{r}\right\rangle$ are zero.

Real part. The inequality holds for every vector in $\mathbb{K}^n$, so it also holds for $-\vec{d}$. The inner product is conjugate-linear in its first argument, and $\overline{-1} = -1$, so

$$0 \le \operatorname{Re}\left\langle A(-\vec{d}), \vec{r}\right\rangle = \operatorname{Re}(-\left\langle A\vec{d}, \vec{r}\right\rangle) = -\operatorname{Re}\left\langle A\vec{d}, \vec{r}\right\rangle.$$

So $\operatorname{Re}\left\langle A\vec{d}, \vec{r}\right\rangle \le 0$. Together with $\operatorname{Re}\left\langle A\vec{d}, \vec{r}\right\rangle \ge 0$ this gives $\operatorname{Re}\left\langle A\vec{d}, \vec{r}\right\rangle = 0$.

Imaginary part. If $\mathbb{K} = \mathbb{R}$, then $\left\langle A\vec{d}, \vec{r}\right\rangle$ is real and we are done. If $\mathbb{K} = \mathbb{C}$, we use the vector $i\vec{d}$ instead. The real-part argument above works for every vector, so it gives $\operatorname{Re}\left\langle A(i\vec{d}), \vec{r}\right\rangle = 0$. Since $\bar{i} = -i$,

$$\left\langle A(i\vec{d}), \vec{r}\right\rangle = \bar{i}\left\langle A\vec{d}, \vec{r}\right\rangle = -i\left\langle A\vec{d}, \vec{r}\right\rangle.$$

For any complex number $a + bi$ with a, b real, $-i(a + bi) = b - ai$, so $\operatorname{Re}(-i(a + bi)) = b$. That is, multiplying by $-i$ turns the imaginary part into the real part. Hence

$$0 = \operatorname{Re}\left\langle A(i\vec{d}), \vec{r}\right\rangle = \operatorname{Re}(-i\left\langle A\vec{d}, \vec{r}\right\rangle) = \operatorname{Im}\left\langle A\vec{d}, \vec{r}\right\rangle.$$

So $\left\langle A\vec{d}, \vec{r}\right\rangle = 0$ for all $\vec{d} \in \mathbb{K}^n$.

If the residual is orthogonal, $\vec{x}$ is a minimizer. Suppose (8.8) holds. Take any $\vec{y} \in \mathbb{K}^n$. We use Lemma 35 with $\vec{\varepsilon} = \vec{y} - \vec{x}$. The mixed term is zero by (8.8), so

$$\left\|A\vec{y} - \vec{b}\right\|^2 = \left\|A\vec{x} - \vec{b}\right\|^2 + \|A(\vec{y} - \vec{x})\|^2 \ge \left\|A\vec{x} - \vec{b}\right\|^2. \tag{8.10}$$

All minimizers. By (8.10), $\vec{y}$ is also a minimizer if and only if $A(\vec{y} - \vec{x}) = \vec{0}$, i.e., $\vec{y} - \vec{x} \in \mathcal{N}(A)$. So the minimizer is unique if and only if $\mathcal{N}(A) = \{\vec{0}\}$, i.e., A has full column rank. □


Thus, if $\vec{x}$ minimizes the squared residual, then

$$\left\langle A\vec{d}, A\vec{x} \right\rangle = \left\langle A\vec{d}, \vec{b} \right\rangle \qquad \text{for all } \vec{d} \in \mathbb{K}^n.$$

Both directions of the theorem are useful:

- Every minimizer satisfies (8.8). So solving (8.8) finds all minimizers.
- To show that a given $\vec{x}$ is a minimizer, it is enough to check (8.8). We do this for the QR solution in Section 8.4.

Geometrically, (8.10) is the Pythagorean theorem.



When the residual $A\vec{x} - \vec{b}$ is orthogonal to $\mathcal{R}(A)$, (8.10) is the Pythagorean theorem. Every other point $A\vec{y}$ in $\mathcal{R}(A)$ is at least as far from $\vec{b}$.

Geometric meaning. The vectors $A\vec{d}$, for all $\vec{d} \in \mathbb{K}^n$, are exactly the vectors in $\mathcal{R}(A)$. So the condition says: the residual is orthogonal to every vector in $\mathcal{R}(A)$. In other words, $A\vec{x}$ is the point of $\mathcal{R}(A)$ closest to $\vec{b}$. It is the orthogonal projection of $\vec{b}$ onto $\mathcal{R}(A)$. This is the picture of Lecture 5, now with a proof.

8.4 What the Optimality Condition Says

What does the optimality condition mean in concrete cases? The condition only tests the residual against vectors of the form $A\vec{d}$. How much this tells us depends on how many vectors have this form.

Invertible square matrices: solving a linear system. Let $m = n$ and let A be invertible. Let $\vec{x}$ be a minimizer. By Theorem 36, the residual is orthogonal to every $A\vec{d}$. Since A is invertible, we can choose $\vec{d} = A^{-1}(A\vec{x} - \vec{b})$, so that $A\vec{d} = A\vec{x} - \vec{b}$. Then

$$0 = \left\langle A\vec{d}, A\vec{x} - \vec{b} \right\rangle = \left\langle A\vec{x} - \vec{b}, A\vec{x} - \vec{b} \right\rangle = \left\| A\vec{x} - \vec{b} \right\|^2, \tag{8.11}$$

so $A\vec{x} = \vec{b}$. So for invertible square matrices, least squares is just solving the linear system, and the smallest residual is zero.

Singular or tall matrices: only some directions are tested. The trick above needs $A\vec{d} = \vec{r}$, where $\vec{r} = A\vec{x} - \vec{b}$ is the residual. If A is singular or $m > n$, then $\mathcal{R}(A)$ is smaller than $\mathbb{K}^m$. If also $\vec{b} \notin \mathcal{R}(A)$, then $\vec{r} \notin \mathcal{R}(A)$, and no such $\vec{d}$ exists. So we only get $\vec{r} \perp \mathcal{R}(A)$, not $\vec{r} = \vec{0}$. For example, $A = \left(\begin{smallmatrix} 1 \\ 0 \end{smallmatrix}\right)$ and $\vec{b} = \left(\begin{smallmatrix} 1 \\ 1 \end{smallmatrix}\right)$ give $x = 1$ and $\vec{r} = \left(\begin{smallmatrix} 0 \\ -1 \end{smallmatrix}\right) \neq \vec{0}$.

Checking the QR solution. Now we can prove the informal QR argument from the start of this lecture. Let $A = \hat{Q}\hat{R}$ be a reduced QR factorization, with $\hat{Q}^*\hat{Q} = I$. Let A have full column rank, so $\hat{R}$ is upper triangular and invertible. Take $\vec{x} = \hat{R}^{-1}\hat{Q}^*\vec{b}$. We check the orthogonality condition, moving $\hat{Q}$ to the right with Lemma 34:

$$\begin{aligned}\left\langle A\vec{d}, A\vec{x} - \vec{b}\right\rangle &= \left\langle \hat{Q}\hat{R}\vec{d}, \hat{Q}\hat{R}\hat{R}^{-1}\hat{Q}^*\vec{b} - \vec{b}\right\rangle \\ &= \left\langle \hat{R}\vec{d}, \hat{Q}^*\hat{Q}\hat{R}\hat{R}^{-1}\hat{Q}^*\vec{b} - \hat{Q}^*\vec{b}\right\rangle \\ &= \left\langle \hat{R}\vec{d}, \hat{Q}^*\vec{b} - \hat{Q}^*\vec{b}\right\rangle = 0.\end{aligned} \tag{8.12}$$

So $\vec{x} = \hat{R}^{-1}\hat{Q}^*\vec{b}$ satisfies the orthogonality condition. By Theorem 36, it is a least squares solution, and since A has full column rank, it is the only one. In practice we never form $\hat{R}^{-1}$. We solve the upper triangular system $\hat{R}\vec{x} = \hat{Q}^*\vec{b}$ by back substitution.

8.5 The Normal Equations

The optimality condition holds "for all $\vec{d}$". We now turn it into one linear system. The trick is to use the transpose: we apply Lemma 34 to both sides and move A to the right.

$$\left\langle A\vec{d}, A\vec{x}\right\rangle = \left\langle A\vec{d}, \vec{b}\right\rangle \ \ \forall \vec{d} \in \mathbb{K}^n \quad \Longleftrightarrow \quad \left\langle \vec{d}, A^*A\vec{x}\right\rangle = \left\langle \vec{d}, A^*\vec{b}\right\rangle \ \ \forall \vec{d} \in \mathbb{K}^n \quad \Longleftrightarrow \quad A^*A\vec{x} = A^*\vec{b}. \tag{8.13}$$

The first step is Lemma 34 applied to each side. For the last step, let $\vec{w} = A^*A\vec{x} - A^*\vec{b}$. The inner product is linear in its second argument, so subtracting the two sides turns the middle condition into $\left\langle \vec{d}, \vec{w}\right\rangle = 0$ for every $\vec{d} \in \mathbb{K}^n$. ($\Rightarrow$) Then $\vec{w}$ is orthogonal to every vector, including itself. Taking $\vec{d} = \vec{w}$ gives $\|\vec{w}\|^2 = 0$, so $\vec{w} = \vec{0}$, that is, $A^*A\vec{x} = A^*\vec{b}$. ($\Leftarrow$) If $\vec{w} = \vec{0}$, then $\left\langle \vec{d}, \vec{w}\right\rangle = 0$ for every $\vec{d}$.

So the orthogonality condition is the same as the normal equations. Together with Theorem 36, we get the following theorem.

Theorem 37 (The Normal Equations)

$\vec{x}$ minimizes $\left\|A\vec{x} - \vec{b}\right\|^2$ if and only if it solves the *normal equations*

$$A^*A\vec{x} = A^*\vec{b}. \tag{8.14}$$

We have turned our system into a square one! The $m \times n$ system $A\vec{x} = \vec{b}$ has become an $n \times n$ system. "Normal" means perpendicular: the equation $A^*(A\vec{x} - \vec{b}) = \vec{0}$ says that the residual is perpendicular to the column space, as in (5.7).

The normal equations have advantages and one serious drawback.

- *Pro:* the system is square, and its matrix A^*A is Hermitian (symmetric in the real case). In fact it is positive semidefinite (Theorem 42). We can solve such systems with the Cholesky factorization of Lecture 9, at half the cost of LU.
- *Con:* the condition number is squared, $\kappa_2(A^*A) = \kappa_2(A)^2$. So forming A^*A loses precision when A is ill-conditioned (see the supplement at the end of this lecture).

8.6 Gram Matrices and Positive Semidefinite Matrices

Solving the normal equations means solving a system with a matrix of the form B^*B. (We use the letter B here, so that A is free for the matrix of the system.) So we study such matrices. We start with a larger class.

Definition 38 (Hermitian and Anti-Hermitian Matrices)

A matrix $A \in \mathbb{K}^{m\times m}$ is *Hermitian* if $A^* = A$, and *anti-Hermitian* (or skew-Hermitian) if $A^* = -A$. For $\mathbb{K} = \mathbb{R}$, they are called *symmetric* and *antisymmetric* (or skew-symmetric).

Proposition 39 (Eigenvalues of (Anti-)Hermitian Matrices)

All eigenvalues of a Hermitian matrix are real. All eigenvalues of an anti-Hermitian matrix are purely imaginary.

Proof. Let $A\vec{v} = \lambda\vec{v}$ with $\vec{v} \neq \vec{0}$. Then $\vec{v}^*A\vec{v} = \lambda \|\vec{v}\|^2$. We take the complex conjugate of the number $\vec{v}^*A\vec{v}$:

$$\overline{\vec{v}^*A\vec{v}} = (\vec{v}^*A\vec{v})^* = \vec{v}^*A^*\vec{v} = \pm\vec{v}^*A\vec{v}, \tag{8.15}$$

with $+$ for Hermitian and $-$ for anti-Hermitian A. So $\vec{v}^*A\vec{v}$ is real in the first case and purely imaginary in the second. The same holds for $\lambda = \vec{v}^*A\vec{v}/\|\vec{v}\|^2$. □

These matrices behave a lot like complex numbers.

- Every square matrix splits into a Hermitian and an anti-Hermitian part, $A = \frac{1}{2}(A + A^*) + \frac{1}{2}(A - A^*)$. This is like splitting a complex number into its real and imaginary parts.
- A second analogy is $z = e^{i\theta}r$ with $r \geq 0$. Its matrix version is the polar decomposition below. It uses the following notion, which we study in detail later in this section: a Hermitian matrix S is *positive semidefinite* if $\vec{v}^*S\vec{v} \geq 0$ for all $\vec{v}$.

Theorem 40 (Polar Decomposition)

For every $A \in \mathbb{K}^{m\times m}$, there are a unitary U and a Hermitian positive semidefinite S such that

$$A = US. \tag{8.16}$$

Proof. Take an SVD $A = W\Sigma V^*$ and insert $V^*V = I$:

$$A = W\Sigma V^* = \underbrace{(WV^*)}_{U}\underbrace{(V\Sigma V^*)}_{S}. \tag{8.17}$$

We check the two factors.

- $U = WV^*$ is a product of unitary matrices, so it is unitary.
- $S = V\Sigma V^*$ is Hermitian, since Σ is real and diagonal. Also $\vec{v}^*S\vec{v} = \sum_i \sigma_i |(V^*\vec{v})_i|^2 \geq 0$. □

Geometrically, S stretches space along the orthogonal directions given by the columns of V. Then U rotates (or reflects) the result. The factor S is always determined by A: since $A^*A = SU^*US = S^2$, S is the positive semidefinite square root of A^*A. If A is invertible, then so is S, and $U = AS^{-1}$ is determined too.

Gram matrices. A matrix of the form B^*B is called a *Gram matrix* (or inner product matrix). Its entries are the inner products of the columns $B_{:,1}, \dots, B_{:,n}$ of B with each other. Using $B\vec{e}_j = B_{:,j}$,

$$(B^*B)_{ij} = \vec{e}_i^* B^* B \vec{e}_j = \langle B\vec{e}_i, B\vec{e}_j \rangle = \langle B_{:,i}, B_{:,j} \rangle . \tag{8.18}$$

We see right away:

- The diagonal entries $(B^*B)_{ii} = \|B_{:,i}\|^2$ are ≥ 0.
- $B^*B = I$ exactly when the columns of B are orthonormal.

Definition 41 (Positive (Semi)definite Matrices)

A Hermitian matrix $A \in \mathbb{K}^{n\times n}$ is

- *positive semidefinite* if $\vec{x}^* A \vec{x} \geq 0$ for all $\vec{x} \in \mathbb{K}^n$;
- *positive definite* if $\vec{x}^* A \vec{x} > 0$ for all $\vec{x} \neq \vec{0}$.

The main result of this section: Gram matrices and positive semidefinite matrices are the same thing.

Theorem 42 (Gram Matrices Are Exactly the Positive Semidefinite Matrices)

1. $A \in \mathbb{K}^{n\times n}$ is Hermitian positive semidefinite if and only if it is a Gram matrix, i.e., $A = B^*B$ for some matrix B.
2. For $B \in \mathbb{K}^{m\times n}$, the Gram matrix B^*B is positive definite if and only if B has full column rank.

Item 1 is false for general Hermitian matrices. For example, $\mathrm{diag}(1, -1)$ is symmetric but not a Gram matrix: its second diagonal entry would have to be $\|B_{:,2}\|^2 \geq 0$. In item 2, the rank condition is on B, not on the square matrix B^*B.

Proof of item 1. Gram matrices are Hermitian positive semidefinite. We have

$$(B^*B)^* = B^*(B^*)^* = B^*B, \qquad \vec{x}^* B^* B \vec{x} = \|B\vec{x}\|^2 \geq 0. \tag{8.19}$$

Hermitian positive semidefinite matrices are Gram matrices. We take a square root of A in three steps.

1. *Diagonalize.* By the spectral theorem, $A = U\Lambda U^*$, where U is unitary and Λ is diagonal with the eigenvalues of A on its diagonal.
2. *The eigenvalues are* ≥ 0. They are real by Proposition 39. If $A\vec{v} = \lambda\vec{v}$ with $\vec{v} \neq \vec{0}$, then $\lambda \|\vec{v}\|^2 = \vec{v}^* A \vec{v} \geq 0$, so $\lambda \geq 0$. (If A is positive definite, the same argument gives $\lambda > 0$.)
3. *Take square roots.* So we can define $\sqrt{\Lambda} = \mathrm{diag}(\sqrt{\lambda_1}, \dots, \sqrt{\lambda_n})$. We insert $U^*U = I$ in the middle:

$$A = U\sqrt{\Lambda}\, U^* U \sqrt{\Lambda}\, U^* = \left(U\sqrt{\Lambda}\, U^*\right)^* \left(U\sqrt{\Lambda}\, U^*\right). \tag{8.20}$$

   In the last step we used that $\sqrt{\Lambda}$ is real and diagonal, so it equals its conjugate transpose.

So $B = U\sqrt{\Lambda}\, U^*$ gives $A = B^*B$. □

The matrix $B = U\sqrt{\Lambda}U^*$ is the *matrix square root* $A^{1/2}$. It is not the only choice: for any unitary V, $(VB)^*(VB) = B^*B$. For example, a QR factorization $B = QR$ (Theorem 31) gives $A = R^*Q^*QR = R^*R$ with R upper triangular. This is the *Cholesky factorization* of A, the topic of the next lecture.

Proof of item 2. We show the equivalent statement with both sides negated:

$$\begin{aligned} B \text{ is column rank deficient} &\iff \exists\, \vec{x} \in \mathbb{K}^n \setminus \{\vec{0}\} : B\vec{x} = \vec{0} \\ &\iff \exists\, \vec{x} \in \mathbb{K}^n \setminus \{\vec{0}\} : \|B\vec{x}\|^2 = 0 \\ &\iff \exists\, \vec{x} \in \mathbb{K}^n \setminus \{\vec{0}\} : \vec{x}^*B^*B\vec{x} = 0. \end{aligned} \tag{8.21}$$

By item 1, $\vec{x}^*B^*B\vec{x} \geq 0$ always. So B^*B is not positive definite exactly when $\vec{x}^*B^*B\vec{x} = 0$ for some $\vec{x} \neq \vec{0}$. This is the last line above. □

What this means for least squares. Putting Theorems 37 and 42 together,

$$\begin{aligned} A \text{ has full column rank} &\iff A^*A \text{ is positive definite} \\ &\iff \text{the normal equations have the unique solution } \vec{x} = (A^*A)^{-1}A^*\vec{b}. \end{aligned} \tag{8.22}$$

This agrees with the uniqueness statement of Theorem 36. We can solve a positive definite system with the Cholesky factorization $A^*A = R^*R$. It costs about $\frac{1}{3}n^3$ flops, half of LU, and needs no pivoting.

8.7 (SUPPLEMENT) Further Topics

8.7.1 Why the Perturbation Argument

In Theorem 36 we moved $\vec{x}$ a little in every direction $\vec{d}$ and used that the value cannot go down. This idea, "at a minimizer, the first-order change in every direction is ≥ 0", works in many other settings:

- General differentiable functions: at a minimizer, the gradient is zero.
- Problems with constraints: we can no longer replace $\vec{d}$ by $-\vec{d}$, so the result stays an inequality. This leads to variational inequalities and the KKT conditions.
- Infinite-dimensional problems, such as finite elements. There, Galerkin orthogonality says that the residual is orthogonal to every test function v. This is the same as "for all $\vec{d}$" here.

8.7.2 A Numerical Warning About the Normal Equations

By Theorem 113, the singular values of A^*A are the squares of those of A. So

$$\kappa_2(A^*A) = \kappa_2(A)^2. \tag{8.23}$$

If A is ill-conditioned, solving $A^*A\vec{x} = A^*\vec{b}$ loses about twice as many digits as necessary.

The normal equations are faster: forming A^*A takes about mn^2 flops and Cholesky another $\frac{1}{3}n^3$. This is about half the cost of Householder QR when $m \gg n$. But the speed is only worth it when A is well-conditioned. So in practice we use a QR factorization and solve $\hat{R}\vec{x} = \hat{Q}^*\vec{b}$ instead [1, Lecture 11].

Lecture 9

Cholesky Factorization

Supplementary reading:

- [1] Lecture 23.
- [2] Section 2.7.1.
- [7] Chapter 10.

9.1 Recap and Motivation

Last lecture we solved the least squares problem $\min_{\vec{x}} \left\|A\vec{x} - \vec{b}\right\|^2$ in two ways.

- *By QR.* With a full QR factorization,

$$\left\|A\vec{x} - \vec{b}\right\|^2 = \left\|\hat{R}\vec{x} - (Q^*\vec{b})_{1:n}\right\|^2 + \left\|(Q^*\vec{b})_{n+1:m}\right\|^2.$$

  We make the first term zero by solving $\hat{R}\vec{x} = (Q^*\vec{b})_{1:n}$. The second term does not depend on $\vec{x}$. It is the squared distance from $\vec{b}$ to $\mathcal{R}(A)$.

- *By the optimality condition.* At the minimizer, the residual is orthogonal to $\mathcal{R}(A)$: $\left\langle A\vec{d}, A\vec{x}\right\rangle = \left\langle A\vec{d}, \vec{b}\right\rangle$ for all $\vec{d}$ (Theorem 36). Written as a linear system, these are the normal equations $A^*A\vec{x} = A^*\vec{b}$ (Theorem 37).

The matrix A^*A of the normal equations is a Gram matrix. We saw that Gram matrices are exactly the positive semidefinite matrices: M is Hermitian positive semidefinite if and only if $M = B^*B$ for some B (Theorem 42).

The factor B in $M = B^*B$ is not unique at all. For any unitary U, $(UB)^*(UB) = B^*U^*UB = B^*B$. Since we have so much freedom, we can ask:

If $A = B^*B$ is positive definite, can we choose B to be upper triangular?

If yes, then $A = R^*R$ with R upper triangular. Then solving $A\vec{x} = \vec{c}$ takes only two triangular solves: first $R^*\vec{y} = \vec{c}$, then $R\vec{x} = \vec{y}$. In this lecture we show that the answer is yes. The resulting factorization is called the *Cholesky factorization*. Even better, we will see that it needs no pivoting and is always backward stable.

Here is our plan.

1. Derive Cholesky from LU by making elimination *symmetric*, and turn it into an algorithm.
2. Prove that the algorithm never breaks down (existence) and that it does not amplify rounding errors (stability).
3. Look at the elimination in blocks, which leads to *Schur complements*.
4. Show that Cholesky and QR compute the same triangular factor.

Conventions. In this lecture $A \in \mathbb{K}^{m\times m}$ is a square matrix. It is not related to the $m \times n$ matrix of least squares. We say that A is *s.p.d.* if it is Hermitian positive definite, i.e., $A = A^*$ and $\vec{x}^* A\vec{x} > 0$ for all $\vec{x} \neq \vec{0}$. (In the real case this means symmetric positive definite, hence the name.) We write the Cholesky factorization as $A = LL^*$ with L lower triangular; this is the same as $A = R^*R$ with $R = L^*$.

Since A is Hermitian, its first row is the conjugate transpose of its first column. So we can split off the first row and column as

$$A = \begin{bmatrix} \alpha & \vec{a}^* \\ \vec{a} & A_{22} \end{bmatrix}, \qquad \alpha = A_{11} \in \mathbb{R}, \quad \vec{a} = A[2{:}m, 1] \in \mathbb{K}^{m-1}, \quad A_{22} = A[2{:}m, 2{:}m]. \tag{9.1}$$

9.2 Deriving Cholesky by Symmetric Elimination

The idea is simple. We already know how to factor a matrix by Gaussian elimination. We run elimination on A. But at every step we do the *same* operation on the rows and on the columns. This way the matrix stays Hermitian. Let us go through one step.

Step 1: one step of LU. The first step of LU subtracts a_i/α times row 1 from row $i + 1$, for every row below the first. In matrix form, this is multiplication from the left by a unit lower triangular L_1:

$$L_1 = \begin{bmatrix} 1 & 0 \\ -\vec{a}/\alpha & I \end{bmatrix}, \qquad L_1 A = \begin{bmatrix} \alpha & \vec{a}^* \\ 0 & A_{22} - \dfrac{\vec{a}\vec{a}^*}{\alpha} \end{bmatrix}. \tag{9.2}$$

The first column is now zero below the diagonal. The lower right block has changed to

$$S := A_{22} - \frac{\vec{a}\vec{a}^*}{\alpha}. \tag{9.3}$$

The matrix S is called a *Schur complement*. We study it in Section 9.6. Note that $\vec{a}\vec{a}^*$ is an outer product. So the update is a rank-one change of A_{22}.

Can we divide by α? Yes: $\alpha = A_{11} = \vec{e}_1^* A\vec{e}_1 > 0$ by positive definiteness.

Step 2: add the matching column operation. If we continued with LU on S, we would get $A = LU$, where L and U are not related. We want $A = LL^*$ instead. So we keep the matrix symmetric. After the row operation L_1 on the left, we apply the matching column operation L_1^* on the right. The result is

$$L_1 A L_1^* = \begin{bmatrix} \alpha & 0 \\ 0 & S \end{bmatrix}. \tag{9.4}$$

Let us check this. Right multiplication by L_1^* subtracts $\overline{a_j}/\alpha$ times column 1 from column $j + 1$.

- It does not change the lower right block, since column 1 of $L_1 A$ is zero below the diagonal.
- It makes the first row zero. Since A is Hermitian, the $(1, j+1)$ entry of $L_1 A$ is $\overline{a_j}$. So it becomes $\overline{a_j} - (\overline{a_j}/\alpha)\alpha = 0$.

This is where symmetry helps. The first row is the conjugate of the first column. So the column operation that clears the row uses the same multipliers as the row operation that cleared the column. One matrix L_1 handles both sides.

Step 3: normalize by $\sqrt{\alpha}$. Finally, we scale the top left entry to 1. We divide the first row and the first column by $\sqrt{\alpha}$. In matrix form, we replace L_1 by $\bar{L}_1 = \operatorname{diag}(1/\sqrt{\alpha}, 1, \ldots, 1) L_1$. Then

$$\bar{L}_1 A \bar{L}_1^* = \begin{bmatrix} 1 & 0 \\ 0 & S \end{bmatrix}, \qquad \bar{L}_1^{-1} = \begin{bmatrix} \sqrt{\alpha} & 0 \\ \vec{a}/\sqrt{\alpha} & I \end{bmatrix}. \tag{9.5}$$

(To invert $\bar{L}_1$, undo the scaling and then flip the sign of the multipliers.) We move $\bar{L}_1$ and $\bar{L}_1^*$ to the other side and get

$$A = \begin{bmatrix} \sqrt{\alpha} & 0 \\ \vec{a}/\sqrt{\alpha} & I \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & S \end{bmatrix} \begin{bmatrix} \sqrt{\alpha} & \vec{a}^*/\sqrt{\alpha} \\ 0 & I \end{bmatrix}. \tag{9.6}$$

Now suppose that S has a Cholesky factorization $S = L_S L_S^*$. We put it into (9.6) and multiply out:

$$A = LL^*, \qquad L = \begin{bmatrix} \sqrt{\alpha} & 0 \\ \vec{a}/\sqrt{\alpha} & L_S \end{bmatrix}. \tag{9.7}$$

In words: *the first column of L is the first column of A divided by $\sqrt{\alpha}$, and the rest of L is the Cholesky factor of the Schur complement S.* So Cholesky is a recursion: compute one column, update the remaining block to S, and repeat on S. Only one thing can go wrong: some later S might not be s.p.d., and then its top left entry might not be positive. We rule this out in Section 9.4.

Problem Solving Strategy 43 (Computing a Cholesky Factorization by Hand)**.** *Repeat for $k = 1, \ldots, m$:*

- *Let d_k be the top left entry of the current matrix.*
- *Column k of L is the current first column divided by $\sqrt{d_k}$.*
- *Replace the current matrix by its Schur complement: delete the first row and column, and subtract $\vec{a}\vec{a}^*/d_k$ from what is left.*

Example 44 (A 3×3 Example)**.** Let

$$A = \begin{bmatrix} 4 & 2 & 2 \\ 2 & 5 & 3 \\ 2 & 3 & 6 \end{bmatrix}. \tag{9.8}$$

Step 1. $d_1 = 4$, so the first column of L is $(4, 2, 2)^\top / 2 = (2, 1, 1)^\top$. The Schur complement is

$$\begin{bmatrix} 5 & 3 \\ 3 & 6 \end{bmatrix} - \frac{1}{4} \begin{bmatrix} 2 \\ 2 \end{bmatrix} \begin{bmatrix} 2 & 2 \end{bmatrix} = \begin{bmatrix} 4 & 2 \\ 2 & 5 \end{bmatrix}. \tag{9.9}$$

Step 2. $d_2 = 4$, so the next column is $(4, 2)^\top / 2 = (2, 1)^\top$. The Schur complement is $5 - 2 \cdot 2/4 = 4$.

Step 3. $d_3 = 4$, so the last entry is $\sqrt{4} = 2$. Altogether,

$$L = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 2 & 0 \\ 1 & 1 & 2 \end{bmatrix}, \tag{9.10}$$

We can check that $LL^\top = A$.

Why the square root? If we eliminate symmetrically without taking square roots, we get $A = \tilde{L}D\tilde{L}^*$. Here $\tilde{L}$ is unit lower triangular and $D = \mathrm{diag}(d_1, \ldots, d_m)$ holds the pivots. LU puts all of D into $U = D\tilde{L}^*$. Cholesky splits it evenly between the two factors:

$$A = (\tilde{L}D^{1/2})(\tilde{L}D^{1/2})^* = LL^*. \tag{9.11}$$

This needs every $d_k > 0$, so that $D^{1/2}$ is real.

9.3 The Cholesky Algorithm

The right-looking algorithm. We now turn the recursion into an in-place algorithm. Step k computes column k of L and then applies the Schur complement update to the lower right block. Since every Schur complement is Hermitian, we only store and update the lower triangle.

Algorithm 12 Cholesky factorization, right-looking, in place (lower triangle).

1: **for** $k = 1, \ldots, m$ **do**
2: $\quad A_{kk} \leftarrow \sqrt{A_{kk}}$
3: $\quad$ **for** $i = k+1, \ldots, m$ **do**
4: $\quad\quad A_{ik} \leftarrow A_{ik}/A_{kk}$ ▷ column k of L
5: $\quad$ **for** $j = k+1, \ldots, m$ **do**
6: $\quad\quad$ **for** $i = j, \ldots, m$ **do**
7: $\quad\quad\quad A_{ij} \leftarrow A_{ij} - A_{ik}\overline{A_{jk}}$ ▷ Schur complement update

In line 2, A_{kk} is the current pivot d_k, the top left entry of the current Schur complement, not an entry of the original A. In the update, A_{ik} and A_{jk} have both been divided by $\sqrt{d_k}$, so we subtract $a_i\overline{a_j}/d_k$, the (i,j) entry of $\vec{a}\vec{a}^*/\alpha$. At the end, the lower triangle of A holds L. In short form:

```
A[:,k] /= sqrt(A[k,k]),   then   A[i,j] -= A[i,k]*conj(A[j,k]).
```

Comparison with LU. Cholesky differs from LU in three ways.

- *Normalization.* Columns are divided by the square root of the pivot, as explained above.
- *Half the work and storage.* Every Schur complement is Hermitian. So we only need to keep one triangle. The other one is its conjugate transpose. The cost drops from about $\frac{2}{3}m^3$ flops for LU to about $\frac{1}{3}m^3$.
- *No pivoting.* We prove in Sections 9.4 and 9.5 that Cholesky without pivoting never breaks down and is stable.

Where positive definiteness enters. The algorithm uses positive definiteness in exactly one place: the square root $\sqrt{A_{kk}}$ in line 2. Every *updated* pivot must be positive. Then it has a real square root, and we can divide by it. Positive definiteness of A directly tells us only that the *original* diagonal entries are positive. To show that the updated pivots are also positive, we need a proof. This is the existence theorem below. The next example shows that a proof is really needed.

Example 45 (A Symmetric Matrix That Is Not Positive Definite)**.** Run Algorithm 12 on

$$A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}. \tag{9.12}$$

For $k = 1$: $\sqrt{1} = 1$, the first column becomes $(1, 2)^\top$, and the update gives $A_{22} = 1 - 2 \cdot 2 = -3$. For $k = 2$ we need $\sqrt{-3}$, so the algorithm fails. Both original diagonal entries are 1. The problem is the updated pivot.


Left-looking and up-looking variants. The right-looking algorithm is not the only possible order of computation. Other orders give especially good memory access patterns. To see the options, we compare entries on both sides of $A = LL^*$ for $i \geq j$:

$$L_{jj} = \sqrt{A_{jj} - \sum_{k<j} |L_{jk}|^2}, \qquad L_{ij} = \frac{1}{L_{jj}} \left(A_{ij} - \sum_{k<j} L_{ik} \overline{L_{jk}} \right) \quad (i > j). \tag{9.13}$$

Any loop order gives a valid algorithm, as long as it evaluates these formulas using only entries of L that are already finished. All variants below do exactly the same floating-point operations. In exact arithmetic they produce the same L. They differ only in the order of the three nested loops over i, j, k.

Variant	Outer loop	Produces per step	Reads	Writes
right-looking	k	column k of L	column k	whole trailing Schur complement
left-looking	j	column j of L	columns $1, \ldots, j-1$	column j only
up-looking	i	row i of L	rows $1, \ldots, i-1$	row i only

The picture below shows one step of each variant.



Left-looking. The right-looking algorithm is *eager*. As soon as a column is finished, it applies that column's update to the whole trailing block. The left-looking algorithm is *lazy*. When it reaches column j, it applies all the updates from the finished columns to its left at once. Then it normalizes the column. So during step j it writes only to column j, and after that step column j is never touched again. In the right-looking variant, by

contrast, every step writes to the whole trailing block. The innermost loop runs down a column. This suits column-major languages (Julia, Fortran, MATLAB).

Algorithm 13 Cholesky factorization, left-looking, in place (lower triangle).

1: **for** $j = 1, \dots, m$ **do**
2: **for** $k = 1, \dots, j-1$ **do**
3: **for** $i = j, \dots, m$ **do**
4: $A_{ij} \leftarrow A_{ij} - A_{ik}\overline{A_{jk}}$
5: $s \leftarrow \sqrt{A_{jj}}$ ▷ save it: line 7 overwrites A_{jj}
6: **for** $i = j, \dots, m$ **do**
7: $A_{ij} \leftarrow A_{ij}/s$

Up-looking. Here L is computed row by row. Row i depends only on its own earlier entries and on the finished rows above it. Row i of L is column i of $R = L^*$. So a column-major code stores the upper triangular factor R instead. Then walking along a row of L becomes walking down a column of R. In *packed storage*, we store the upper triangle column by column. Column i has length i and comes right after column $i-1$:

$$\begin{bmatrix} 1 & 2 & 4 & 7 & 11 \\ & 3 & 5 & 8 & 12 \\ & & 6 & 9 & 13 \\ & & & 10 & 14 \\ & & & & 15 \end{bmatrix}. \tag{9.14}$$

Computing column i uses only earlier data. So when A grows by one row and one column, we just append one new segment at the end of the factor. Nothing computed before has to change.

Algorithm 14 Cholesky factorization, up-looking, in place (lower triangle).

1: **for** $i = 1, \dots, m$ **do**
2: **for** $j = 1, \dots, i$ **do**
3: **for** $k = 1, \dots, j-1$ **do**
4: $A_{ij} \leftarrow A_{ij} - A_{ik}\overline{A_{jk}}$
5: **if** $j < i$ **then**
6: $A_{ij} \leftarrow A_{ij}/A_{jj}$ ▷ A_{jj} already holds L_{jj}
7: **else**
8: $A_{ii} \leftarrow \sqrt{A_{ii}}$

In Algorithms 13 and 14 it is tempting to divide by $\sqrt{A_{jj}}$ in every case. This is wrong.

- In the left-looking loop, A_{jj} itself is overwritten when $i = j$. This is why that algorithm saves s first.
- In the up-looking loop, for $j < i$ the entry A_{jj} already holds L_{jj}. This is why that algorithm divides by A_{jj}.

9.4 Existence: Cholesky Never Breaks Down

We have seen that LU without pivoting can meet a zero pivot. For example, $\left[\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right]$ fails at the very first step. Cholesky needs even more. The pivots must be *real and positive*, because we take their square roots. In this section we show that for s.p.d. matrices this always holds.

Here is the idea of the proof. After k steps of symmetric elimination, our current matrix is A times a nonsingular matrix on the right, and times its conjugate transpose on the left. We first show that such a "sandwich" of an s.p.d. matrix is again s.p.d. Then all its diagonal entries are positive. In particular, the next pivot is positive.

Lemma 46 (Congruence Preserves Positive Definiteness)

Let $A \in \mathbb{K}^{m\times m}$ be s.p.d. and $B \in \mathbb{K}^{m\times m}$ be nonsingular. Then B^*AB is s.p.d.

Proof. First, B^*AB is Hermitian, since $(B^*AB)^* = B^*A^*B = B^*AB$. Next we show positive definiteness. Let $\vec{x} \neq \vec{0}$. Since B is nonsingular, $B\vec{x} \neq \vec{0}$. Hence

$$\vec{x}^* B^* AB\vec{x} = (B\vec{x})^* A(B\vec{x}) > 0, \tag{9.15}$$

as desired. □

The map $A \mapsto B^*AB$ is called a *congruence*. We do need B to be nonsingular. If $B\vec{x} = \vec{0}$ for some $\vec{x} \neq \vec{0}$, then $\vec{x}^* B^* AB\vec{x} = 0$. Then we only get positive semidefiniteness.

Theorem 47 (Existence of the Cholesky Factorization)

Let $A \in \mathbb{K}^{m\times m}$ be s.p.d. Then Cholesky without pivoting meets only real, positive pivots $d_1, \dots, d_m$. Hence it produces $A = LL^*$, where L is lower triangular with positive diagonal. Also, the pivots have the lower bound

$$d_k \geq \frac{1}{\|A^{-1}\|_2} = \lambda_{\min}(A), \qquad k = 1, \dots, m. \tag{9.16}$$

Proof. We show by induction on $k = 0, \dots, m-1$ that the pivot d_{k+1} is real and positive.

Suppose that the first k pivots are positive. Then the elimination matrices $L_1, \dots, L_k$ of the first k steps are defined. (For $k = 0$ there is nothing to assume.) After k symmetric steps, we have

$$S \coloneqq L_k \cdots L_1 A L_1^* \cdots L_k^* = \begin{bmatrix} \operatorname{diag}(d_1, \dots, d_k) & 0 \\ 0 & S^{(k)} \end{bmatrix}, \tag{9.17}$$

where $S^{(k)}$ is the current Schur complement. Its top left entry is the next pivot, $d_{k+1} = S_{k+1,k+1}$.

Let $B = L_1^* \cdots L_k^*$, so that $S = B^*AB$. Each L_i is unit lower triangular, so it has determinant 1. Hence B is nonsingular. By Lemma 46, S is s.p.d. Now:

- *The pivot is real.* S is Hermitian, so $S_{k+1,k+1} = \overline{S_{k+1,k+1}}$.
- *The pivot is positive.* A standard basis vector picks out a diagonal entry, $\vec{e}_j^* S\vec{e}_j = S_{jj}$. So by positive definiteness,

$$d_{k+1} = S_{k+1,k+1} = \vec{e}_{k+1}^* S\vec{e}_{k+1} > 0. \tag{9.18}$$

This completes the induction. With all pivots positive, the construction of the previous section gives $A = LL^*$ with $L_{kk} = \sqrt{d_k} > 0$.

The lower bound (9.16) needs a fact about Schur complements. We prove it in Section 9.6. □

Note that "real" and "positive" come from different assumptions. The pivot is real because S is Hermitian. It is positive because S is positive definite. Along the way we also proved a useful fact: *every diagonal entry of a positive definite matrix is positive*. The lower bound (9.16) says even more than positivity. The pivots stay a fixed distance away from zero, for every k. So tiny pivots cannot occur.

Corollary 48 (Uniqueness of the Cholesky Factorization)**.** *An s.p.d. matrix has exactly one factorization $A = LL^*$, where L is lower triangular with positive diagonal.*

Proof. Existence is Theorem 47. Suppose $A = LL^* = MM^*$ with both factors of this form. Then $M^{-1}L = M^*L^{-*}$. The left side is lower triangular, and the right side is upper triangular. So both sides equal a diagonal matrix Δ. Its entries L_{kk}/M_{kk} are positive. Also,

$$\Delta\Delta^* = (M^{-1}L)(M^{-1}L)^* = M^{-1}LL^*M^{-*} = M^{-1}MM^*M^{-*} = I, \tag{9.19}$$

so $|\Delta_{kk}| = 1$. A positive number of absolute value 1 is 1, so $\Delta = I$ and $L = M$. □

9.5 Rounding Errors

Existence tells us that the algorithm runs to completion in exact arithmetic. We now ask what happens in floating point.

The key quantity for LU. Recall from the error analysis of LU (Theorem 16) that the computed factors satisfy $\tilde{L}\tilde{U} = A + \delta A$ with

$$|\delta A| \le cm\varepsilon_{\text{mach}} \left|\tilde{L}\right| \left|\tilde{U}\right| \quad \text{(entrywise)}. \tag{9.20}$$

So the key quantity is $|L|\,|U|$. If it is of the same size as $|A|$, the algorithm is backward stable. If it is much larger, errors are amplified. Without pivoting, $|L|\,|U|$ can be arbitrarily large. For example, for $A = \left[\begin{smallmatrix} \delta & 1 \\ 1 & 1 \end{smallmatrix}\right]$ with small δ, L contains the entry $1/\delta$.

No element growth in Cholesky. For Cholesky, $U = L^*$, so the key quantity is $|L|\,|L^*|$. We will show that it is never larger than the largest entry of A.

Proposition 49 (No Growth in the Cholesky Factor)

Let $A = LL^*$ be s.p.d. Then for all i, j,

$$|L_{ij}| \le \sqrt{A_{ii}}, \qquad (|L|\,|L^*|)_{ij} \le \sqrt{A_{ii}A_{jj}} \le \max_{k,l} |A_{kl}|\,. \tag{9.21}$$

Proof. We go in four steps.

Step 1: the diagonal of A. The (i, i) entry of $A = LL^*$ is

$$A_{ii} = \sum_k L_{ik}\overline{L_{ik}} = \sum_k |L_{ik}|^2\,.$$

So A_{ii} is the squared length of row i of L. Each $|L_{ij}|^2$ is one term of this sum, so $|L_{ij}|^2 \le A_{ii}$. This is the first claim.

Step 2: an entry of $|L|\,|L^*|$. By the rule for matrix products, with $|L^*|_{kj} = |L_{jk}|$,

$$(|L|\,|L^*|)_{ij} = \sum_k |L_{ik}|\,|L_{jk}|\,.$$

Step 3: Cauchy–Schwarz. For real numbers a_k, b_k, the Cauchy–Schwarz inequality says

$$\sum_k a_k b_k \le \left(\sum_k a_k^2\right)^{1/2} \left(\sum_k b_k^2\right)^{1/2}.$$

We take $a_k = |L_{ik}|$ and $b_k = |L_{jk}|$. By Step 1, the two sums on the right are A_{ii} and A_{jj}. So

$$\sum_k |L_{ik}|\,|L_{jk}| \le \left(\sum_k |L_{ik}|^2\right)^{1/2} \left(\sum_k |L_{jk}|^2\right)^{1/2} = \sqrt{A_{ii}}\sqrt{A_{jj}}.$$

Step 4: compare with the largest entry. The geometric mean of two numbers is at most the larger one. Both A_{ii} and A_{jj} are entries of A. So

$$\sqrt{A_{ii}A_{jj}} \le \max(A_{ii}, A_{jj}) \le \max_{k,l} |A_{kl}|\,.$$

Steps 2–4 together give the second claim. □

The same chain also shows where the large entries of an s.p.d. matrix are. Since $A_{ij} = \sum_k L_{ik}\overline{L_{jk}}$, we have $|A_{ij}| \le \sum_k |L_{ik}|\,|L_{jk}| \le \sqrt{A_{ii}}\sqrt{A_{jj}}$. So the largest entry of A is on its diagonal: $\max_{i,j} |A_{ij}| = \max_i A_{ii}$.

Here $\max_{k,l} |A_{kl}| = \|\mathrm{vec}(A)\|_\infty$ is the largest entry of A. (The vector $\mathrm{vec}(A) \in \mathbb{K}^{m^2}$ stacks the columns of A.) It is *not* the induced ∞-norm (maximum row sum) of Proposition 103. In words: the growth factor of Cholesky is 1. We put this into the backward error bound and get the main result.

Theorem 50 (Backward Stability of Cholesky)

Let $A \in \mathbb{K}^{m\times m}$ be s.p.d. Suppose Cholesky runs to completion in floating-point arithmetic and produces $\tilde{L}$. Then, to first order in $\varepsilon_{\mathrm{mach}}$,

$$\tilde{L}\tilde{L}^* = A + \delta A, \qquad |\delta A_{ij}| \le cm\varepsilon_{\mathrm{mach}} \max_{k,l} |A_{kl}| \tag{9.22}$$

for a modest constant c. That is, Cholesky without pivoting is backward stable.

Proof sketch. The analysis of Theorem 16 applies with $\tilde{U} = \tilde{L}^*$ and gives $|\delta A| \le cm\varepsilon_{\mathrm{mach}} \left|\tilde{L}\right| \left|\tilde{L}^*\right|$ [7, Theorem 10.3]. Strictly speaking, Proposition 49 is about the exact factor. The computed factor satisfies $\sum_k \left|\tilde{L}_{ik}\right|^2 = A_{ii} + \delta A_{ii}$. This differs by a term of order $\varepsilon_{\mathrm{mach}}$, which changes the bound only at second order. Hence $\left|\tilde{L}\right| \left|\tilde{L}^*\right| \le \max_{k,l} |A_{kl}|$ to first order, as desired. □

So Theorems 47 and 50 answer two different questions. The first says that the algorithm runs to completion in exact arithmetic. The second says that in floating point its answer is accurate. In LU, we need pivoting to fix both problems. For s.p.d. matrices, both hold automatically. (In floating point, completion itself is guaranteed as long as $\kappa_2(A)$ is not too close to $1/\varepsilon_{\mathrm{mach}}$ [7, Theorem 10.7].)

Symmetric indefinite matrices. What if A is Hermitian but not positive definite? A Hermitian matrix is *indefinite* if $\vec{x}^* A\vec{x}$ takes both positive and negative values. Equivalently, it has both positive and negative eigenvalues. For such a matrix, pivots can be zero or negative, and Cholesky fails. Instead, we compute

$$PAP^\top = LDL^\top, \tag{9.23}$$

where P is a permutation and D is block diagonal with 1×1 and 2×2 blocks (LDL^* in the complex case).

Why do we use the symmetric permutation $PAP^\top$, and not PA as in LU?

- The product $PAP^\top$ swaps rows and columns in the same way. So it keeps the matrix symmetric, and we can still work on one half.
- PA swaps only rows. This destroys symmetry.

But a symmetric permutation has a limit. It only moves diagonal entries to other diagonal positions. It can never bring an off-diagonal entry onto the diagonal. For example, consider

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}. \tag{9.24}$$

Its diagonal is zero, so every 1×1 pivot is zero no matter how we permute. The solution is to take the whole 2×2 block as one pivot. This block is invertible, so it becomes one block of D, with $L = I$.

In general, each step moves an invertible 1×1 or 2×2 block E to the top left. Then it factors

$$PAP^\top = \begin{bmatrix} E & C^* \\ C & B \end{bmatrix} = \begin{bmatrix} I & 0 \\ CE^{-1} & I \end{bmatrix} \begin{bmatrix} E & 0 \\ 0 & B - CE^{-1}C^* \end{bmatrix} \begin{bmatrix} I & E^{-1}C^* \\ 0 & I \end{bmatrix}, \tag{9.25}$$

and repeats on the Schur complement $B - CE^{-1}C^*$, just as in the derivation of Cholesky. In practice, we choose the blocks by *Bunch–Kaufman pivoting* (LAPACK `sytrf`). If a diagonal entry is too small compared with the off-diagonal entries in its column, it uses a 2×2 block instead. This avoids small pivots. Its stability is about the same as LU with partial pivoting. It costs about $\frac{1}{3}m^3$ flops.

9.6 Schur Complements

So far we eliminated one row and column at a time. Let us now do the same thing with blocks. This leads to the Schur complement. It appears everywhere in numerical linear algebra (and also in statistics and optimization).

Block Cholesky. We partition $A = LL^*$ into 2×2 blocks. Here L_{11} and L_{22} are square and lower triangular:

$$L = \begin{bmatrix} L_{11} & 0 \\ L_{21} & L_{22} \end{bmatrix}, \qquad A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}, \qquad LL^* = \begin{bmatrix} L_{11}L_{11}^* & L_{11}L_{21}^* \\ L_{21}L_{11}^* & L_{21}L_{21}^* + L_{22}L_{22}^* \end{bmatrix}. \tag{9.26}$$

Comparing blocks, we have

$$L_{11}L_{11}^* = A_{11}, \qquad L_{21}L_{11}^* = A_{21}, \qquad L_{21}L_{21}^* + L_{22}L_{22}^* = A_{22}. \tag{9.27}$$

The top right block gives the conjugate transpose of the second equation, so it gives nothing new.

We want to find $L_{22}L_{22}^*$. The second equation gives

$$L_{21} = A_{21}L_{11}^{-*}.$$

Using $A_{21}^* = A_{12}$, we get

$$L_{21}L_{21}^* = A_{21}L_{11}^{-*}L_{11}^{-1}A_{21}^* = A_{21}(L_{11}L_{11}^*)^{-1}A_{21}^* = A_{21}A_{11}^{-1}A_{12}. \tag{9.28}$$

We put this into the third equation and get

$$L_{22}L_{22}^* = A_{22} - A_{21}A_{11}^{-1}A_{12}. \tag{9.29}$$

The matrix on the right has a name.

Definition 51 (Schur Complement)

Let $A = \left[\begin{smallmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{smallmatrix}\right]$ with A_{11} square and invertible. The *Schur complement* of A_{11} in A is

$$A/A_{11} \coloneqq A_{22} - A_{21}A_{11}^{-1}A_{12}. \tag{9.30}$$

The Schur complement S from Step 1 of Section 9.2 is the special case where $A_{11} = \alpha$ is 1×1. Then A_{11}^{-1} is just division by a scalar.

Note that the Schur complement is *defined* as $A_{22} - A_{21}A_{11}^{-1}A_{12}$. The block equations show that it is *equal* to $L_{22}L_{22}^*$. This tells us two things.

1. L_{22} is the Cholesky factor of the Schur complement. So the algorithm can recurse on blocks instead of single columns.
2. The Schur complement equals $L_{22}L_{22}^*$ with L_{22} invertible, so it is s.p.d. This gives a second proof that the recursion never breaks down. (The first proof was the congruence argument.)

The Schur complement is also exactly what the column-by-column elimination of Section 9.4 computes. Let A_{11} be the leading $k \times k$ block. Then after k steps of symmetric elimination,

$$L_k \cdots L_1 A L_1^* \cdots L_k^* = \begin{bmatrix} \operatorname{diag}(d_1, \ldots, d_k) & 0 \\ 0 & A/A_{11} \end{bmatrix}. \tag{9.31}$$

Indeed, both come from eliminating the first k rows and columns of A. The block equations above show that what is left in the lower right is $L_{22}L_{22}^* = A/A_{11}$.

In other words, *k steps of elimination produce the Schur complement of the leading $k \times k$ block.*

The inverse and the Schur complement. The following lemma is the key tool for the pivot bound. It relates the Schur complement to a block of the inverse.

Lemma 52 (Blocks of the Inverse)

Let $A = \left[\begin{smallmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{smallmatrix}\right]$ be invertible with A_{11} invertible, let $S = A_{22} - A_{21}A_{11}^{-1}A_{12}$ be invertible, and partition $\Theta \coloneqq A^{-1}$ in the same way. Then

$$\Theta_{22} = S^{-1}. \tag{9.32}$$

Proof. The idea is to compute a block LU factorization of A and invert it. Block elimination gives

$$A = \begin{bmatrix} I & 0 \\ A_{21}A_{11}^{-1} & I \end{bmatrix} \begin{bmatrix} A_{11} & 0 \\ 0 & S \end{bmatrix} \begin{bmatrix} I & A_{11}^{-1}A_{12} \\ 0 & I \end{bmatrix}. \tag{9.33}$$

(Multiply it out to check.) To invert a unit block triangular matrix, we negate its off-diagonal block. So

$$A^{-1} = \begin{bmatrix} I & -A_{11}^{-1}A_{12} \\ 0 & I \end{bmatrix} \begin{bmatrix} A_{11}^{-1} & 0 \\ 0 & S^{-1} \end{bmatrix} \begin{bmatrix} I & 0 \\ -A_{21}A_{11}^{-1} & I \end{bmatrix}. \tag{9.34}$$

To read off the lower right block of the product, we only need two pieces:

- the second block row of the left factor, which is $\begin{bmatrix} 0 & I \end{bmatrix}$;
- the second block column of the right factor, which is $\begin{bmatrix} 0 & I \end{bmatrix}^\top$.

So the lower right block of A^{-1} is

$$\begin{bmatrix} 0 & I \end{bmatrix} \begin{bmatrix} A_{11}^{-1} & 0 \\ 0 & S^{-1} \end{bmatrix} \begin{bmatrix} 0 \\ I \end{bmatrix} = S^{-1}.$$

□

The proof uses only that A_{11} and S are invertible. It needs neither symmetry nor definiteness.

In words: *to get the Schur complement, invert, take the submatrix, and invert again*:

$$A/A_{11} = \left((A^{-1})_{22}\right)^{-1}. \tag{9.35}$$

Watch the inverse here. It is easy to remember the lemma wrongly as $\Theta_{22} = A_{22} - A_{21}A_{11}^{-1}A_{12}$. But the block of the inverse is S^{-1}, not S. The slogan "invert, take submatrix, invert" helps us remember the correct version.

Proof of the pivot bound. We can now finish the proof of Theorem 47. The plan has three parts.

- The k-th pivot is a diagonal entry of a Schur complement S'.
- A diagonal entry is at least $\lambda_{\min}(S')$.
- By Lemma 52, $(S')^{-1}$ is a submatrix of A^{-1}. So it cannot be larger than A^{-1}.

Proof of (9.16). Fix k. Let S' be the Schur complement of the leading $(k-1) \times (k-1)$ block of A (for $k = 1$, $S' = A$). As shown above, S' is the matrix left over after $k-1$ steps of elimination. So the k-th pivot is its top left entry, $d_k = \vec{e}_1^* S' \vec{e}_1$. We bound d_k in three steps.

1. *A diagonal entry is at least the smallest eigenvalue.* S' is s.p.d., so by the Rayleigh quotient bound $\vec{x}^* S' \vec{x} \geq \lambda_{\min}(S') \|\vec{x}\|^2$,
$$d_k = \vec{e}_1^* S' \vec{e}_1 \geq \lambda_{\min}(S') = \frac{1}{\|(S')^{-1}\|_2}, \tag{9.36}$$
where the last equality holds because the singular values of an s.p.d. matrix are its eigenvalues (Theorem 113).

2. *The inverse of the Schur complement is a submatrix of A^{-1}.* By Lemma 52, $(S')^{-1} = \Theta[k{:}m, k{:}m]$ with $\Theta = A^{-1}$.

3. *A submatrix is no larger than the whole matrix.* We have $\Theta[k{:}m, k{:}m] = P^\top \Theta P$, where $P = I[:, k{:}m]$ has orthonormal columns. So $\|P\|_2 = 1$ and
$$\left\|(S')^{-1}\right\|_2 = \left\|P^\top \Theta P\right\|_2 \leq \|\Theta\|_2 = \left\|A^{-1}\right\|_2.$$

Putting the three steps together,
$$d_k \geq \frac{1}{\|(S')^{-1}\|_2} \geq \frac{1}{\|A^{-1}\|_2} = \lambda_{\min}(A), \tag{9.37}$$
where the last equality again uses that A is s.p.d. □

9.7 Cholesky and QR

We end with a link back to least squares. The Cholesky factorization and the QR factorization are two views of the same thing. Let $B \in \mathbb{K}^{m\times n}$ have full column rank. Then B^*B is s.p.d. (Theorem 42). If $B = QR$, then
$$B^*B = R^*Q^*QR = R^*R. \tag{9.38}$$
So the R of QR is a Cholesky factor of B^*B. The next theorem states this exactly, and also gives the converse.

Theorem 53 (QR of B Is Cholesky of B^*B)

Let $B \in \mathbb{K}^{m\times n}$ have full column rank.

1. If $B = \hat{Q}\hat{R}$ is a reduced QR factorization with $r_{jj} > 0$, then $B^*B = \hat{R}^*\hat{R}$ is the Cholesky factorization of B^*B.
2. Conversely, if $B^*B = R^*R$ is the Cholesky factorization, then $B = (BR^{-1})R$ is a reduced QR factorization: BR^{-1} has orthonormal columns.

Proof. Item 1. We have $B^*B = \hat{R}^*\hat{Q}^*\hat{Q}\hat{R} = \hat{R}^*\hat{R}$, and $\hat{R}^*$ is lower triangular with positive diagonal. By Corollary 48, this is *the* Cholesky factorization.

Item 2. R is invertible because its diagonal is positive. So
$$(BR^{-1})^*(BR^{-1}) = R^{-*}B^*BR^{-1} = R^{-*}R^*RR^{-1} = I, \tag{9.39}$$
as desired. □

So QR of B and Cholesky of B^*B compute the same triangular factor. This has three consequences.

- *Uniqueness of QR.* The uniqueness of Cholesky (Corollary 48) gives the uniqueness of reduced QR with $r_{jj} > 0$ (Theorem 33). Indeed, $\hat{R}$ must be the Cholesky factor of B^*B, and then $\hat{Q} = B\hat{R}^{-1}$.
- *Least squares.* Solving the normal equations with Cholesky computes the same R as QR. But it goes through B^*B, whose condition number is $\kappa_2(B)^2$. The two methods agree in exact arithmetic, but not in floating point.
- *CholeskyQR.* Item 2 is itself an algorithm: form B^*B, factor it, and set $\hat{Q} = BR^{-1}$. It is fast, and it is mostly matrix multiplications. But for the same reason (it goes through B^*B), the computed $\hat{Q}$ loses orthogonality like $\varepsilon_{\text{mach}}\kappa_2(B)^2$. This is worse than modified Gram–Schmidt.

9.8 (SUPPLEMENT) Further Topics

9.8.1 Over $\mathbb{R}$, "Positive Definite" Does Not Imply Symmetric

If $A = B^*B$ with B real, then A is automatically symmetric. But suppose we only ask that $\vec{x}^\top A\vec{x} > 0$ for all real $\vec{x} \neq \vec{0}$. Then A need not be symmetric. For example, $A = \left[\begin{smallmatrix} 1 & 1 \\ -1 & 1 \end{smallmatrix}\right]$ has $\vec{x}^\top A\vec{x} = x_1^2 + x_2^2$.

The reason is this. Every real square matrix is the sum of a symmetric part and a skew-symmetric part K. Since $\vec{x}^\top K\vec{x} = 0$, the quadratic form only sees the symmetric part. This is why, over $\mathbb{R}$, symmetry is stated as a separate condition in "s.p.d.".

Over $\mathbb{C}$, symmetry is automatic. Suppose $\vec{x}^* A\vec{x}$ is real for all complex $\vec{x}$. Then $\vec{x}^*(A - A^*)\vec{x} = 0$ for all $\vec{x}$. Substituting $\vec{x} = \vec{e}_j + \vec{e}_k$ and $\vec{x} = \vec{e}_j + i\vec{e}_k$ (polarization) gives $A = A^*$. The example above, at $\vec{x} = (1, i)^\top$, gives $2 + 2i$. So it is not positive definite over $\mathbb{C}$.

9.8.2 Cholesky as a Test for Positive Definiteness

Congruences by unit lower triangular matrices do not change the determinants of the leading principal submatrices. So $\det A[1{:}k, 1{:}k] = d_1 \cdots d_k$. Hence the pivots are ratios of consecutive leading principal minors:

$$d_k = \frac{\det A[1{:}k, 1{:}k]}{\det A[1{:}k-1, 1{:}k-1]}. \tag{9.40}$$

Combined with Sylvester's criterion (a Hermitian matrix is positive definite if and only if all its leading principal minors are positive), this gives

$$\begin{aligned} A \text{ is positive definite} &\iff \text{all leading principal minors are } > 0 \\ &\iff \text{all pivots are } > 0 \iff \text{Cholesky completes.} \end{aligned} \tag{9.41}$$

So Cholesky is the standard test for positive definiteness. It is much cheaper than computing eigenvalues. For example, `numpy.linalg.cholesky` raises `LinAlgError` exactly at this point when the input is not positive definite.

9.8.3 Blocked Cholesky and Hardware

Libraries (LAPACK and cuSOLVER `potrf`) use a blocked version, as for LU in Section 2.9.3. We take A_{11} to be a block of size, say, 256×256, and factor it. Then:

- $L_{21} = A_{21} L_{11}^{-*}$ is a triangular solve with many right-hand sides (TRSM);
- the Schur complement update $A_{22} - L_{21} L_{21}^*$ is a matrix multiplication (SYRK/GEMM).

So almost all flops are Level 3 BLAS operations, and these run close to peak speed. This continues our discussion of memory access in the left- and up-looking variants.

9.8.4 The Normal Equations, Again

Backward stability of Cholesky means this: the computed $\tilde{R}$ is the exact Cholesky factor of $A^*A + \delta$ with a small δ. But when we solve least squares through the normal equations, the matrix we factor is A^*A. So the error in $\vec{x}$ is amplified by $\kappa(A^*A) = \kappa(A)^2$. A stable Cholesky therefore does not make "normal equations plus Cholesky" a stable least squares solver. For ill-conditioned A, QR is still the best method (Lecture 8).

Lecture 10

Eigen-like Decompositions, Computing a Single Eigenvalue

Supplementary reading:

- [1] Lectures 24, 25, and 27.
- [2] Sections 4.4.1–4.4.2 and 5.3.2.
- [5] Sections 7.3.1 and 8.2.

10.1 Recap and Motivation

Last lecture we factored an s.p.d. matrix as $A = LL^*$, using symmetric elimination $A \mapsto L_1 A L_1^*$. We showed three things.

- No pivoting is needed.
- There is no element growth.
- The steps of elimination compute Schur complements (Lemma 52).

The last statement is easy to get backwards, so we recall it exactly. Let $\Theta = A^{-1}$. Then the Schur complement $A_{22} - A_{21}A_{11}^{-1}A_{12}$ equals Θ_{22}^{-1}, not Θ_{22}.

From elimination to similarity transformations. So far we have seen LU, Cholesky, and QR. All of them are based on *elimination* or *projection*.

- LU and Cholesky are Gaussian elimination: lower triangular matrices zero out the entries below the pivots, column by column.
- QR uses projection. Gram–Schmidt subtracts from each column its projections onto the previous columns. Householder reflects each column onto the direction of $\vec{e}_1$.

These factorizations have two things in common.

1. Their purpose is to *solve equations*. They use one-sided transformations ($A \mapsto LA$, $A \mapsto Q^*A$) or congruences ($A \mapsto LAL^*$). None of these preserve eigenvalues. This is why the pivots of Cholesky are not the eigenvalues: a congruence is not a similarity.
2. They finish in finitely many steps, about $O(m^3)$. Without rounding, they are exact.

Starting with this lecture we study a second family: the eigen, polar, Schur, and singular value decompositions. All of them are closely tied to the *eigenpairs* $(\lambda_i, \vec{x}_i)$ of a matrix. They use *similarity transformations* $A \mapsto X^{-1}AX$, which preserve eigenvalues. In this way they show the structure of A as a linear map.

Factorization	Form	Exists for	Transformation
Eigen	$A = X\Lambda X^{-1}$	diagonalizable (nondefective) A	general X, possibly ill-conditioned
Polar	$A = QP$, $P = (A^*A)^{1/2} \succeq 0$	every square A	Q unitary
Schur	$A = QTQ^*$, T upper triangular	every square A	Q unitary
SVD	$A = U\Sigma V^*$	every A	U, V unitary

A few remarks on the table.

- Only the eigendecomposition may fail to exist. The other three exist for every (square) matrix. They use only unitary transformations, which have condition number 1 and do not amplify errors.
- A *normal* matrix ($A^*A = AA^*$, which includes Hermitian matrices) can be diagonalized by a unitary matrix, $A = Q\Lambda Q^*$.
- The polar decomposition (Theorem 40) is the matrix version of $z = e^{i\theta}r$. Here Q is a "rotation or reflection" and P is a "stretch". It follows directly from the SVD:

$$A = U\Sigma V^* = (UV^*)(V\Sigma V^*), \qquad Q = UV^*, \quad P = V\Sigma V^*. \tag{10.1}$$

  The factor Q is the unitary matrix closest to A in the Frobenius norm (the orthogonal Procrustes problem).
- The Schur *decomposition* has nothing directly to do with the Schur *complement* of Lecture 9.

We start with the eigendecomposition of Hermitian matrices. Here is our plan.

1. Recall the basic facts about eigenpairs of Hermitian matrices and introduce the Rayleigh quotient.
2. Show that eigenvalues *cannot* be computed exactly in finitely many steps, so every eigenvalue algorithm must be iterative.
3. Develop three iterative methods, each improving on the previous one: power iteration, inverse iteration, and Rayleigh quotient iteration.

10.2 Eigenpairs of Hermitian Matrices

Problem. Given a Hermitian $A \in \mathbb{K}^{m\times m}$, find $\lambda_1, \dots, \lambda_m \in \mathbb{R}$ and $\vec{x}_1, \dots, \vec{x}_m \in \mathbb{K}^m$ with

$$A\vec{x}_i = \lambda_i \vec{x}_i, \qquad i = 1, \dots, m. \tag{10.2}$$

Recall that $\langle \vec{x}, \vec{y} \rangle = \vec{x}^*\vec{y}$ is conjugate-linear in its first argument.

Real eigenvalues and the Rayleigh quotient. Proposition 39 showed that the eigenvalues of a Hermitian matrix are real. With inner products, the argument is the following chain:

$$\lambda_i = \frac{\langle \vec{x}_i, A\vec{x}_i \rangle}{\|\vec{x}_i\|^2} = \frac{\langle A\vec{x}_i, \vec{x}_i \rangle}{\|\vec{x}_i\|^2} = \overline{\lambda_i}. \tag{10.3}$$

We check each step. We use two facts. First, the inner product $\langle \vec{x}, \vec{y} \rangle = \vec{x}^* \vec{y}$ is linear in its second argument and conjugate-linear in its first. Second, $A = A^*$.

- *The first equality.* We put $A\vec{x}_i = \lambda_i \vec{x}_i$ into the second argument. There the scalar comes out unchanged:

$$\langle \vec{x}_i, A\vec{x}_i \rangle = \vec{x}_i^* (\lambda_i \vec{x}_i) = \lambda_i \, \vec{x}_i^* \vec{x}_i = \lambda_i \, \|\vec{x}_i\|^2 . \tag{10.4}$$

- *The second equality.* We move A to the other side of the inner product. For this we use $(A\vec{x})^* = \vec{x}^* A^*$ and $A = A^*$:

$$\langle \vec{x}_i, A\vec{x}_i \rangle = \vec{x}_i^* A \vec{x}_i = \vec{x}_i^* A^* \vec{x}_i = (A\vec{x}_i)^* \vec{x}_i = \langle A\vec{x}_i, \vec{x}_i \rangle . \tag{10.5}$$

  In general $\langle \vec{x}, A\vec{y} \rangle = \langle A^*\vec{x}, \vec{y} \rangle$. So for a Hermitian A, we can move A freely between the two arguments.

- *The third equality.* We put $A\vec{x}_i = \lambda_i \vec{x}_i$ in again, but now into the first argument. There the conjugate transpose turns λ_i into $\overline{\lambda_i}$:

$$\langle A\vec{x}_i, \vec{x}_i \rangle = (\lambda_i \vec{x}_i)^* \vec{x}_i = \overline{\lambda_i} \, \vec{x}_i^* \vec{x}_i = \overline{\lambda_i} \, \|\vec{x}_i\|^2 . \tag{10.6}$$

Thus $\lambda_i = \overline{\lambda_i}$, i.e., $\lambda_i \in \mathbb{R}$.

Here is a shorter argument. The number $\vec{x}^* A \vec{x}$ is a 1×1 matrix that equals its own conjugate transpose:

$$(\vec{x}^* A \vec{x})^* = \vec{x}^* A^* \vec{x} = \vec{x}^* A \vec{x}.$$

So it is real for *every* $\vec{x}$. Dividing by the real number $\|\vec{x}_i\|^2 > 0$ gives $\lambda_i \in \mathbb{R}$.

Even for a real symmetric A, the argument must allow $\vec{x}_i \in \mathbb{C}^m$. We do not know yet that λ_i is real, so we cannot assume that its eigenvector is real. If we take $\vec{x}_i \in \mathbb{R}^m$ from the start, we assume what we want to prove.

The first expression makes sense for *any* nonzero vector, not just for eigenvectors. This leads to the following definition. It will be our main tool for estimating eigenvalues.

Definition 54 (Rayleigh Quotient)

Let $A \in \mathbb{K}^{m \times m}$. The *Rayleigh quotient* of a nonzero $\vec{x} \in \mathbb{K}^m$ is

$$r(\vec{x}) := \frac{\vec{x}^* A \vec{x}}{\vec{x}^* \vec{x}}. \tag{10.7}$$

If A is Hermitian, the Rayleigh quotient is always real. At an eigenvector it equals the eigenvalue. So if we have a good approximate eigenvector, its Rayleigh quotient is a natural guess for the eigenvalue. We will see that this guess is surprisingly accurate.

Proposition 55 (Orthogonal Eigenvectors)

Let A be Hermitian, $A\vec{x}_i = \lambda_i \vec{x}_i$ and $A\vec{x}_j = \lambda_j \vec{x}_j$ with $\lambda_i \neq \lambda_j$. Then $\vec{x}_i \perp \vec{x}_j$.

Proof. We have

$$\lambda_i \langle \vec{x}_i, \vec{x}_j \rangle = \langle A\vec{x}_i, \vec{x}_j \rangle = \langle \vec{x}_i, A\vec{x}_j \rangle = \lambda_j \langle \vec{x}_i, \vec{x}_j \rangle . \tag{10.8}$$

The first equality uses $\langle \lambda_i \vec{x}_i, \vec{x}_j \rangle = \overline{\lambda_i} \langle \vec{x}_i, \vec{x}_j \rangle$ and $\overline{\lambda_i} = \lambda_i$ (Proposition 39). The second uses $A = A^*$. Thus $(\lambda_i - \lambda_j) \langle \vec{x}_i, \vec{x}_j \rangle = 0$. Since $\lambda_i \neq \lambda_j$, we get $\langle \vec{x}_i, \vec{x}_j \rangle = 0$, as desired. □

The proof uses that the eigenvalues are real. So we must prove the two facts in this order.

For a repeated eigenvalue, we can make the eigenvectors in its eigenspace orthonormal by Gram–Schmidt. So orthogonality is never the problem. What we have *not* yet proved is *existence*: why does a Hermitian matrix have m linearly independent eigenvectors? This is the spectral theorem $A = Q\Lambda Q^*$. The usual proof first shows the Schur decomposition $A = QTQ^*$ for every square matrix. Then $T = Q^*AQ$ is both Hermitian and upper triangular, so it is diagonal. We do this in Lecture 11.

10.3 A Negative Result: Eigenvalues Cannot Be Computed in Finitely Many Steps

LU and its relatives can be computed exactly (up to rounding) in $O(m^3)$ arithmetic operations. Can we do the same for eigenvalues? The answer is no. The reason comes from algebra.

Theorem 56 (Abel–Ruffini)

For every $m \geq 5$, there is a polynomial of degree m with rational coefficients whose roots cannot be expressed in terms of its coefficients using finitely many operations $+$, $-$, $\times$, $\div$, and k-th roots.

Why is this a problem for eigenvalues? We can turn any polynomial into a matrix whose eigenvalues are its roots. So if we could compute eigenvalues exactly, we could find the roots of any polynomial exactly. This contradicts Abel. Let us make this precise.

Lemma 57 (Companion Matrix)

Let $p(z) = z^m + a_{m-1}z^{m-1} + \cdots + a_1 z + a_0$ be monic, and let

$$A = \begin{bmatrix} 0 & 1 & & & \\ & 0 & 1 & & \\ & & \ddots & \ddots & \\ & & & 0 & 1 \\ -a_0 & -a_1 & \cdots & -a_{m-2} & -a_{m-1} \end{bmatrix}, \qquad \vec{x} = \begin{bmatrix} 1 \\ z \\ z^2 \\ \vdots \\ z^{m-1} \end{bmatrix}. \tag{10.9}$$

If $p(z) = 0$, then $A\vec{x} = z\vec{x}$, so z is an eigenvalue of A with eigenvector $\vec{x} \neq \vec{0}$.

Proof. We check $A\vec{x} = z\vec{x}$ row by row.

- *The first $m-1$ rows.* Row i of A picks out the next entry of $\vec{x}$: $(A\vec{x})_i = x_{i+1} = z^i = z \cdot z^{i-1} = zx_i$.
- *The last row.* Using $p(z) = 0$,

$$(A\vec{x})_m = -\sum_{i=0}^{m-1} a_i z^i = z^m - \underbrace{p(z)}_{=0} = z \cdot z^{m-1} = zx_m. \tag{10.10}$$

Finally $\vec{x} \neq \vec{0}$ since $x_1 = 1$. □

If we expand $\det(zI - A)$ along the last row, we get more: $\det(zI - A) = p(z)$. So the eigenvalues of A are *exactly* the roots of p, counted with multiplicity. If p is not monic, we first divide by the leading coefficient a_m. The last row is then $-a_i/a_m$.

Theorem 58 (Eigenvalue Algorithms Must Be Iterative)

For $m \geq 5$, there is no algorithm that computes the eigenvalues of every $m \times m$ matrix exactly in finitely many operations $+$, $-$, $\times$, $\div$, and k-th roots.

Proof. Suppose such an algorithm exists. Given any monic polynomial p of degree m, form its companion matrix. This needs no arithmetic at all, since its entries are 0, 1, and the coefficients $-a_i$. By Lemma 57 and the remark after it, the algorithm applied to this matrix returns the roots of p in finitely many operations. This contradicts Theorem 56. □

So every eigenvalue algorithm must be *iterative*: it can only produce a sequence that converges to the eigenvalues.

The direction of the reduction. We must be careful about which way the argument goes.

- The familiar direction "matrix → characteristic polynomial → roots" only shows that the eigenvalue problem is *no harder* than root finding. It proves no impossibility.
- To show that the eigenvalue problem is *at least as hard* as root finding, we need the opposite direction: every polynomial can be written as an eigenvalue problem.

Also, this step must be free. The companion matrix is built from 0, 1, and the coefficients, without any computation. If we used $\operatorname{diag}(z_1, \ldots, z_m)$ instead, we would need to know the roots already, so the argument would be circular.

Where does the companion matrix come from? It is the matrix of the linear map "multiply by z" on the quotient space $\mathbb{C}[z]/(p)$, in the basis $1, z, \ldots, z^{m-1}$. The ones on the superdiagonal move each basis vector to the next one. The power z^m falls outside the basis. The last row uses $p(z) = 0$ to write it in terms of lower powers.

The conclusion also holds for Hermitian matrices. Take a polynomial whose roots are all real. A Sturm sequence (built with arithmetic operations and square roots) gives a real symmetric tridiagonal matrix with this characteristic polynomial.

Iterative does not mean slow. The standard approach has two phases.

1. Finitely many unitary similarity transformations reduce A to Hessenberg form (tridiagonal if A is Hermitian). This phase is a direct method.
2. An iteration runs on the reduced matrix and converges very fast.

The total cost is still $O(m^3)$. The reduction above is also used in practice, in the other direction. The function `roots` in MATLAB and NumPy finds the roots of a polynomial by forming its companion matrix and calling an eigenvalue solver.

10.4 Power Iteration

We cannot hope for exact computation. So how can we *approximate* the eigenpairs of A? The simplest idea is to multiply a vector by A again and again. To see why this works, we first look at what multiplication by A does in the eigenbasis.

Multiplying by A in the eigenbasis. Let $A = X\Lambda X^{-1}$ be diagonalizable. We write any vector as a combination of eigenvectors, $\vec{z} = X\vec{y} = \sum_i y_i \vec{x}_i$. Here $\vec{y} = X^{-1}\vec{z}$ holds the coordinates of $\vec{z}$ in the eigenbasis. Then

$$A\vec{z} = X\Lambda\vec{y}, \qquad A^2\vec{z} = X\Lambda^2\vec{y}, \qquad A^k\vec{z} = X\Lambda^k\vec{y} = \sum_{i=1}^{m} \lambda_i^k y_i \vec{x}_i. \tag{10.11}$$

In words: *in the eigenbasis, multiplication by A multiplies each coordinate by its own eigenvalue. The directions do not mix.*

Now assume that the largest eigenvalue strictly dominates, $|\lambda_1| > |\lambda_2| \geq \cdots \geq |\lambda_m|$. We factor out λ_1^k:

$$A^k\vec{z} = \lambda_1^k \left(y_1\vec{x}_1 + \sum_{i=2}^{m} \left(\frac{\lambda_i}{\lambda_1}\right)^k y_i \vec{x}_i \right). \tag{10.12}$$

Every ratio $|\lambda_i/\lambda_1|$ with $i \geq 2$ is less than 1. So $(\lambda_i/\lambda_1)^k \to 0$, and the *direction* of $A^k\vec{z}$ approaches $\vec{x}_1$. Informally, $\Lambda^k \approx \lambda_1^k \operatorname{diag}(1, 0, \ldots, 0)$ and $A^k\vec{z} \approx \lambda_1^k y_1 \vec{x}_1$. These approximations hold in the relative sense, $\Lambda^k/\lambda_1^k \to \operatorname{diag}(1, 0, \ldots, 0)$. The vector $A^k\vec{z}$ itself may blow up or go to zero.

The condition $y_1 \neq 0$. All of this needs $y_1 \neq 0$. If the starting vector has no component along $\vec{x}_1$, multiplying by A will never create one. What does $y_1 \neq 0$ mean concretely?

The coordinate y_1 is the inner product of the first row of X^{-1} with $\vec{z}$. From $X^{-1}A = \Lambda X^{-1}$, the i-th row $\vec{w}_i^*$ of X^{-1} satisfies $\vec{w}_i^* A = \lambda_i \vec{w}_i^*$. So it is a *left eigenvector*. Hence

$$y_1 \neq 0 \iff \langle \vec{w}_1, \vec{z} \rangle \neq 0, \tag{10.13}$$

i.e., the starting vector must not be orthogonal to the left eigenvector of λ_1. If A is Hermitian, then $X^{-1} = Q^*$, and the left and right eigenvectors are the same. Then the condition simply says that $\vec{z}$ is not orthogonal to $\vec{x}_1$.

What if $y_1 = 0$ anyway? In exact arithmetic the direction $\vec{x}_1$ never appears. The iteration then converges to $\vec{x}_j$ for the first j with $y_j \neq 0$, and it does not notice the problem. In practice this is not a concern, for two reasons.

- A random starting vector lies on the hyperplane $\vec{w}_1^{\perp}$ with probability 0.
- Even if it does, rounding errors add an $O(\varepsilon_{\text{mach}})$ component along $\vec{x}_1$. This component then grows step by step.

The algorithm. We compute $\vec{z}_k = A^k\vec{z} / \left\| A^k\vec{z} \right\|$ as an approximation to $\vec{x}_1$. We use the Rayleigh quotient $\langle \vec{z}_k, A\vec{z}_k \rangle$ as an approximation to λ_1. We choose $\vec{z} \sim N(0, I)$ to make sure that $y_1 \neq 0$. We normalize at every step to avoid overflow.

Algorithm 15 Power iteration.

1: $\vec{q} \leftarrow$ vector with i.i.d. standard normal entries; $\vec{q} \leftarrow \vec{q}/\|\vec{q}\|$
2: **for** $k = 1, 2, \ldots$ **do**
3: $\quad \vec{z} \leftarrow A\vec{q}$
4: $\quad \lambda \leftarrow \vec{q}^* \vec{z}$ ▷ Rayleigh quotient, reuses $\vec{z}$
5: $\quad$ **if** $\|\vec{z} - \lambda\vec{q}\| \le \mathrm{tol} \cdot |\lambda|$ **then**
6: $\quad\quad$ **return** $\lambda, \vec{q}$
7: $\quad \vec{q} \leftarrow \vec{z}/\|\vec{z}\|$

Some details of the algorithm.

- *The random start.* A standard normal vector has a rotation-invariant distribution. After normalization it is uniformly distributed on the unit sphere. It satisfies $y_1 \neq 0$ almost surely.
- *The order of lines 4 and 7.* Line 4 must come before line 7, because $\vec{z} = A\vec{q}$ belongs to the old $\vec{q}$.
- *The order in $\vec{q}^*\vec{z}$.* In the complex case, $\vec{z}^*\vec{q} = \overline{\vec{q}^* A\vec{q}}$. This agrees with $\vec{q}^*\vec{z}$ only when $\vec{q}^* A\vec{q}$ is real. For real A either order works.
- *Cost.* Each step costs one matrix-vector product: $O(m^2)$ for a dense matrix and $O(\mathrm{nnz})$ for a sparse one. So power iteration suits large matrices that we can only multiply by (PageRank, for example).

Why normalize. The length $\left\|A^k\vec{z}\right\|$ changes like $|\lambda_1|^k$. Double precision covers only about $10^{\pm 308}$.

- With $|\lambda_1| = 10$, the iterates overflow to `Inf` after about 300 steps.
- With $|\lambda_1| = 0.1$, they underflow to 0.

The number of steps needed for convergence depends on $|\lambda_2/\lambda_1|$, not on $|\lambda_1|$. So overflow or underflow can easily happen before convergence. Normalizing only at the end is too late.

Normalizing does not change the result. Since $A(c\vec{v}) = c \cdot A\vec{v}$, scaling at one step does not affect the directions at later steps. In exact arithmetic, $\vec{q}^{(k)} = A^k\vec{q}^{(0)}/\left\|A^k\vec{q}^{(0)}\right\|$. We also get two small benefits: the Rayleigh quotient needs no division by $\|\vec{q}\|^2$, and the stopping test is not affected by scaling.

Why the residual. The stopping test uses the residual $\|A\vec{q} - \lambda\vec{q}\|$, not the change $\left\|\vec{q}^{(k)} - \vec{q}^{(k-1)}\right\|$. The reason: if $\lambda_1 < 0$, then $\vec{q}$ changes sign at every step. (If λ_1 is complex, $\vec{q}$ is multiplied by a phase at every step.) So the *direction* converges, but the vector does not. For Hermitian matrices the residual also bounds the eigenvalue error directly.

Proposition 59 (Residual Bound)

Let A be Hermitian, $\vec{q}$ a unit vector, and $\lambda \in \mathbb{R}$. Then A has an eigenvalue λ_j with

$$|\lambda_j - \lambda| \le \|A\vec{q} - \lambda\vec{q}\|\,. \tag{10.14}$$

Proof. Expand $\vec{q} = \sum_j c_j\vec{x}_j$ in an orthonormal eigenbasis, so that $\sum_j |c_j|^2 = 1$. Then $A\vec{q} - \lambda\vec{q} =$

$\sum_j (\lambda_j - \lambda) c_j \vec{x}_j$, and by orthonormality

$$\|A\vec{q} - \lambda\vec{q}\|^2 = \sum_j |\lambda_j - \lambda|^2 |c_j|^2 \geq \min_j |\lambda_j - \lambda|^2 \sum_j |c_j|^2 = \min_j |\lambda_j - \lambda|^2 , \tag{10.15}$$

as desired. □

Rate of convergence. We now study how fast power iteration converges, for Hermitian A. We track two errors: the error in the vector and the error in the eigenvalue. The key point is that the second error is much smaller than the first. The reason is that the Rayleigh quotient of a Hermitian matrix is *quadratically* accurate.

Lemma 60 (Quadratic Accuracy of the Rayleigh Quotient)

Let A be Hermitian with orthonormal eigenvectors $\vec{x}_1, \ldots, \vec{x}_m$, and let $\vec{q}$ be a unit vector. Then for every J,

$$r(\vec{q}) - \lambda_J = \sum_{j \neq J} (\lambda_j - \lambda_J) |c_j|^2 , \qquad c_j = \langle \vec{x}_j, \vec{q} \rangle , \tag{10.16}$$

and hence $|r(\vec{q}) - \lambda_J| \leq (\lambda_{\max} - \lambda_{\min}) \|\vec{q} - \vec{x}_J\|^2$.

Proof. Since $\vec{q} = \sum_j c_j \vec{x}_j$ and $\sum_j |c_j|^2 = 1$,

$$r(\vec{q}) = \vec{q}^* A \vec{q} = \sum_j \lambda_j |c_j|^2 = \lambda_J \underbrace{\sum_j |c_j|^2}_{=1} + \sum_{j \neq J} (\lambda_j - \lambda_J) |c_j|^2 . \tag{10.17}$$

For the bound, we use two facts. First, $|\lambda_j - \lambda_J| \leq \lambda_{\max} - \lambda_{\min}$. Second,

$$\sum_{j \neq J} |c_j|^2 = \|\vec{q} - c_J \vec{x}_J\|^2 \leq \|\vec{q} - \vec{x}_J\|^2 , \tag{10.18}$$

since $c_J \vec{x}_J$ is the orthogonal projection of $\vec{q}$ onto $\mathrm{span}(\vec{x}_J)$. This is the point of that line closest to $\vec{q}$. □

Why "quadratic"? For $j \neq J$, the coefficients c_j measure the vector error. The right side is quadratic in them. Another way to see it: the Rayleigh quotient of a Hermitian matrix is *stationary* at eigenvectors. In the real case,

$$\nabla r(\vec{x}) = \frac{2}{\vec{x}^\top \vec{x}} (A\vec{x} - r(\vec{x})\vec{x}),$$

which is zero exactly at eigenvectors. So an ε error in the vector moves the Rayleigh quotient only by $O(\varepsilon^2)$. We come back to this picture in Section 10.7.1.

Theorem 61 (Convergence of Power Iteration)

Let A be Hermitian, with orthonormal eigenvectors $\vec{x}_1, \ldots, \vec{x}_m$ and eigenvalues $|\lambda_1| > |\lambda_2| \geq \cdots \geq |\lambda_m|$. Let the unit starting vector satisfy $c_1 = \left\langle \vec{x}_1, \vec{q}^{(0)} \right\rangle \neq 0$. Then there are unimodular scalars s_k ($s_k = \pm 1$ in the real case) such that the iterates of Algorithm 15 satisfy

$$\left\| \vec{q}^{(k)} - s_k \vec{x}_1 \right\| = O\left(\left| \frac{\lambda_2}{\lambda_1} \right|^k \right) , \qquad \left| r(\vec{q}^{(k)}) - \lambda_1 \right| = O\left(\left| \frac{\lambda_2}{\lambda_1} \right|^{2k} \right) . \tag{10.19}$$

The plan of the proof has three steps.

1. Write $A^k\vec{q}^{(0)}$ as $\lambda_1^k c_1$ times "$\vec{x}_1$ plus a small error".
2. Show that normalizing keeps the error small.
3. Square the error with Lemma 60.

Proof. Step 1: the unnormalized iterate. By (10.12) with $y_j = c_j$, we have $A^k\vec{q}^{(0)} = \lambda_1^k c_1(\vec{x}_1 + \vec{e}_k)$, where

$$\vec{e}_k = \sum_{j=2}^{m} \left(\frac{\lambda_j}{\lambda_1}\right)^k \frac{c_j}{c_1}\vec{x}_j, \qquad \|\vec{e}_k\|^2 = \sum_{j=2}^{m} \left|\frac{\lambda_j}{\lambda_1}\right|^{2k} \frac{|c_j|^2}{|c_1|^2} \le \left|\frac{\lambda_2}{\lambda_1}\right|^{2k} \frac{1}{|c_1|^2}, \tag{10.20}$$

using $\sum_j |c_j|^2 = 1$.

Step 2: normalization. Let $s_k = \lambda_1^k c_1 / \left|\lambda_1^k c_1\right|$ and $\vec{u} = \vec{x}_1 + \vec{e}_k$. Then $\vec{q}^{(k)} = s_k\vec{u}/\|\vec{u}\|$. Since $\|\vec{x}_1\| = 1$, the triangle inequality gives

$$\left\|\frac{\vec{u}}{\|\vec{u}\|} - \vec{x}_1\right\| \le \left\|\frac{\vec{u}}{\|\vec{u}\|} - \vec{u}\right\| + \|\vec{u} - \vec{x}_1\| = \underbrace{|1 - \|\vec{u}\||}_{\le \|\vec{u}-\vec{x}_1\|} + \|\vec{u} - \vec{x}_1\| \le 2\|\vec{e}_k\|, \tag{10.21}$$

so $\left\|\vec{q}^{(k)} - s_k\vec{x}_1\right\| \le 2\left|\lambda_2/\lambda_1\right|^k / |c_1|$.

Step 3: the eigenvalue. The vector $s_k\vec{x}_1$ is a unit eigenvector for λ_1, so Lemma 60 gives

$$\left|r(\vec{q}^{(k)}) - \lambda_1\right| \le (\lambda_{\max} - \lambda_{\min})\left\|\vec{q}^{(k)} - s_k\vec{x}_1\right\|^2 \le \frac{4(\lambda_{\max} - \lambda_{\min})}{|c_1|^2}\left|\frac{\lambda_2}{\lambda_1}\right|^{2k}, \tag{10.22}$$

as desired. □

The eigenvalue error is often quoted as $O(|\lambda_2/\lambda_1|^k)$, without the factor 2 in the exponent. That is the right rate for a general diagonalizable matrix. For non-Hermitian A:

- The first estimate still holds, with a constant that also depends on the condition number of X.
- The second estimate does not hold. The eigenvectors are no longer orthogonal, so the expansion of $r(\vec{q})$ has mixed terms that are linear in the c_j. The eigenvalue error is then only $O(|\lambda_2/\lambda_1|^k)$.

For Hermitian A we get the square.

How many steps does this take? With linear convergence, reaching an error ε takes about $k \approx \log\varepsilon / \log|\lambda_2/\lambda_1|$ steps. For a vector accuracy of 10^{-8}, this is about 27 steps when the ratio is 0.5, 175 when it is 0.9, and 1800 when it is 0.99. So the speed depends on λ_2/λ_1. The natural question is: what happens when this ratio is close to 1 in modulus? There are two very different cases: $\lambda_2 \approx \lambda_1$ and $\lambda_2 \approx -\lambda_1$.

When $\lambda_2 \approx \lambda_1$: the eigenvalue still converges. In short, in this case *the eigenvalue converges well enough, but the eigenvector converges slowly.*

Suppose $|\lambda_1| \sim |\lambda_2| \gg |\lambda_3|$. In the space spanned by $\vec{x}_1, \vec{x}_2, \vec{x}_3$, the iterates move in two phases.

1. They first fall quickly into the plane $\mathrm{span}(\vec{x}_1, \vec{x}_2)$, at the rate $|\lambda_3/\lambda_1|^k$.
2. Then they turn slowly toward $\vec{x}_1$ within the plane, at the rate $|\lambda_2/\lambda_1|^k$.

After the first phase, $\vec{q} \approx c_1\vec{x}_1 + c_2\vec{x}_2$. The Rayleigh quotient $\lambda_1 |c_1|^2 + \lambda_2 |c_2|^2$ is then a weighted average of λ_1 and λ_2. Its error is at most $|\lambda_1 - \lambda_2|$, which is already small.

We can also see this from the formula $\sum_{j\geq 2}(\lambda_j - \lambda_1)|c_j|^2$ of Lemma 60.

- For directions far from λ_1, $|c_j|^2$ decays quickly.
- For directions close to λ_1, $|c_j|^2$ decays slowly, but the factor $\lambda_j - \lambda_1$ is small.

So one of the two factors is always small.


In the extreme case $\lambda_2 = \lambda_1$, $\vec{q}$ converges to some vector in $\mathrm{span}(\vec{x}_1, \vec{x}_2)$. Which vector depends on the starting vector. Every nonzero vector in this eigenspace is an eigenvector for λ_1. So the algorithm still succeeds. The component of $\vec{q}$ outside this eigenspace decays like $|\lambda_3/\lambda_1|^k$, and the eigenvalue error like $|\lambda_3/\lambda_1|^{2k}$.

This matches the conditioning of the problem. Eigenvalues of Hermitian matrices are always well-conditioned (Weyl: $|\delta\lambda| \leq \|\delta A\|_2$). The sensitivity of an eigenvector, on the other hand, is about $\|\delta A\|/\mathrm{gap}$. When the gap is small, $\vec{x}_1$ itself is ill-conditioned, so we should not expect to compute it quickly.

When $\lambda_2 \approx -\lambda_1$: the real failure. Now only the plane matters. Let $\vec{q} = c_1\vec{x}_1 + c_2\vec{x}_2$. One multiplication by A gives $\lambda_1(c_1\vec{x}_1 - c_2\vec{x}_2)$. Geometrically this is a *reflection across the $\vec{x}_1$ axis*. The iterates jump back and forth between two directions that are symmetric about $\vec{x}_1$. If $\lambda_2 = -\lambda_1$ exactly, $|c_2/c_1|$ never changes, and there is no convergence at all. The Rayleigh quotient

$$\vec{q}^* A\vec{q} \approx \lambda_1 \left(|c_1|^2 - |c_2|^2\right) \tag{10.23}$$

is fixed by the starting vector. It may lie anywhere in $[-\lambda_1, \lambda_1]$.



Left: with $\lambda_2/\lambda_1 = 0.8$ the iterates turn slowly but steadily toward $\vec{x}_1$. Right: with $\lambda_2 = -\lambda_1$ each step reflects $\vec{q}$ across the $\vec{x}_1$ axis, and the iteration never converges.

So the rate depends only on $|\lambda_2/\lambda_1|$. But the truly bad case is eigenvalues with *similar modulus but far-apart values*. Close values do no harm. The cure is to change the spectrum. This is the topic of the next section.

10.5 Matrix Functions: Changing the Spectrum

Power iteration finds the eigenvalue of largest modulus. It is fast when that eigenvalue stands out. So we ask: *how can we change the eigenvalues of a matrix, so that the one we want stands out?* The answer is to apply a function to the matrix. For every polynomial p, $p(A)\vec{x}_i = p(\lambda_i)\vec{x}_i$. This gives the following table.

Transformation	Eigenvalues	Eigenvectors
$A \to A^k$	$\lambda \to \lambda^k$	unchanged
$A \to \alpha A$	$\lambda \to \alpha\lambda$	unchanged
$A \to A + \beta I$	$\lambda \to \lambda + \beta$	unchanged
$A \to p(A)$	$\lambda \to p(\lambda)$	unchanged

The important column is the last one. Power iteration on $p(A)$ produces eigenvectors of A. From these, the Rayleigh quotient of A gives the eigenvalues of A.

From this point of view, power iteration itself filters the spectrum with $p(z) = z^k$. It amplifies eigenvalues of large modulus and damps small ones. This is called *polynomial filtering*. A different p filters differently. Scaling by α does not change the ratios; it acts like normalization. The useful operation is the shift.

Shifting. As an example, suppose a cluster of eigenvalues sits on the negative axis, and an isolated eigenvalue sits on the positive axis, with similar moduli. Then the ratio for plain power iteration is close to 1. Iterating with $A + I$ shifts the whole spectrum to the right. The cluster moves toward the origin, and the isolated eigenvalue moves farther out. So the ratio drops.

How should we choose the shift in general? Let A have real eigenvalues $\lambda_1 > \lambda_2 \geq \cdots \geq \lambda_m$, ordered by value. Let λ_1 be the one we want. Power iteration on $A + \beta I$ converges at the rate

$$\max_{j \geq 2} \frac{|\lambda_j + \beta|}{|\lambda_1 + \beta|}. \tag{10.24}$$

The best shift moves the midpoint of $[\lambda_m, \lambda_2]$ to the origin. This is $\beta = -\frac{\lambda_2 + \lambda_m}{2}$, and it gives the ratio

$$\frac{(\lambda_2 - \lambda_m)/2}{\lambda_1 - \frac{\lambda_2 + \lambda_m}{2}}. \tag{10.25}$$

Why is this the best choice?

- For larger β, the ratio is $(\lambda_2 + \beta)/(\lambda_1 + \beta)$, which increases with β.
- For smaller β, it is $-(\lambda_m + \beta)/(\lambda_1 + \beta)$, which decreases with β.

The bad case $\lambda_2 \approx -\lambda_1$ of the previous section is solved in the same way. With $\lambda_1 > 0$ and any $\beta > 0$, the ratio becomes $|\beta - \lambda_1| / (\beta + \lambda_1) < 1$.

A shift can only move the spectrum left or right. To choose it, we need a rough idea of where the spectrum lies. Two generalizations go further.

- We can design higher-degree polynomials p that are large at λ_1 and uniformly small on the interval that contains the rest of the spectrum (Chebyshev acceleration).
- If we replace polynomials by rational functions, we get inverse iteration. This is the topic of the next section.


10.6 Inverse Iteration

Power iteration can only find eigenvalues at the edge of the spectrum. What if we want an eigenvalue in the *middle*: the one closest to a given $\mu \in \mathbb{R}$? The idea is to use $(A - \mu I)^{-1}$, i.e., the function

$$f(z) = \frac{1}{z - \mu}, \qquad (A - \mu I)^{-1}\vec{x}_j = \frac{1}{\lambda_j - \mu}\vec{x}_j. \tag{10.26}$$

Let us see what this does to the spectrum.

- The eigenvalue λ_J closest to μ is mapped to $1/(\lambda_J - \mu)$, which has the largest modulus.
- The second closest is mapped a bit closer to the origin.
- All the far ones are squeezed toward the origin. Those with $\lambda_j < \mu$ go to the negative axis, and those with $\lambda_j > \mu$ go to the positive axis.



The map $\lambda \mapsto 1/(\lambda - \mu)$ sends the eigenvalue closest to μ far away and squeezes all the others toward 0.

So power iteration on $(A - \mu I)^{-1}$ finds $\vec{x}_J$. Let λ_K be the eigenvalue second closest to μ.

Theorem 62 (Convergence of Inverse Iteration)

Let A be Hermitian. Let λ_J be the eigenvalue closest to μ and λ_K the second closest, with $|\lambda_J - \mu| < |\lambda_K - \mu|$, and let $\left\langle \vec{x}_J, \vec{q}^{(0)} \right\rangle \neq 0$. Then there are unimodular s_k such that the iterates of Algorithm 16 satisfy

$$\left\| \vec{q}^{(k)} - s_k \vec{x}_J \right\| = O\left(\left| \frac{\lambda_J - \mu}{\lambda_K - \mu} \right|^k \right), \qquad \left| \lambda^{(k)} - \lambda_J \right| = O\left(\left| \frac{\lambda_J - \mu}{\lambda_K - \mu} \right|^{2k} \right). \tag{10.27}$$

Proof. The matrix $(A - \mu I)^{-1}$ is Hermitian. It has the same orthonormal eigenvectors as A, and eigenvalues $1/(\lambda_j - \mu)$. The largest in modulus is $1/(\lambda_J - \mu)$ and the second largest is $1/(\lambda_K - \mu)$. Up to normalization, the iterates of Algorithm 16 are those of Algorithm 15 applied to $(A - \mu I)^{-1}$. So the first estimate is Theorem 61 with the ratio

$$\left| \frac{(\lambda_K - \mu)^{-1}}{(\lambda_J - \mu)^{-1}} \right| = \left| \frac{\lambda_J - \mu}{\lambda_K - \mu} \right|. \tag{10.28}$$

The eigenvalue estimate $\lambda^{(k)} = r(\vec{q}^{(k)})$ is the Rayleigh quotient of A itself. Lemma 60 applied to A squares the vector error. □

The closer μ is to λ_J, the smaller the ratio. In other words, *the rate of convergence is now controlled by our choice of* μ, not only by the spectrum of the matrix.

Algorithm 16 Inverse iteration with a fixed shift μ.

1: factor $A - \mu I = PLU$ ▷ once, $O(m^3)$
2: $\vec{q} \leftarrow$ random unit vector
3: **for** $k = 1, 2, \ldots$ **do**
4: solve $(A - \mu I)\vec{z} = \vec{q}$ with the factorization ▷ $O(m^2)$
5: $\vec{q} \leftarrow \vec{z} / \|\vec{z}\|$
6: $\vec{w} \leftarrow A\vec{q}, \quad \lambda \leftarrow \vec{q}^* \vec{w}$ ▷ Rayleigh quotient of A
7: **if** $\|\vec{w} - \lambda\vec{q}\| \leq \text{tol} \cdot |\lambda|$ **then**
8: **return** $\lambda, \vec{q}$

In MATLAB notation, line 4 is `z = (A - mu*I) \ q`. Some remarks on implementing it.

- *Factor once.* Since μ is fixed, we compute the factorization once, outside the loop. If we solved from scratch in every step, each step would cost $O(m^3)$ instead of $O(m^2)$.
- *Which factorization.* For Hermitian A we can use a symmetric indefinite factorization LDL^* (Bunch–Kaufman, Section 9.5). We cannot use Cholesky: with μ in the middle of the spectrum, $A - \mu I$ is indefinite.
- *The eigenvalue estimate.* We use the Rayleigh quotient of A itself, which gives λ_J directly. We do not need to recover it from $1/(\lambda_J - \mu)$.
- *Normalization* is especially important here, since $\|\vec{z}\| \approx 1/|\lambda_J - \mu|$ can be huge.

Ill-conditioning is harmless. The closer μ is to λ_J, the closer $A - \mu I$ is to singular, and the larger the error in the solve. Is this a problem? No. This error lies almost entirely along $\vec{x}_J$, which is exactly the direction we want, and normalization removes it. An exact zero pivot (when μ is exactly an eigenvalue) almost never happens in floating point. If it does, it is enough to perturb μ by about $\varepsilon_{\text{mach}}$.

10.7 Rayleigh Quotient Iteration

In some problems we know a good μ in advance. But what if we don't? And even with a good fixed μ, inverse iteration still converges only linearly. The idea of Rayleigh quotient iteration (RQI) is to *choose μ to be our current guess of the eigenvalue*: the Rayleigh quotient of the current vector.

Algorithm 17 Rayleigh quotient iteration.

1: $\vec{q} \leftarrow$ unit starting vector; $\mu \leftarrow \vec{q}^* A \vec{q}$ (or a given guess)
2: **for** $k = 1, 2, \ldots$ **do**
3: solve $(A - \mu I)\vec{z} = \vec{q}$
4: $\vec{q} \leftarrow \vec{z} / \|\vec{z}\|$
5: $\mu \leftarrow \vec{q}^* A \vec{q}$

This creates a positive feedback loop.

- A more accurate μ gives a smaller inverse iteration ratio, so $\vec{q}$ improves more.
- A better $\vec{q}$ makes the Rayleigh quotient μ more accurate.

In short: the Rayleigh quotient turns good vectors into better eigenvalues, and inverse iteration turns good eigenvalues into good vectors.

10.7.1 The Rayleigh Quotient Near an Eigenvector

To analyze RQI, we first need to know exactly how close $r(\vec{q})$ is to λ_J when $\vec{q}$ is close to $\vec{x}_J$. We already know the answer from Lemma 60. Here we derive it again in a different way, with calculus. This gives a nice picture.

Take $A \in \mathbb{R}^{m\times m}$ symmetric. (The complex case is covered by Lemma 60.) We view $r(\vec{x}) = \vec{x}^\top A\vec{x}/\vec{x}^\top\vec{x}$ as a function on $\mathbb{R}^m \setminus \{\vec{0}\}$. In inverse iteration and RQI the eigenvalue estimate is $\mu_k = r(\vec{q}^{(k)})$. For an eigenpair, $r(\vec{x}_J) = \lambda_J$, so $\vec{q} \approx \vec{x}_J$ implies $r(\vec{q}) \approx \lambda_J$. How close? We compute the Taylor expansion around $\vec{x}_J$, with $\vec{e} = \vec{y} - \vec{x}_J$:

$$r(\vec{y}) = r(\vec{x}_J) + \nabla r(\vec{x}_J)^\top \vec{e} + \tfrac{1}{2}\vec{e}^\top H(\vec{x}_J)\vec{e} + O(\|\vec{e}\|^3), \tag{10.29}$$

where H is the Hessian of r. By the quotient rule, the gradient is

$$\nabla r(\vec{x}) = \frac{2A\vec{x}\,(\vec{x}^\top\vec{x}) - 2\vec{x}\,(\vec{x}^\top A\vec{x})}{(\vec{x}^\top\vec{x})^2} = \frac{2}{\vec{x}^\top\vec{x}}(A\vec{x} - r(\vec{x})\vec{x}). \tag{10.30}$$


So $\nabla r(\vec{x}) = \vec{0}$ if and only if $\vec{x}$ is an eigenvector: *the eigenvectors are exactly the critical points of* r. Therefore the linear term in the Taylor expansion drops out, and


$$r(\vec{y}) = \lambda_J + O(\|\vec{y} - \vec{x}_J\|^2). \tag{10.31}$$

We can go one step further and compute the quadratic term. We differentiate ∇r once more and use $\nabla r(\vec{x}_J) = \vec{0}$. For $\|\vec{x}_J\| = 1$, this gives

$$H(\vec{x}_J) = 2(A - \lambda_J I), \qquad \tfrac{1}{2}\vec{e}^\top H(\vec{x}_J)\vec{e} = \vec{e}^\top(A - \lambda_J I)\vec{e} = \sum_{j\neq J}(\lambda_j - \lambda_J)c_j^2, \quad c_j = \vec{x}_j^\top\vec{e}, \tag{10.32}$$

which is the expression of Lemma 60 again. The Hessian is indefinite unless λ_J is the largest or the smallest eigenvalue. So $\vec{x}_{\max}$ and $\vec{x}_{\min}$ are the maximum and minimum of r. Every other eigenvector is a saddle point.





The Rayleigh quotient on the unit circle $\vec{q} = \cos\theta\,\vec{x}_1 + \sin\theta\,\vec{x}_2$, for $\lambda_1 = 3$, $\lambda_2 = 1$, is $r(\theta) = \lambda_2 + (\lambda_1 - \lambda_2)\cos^2\theta$. The derivative is zero at the eigenvectors. So an angle error ε moves r by only $(\lambda_1 - \lambda_2)\sin^2\varepsilon \approx (\lambda_1 - \lambda_2)\varepsilon^2$.

In RQI the next vector is $\vec{q}^{(k+1)} \propto (A - r(\vec{q}^{(k)})I)^{-1}\vec{q}^{(k)}$. The good news: *if* $\vec{q}$ *is close to* $\vec{x}_J$*, then* μ *is even closer to* λ_J. A vector error ε gives an eigenvalue error $O(\varepsilon^2)$.

Analogy with Newton's method. RQI is similar to Newton's method, and we can make this precise. An eigenpair is a root of

$$F(\vec{x},\lambda)=\begin{bmatrix}(A-\lambda I)\vec{x}\\ \frac{1}{2}(1-\vec{x}^\top\vec{x})\end{bmatrix},\qquad F'(\vec{x},\lambda)=\begin{bmatrix}A-\lambda I & -\vec{x}\\ -\vec{x}^\top & 0\end{bmatrix}. \tag{10.33}$$

The first block row of the Newton step $F'(\vec{x},\lambda)(\Delta\vec{x},\Delta\lambda)=-F(\vec{x},\lambda)$ reads $(A-\lambda I)\Delta\vec{x}-\Delta\lambda\,\vec{x}=-(A-\lambda I)\vec{x}$, i.e.,

$$\vec{x}+\Delta\vec{x}=\Delta\lambda\,(A-\lambda I)^{-1}\vec{x}. \tag{10.34}$$

So the Newton update of the vector is parallel to one step of inverse iteration with shift λ. RQI takes this direction and normalizes it. Then it replaces Newton's $\lambda+\Delta\lambda$ by the Rayleigh quotient of the new vector. Newton's method converges quadratically, because the error of the linearization is quadratic. For symmetric A, the stationarity of r adds one more order.

10.7.2 Cubic Convergence

Theorem 63 (Convergence of Rayleigh Quotient Iteration [1, Thm. 27.3])

Let A be Hermitian. Suppose RQI converges to an eigenpair $(\lambda_J,\vec{x}_J)$. Then for $\vec{q}^{(k)}$ close enough to $\pm\vec{x}_J$ (in the complex case, to a unimodular multiple of $\vec{x}_J$),

$$\left\|\vec{q}^{(k+1)}-(\pm\vec{x}_J)\right\|=O\left(\left\|\vec{q}^{(k)}-(\pm\vec{x}_J)\right\|^3\right),\qquad \left|\mu^{(k+1)}-\lambda_J\right|=O\left(\left|\mu^{(k)}-\lambda_J\right|^3\right). \tag{10.35}$$

For a general (non-Hermitian) matrix, the convergence is quadratic instead.

Why cubic: the heuristic. Before the proof, let us see where the exponent 3 comes from. Suppose $\left\|\vec{q}^{(k)}-\vec{x}_J\right\|=\varepsilon$.

1. One step of inverse iteration multiplies the vector error by $|\lambda_J-\mu|\,/\,|\lambda_K-\mu|$. The denominator is roughly the spectral gap and stays away from 0. So the new error is about $\varepsilon\,|\lambda_J-\mu|$.
2. For a general matrix the Rayleigh quotient is only first-order accurate, $|\mu-\lambda_J|=O(\varepsilon)$. So the new error is $O(\varepsilon^2)$: quadratic convergence.
3. For Hermitian A, Lemma 60 gives $|\mu-\lambda_J|=O(\varepsilon^2)$. So the new error is $O(\varepsilon^3)$: cubic convergence.

In the cubic case the number of correct digits triples at every step: $10^{-2}\to10^{-6}\to10^{-18}$. The next proposition makes this precise.

Proposition 64 (One Step of RQI)

Let A be Hermitian with orthonormal eigenvectors $\vec{x}_1, \dots, \vec{x}_m$, let λ_J be simple, and let

$$g \coloneqq \min_{j \neq J} |\lambda_j - \lambda_J| > 0, \qquad s_A \coloneqq \lambda_{\max} - \lambda_{\min}. \tag{10.36}$$

Let $\vec{q}$ be a unit vector with $\|\vec{q} - \vec{x}_J\| \leq \varepsilon$, where $\varepsilon \leq \frac{1}{2}$ and $s_A \varepsilon^2 \leq g/2$. Let $\mu = r(\vec{q}) \neq \lambda_J$, and let $\vec{q}^+ = (A - \mu I)^{-1}\vec{q} / \left\|(A - \mu I)^{-1}\vec{q}\right\|$. Then there is a unimodular s with

$$\left\|\vec{q}^+ - s\vec{x}_J\right\| \leq \frac{4 s_A}{g}\varepsilon^3. \tag{10.37}$$

Proof. Step 1: split $\vec{q}$. Write $\vec{q} = \alpha\vec{x}_J + \beta\vec{v}$ with $\alpha = \langle \vec{x}_J, \vec{q}\rangle$, $\vec{v} \perp \vec{x}_J$, and $\|\vec{v}\| = 1$. The point $\alpha\vec{x}_J$ is the point of $\operatorname{span}(\vec{x}_J)$ closest to $\vec{q}$. So $|\beta| = \|\vec{q} - \alpha\vec{x}_J\| \leq \|\vec{q} - \vec{x}_J\| \leq \varepsilon$. Also $|\alpha - 1| = |\langle \vec{x}_J, \vec{q} - \vec{x}_J\rangle| \leq \varepsilon$, so $|\alpha| \geq \frac{1}{2}$.

Step 2: apply $(A - \mu I)^{-1}$. The vector $\vec{x}_J$ is an eigenvector of $(A - \mu I)^{-1}$ with eigenvalue $1/(\lambda_J - \mu)$, so

$$(A - \mu I)^{-1}\vec{q} = \frac{\alpha}{\lambda_J - \mu}\vec{x}_J + \beta(A - \mu I)^{-1}\vec{v}. \tag{10.38}$$

We bound the second term. The subspace $\operatorname{span}\{\vec{x}_j : j \neq J\}$ contains $\vec{v}$ and is invariant under $(A - \mu I)^{-1}$. So $(A - \mu I)^{-1}\vec{v} \perp \vec{x}_J$, and $\left\|(A - \mu I)^{-1}\vec{v}\right\| \leq \max_{j \neq J} 1/|\lambda_j - \mu|$. By Lemma 60, $|\mu - \lambda_J| \leq s_A\varepsilon^2 \leq g/2$. Hence for $j \neq J$

$$|\lambda_j - \mu| \geq |\lambda_j - \lambda_J| - |\lambda_J - \mu| \geq g/2. \tag{10.39}$$

Step 3: factor out the main term. With $c = \alpha/(\lambda_J - \mu)$,

$$(A - \mu I)^{-1}\vec{q} = c(\vec{x}_J + \vec{d}), \qquad \vec{d} = \frac{\beta(\lambda_J - \mu)}{\alpha}(A - \mu I)^{-1}\vec{v} \perp \vec{x}_J, \tag{10.40}$$

and

$$\left\|\vec{d}\right\| \leq \frac{|\beta|\,|\lambda_J - \mu|}{|\alpha|} \cdot \frac{2}{g} \leq \frac{4}{g}\,\varepsilon\,|\lambda_J - \mu| \leq \frac{4 s_A}{g}\varepsilon^3. \tag{10.41}$$

In O-notation, the middle bound is $O(\varepsilon\,|\lambda_J - \mu|)$. The last step uses $|\lambda_J - \mu| = O(\varepsilon^2)$.

Step 4: normalize. With $s = c/|c|$, we have $\vec{q}^+ = s\vec{u}$, where $\vec{u} = (\vec{x}_J + \vec{d})/\left\|\vec{x}_J + \vec{d}\right\|$. Let $t = \left\|\vec{d}\right\|$. Since $\vec{d} \perp \vec{x}_J$, we have $\left\|\vec{x}_J + \vec{d}\right\| = \sqrt{1 + t^2}$, and

$$\|\vec{u} - \vec{x}_J\|^2 = \left(1 - \frac{1}{\sqrt{1 + t^2}}\right)^2 + \frac{t^2}{1 + t^2} = 2 - \frac{2}{\sqrt{1 + t^2}} \leq t^2, \tag{10.42}$$

using $1/\sqrt{1 + t^2} \geq 1 - t^2/2$. Hence $\left\|\vec{q}^+ - s\vec{x}_J\right\| = \|\vec{u} - \vec{x}_J\| \leq t$, as desired. □

Step 4 is the rigorous form of the estimate $\frac{1}{a+\varepsilon} - \frac{1}{a} \approx -\frac{\varepsilon}{a^2}$. If we divide by a length that changes by a small amount, the result changes only by the same order. So normalizing does not destroy the $O(\varepsilon^3)$ bound.

From one step to the theorem. Since $s\vec{x}_J$ is again a unit eigenvector for λ_J, we can apply the proposition at every step. Let $\varepsilon_k = \left\|\vec{q}^{(k)} - s_k\vec{x}_J\right\|$. Then the proposition gives $\varepsilon_{k+1} \le C\varepsilon_k^3$ with $C = 4s_A/g$. The eigenvalue error satisfies $\left|\mu^{(k)} - \lambda_J\right| \le s_A\varepsilon_k^2$, so this bound also shrinks cubically: $s_A\varepsilon_{k+1}^2 \le s_A C^2\varepsilon_k^6$. This proves the Hermitian case of Theorem 63 in terms of these error bounds; for the statement directly in terms of $\mu^{(k)}$, see [1, Thm. 27.3]. Here "close enough" means $\varepsilon \le \frac{1}{2}$ and $s_A\varepsilon^2 \le g/2$, together with $C\varepsilon^2 < 1$ so that the errors decrease. If $\mu = \lambda_J$ exactly, the iteration has already converged.

This is a *local* statement. It applies only once $\vec{q}^{(k)}$ is already close enough to some eigenvector.

Cost and limitations.

- *Cost.* Since μ changes at every step, we cannot reuse the factorization. Each step costs $O(m^3)$ for a dense matrix. In practice we first reduce A to tridiagonal (Hermitian) or Hessenberg form. After that, each solve costs only $O(m)$ or $O(m^2)$. The shifted QR algorithm is essentially RQI in another form.
- *Near-singular systems.* As RQI converges, $A - \mu I$ gets closer and closer to singular. As for inverse iteration, this ill-conditioning is harmless. An exact zero pivot means that the iteration has converged, and we can stop.
- *Which eigenvalue.* The eigenvalue that RQI converges to depends on the starting vector. We cannot choose it in advance. For Hermitian matrices, one can show that RQI converges for almost every starting vector. A common practice: first run a few steps of inverse iteration with a fixed shift, to move the vector toward the target. Then switch to RQI to speed up.

10.8 (SUPPLEMENT) Further Topics

10.8.1 The Jordan Normal Form

A diagonalizable matrix needs m linearly independent eigenvectors. This can fail. For example, take

$$J = \begin{bmatrix} \lambda & 1 \\ 0 & \lambda \end{bmatrix}. \tag{10.43}$$

Its eigenvalue λ has algebraic multiplicity 2. But $\mathcal{N}(J - \lambda I) = \operatorname{span}(\vec{e}_1)$, so its geometric multiplicity is 1. A matrix is called *defective* if it has an eigenvalue whose geometric multiplicity is smaller than its algebraic multiplicity.

Every square matrix is similar to a Jordan normal form: $A = XJX^{-1}$ with $J = \operatorname{diag}(J_1, \ldots, J_p)$. Each Jordan block is $J_i = \lambda_i I + N$, where N is the nilpotent matrix with ones on the superdiagonal. The matrix J is unique up to the order of the blocks.

- The algebraic multiplicity of an eigenvalue is the total size of its blocks. The geometric multiplicity is the number of its blocks. So A is diagonalizable if and only if all blocks are 1×1.
- Take the columns of X that belong to a $k \times k$ block. They satisfy $(A - \lambda I)\vec{x}_j = \vec{x}_{j-1}$. Here $\vec{x}_1$ is an eigenvector, and the others are *generalized eigenvectors*.

In theory, the Jordan form explains the transient (temporary) growth of defective matrices. We have $J_i^n = \sum_{j=0}^{k-1} \binom{n}{j} \lambda^{n-j} N^j$. So even with $|\lambda| < 1$, $\|A^n\|$ may first grow like $n^{k-1}\,|\lambda|^n$. The factor t in the solutions of ODEs with repeated roots comes from the same place.

Numerically, however, we do not use the Jordan form, because it is not continuous under perturbation.

- For example, $\left[\begin{smallmatrix}\lambda & 1\\ \varepsilon & \lambda\end{smallmatrix}\right]$ has eigenvalues $\lambda \pm \sqrt{\varepsilon}$, so it is already diagonalizable.
- A $k \times k$ block amplifies perturbations like $\varepsilon^{1/k}$.
- In floating point, we cannot decide whether a matrix is defective, or how large its blocks are.
- $\kappa(X) \to \infty$ as a matrix approaches a defective one.

This is why we give up on diagonal form and use the upper triangular Schur form instead.

10.8.2 Why Not Go Through the Characteristic Polynomial

Even in low degrees, where root formulas exist, we do not compute $\det(zI - A)$ and then its roots. The reason: going from a matrix to the coefficients of its polynomial is extremely ill-conditioned. Take the Wilkinson polynomial $\prod_{i=1}^{20}(z - i)$. If we change the coefficient of z^{19} by $2^{-23} \approx 10^{-7}$, several roots move into the complex plane, by $O(1)$. Yet the eigenvalues of $\mathrm{diag}(1, \dots, 20)$ are perfectly well-conditioned under perturbations of the matrix. So all iterative methods of this lecture work directly on the matrix, using only matrix-vector products or linear solves. They use $p(A)\vec{x}_i = p(\lambda_i)\vec{x}_i$, and do not approximate roots of a polynomial.

10.8.3 A Gap-Independent Eigenvalue Bound

Let A be s.p.d., with eigenvalues $\lambda_1 \geq \lambda_2 \geq \cdots > 0$ and orthonormal eigenvectors. Let $t_j = \lambda_j/\lambda_1 \in (0, 1]$. The Rayleigh quotient of power iteration satisfies

$$\frac{\lambda_1 - r(\vec{q}^{(k)})}{\lambda_1} = \frac{\sum_{j\geq 2}(1 - t_j)t_j^{2k}\,|y_j|^2}{\sum_j t_j^{2k}\,|y_j|^2} \leq \sum_{j\geq 2}(1 - t_j)t_j^{2k}\frac{|y_j|^2}{|y_1|^2} \leq \frac{1}{2k+1}\cdot\frac{\|\vec{y}\|^2}{|y_1|^2}. \tag{10.44}$$

For the last step: on $[0, 1]$, the function $(1 - t)t^{2k}$ has its maximum at $t = \frac{2k}{2k+1}$, and this maximum is at most $\frac{1}{2k+1}$. The bound does not depend on λ_2/λ_1. However small the gap, the relative eigenvalue error decreases at least like $O(1/k)$.

10.8.4 Matrix Functions in General

A matrix function does not act entrywise. (In Julia, `exp(A)` is the matrix exponential, while `exp.(A)` acts entrywise.) Polynomials extend to power series, such as $e^A = \sum_k A^k/k!$. For diagonalizable A, we can use

$$f(A) = Xf(\Lambda)X^{-1}, \qquad f(\Lambda) = \mathrm{diag}(f(\lambda_1), \dots, f(\lambda_m)) \tag{10.45}$$

as the definition for a general f. For example, an s.p.d. matrix has the square root $\sqrt{A} = Q\,\mathrm{diag}(\sqrt{\lambda_i})Q^*$. This is how $P = (A^*A)^{1/2}$ in the polar decomposition is defined. For a Jordan block, $f(J)$ is upper triangular, with $f'(\lambda), f''(\lambda)/2!, \dots$ on the successive superdiagonals. All methods of this lecture are choices of f:

$f(z)$	Amplified eigenvalue	Method
z	largest modulus	power iteration
$z + \sigma$	breaks symmetry of moduli	shifted power iteration
$(z - \mu)^{-1}$	closest to μ	inverse iteration
$(z - \mu_k)^{-1}$, μ_k updated	adaptive	Rayleigh quotient iteration
well-designed polynomial $p_k(z)$ of degree k	small on the rest of the spectrum	Chebyshev acceleration, Krylov methods

The last row is the natural generalization of power iteration. The vector $A^k\vec{q}$ is just one of many polynomials of degree k in A, applied to $\vec{q}$. Krylov subspace methods (Lanczos, Arnoldi) search for the best one in $\operatorname{span}(\vec{q}, A\vec{q}, \ldots, A^k\vec{q})$.

Numerically, when X is ill-conditioned, the error of $Xf(\Lambda)X^{-1}$ is amplified by $\kappa(X)$.

- For Hermitian A, $X = Q$ is unitary, and the formula is safe.
- For general matrices we use the Schur decomposition $f(A) = Qf(T)Q^*$ and a recurrence on the triangular T (Schur–Parlett).
- We compute e^A by scaling and squaring, $e^A = (e^{A/2^s})^{2^s}$, with a Padé approximant on the small matrix.

See Higham [8].

10.8.5 Why Iterative Approximation Is Reliable

Eigenvalues cannot be computed exactly in finitely many steps. Still, numerically, iterative approximation is as good as direct methods. There are three reasons.

- *Why the vector converges.* Every vector is a combination of eigenvectors. In $f(A)^k\vec{q} = \sum_j f(\lambda_j)^k y_j \vec{x}_j$, each direction is multiplied by its own $f(\lambda_j)^k$, independently of the others. Because of exponential growth, the direction with the largest $|f(\lambda)|$ dominates, and the convergence is geometric. The three algorithms differ only in the choice of f.
- *Why the eigenvalue can be read off.* The Rayleigh quotient equals the eigenvalue at an eigenvector. For Hermitian matrices it is also stationary there. So a roughly correct vector already gives a quadratically accurate eigenvalue.
- *Why an approximation is good enough.* For any unit vector $\vec{q}$ and number μ, let $\vec{r} = A\vec{q} - \mu\vec{q}$ and $\delta A = -\vec{r}\vec{q}^*$. Then

$$(A + \delta A)\vec{q} = A\vec{q} - \vec{r} = \mu\vec{q}, \qquad \|\delta A\|_2 = \|\vec{r}\| \, \|\vec{q}\| = \|\vec{r}\| \,. \tag{10.46}$$

  So an approximate eigenpair with residual $\|\vec{r}\|$ is an *exact* eigenpair of a matrix that differs from A by $\|\vec{r}\|$. The input A already has errors of order $\varepsilon_{\text{mach}}$. So once $\|\vec{r}\| \approx \varepsilon_{\text{mach}} \|A\|$, the result is as good as an exact computation on a matrix that we cannot tell apart from A. This is exactly backward stability (Definition 119), the same standard as for LU and QR. Hermitian eigenvalues are well-conditioned (Weyl). So a small backward error also guarantees a small forward error. This is the basic reason why we use the residual as the stopping criterion.

10.8.6 A Preview of Subspace Iteration

Look again at the example $|\lambda_1| \sim |\lambda_2| \gg |\lambda_3|$. What converges quickly there is not a vector but the subspace $\operatorname{span}(\vec{x}_1, \vec{x}_2)$. It converges at the rate $|\lambda_3/\lambda_2|$, no matter how close λ_1 and λ_2 are. This suggests the following method.

- Iterate several vectors at once, and keep them orthonormal with a QR factorization at each step. This quickly gives an invariant subspace.
- Solve a small eigenvalue problem inside this subspace. This separates the eigenpairs.

This is simultaneous (subspace) iteration, the method that led to the QR algorithm. When $\lambda_2 \approx -\lambda_1$, two consecutive iterates $\vec{q}^{(k)}, \vec{q}^{(k+1)}$ also span this invariant subspace, and the same approach separates $\pm\lambda_1$.

Lecture 11

Schur Factorization

Supplementary reading:

- [1] Lecture 24.
- [2] Sections 4.2–4.3.
- [4] Sections 5A, 5C, 6B, 7B, and 8A–8C (linear algebra background).

11.1 Recap and Motivation

Last lecture we started on eigenvalue problems.

- *A negative result.* We cannot compute eigenvalues in finitely many steps (Theorem 58). The reason: the companion matrix turns every polynomial into an eigenvalue problem, and by Abel there is no finite formula for the roots. So every eigenvalue algorithm is iterative.
- *Power iteration.* We expand the starting vector in the eigenbasis as $\vec{z} = X\vec{y}$. Then $A^k\vec{z} = X\Lambda^k\vec{y}$. After normalization, Λ^k keeps only the dominant eigenvalue.
- *Matrix functions.* We can move the spectrum around. A shift $A + \sigma I$ can make the wanted eigenvalue dominant. Inverse iteration and Rayleigh quotient iteration use $f(z) = (z - \mu)^{-1}$.

All of these methods compute *one* eigenpair. In this lecture we ask two questions.

1. How do we get *all* the eigenpairs, efficiently?
2. What object should "all eigenpairs" be? The obvious answer is the eigenvalue decomposition. We will see that it does not always exist, and that it is numerically unstable. The right answer is the *Schur factorization*.

11.2 Deflation

We have a fast way to compute *one* eigenpair. What if we want all of them? Suppose we have already computed the largest eigenpair $(\lambda_1, \vec{x}_1)$. If we start power iteration again from a new random vector, it converges to $\vec{x}_1$ again. How do we avoid computing the same thing again?

The idea: we change the matrix so that λ_1 is removed. The new matrix has the same eigenpairs, except that λ_1 becomes 0. For Hermitian A, this change is a rank-one update. It is called Hotelling deflation.

Proposition 65 (Hotelling Deflation)

Let $A \in \mathbb{K}^{m\times m}$ be Hermitian with eigenpairs $(\lambda_i, \vec{x}_i)$, where the $\vec{x}_i$ are pairwise orthogonal. Let

$$B \coloneqq A - \lambda_1 \frac{\vec{x}_1 \vec{x}_1^*}{\vec{x}_1^* \vec{x}_1}. \tag{11.1}$$

Then $B\vec{x}_1 = \vec{0}$ and $B\vec{x}_i = \lambda_i \vec{x}_i$ for $i = 2, \dots, m$. In particular, the eigenvalues of B are $0, \lambda_2, \dots, \lambda_m$.

Proof. We check the two cases.

- For $i = 1$,
$$B\vec{x}_1 = A\vec{x}_1 - \lambda_1 \vec{x}_1 \frac{\vec{x}_1^* \vec{x}_1}{\vec{x}_1^* \vec{x}_1} = \lambda_1 \vec{x}_1 - \lambda_1 \vec{x}_1 = \vec{0}. \tag{11.2}$$
- For $i \geq 2$, orthogonality gives $\vec{x}_1^* \vec{x}_i = 0$. So the correction term is zero, and $B\vec{x}_i = A\vec{x}_i = \lambda_i \vec{x}_i$. □

The proof uses only one property of Hermitian matrices: the eigenvectors are orthogonal. For distinct eigenvalues this holds automatically (Proposition 55). For a repeated eigenvalue, we can choose orthogonal eigenvectors inside its eigenspace by Gram–Schmidt. Also note that $\vec{x}_1 \vec{x}_1^* / (\vec{x}_1^* \vec{x}_1)$ is the orthogonal projector onto $\operatorname{span}(\vec{x}_1)$. This is why $\vec{x}_1$ does not need to be normalized.

We then apply power iteration to B. It converges to the eigenpair with the next largest modulus. By repeating this, we can compute the eigenpairs one by one.

Implementation. We never form B, because if A is sparse, B is dense. Instead we compute

$$B\vec{z} = A\vec{z} - \lambda_1 \vec{x}_1 \frac{\vec{x}_1^* \vec{z}}{\vec{x}_1^* \vec{x}_1}. \tag{11.3}$$

This costs one product with A plus $O(m)$ work.

There is another way to see what B does. Let $P \coloneqq I - \vec{x}_1 \vec{x}_1^* / (\vec{x}_1^* \vec{x}_1)$. Then one can check that $B = AP = PA$. So power iteration on B is power iteration on A, with the $\vec{x}_1$ component removed by P. Could we instead project the starting vector only once, and then iterate with A?

- In exact arithmetic, yes.
- In floating point, no. Rounding brings back a component of size $O(\varepsilon_{\text{mach}})$ along $\vec{x}_1$. This component grows like $|\lambda_1/\lambda_2|^k$ and finally dominates.

So we must apply the projection at every step. This is exactly what multiplication by $B = AP$ does.

Limitations.

- The computed $(\lambda_1, \vec{x}_1)$ is only approximate. Its error goes into B, and so into every eigenpair we compute after it. If we deflate many times in a row, these errors add up.

- For non-Hermitian A, the eigenvectors are not orthogonal, so the proof above does not work. See Section 11.9 for a version that works. See Section 11.7 for the stable way of deflating, which uses only unitary matrices.

11.3 Subspace Iteration: Motivation

So far we can compute single eigenvalues. With deflation we can, in principle, compute all eigenpairs one by one. The problem is cost. Surprisingly, the main cost is not the number of flops but memory traffic.

- Computing N eigenpairs needs at least N matrix-vector products. In practice it needs many more, since each eigenpair needs many iterations.
- These matrix-vector products happen one after another (*serially*). We cannot start the $(k+1)$-st eigenpair until the k-th has converged and been deflated.
- Each matrix-vector product must load the whole matrix from RAM. It does $2m^2$ flops on m^2 entries of A. So its computational intensity is $q \approx 2$ (Section 2.9.2). It is a Level 2 operation, and memory bandwidth limits its speed.

So we load the matrix from RAM at least N times, one after another.

The alternative is to run power iteration on *many vectors at the same time*. With $Z \in \mathbb{K}^{m \times n}$, the product AZ does $2m^2 n$ flops but reads A only once. So its intensity grows proportionally to n. Matrix-vector products become matrix-matrix products (Level 3). With the tiling of Lecture 2, these run near peak speed. This is *subspace iteration*.

But simply iterating $Z \mapsto AZ$ is not enough. Then every column is an independent power iteration, so every column converges to the same direction $\vec{x}_1$. We must keep the columns linearly independent. For example, we can orthonormalize them with a QR factorization after each multiplication. We give the algorithm and its convergence in a later lecture. The end of Lecture 10 gave a preview.

In this lecture we answer a more basic question: *when we iterate a whole set of vectors, what does the iteration converge to?* The answer (Section 11.7) is an orthonormal basis of nested invariant subspaces. This is a Schur factorization. We build up to it in the rest of this chapter.

11.4 Eigenvalue Problems as Matrix Factorizations

To ask for "all eigenpairs at once", we need to write them as one object. The first candidate is the eigenvalue decomposition $A = X \Lambda X^{-1}$. When does it exist? We start with the simplest sufficient condition: distinct eigenvalues.

Lemma 66 (Eigenvectors for Distinct Eigenvalues)

Eigenvectors $\vec{x}_1, \ldots, \vec{x}_k$ of A corresponding to distinct eigenvalues $\lambda_1, \ldots, \lambda_k$ are linearly independent [4, 5.11].

Proof. Suppose not. Choose a shortest nontrivial dependency $\sum_{i \in S} c_i \vec{x}_i = \vec{0}$, where all $c_i \neq 0$. Since eigenvectors are nonzero, S has at least two elements. The idea is to remove one term by applying $A - \lambda_j I$.

Fix some $j \in S$. Then

$$\sum_{i \in S \setminus \{j\}} c_i(\lambda_i - \lambda_j)\vec{x}_i = \vec{0}. \tag{11.4}$$

All coefficients $c_i(\lambda_i - \lambda_j)$ are nonzero, because the eigenvalues are distinct. So this is a shorter nontrivial dependency. This is a contradiction. □

Proposition 67 (Distinct Eigenvalues Give an Eigenvalue Decomposition)

If $A \in \mathbb{K}^{m \times m}$ has m distinct eigenvalues $\lambda_1, \ldots, \lambda_m$ with eigenvectors $\vec{x}_1, \ldots, \vec{x}_m$, then

$$A = X\Lambda X^{-1}, \qquad X = \begin{bmatrix} \vec{x}_1 & \cdots & \vec{x}_m \end{bmatrix}, \quad \Lambda = \operatorname{diag}(\lambda_1, \ldots, \lambda_m). \tag{11.5}$$

Proof. Written column by column, the m equations $A\vec{x}_i = \lambda_i \vec{x}_i$ are $AX = X\Lambda$. By Lemma 66, the columns of X are linearly independent. So X is invertible, and we can multiply by X^{-1} on the right. □

Distinct eigenvalues are sufficient but not necessary. For example, I has the single eigenvalue 1, and it is diagonal. Theorem 72 gives the exact condition.

For self-adjoint matrices we can say more: we can choose the eigenvectors orthonormal. First we recall the term.

Definition 68 (Adjoint and Self-Adjoint)

Let $\langle \vec{x}, \vec{y} \rangle = \vec{x}^* \vec{y}$ be the standard inner product on $\mathbb{K}^m$. The *adjoint* of $A \in \mathbb{K}^{m \times m}$ is the unique matrix A^* with

$$\langle A\vec{x}, \vec{y} \rangle = \langle \vec{x}, A^* \vec{y} \rangle \quad \text{for all } \vec{x}, \vec{y} \in \mathbb{K}^m. \tag{11.6}$$

A is *self-adjoint* if $A^* = A$.

Since $\langle A\vec{x}, \vec{y} \rangle = (A\vec{x})^* \vec{y} = \vec{x}^* (\overline{A}^\top \vec{y})$, the adjoint for the standard inner product is the conjugate transpose. So self-adjoint means *symmetric* for $\mathbb{K} = \mathbb{R}$ and *Hermitian* for $\mathbb{K} = \mathbb{C}$. For example, $\left[\begin{smallmatrix} 2 & i \\ -i & 3 \end{smallmatrix}\right]$ is Hermitian but not symmetric. Why use a new word? The name depends only on the inner product, not on a matrix representation. So it also applies to operators on infinite-dimensional spaces, for example $-\mathrm{d}^2/\mathrm{d}x^2$ with zero boundary values. It also covers the real and complex cases at once. Below we keep saying "Hermitian", as in Lecture 10.

Theorem 69 (Spectral Theorem)

If $A \in \mathbb{K}^{m \times m}$ is self-adjoint, there is a unitary X with

$$A = X\Lambda X^*, \tag{11.7}$$

and Λ is real [1, Thm. 24.7], [4, 7.29, 7.31].

We postpone the proof to Corollary 79. There it follows from the Schur factorization in two lines. A few remarks.

- The theorem says that X *can be chosen* orthogonal. If the eigenvalues are distinct, the eigenvectors are orthogonal automatically (Proposition 55), and we only need to normalize them. If an eigenvalue

is repeated, its eigenspace also contains eigenvectors that are not orthogonal. Then we must pick an orthonormal basis of the eigenspace.

- Compared with Proposition 67, X is free to invert ($X^{-1} = X^*$). Also $\kappa_2(X) = 1$, so changing to and from eigenvector coordinates does not amplify errors.

11.5 Defective Matrices and Invariant Subspaces


Are all matrices diagonalizable? Consider

$$J = \begin{bmatrix} \lambda & 1 \\ 0 & \lambda \end{bmatrix}. \tag{11.8}$$

What are its eigenpairs? Its only eigenvalue is λ. The equation $(J - \lambda I)\vec{x} = \left[\begin{smallmatrix} 0 & 1 \\ 0 & 0 \end{smallmatrix}\right]\vec{x} = \vec{0}$ forces $x_2 = 0$. So, up to scaling, $(\lambda, \vec{e}_1)$ is the only eigenpair. There is no second independent eigenvector. So there is no invertible X with $J = X\Lambda X^{-1}$. Such matrices are called *defective*.

To see what happens, take $\lambda = 0$, so that $J = N := \left[\begin{smallmatrix} 0 & 1 \\ 0 & 0 \end{smallmatrix}\right]$. Then $N\vec{e}_2 = \vec{e}_1$ and $N\vec{e}_1 = \vec{0}$. So the matrix pushes $\vec{e}_2$ to $\vec{e}_1$, and $\vec{e}_1$ to $\vec{0}$. This is a chain $\vec{e}_2 \mapsto \vec{e}_1 \mapsto \vec{0}$. Hence $N^2 = 0$, although $N \neq 0$.

This gives another short proof that N is not diagonalizable. Suppose $N = X\Lambda X^{-1}$. The diagonal of Λ holds the eigenvalues of N, which are all 0. So $\Lambda = 0$ and $N = 0$, a contradiction.



The nilpotent matrix N maps $\vec{e}_2$ to $\vec{e}_1$ and $\vec{e}_1$ to $\vec{0}$. Only $\mathrm{span}(\vec{e}_1)$ is an eigenspace.

11.5.1 Algebraic and Geometric Multiplicity

We follow the terms of [1, Lecture 24]. The characteristic polynomial $p_A(z) := \det(zI - A)$ is monic of degree m. Its roots are the eigenvalues. Similar matrices have the same characteristic polynomial, since

$$\det(zI - X^{-1}AX) = \det(X^{-1}(zI - A)X) = p_A(z). \tag{11.9}$$

There are two ways to count how often an eigenvalue occurs. A matrix is defective exactly when the two counts differ.

Definition 70 (Multiplicities, Defective Matrices)

Let λ be an eigenvalue of A.

- The *eigenspace* is $E_\lambda := \mathcal{N}(A - \lambda I)$. The *geometric multiplicity* of λ is $\dim E_\lambda$. This is the largest number of linearly independent eigenvectors for λ.
- The *algebraic multiplicity* of λ is its multiplicity as a root of p_A. The eigenvalue is *simple* if its algebraic multiplicity is 1.
- λ is *defective* if its algebraic multiplicity is larger than its geometric multiplicity. A is *defective* if it has a defective eigenvalue.

For the matrix J above, $p_J(z) = (z - \lambda)^2$, so the algebraic multiplicity is 2. But $E_\lambda = \operatorname{span}(\vec{e}_1)$, so the geometric multiplicity is 1. The difference can only go in one direction.

Theorem 71 (Algebraic Multiplicity Bounds Geometric Multiplicity)

For every eigenvalue, the algebraic multiplicity is at least the geometric multiplicity [1, Thm. 24.4].

Proof. The idea is to change to a basis that starts with the eigenspace. Let $n = \dim E_\lambda$. Let the columns of $\hat{V} \in \mathbb{K}^{m \times n}$ be an orthonormal basis of E_λ, and extend $\hat{V}$ to a unitary $V = [\hat{V}\ \tilde{V}]$. Since $A\hat{V} = \lambda\hat{V}$ and $\tilde{V}^*\hat{V} = 0$,

$$V^*AV = \begin{bmatrix} \lambda I_n & C \\ 0 & D \end{bmatrix}, \qquad C = \hat{V}^*A\tilde{V}, \quad D = \tilde{V}^*A\tilde{V}. \tag{11.10}$$

The determinant of a block triangular matrix is the product of the determinants of its diagonal blocks. So $p_A(z) = (z - \lambda)^n \det(zI - D)$. Thus λ is a root of multiplicity at least n. □

Theorem 72 (Diagonalizable Means Nondefective)

$A \in \mathbb{K}^{m \times m}$ has an eigenvalue decomposition $A = X\Lambda X^{-1}$ if and only if A is nondefective [1, Thm. 24.5].

Proof. ($\Rightarrow$) Let $A = X\Lambda X^{-1}$. Then A and Λ have the same eigenvalues with the same algebraic multiplicities. Also, X maps each eigenspace of Λ onto the matching eigenspace of A, so the geometric multiplicities are also equal. A diagonal matrix is clearly nondefective.

($\Leftarrow$) Let A be nondefective, and choose a basis of each eigenspace. The geometric multiplicities add up to the algebraic ones, and these add up to m. So we have m vectors. They are linearly independent. Indeed, take a relation $\sum_\lambda \vec{u}_\lambda = \vec{0}$ with $\vec{u}_\lambda \in E_\lambda$. The nonzero $\vec{u}_\lambda$ would be eigenvectors for distinct eigenvalues, which contradicts Lemma 66. So every $\vec{u}_\lambda = \vec{0}$. Then all coefficients are zero, because each basis is independent. These m vectors form an invertible X with $AX = X\Lambda$. □

11.5.2 Invariant Subspaces

A defective matrix does not have enough eigenvectors to fill $\mathbb{K}^m$. But intuitively, all vectors still show "λ-growth" under J. Our idea is to generalize the λ-eigenspace to an *invariant subspace with λ-growth*. This is a subspace that A maps into itself. On it, A need not act as a multiple of the identity. For J, the whole plane $\operatorname{span}(\vec{e}_1, \vec{e}_2)$ is mapped into itself. On it, J acts as $\lambda I + N$, not as λI.

Definition 73 (Invariant Subspace)

Let $X = [\vec{x}_1 \ \cdots \ \vec{x}_n] \in \mathbb{K}^{m\times n}$ have linearly independent columns. The space $\mathcal{R}(X)$ spanned by its columns is *A-invariant* if and only if there is $B \in \mathbb{K}^{n\times n}$ with

$$AX = XB. \tag{11.11}$$

Let us unpack the definition.

- Read column by column, the definition says $A\vec{x}_j = X\vec{b}_j$. So A maps every basis vector into $\mathcal{R}(X)$. So the definition is equivalent to $A\,\mathcal{R}(X) \subseteq \mathcal{R}(X)$. This is the basis-free definition in [4, 5.2] and [2, Def. 4.4].
- The matrix B is the matrix of A restricted to $\mathcal{R}(X)$, in the basis X. Since X has independent columns, B is unique.

The following table shows that the definition covers every case we have seen so far.

Situation	X	B
eigenpair	$\vec{x}_1$ $(n = 1)$	λ_1
eigenspace	a basis of E_λ	λI
diagonalizable A	all eigenvectors $(n = m)$	Λ
the matrix J above	I_2	J

So $AX = XB$ generalizes both $A\vec{x} = \lambda\vec{x}$ and $AX = X\Lambda$, and B need not be diagonal. The eigenvalues of B are still eigenvalues of A.

Proposition 74 (Eigenvalues on an Invariant Subspace)

Let $AX = XB$ with X of full column rank. Then every eigenvalue of B is an eigenvalue of A. If the columns of $X = Q$ are orthonormal, then $B = Q^*AQ$ [2, Prop. 4.3].

Proof. Suppose $B\vec{v} = \lambda\vec{v}$ with $\vec{v} \neq \vec{0}$. Then $A(X\vec{v}) = XB\vec{v} = \lambda X\vec{v}$, and $X\vec{v} \neq \vec{0}$ because X has independent columns. If $Q^*Q = I$, we multiply $AQ = QB$ by Q^* on the left and get $B = Q^*AQ$. □

For $n = 1$ and $\|\vec{q}\| = 1$, $Q^*AQ = \vec{q}^*A\vec{q}$ is the Rayleigh quotient (Definition 54). So Q^*AQ is a block version of the Rayleigh quotient.

The meaning of "λ-growth". Iterating $AX = XB$ gives $A^kX = XB^k$. So on an invariant subspace, the powers of A (and so power iteration) behave like the powers of the small matrix B. The eigenvalues of B control their growth. For the 2×2 block,

$$J^k = \begin{bmatrix} \lambda^k & k\lambda^{k-1} \\ 0 & \lambda^k \end{bmatrix}. \tag{11.12}$$

This grows like $|\lambda|^k$, up to a factor that is polynomial in k. Note: even for $|\lambda| < 1$, $\left\|J^k\right\|$ can first increase and then decay (see the supplement of Lecture 10).

11.6 The Jordan Normal Form

If we collect invariant subspaces on which A acts like a single eigenvalue, we get the Jordan normal form. It splits the whole space into A-invariant subspaces.

Theorem 75 (Jordan Normal Form)

For every $A \in \mathbb{K}^{m\times m}$ there is a nonsingular $S \in \mathbb{C}^{m\times m}$ with

$$A = SJS^{-1}, \qquad J = \begin{bmatrix} J_1 & & \\ & \ddots & \\ & & J_k \end{bmatrix}, \qquad J_\ell = \begin{bmatrix} \lambda_\ell & 1 & & \\ & \lambda_\ell & \ddots & \\ & & \ddots & 1 \\ & & & \lambda_\ell \end{bmatrix} \in \mathbb{C}^{n_\ell \times n_\ell}. \tag{11.13}$$

J is unique up to the order of the blocks [2, Thm. 4.1], [4, 8.46].

Reading the theorem.

- S and J are complex in general, even for real A, because the eigenvalues may be complex.
- The λ_ℓ need not be distinct: one eigenvalue may have several blocks. For each eigenvalue, the algebraic multiplicity is the total size of its blocks. The geometric multiplicity is the number of its blocks (by Proposition 76 below).
- A is diagonalizable if and only if all $n_\ell = 1$. Then $J = \Lambda$ and $S = X$.
- We split $S = [S_1 \ \cdots \ S_k]$ into column blocks that match the blocks of J. Then $AS_\ell = S_\ell J_\ell$. So each block gives an invariant subspace in the sense of Definition 73, with $B = J_\ell$.

Idea of the proof. The proof is pure linear algebra. We do not give it in full; see [4, Chapter 8]. Here is the outline.

- A nonzero $\vec{v}$ with $(A - \lambda I)^p \vec{v} = \vec{0}$ for some $p \geq 1$ is a *generalized eigenvector* [4, 8.8].
- The space $\mathbb{K}^m$ (over $\mathbb{C}$) is the direct sum of the generalized eigenspaces $\mathcal{N}(A - \lambda I)^m$. On each of them, $A - \lambda I$ is nilpotent [4, 8.14], that is, some power of it is zero.
- The columns of S that belong to one block form a *Jordan chain*: $(A - \lambda I)\vec{s}_j = \vec{s}_{j-1}$, $(A - \lambda I)\vec{s}_1 = \vec{0}$. This is the chain $\vec{e}_2 \mapsto \vec{e}_1 \mapsto \vec{0}$ of N above. Only $\vec{s}_1$ is an eigenvector.

Proposition 76 (One Eigenvector per Jordan Block)

Every Jordan block J_ℓ has, up to scaling, only one eigenvector, $\vec{e}_1$ [2, Prop. 4.2].

Proof. The only eigenvalue of J_ℓ is λ_ℓ. The matrix $J_\ell - \lambda_\ell I$ has ones on the superdiagonal and zeros elsewhere. So it shifts the entries up: $(J_\ell - \lambda_\ell I)\vec{x} = (x_2, \ldots, x_{n_\ell}, 0)^\top$. This is zero if and only if $x_2 = \cdots = x_{n_\ell} = 0$, that is, $\vec{x} \in \operatorname{span}(\vec{e}_1)$. □

11.6.1 The Jordan Form Is Not Numerically Stable

The Jordan form exists for every matrix. So why not compute it? The problem is that it is not (numerically) stable. This shows in two ways.

- It is discontinuous in the input A.
- The transformation S can be arbitrarily ill-conditioned.

Example 77 (A Tiny Perturbation Changes the Jordan Form)**.** Perturb $N = \left[\begin{smallmatrix} 0 & 1 \\ 0 & 0 \end{smallmatrix}\right]$ to

$$A_\varepsilon = \begin{bmatrix} \varepsilon & 1 \\ 0 & 2\varepsilon \end{bmatrix}, \qquad \varepsilon > 0. \tag{11.14}$$

A_ε is upper triangular with distinct eigenvalues ε and 2ε. So it is diagonalizable. Let us compute its eigenvectors.

- For ε, the eigenvector is $\vec{e}_1$.
- For 2ε, the equation $(A_\varepsilon - 2\varepsilon I)\vec{x} = \left[\begin{smallmatrix} -\varepsilon & 1 \\ 0 & 0 \end{smallmatrix}\right] \vec{x} = \vec{0}$ gives the eigenvector $(1, \varepsilon)^\top$.

Hence

$$A_\varepsilon = S_\varepsilon \begin{bmatrix} \varepsilon & 0 \\ 0 & 2\varepsilon \end{bmatrix} S_\varepsilon^{-1}, \qquad S_\varepsilon = \begin{bmatrix} 1 & 1 \\ 0 & \varepsilon \end{bmatrix}. \tag{11.15}$$

As $\varepsilon \to 0$, $A_\varepsilon \to N$. But the Jordan form $\operatorname{diag}(\varepsilon, 2\varepsilon)$ tends to the zero matrix, while the Jordan form of N is $N \neq 0$. So the Jordan form is a *discontinuous* function of A. More generally, take an $n \times n$ Jordan block with eigenvalue 0 and add $\operatorname{diag}(\varepsilon, 2\varepsilon, \ldots, n\varepsilon)$. Its Jordan form becomes $\operatorname{diag}(\varepsilon, \ldots, n\varepsilon)$ [2, Example 4.2].

In floating point, perturbations of size $\varepsilon_{\text{mach}} \|A\|$ are always present. So we cannot answer the question "does A have a nontrivial Jordan block?" by computation.

S can be arbitrarily ill-conditioned. The two columns of S_ε are nearly parallel: the angle θ between them satisfies $\tan\theta = \varepsilon$. We normalize both columns to unit length. Then the singular values are $\sqrt{1 \pm \cos\theta}$, so

$$\kappa_2(S_\varepsilon) = \sqrt{\frac{1+\cos\theta}{1-\cos\theta}} = \cot\frac{\theta}{2} \approx \frac{2}{\varepsilon}. \tag{11.16}$$

Could a better scaling of the columns help? No. Let $\vec{u}, \vec{v}$ be the unit columns, and let $SD = [a\vec{u}\; b\vec{v}]$ with $a, b \neq 0$.

- For $\vec{w} = (1/a, -1/b)^\top$ we get $SD\vec{w} = \vec{u} - \vec{v}$ and $\|\vec{w}\| \geq 1/\min(|a|, |b|)$. So $\sigma_{\min}(SD) \leq \min(|a|, |b|)\, \|\vec{u} - \vec{v}\|$.
- Also $\sigma_{\max}(SD) \geq \max(|a|, |b|)$, the norm of the larger column.

Hence $\kappa_2(SD) \geq 1/\|\vec{u} - \vec{v}\| = 1/(2\sin(\theta/2)) \approx 1/\varepsilon$. Since A_ε is diagonalizable, this shows one more thing. The eigenvector matrix X of a diagonalizable matrix can be arbitrarily ill-conditioned when the matrix is close to a defective one.

Why an ill-conditioned S is a serious problem. Suppose we are lucky: we compute S exactly, and J with a tiny error δJ, $\|\delta J\| = O(\varepsilon_{\text{mach}})\,\|A\|$. The backward error δA with $S^{-1}(A+\delta A)S = J + \delta J$ is $\delta A = S\,\delta J\,S^{-1}$. All we can say is

$$\|\delta A\| \le \kappa(S)\,\|\delta J\| = O(\varepsilon_{\text{mach}})\,\kappa(S)\,\|A\|\,. \tag{11.17}$$

This can be much larger than $\varepsilon_{\text{mach}}\,\|A\|$. So we cannot compute the Jordan form in a backward stable way [2, Example 4.3]. The same factor $\kappa(S)$ multiplies the error of any computation of the form $f(A) = Sf(J)S^{-1}$.

The conclusion: we should use only *unitary* similarity transformations, so that $\kappa_2 = 1$. Then we cannot reach a form as simple as J. But we get something almost as good.

11.7 The Schur Factorization

Theorem 78 (Schur Factorization)

For every $A \in \mathbb{K}^{m\times m}$ there exist a unitary Q and an upper triangular T with

$$A = QTQ^*. \tag{11.18}$$

The eigenvalues of A are the diagonal entries of T [1, Thm. 24.9], [2, Thm. 4.2], [4, 6.38].

The idea of the proof: every matrix has at least one eigenpair. We rotate it into the first column by a unitary similarity. Then we recurse on the bottom right part.

Proof. We use induction on m. For $m = 1$, take $Q = [1]$ and $T = A$. Now let $m \ge 2$.

Step 1: find one eigenpair. By the fundamental theorem of algebra, p_A has a root $\lambda \in \mathbb{C}$. So there is $\vec{u}$ with $A\vec{u} = \lambda\vec{u}$, and we normalize it to $\|\vec{u}\|_2 = 1$.

Step 2: rotate it into the first column. Extend $\vec{u}$ to a square unitary matrix $U = [\vec{u}\;\tilde{U}]$, with $\tilde{U} \in \mathbb{C}^{m\times(m-1)}$. For example, we can take a Householder reflector that maps $\vec{e}_1$ to a multiple of $\vec{u}$ (Lecture 6). Then

$$U^*AU = \begin{bmatrix} \vec{u}^*A\vec{u} & \vec{u}^*A\tilde{U} \\ \tilde{U}^*A\vec{u} & \tilde{U}^*A\tilde{U} \end{bmatrix} = \begin{bmatrix} \lambda & \vec{u}^*A\tilde{U} \\ \vec{0} & A_2 \end{bmatrix}, \qquad A_2 \coloneqq \tilde{U}^*A\tilde{U}, \tag{11.19}$$

because $\vec{u}^*A\vec{u} = \lambda\vec{u}^*\vec{u} = \lambda$ and $\tilde{U}^*A\vec{u} = \lambda\tilde{U}^*\vec{u} = \vec{0}$ by orthogonality.

Step 3: recurse on the bottom right part. By the induction hypothesis, $A_2 = \tilde{Q}\tilde{T}\tilde{Q}^*$ with $\tilde{Q}$ unitary and $\tilde{T}$ upper triangular. Set

$$Q \coloneqq U\begin{bmatrix} 1 & \\ & \tilde{Q} \end{bmatrix}. \tag{11.20}$$

This is a product of unitary matrices. Then

$$Q^*AQ = \begin{bmatrix} 1 & \\ & \tilde{Q}^* \end{bmatrix}\begin{bmatrix} \lambda & \vec{u}^*A\tilde{U} \\ \vec{0} & A_2 \end{bmatrix}\begin{bmatrix} 1 & \\ & \tilde{Q} \end{bmatrix} = \begin{bmatrix} \lambda & \vec{u}^*A\tilde{U}\tilde{Q} \\ \vec{0} & \tilde{T} \end{bmatrix} \tag{11.21}$$

is upper triangular. We call it T.

Step 4: the eigenvalues. A and T are similar, so they have the same characteristic polynomial. Since T is triangular, $p_T(z) = \prod_i (z - t_{ii})$. □

Step 1 uses the fundamental theorem of algebra. This is where $\mathbb{C}$ enters. Over $\mathbb{R}$, a matrix may have no eigenvalue at all (for example, a rotation of the plane). So Q and T are complex in general, even for real A.

Remarks.

- *Comparison with Jordan.* Both forms exist for every matrix. Jordan uses an S that may be ill-conditioned, and it is discontinuous. Schur uses a unitary Q with $\kappa_2(Q) = 1$. The price is that T is only triangular: its strictly upper part may be dense.
- *Non-uniqueness.* In Step 1 we can pick any eigenvalue. So the eigenvalues can appear on the diagonal of T in any order.
- *Nested invariant subspaces.* We write $AQ = QT$ and keep the first k columns. Since T is upper triangular, this gives $AQ_{:,1:k} = Q_{:,1:k}T_{1:k,1:k}$. So $\operatorname{span}(\vec{q}_1) \subset \operatorname{span}(\vec{q}_1, \vec{q}_2) \subset \cdots$ is a nested sequence of invariant subspaces (Definition 73), with orthonormal bases. Conversely, if all leading spans of a basis are invariant, the basis gives an upper triangular matrix [4, 5.39]. This is the goal of subspace iteration with orthonormalization (Section 11.3).
- *The proof is deflation.* Each step puts one eigenvalue in the top left corner by a unitary similarity. The other eigenvalues are those of A_2. This answers the question of Section 11.2 for all matrices, Hermitian or not, with only unitary transformations. But it is not yet an algorithm: each step needs an eigenpair, and we must compute that eigenpair iteratively.

As promised, the spectral theorem now follows in two lines.

Corollary 79 (Normal Matrices and the Spectral Theorem)**.** *If A is normal ($A^*A = AA^*$), then the triangular factor T of any Schur factorization of A is diagonal, so $A = Q\Lambda Q^*$. If A is Hermitian, Λ is also real. In particular, Theorem 69 holds [1, Thm. 24.8].*

Proof. Hermitian case. $T = Q^*AQ$ satisfies $T^* = Q^*A^*Q = T$. A Hermitian upper triangular matrix is diagonal, and its diagonal is real.

Normal case. This is the forward direction of Theorem 84, proved in Lecture 12. □

11.7.1 The Real Schur Form

For real matrices, we prefer to stay with real numbers. There are two reasons.

- *Cost.* A complex multiplication costs four real multiplications and two additions. Complex storage is twice as large.
- *Structure.* A real matrix has its non-real eigenvalues in conjugate pairs $\lambda, \overline{\lambda}$. Complex arithmetic with rounding would break this exact symmetry. Real arithmetic keeps it automatically.


Theorem 80 (Real Schur Form)

If $A \in \mathbb{R}^{m\times m}$, there exist an orthogonal Q and a block upper triangular T with 1×1 or 2×2 diagonal blocks such that

$$A = QTQ^\top. \tag{11.22}$$

The 1×1 blocks are the real eigenvalues of A. Each 2×2 block has a pair of complex conjugate eigenvalues of A [2, Thm. 4.3].

Such a T is called *quasi-upper triangular*.



The quasi-upper triangular factor of the real Schur form. Shaded entries may be nonzero. Each diagonal block holds a real eigenvalue or a conjugate pair.

The key idea: the complex eigenvalues of a real matrix come in conjugate pairs. Each pair gives a *real* two-dimensional invariant subspace.

Lemma 81 (A Conjugate Pair Gives a Real Invariant Plane)

Let $A \in \mathbb{R}^{m\times m}$, and let $A\vec{x} = \lambda\vec{x}$ with $\lambda = \alpha + i\beta$, $\beta \neq 0$, and $\vec{x} = \vec{a} + i\vec{b}$ with $\vec{a}, \vec{b} \in \mathbb{R}^m$. Then $\vec{a}, \vec{b}$ are linearly independent and

$$A\begin{bmatrix}\vec{a} & \vec{b}\end{bmatrix} = \begin{bmatrix}\vec{a} & \vec{b}\end{bmatrix}\begin{bmatrix}\alpha & \beta\\ -\beta & \alpha\end{bmatrix}, \tag{11.23}$$

where the 2×2 matrix has eigenvalues $\alpha \pm i\beta$.

Proof. The identity. Since A is real, we can compare the real and imaginary parts of $A(\vec{a} + i\vec{b}) = (\alpha + i\beta)(\vec{a} + i\vec{b})$. This gives

$$A\vec{a} = \alpha\vec{a} - \beta\vec{b}, \qquad A\vec{b} = \beta\vec{a} + \alpha\vec{b}. \tag{11.24}$$

This is the matrix identity, column by column.

Independence. Suppose $\vec{a}, \vec{b}$ were dependent. Then $\vec{x}$ would be a complex multiple of a single real vector $\vec{r} \neq \vec{0}$, so $A\vec{r} = \lambda\vec{r}$. The left side is real, so λ must be real. This is a contradiction.

The eigenvalues. The eigenvalues of $\left[\begin{smallmatrix}\alpha & \beta\\ -\beta & \alpha\end{smallmatrix}\right]$ are the roots of $(z-\alpha)^2 + \beta^2$, that is, $\alpha \pm i\beta$. □

The 2×2 matrix is the standard real representation of the complex number $\alpha + i\beta$, acting by multiplication.

Proof of Theorem 80. We use induction as in Theorem 78, but with real arithmetic. There are two cases.

A real eigenvalue. It has a real eigenvector. The step is the same as before, with a 1×1 block.

No real eigenvalue. Take $\lambda = \alpha + i\beta$ with $\beta \neq 0$, and form $[\vec{a}\ \vec{b}]$ as in Lemma 81. Compute its QR factorization $[\vec{a}\ \vec{b}] = \hat{Q}R$, with $\hat{Q} \in \mathbb{R}^{m\times 2}$ and R invertible. Then $\mathcal{R}(\hat{Q}) = \operatorname{span}(\vec{a}, \vec{b})$ is invariant. We

have $A\hat{Q} = \hat{Q}\hat{B}$ with $\hat{B} = R \left[\begin{smallmatrix} \alpha & \beta \\ -\beta & \alpha \end{smallmatrix}\right] R^{-1}$, and $\hat{B}$ still has eigenvalues $\alpha \pm i\beta$. Extend $\hat{Q}$ to an orthogonal $U = [\hat{Q}\ \tilde{Q}]$. By Proposition 74, $\hat{Q}^\top A\hat{Q} = \hat{B}$. Also $\tilde{Q}^\top A\hat{Q} = \tilde{Q}^\top \hat{Q}\hat{B} = 0$. So

$$U^\top AU = \begin{bmatrix} \hat{B} & \hat{Q}^\top A\tilde{Q} \\ 0 & \tilde{Q}^\top A\tilde{Q} \end{bmatrix}, \tag{11.25}$$

and we recurse on $\tilde{Q}^\top A\tilde{Q}$. □

A 2×2 block with a non-real conjugate pair cannot be made triangular over $\mathbb{R}$. Its triangular form would show the non-real eigenvalues on a real diagonal.

11.8 Computing Eigenvectors from the Schur Form

The Schur factorization gives us the eigenvalues. What about the eigenvectors? Suppose we know a Schur factorization $A = QTQ^*$. Then

$$T\vec{x} = \lambda\vec{x} \iff (QTQ^*)(Q\vec{x}) = \lambda Q\vec{x}. \tag{11.26}$$

So $\vec{x}$ is an eigenvector of T if and only if $Q\vec{x}$ is an eigenvector of A. So we need eigenvectors of the triangular matrix T. We will see that this is just a triangular solve.

Proposition 82 (Eigenvectors of a Triangular Matrix)

Let T be upper triangular and let $\lambda = t_{kk}$ be a simple eigenvalue. Partition

$$T - \lambda I = \begin{bmatrix} T_{11} - \lambda I & \vec{t}_{12} & T_{13} \\ & 0 & \vec{t}_{23}^* \\ & & T_{33} - \lambda I \end{bmatrix}. \tag{11.27}$$

Here T_{11} is $(k-1) \times (k-1)$, the middle block is 1×1, and T_{33} is $(m-k) \times (m-k)$. Then

$$\vec{x} = \begin{bmatrix} (\lambda I - T_{11})^{-1}\vec{t}_{12} \\ 1 \\ \vec{0} \end{bmatrix} \tag{11.28}$$

is an eigenvector of T for λ, and $Q\vec{x}$ is an eigenvector of A [2, §4.2.1].

Proof. We partition $\vec{x} = (\vec{x}_1, x_2, \vec{x}_3)$ in the same way. Then $(T - \lambda I)\vec{x} = \vec{0}$ reads

$$\begin{bmatrix} (T_{11} - \lambda I)\vec{x}_1 + \vec{t}_{12}x_2 + T_{13}\vec{x}_3 \\ \vec{t}_{23}^*\vec{x}_3 \\ (T_{33} - \lambda I)\vec{x}_3 \end{bmatrix} = \vec{0}. \tag{11.29}$$

Since λ is simple, it does not appear on the diagonal of T_{11} or T_{33}. So $T_{11} - \lambda I$ and $T_{33} - \lambda I$ are nonsingular triangular matrices. We solve from the bottom up.

- The third row gives $\vec{x}_3 = \vec{0}$. Then the second row holds automatically.
- The scalar x_2 is free, and we choose $x_2 = 1$.
- The first row becomes $(T_{11} - \lambda I)\vec{x}_1 = -\vec{t}_{12}$, that is, $\vec{x}_1 = (\lambda I - T_{11})^{-1}\vec{t}_{12}$. □

Cost. Computing $\vec{x}_1$ is one triangular solve of size $k-1$, about $(k-1)^2$ flops. Computing $Q\vec{x}$ uses only the first k columns of Q, about $2mk$ flops. Summing over $k = 1, \dots, m$ gives roughly $\frac{1}{3}m^3 + m^3$ flops for all eigenvectors. This is the same order $O(m^3)$ as computing the Schur form itself. For $k = 1$, $\vec{x} = \vec{e}_1$ and $Q\vec{x} = \vec{q}_1$. So the first Schur vector is always an eigenvector: it is the $\vec{u}$ chosen in Step 1 of the proof of Theorem 78.

Remarks.

- *Repeated eigenvalues.* If $t_{jj} = \lambda$ for some $j < k$, then $T_{11} - \lambda I$ is singular, and there may be no eigenvector with $x_k = 1$. This is the defective case of Section 11.5 again.
- *Sensitivity.* The Schur form can be computed stably. But if λ is close to some other diagonal entry of T_{11}, then $(\lambda I - T_{11})^{-1}$ is large, and $\vec{x}$ is very sensitive to perturbations. This is not a flaw of the method. The eigenvector itself is ill-conditioned. Compare Example 77, where two eigenvectors become nearly parallel.
- *Real Schur form.* From the real Schur form, we solve a quasi-triangular system instead. Computing complex eigenvectors in real arithmetic needs a bit more care. LAPACK does it in the routine `xTREVC` [2, §4.2.1].

11.9 (SUPPLEMENT) Further Topics

11.9.1 Deflation for Non-Hermitian Matrices

Hotelling deflation (Proposition 65) needs orthogonal eigenvectors. Now let A be general, with $A\vec{x}_1 = \lambda_1\vec{x}_1$, and let $\vec{w} \in \mathbb{K}^m$ be arbitrary. Then the eigenvalues of $A - \vec{x}_1\vec{w}^*$ are $\lambda_1 - \vec{w}^*\vec{x}_1, \lambda_2, \dots, \lambda_m$, counted with multiplicity (Brauer's theorem). The proof is one step of the Schur proof. Let $\hat{\vec{x}}_1 = \vec{x}_1 / \|\vec{x}_1\|$, and let $U = [\hat{\vec{x}}_1\ \tilde{U}]$ be unitary. Then

$$U^*(A - \vec{x}_1\vec{w}^*)U = \begin{bmatrix} \lambda_1 & * \\ \vec{0} & A_2 \end{bmatrix} - \begin{bmatrix} \|\vec{x}_1\| \\ \vec{0} \end{bmatrix} \vec{w}^*U = \begin{bmatrix} \lambda_1 - \vec{w}^*\vec{x}_1 & * \\ \vec{0} & A_2 \end{bmatrix}. \tag{11.30}$$

Here we used $U^*\vec{x}_1 = \|\vec{x}_1\|\,\vec{e}_1$. Also, the $(1,1)$ entry of the correction is $\|\vec{x}_1\|\,\vec{w}^*\hat{\vec{x}}_1 = \vec{w}^*\vec{x}_1$.

- Choose $\vec{w} = \overline{\lambda_1}\vec{v}$ with $\vec{v}^*\vec{x}_1 = 1$, so that $\vec{x}_1\vec{w}^* = \lambda_1\vec{x}_1\vec{v}^*$ and $\vec{w}^*\vec{x}_1 = \lambda_1$. This moves λ_1 to 0 and keeps the rest of the spectrum (Wielandt deflation).
- The choice $\vec{v} = \vec{x}_1 / \|\vec{x}_1\|^2$ gives back Hotelling's formula.

Unlike in the Hermitian case, the eigenvectors change. Let $B = A - \lambda_1\vec{x}_1\vec{v}^*$, and let $B\vec{w}_i = \lambda_i\vec{w}_i$ with $\lambda_i \neq \lambda_1$. A direct computation shows that $\vec{x}_i = \vec{w}_i + c\,\vec{x}_1$ with $c = \lambda_1\vec{v}^*\vec{w}_i/(\lambda_i - \lambda_1)$ is an eigenvector of A.

11.9.2 How Ill-Conditioned Eigenvalues Near a Jordan Block Are

In Example 77, the eigenvalues moved by only $O(\varepsilon)$, while the Jordan structure and S broke down. A different perturbation moves the eigenvalues themselves much more. Put ε in the bottom left corner of an $n \times n$ Jordan block with eigenvalue 0. The result is the companion matrix (Lemma 57) of $z^n - \varepsilon$. Its eigenvalues are the n complex n-th roots of ε, all of modulus $\varepsilon^{1/n}$ [2, Example 4.4]. For $n = 10$ and $\varepsilon = 10^{-10}$, a perturbation of size 10^{-10} moves the eigenvalues by 0.1. A multiple eigenvalue in a nontrivial Jordan block has infinite condition number.

11.9.3 The Bauer–Fike Theorem

We can make the role of $\kappa(X)$ quantitative. Let $A = X\Lambda X^{-1}$ be diagonalizable, and let μ be an eigenvalue of $A + E$. Then

$$\min_i |\mu - \lambda_i| \le \kappa_2(X) \|E\|_2 . \tag{11.31}$$

To see this, assume μ is not an eigenvalue of A. Since $A + E - \mu I$ is singular, so is $X^{-1}(A + E - \mu I)X = (\Lambda - \mu I) + X^{-1}EX$. Hence $I + (\Lambda - \mu I)^{-1}X^{-1}EX$ is also singular. A matrix $I + M$ can only be singular if $\|M\| \ge 1$. So

$$1 \le \left\|(\Lambda - \mu I)^{-1}\right\| \left\|X^{-1}\right\| \|E\| \|X\| = \frac{\kappa_2(X)\|E\|_2}{\min_i |\lambda_i - \mu|}. \tag{11.32}$$

For normal matrices, X can be unitary. Then $\kappa_2(X) = 1$, and the eigenvalues move by at most $\|E\|_2$. This is the eigenvalue version of the remark after Theorem 69. When A approaches a defective matrix, $\kappa_2(X) \to \infty$ and the bound becomes useless. This agrees with the $\varepsilon^{1/n}$ behavior above. See [2, §4.3].

11.9.4 From the Schur Form to the QR Algorithm

Practical eigenvalue solvers compute the Schur factorization. Consider subspace iteration with $n = m$ vectors and a QR factorization after each multiplication by A. Under suitable conditions, it produces a $Q^{(k)}$ whose leading columns converge to the nested invariant subspaces of Section 11.7. Then $(Q^{(k)})^* A Q^{(k)}$ converges to an upper triangular T. If we rewrite this iteration in terms of $(Q^{(k)})^* A Q^{(k)}$ alone, we get the QR algorithm. We combine it with a first reduction to Hessenberg form and with shifts (the Rayleigh quotient idea of Lecture 10). Then it computes the Schur form in $O(m^3)$ flops. The real version produces the real Schur form in real arithmetic. See [1, Lectures 25 and 28].

Lecture 12

Orthogonal Iteration

Supplementary reading:

- [1] Lectures 24 and 28.
- [2] Section 4.4.3.

12.1 Recap

Last lecture (Lecture 11) we moved from computing one eigenpair to a new question: what should "all eigenpairs" mean as a single object?

- *Deflation* (Section 11.2). Once we know $(\lambda_1, \vec{x}_1)$, power iteration on $A - \lambda_1 \vec{x}_1 \vec{x}_1^* / (\vec{x}_1^* \vec{x}_1)$ finds the next eigenpair (for Hermitian A, Proposition 65).
- *The need for subspace iteration* (Section 11.3). Deflation computes the eigenpairs one after another. Each one needs many Level 2 matrix-vector products. If we iterate on a block of vectors at once, these become Level 3 matrix-matrix products.
- *Eigenvalues as matrix factorizations* (Section 11.4). All eigenpairs together form the eigenvalue decomposition $A = X \Lambda X^{-1}$. But it does not always exist. For example, $N = \left[\begin{smallmatrix} 0 & 1 \\ 0 & 0 \end{smallmatrix}\right]$ sends $\vec{e}_2 \mapsto \vec{e}_1 \mapsto \vec{0}$ and has only one eigenvector.
- *Defective matrices and Jordan blocks* (Sections 11.5 and 11.6). Every matrix has a Jordan form $A = SJS^{-1}$. Here $J = \operatorname{diag}(J_1, \ldots, J_k)$, and each $J_\ell \in \mathbb{C}^{n_\ell \times n_\ell}$ has λ_ℓ on the diagonal and ones on the superdiagonal. The Jordan form is not continuous in A, and S can be arbitrarily ill-conditioned (Example 77).
- *Schur factorization* (Section 11.7). Every matrix can be written as $A = QTQ^*$ with Q unitary and T upper triangular (Theorem 78). This is the stable replacement for the Jordan form. Eigenvalue algorithms compute this factorization.

12.2 From Schur Factorization to Diagonalization

The Schur factorization $A = QTQ^*$ always exists, but T is only upper triangular. When is it a diagonalization? That is, when is T diagonal, so that $A = Q \Lambda Q^*$ with Q unitary?

Definition 83 (Normal Matrix)

$A \in \mathbb{K}^{m\times m}$ is *normal* if $AA^* = A^*A$.

Hermitian ($A^* = A$), skew-Hermitian ($A^* = -A$), and unitary ($A^* = A^{-1}$) matrices are normal. In each case A^* is a function of A, so it commutes with A. The nilpotent matrix $N = \left[\begin{smallmatrix} 0 & 1 \\ 0 & 0 \end{smallmatrix}\right]$ is not normal:

$$NN^* = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \neq \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} = N^*N. \tag{12.1}$$

Theorem 84 (Normal Matrices Are Unitarily Diagonalizable)

$A \in \mathbb{K}^{m\times m}$ is normal if and only if it has an orthogonal eigenvalue decomposition $A = Q\Lambda Q^*$ with Q unitary and Λ diagonal. In this case the triangular factor T of every Schur factorization $A = QTQ^*$ is diagonal.

Over $\mathbb{C}$, an "orthogonal" eigenvalue decomposition means that the eigenvector matrix Q is unitary.

Proof. If $A = Q\Lambda Q^*$, then A is normal. We have $A^* = Q\overline{\Lambda}Q^*$. Diagonal matrices commute, so

$$AA^* = Q\Lambda\overline{\Lambda}Q^* = Q\overline{\Lambda}\Lambda Q^* = A^*A. \tag{12.2}$$

If A is normal, then T is diagonal. Take a Schur factorization $A = QTQ^*$ (Theorem 78). Then


$$QTT^*Q^* = AA^* = A^*A = QT^*TQ^*. \tag{12.3}$$

We multiply by Q^* on the left and by Q on the right. This gives $TT^* = T^*T$, so the triangular factor is normal too. Now we compare the $(1,1)$ entries. Since T is upper triangular, its first column is $t_{11}\vec{e}_1$. But its first row is full:

$$(T^*T)_{11} = \sum_{i=1}^{m} |t_{i1}|^2 = |t_{11}|^2, \qquad (TT^*)_{11} = \sum_{i=1}^{m} |t_{1i}|^2. \tag{12.4}$$

Setting them equal gives $\sum_{i=2}^{m} |t_{1i}|^2 = 0$. So $t_{1i} = 0$ for $i \geq 2$, and $T = \operatorname{diag}(t_{11}, T_2)$ with T_2 upper triangular. The lower right blocks of $TT^* = T^*T$ give $T_2T_2^* = T_2^*T_2$, so T_2 is normal again. By induction on m, T is diagonal. So $A = QTQ^*$ is an orthogonal eigenvalue decomposition. □

In pictures, using the shapes of the factors, the key identity says: lower times upper equals upper times lower.



The $(1,1)$ entry of the left side uses only the first column of T, which is just t_{11}. The $(1,1)$ entry of the right side uses the whole first row. For the two to be equal, the rest of the first row must be zero.

12.3 Orthogonal Iteration

We now come back to the question from Section 11.3: how do we run power iteration on several vectors at once, without all of them collapsing onto $\vec{x}_1$? In this section we assume that $A \in \mathbb{R}^{m\times m}$ is diagonalizable, with

$$|\lambda_1| > |\lambda_2| > \cdots > |\lambda_m|. \tag{12.5}$$

The non-real eigenvalues of a real matrix come in conjugate pairs with equal modulus. So the strict ordering forces all eigenvalues to be real, and then all eigenvectors are real too.

Suppose we want to compute $r = 2$ eigenpairs at once. We start by applying power iteration to two vectors at the same time.

- *Problem.* If we run them independently, both vectors converge to the same vector $\vec{x}_1$ (up to sign). The reason is that almost every starting vector has a component along $\vec{x}_1$.
- *Idea.* After each multiplication, we project $\vec{q}_2$ onto the orthogonal complement of $\vec{q}_1$. Then $\vec{q}_2$ stays orthogonal to $\vec{q}_1$, so it cannot also converge to $\vec{x}_1$.

Algorithm 18 Orthogonal iteration with two vectors.

1: $\vec{q}_1, \vec{q}_2 \leftarrow$ random vectors
2: **while** not converged **do**
3: $\quad \vec{q}_1 \leftarrow A\vec{q}_1, \quad \vec{q}_2 \leftarrow A\vec{q}_2$
4: $\quad \vec{q}_2 \leftarrow \left(I - \dfrac{\vec{q}_1\vec{q}_1^*}{\vec{q}_1^*\vec{q}_1}\right)\vec{q}_2$ ▷ project out $\vec{q}_1$
5: $\quad \vec{q}_1 \leftarrow \vec{q}_1 / \|\vec{q}_1\|, \quad \vec{q}_2 \leftarrow \vec{q}_2 / \|\vec{q}_2\|$

The last two lines are one step of Gram–Schmidt on $[A\vec{q}_1 \; A\vec{q}_2]$. In other words, they compute a reduced QR factorization of a two-column matrix (Lecture 5). The first column is plain power iteration, so $\vec{q}_1 \to \pm\vec{x}_1 / \|\vec{x}_1\|$. Compare this with deflation (Section 11.2):

- In deflation, we projected out the *converged* $\vec{x}_1$ at every step.
- Here, we project out the current approximation $\vec{q}_1$. So both vectors are iterated at the same time.

12.3.1 What Do the Two Vectors Converge To?

We write $\vec{q}_i^{(k)}$ for the iterates after k steps. The projection never changes the first vector, so

$$\vec{q}_1^{(k)} = \frac{A^k \vec{q}_1^{(0)}}{\left\|A^k \vec{q}_1^{(0)}\right\|}. \tag{12.6}$$

So $\vec{q}_1$ does plain power iteration (Algorithm 15). It converges to $\vec{x}_1$ (normalized, up to sign) at the rate $|\lambda_2/\lambda_1|$.

For $\vec{q}_2$, assume that $\vec{q}_1 \approx \vec{x}_1$ already, with $\|\vec{x}_1\| = 1$. Then the projection in Algorithm 18 is (approximately) a fixed matrix. Up to normalization,

$$\vec{q}_2^{(k)} \approx (I - \vec{x}_1\vec{x}_1^*)A\vec{q}_2^{(k-1)} = (I - \vec{x}_1\vec{x}_1^*)A(I - \vec{x}_1\vec{x}_1^*)\vec{q}_2^{(k-1)}. \tag{12.7}$$

The equality holds because $\vec{q}_2^{(k-1)}$ was already projected onto the orthogonal complement of $\vec{x}_1$ in the previous step. So applying the projector once more changes nothing. Thus, once $\vec{q}_1$ has converged, $\vec{q}_2$ does power iteration with the matrix $(I - \vec{x}_1\vec{x}_1^*)A(I - \vec{x}_1\vec{x}_1^*)$.

The normal case. For normal A, this matrix is exactly the Hotelling deflation of Proposition 65:

$$(I - \vec{x}_1\vec{x}_1^*)A(I - \vec{x}_1\vec{x}_1^*) = A - 2\lambda_1\vec{x}_1\vec{x}_1^* + \lambda_1\vec{x}_1\vec{x}_1^* = A - \lambda_1\vec{x}_1\vec{x}_1^*. \tag{12.8}$$

To see the first equality, we expand the product into four terms:

$$A - \vec{x}_1(\vec{x}_1^* A) - (A\vec{x}_1)\vec{x}_1^* + \vec{x}_1(\vec{x}_1^* A\vec{x}_1)\vec{x}_1^*. \tag{12.9}$$

We have $A\vec{x}_1 = \lambda_1\vec{x}_1$ and $\vec{x}_1^* A\vec{x}_1 = \lambda_1$. For the remaining term we need normality. It makes $\vec{x}_1$ also a *left* eigenvector: $\vec{x}_1^* A = \lambda_1\vec{x}_1^*$. To see this, note that $B \coloneqq A - \lambda_1 I$ is normal, since A is. For any normal B and any $\vec{v}$,

$$\|B\vec{v}\|^2 = \vec{v}^* B^* B\vec{v} = \vec{v}^* BB^*\vec{v} = \|B^*\vec{v}\|^2. \tag{12.10}$$

So $B\vec{x}_1 = \vec{0}$ implies $B^*\vec{x}_1 = \vec{0}$, i.e., $A^*\vec{x}_1 = \overline{\lambda_1}\vec{x}_1$. Taking the conjugate transpose gives $\vec{x}_1^* A = \lambda_1\vec{x}_1^*$. So three of the four terms are equal to $\pm\lambda_1\vec{x}_1\vec{x}_1^*$.

So for normal A, $\vec{q}_2$ does power iteration on the deflated matrix $A - \lambda_1\vec{x}_1\vec{x}_1^*$. Its eigenvalues are $0, \lambda_2, \ldots, \lambda_m$. So, once $\vec{q}_1$ has converged, $\vec{q}_2$ converges to $\vec{x}_2$ at the rate $|\lambda_3/\lambda_2|$. Overall, its error decays like $\max(|\lambda_2/\lambda_1|, |\lambda_3/\lambda_2|)^k$. Orthogonal iteration does deflation automatically. It does not wait for $\vec{x}_1$ to converge first.

The general case. Without normality, in general $\vec{x}_1^* A \neq \lambda_1\vec{x}_1^*$. Then the deflated matrix is no longer $A - \lambda_1\vec{x}_1\vec{x}_1^*$. We can still find its eigenpairs, by applying it to the eigenvectors of A. Write $P \coloneqq I - \vec{x}_1\vec{x}_1^*$. For every i,

$$PAP\vec{x}_i = P\big(\lambda_i\vec{x}_i - \lambda_1\vec{x}_1(\vec{x}_1^*\vec{x}_i)\big) = \lambda_i P\vec{x}_i, \tag{12.11}$$

since $P\vec{x}_1 = \vec{0}$. We can read the left side in two ways.

- As $[PA](P\vec{x}_i)$, the matrix PA applied to the vector $P\vec{x}_i$. Then the identity says directly that $PA(P\vec{x}_i) = \lambda_i(P\vec{x}_i)$.
- As PAP applied to $\vec{x}_i$. Using $P^2 = P$, we also get $PAP(P\vec{x}_i) = \lambda_i(P\vec{x}_i)$.

So the eigenpairs of $PA = (I - \vec{x}_1\vec{x}_1^*)A$ (and of PAP) are

$$(0, \vec{x}_1) \qquad \text{and} \qquad (\lambda_i, P\vec{x}_i), \quad i = 2, \ldots, m. \tag{12.12}$$

Here $P\vec{x}_i = \vec{x}_i - (\vec{x}_1^*\vec{x}_i)\vec{x}_1 \neq \vec{0}$ for $i \geq 2$, because the $\vec{x}_i$ are linearly independent. In words: the projection keeps the eigenvalues $\lambda_2, \ldots, \lambda_m$. But it replaces each eigenvector $\vec{x}_i$ by its component orthogonal to $\vec{x}_1$. In the normal case $\vec{x}_i \perp \vec{x}_1$, so $P\vec{x}_i = \vec{x}_i$. This agrees with Equation (12.8).

Conclusion. Once $\vec{q}_1$ has converged, $\vec{q}_2$ does power iteration with PA. The dominant eigenpair of PA is $(\lambda_2, P\vec{x}_2)$, and its next largest eigenvalue in modulus is λ_3. So, normalized and up to sign,

$$\vec{q}_2^{(k)} \longrightarrow (I - \vec{x}_1\vec{x}_1^*)\vec{x}_2, \tag{12.13}$$

at the rate $|\lambda_3/\lambda_2|$. This is *not* $\vec{x}_2$, unless $\vec{x}_1 \perp \vec{x}_2$ (as for normal A). In general the two vectors cannot both be eigenvectors: $\vec{q}_1 \perp \vec{q}_2$ by construction, but $\vec{x}_1, \vec{x}_2$ need not be orthogonal.

What does converge is the subspace. Since $P\vec{x}_2 = \vec{x}_2 - (\vec{x}_1^*\vec{x}_2)\vec{x}_1$, we get

$$\operatorname{span}(\vec{q}_1, \vec{q}_2) \longrightarrow \operatorname{span}(\vec{x}_1, \vec{x}_2). \tag{12.14}$$

This is an invariant subspace of A. The vectors $\vec{q}_1, \vec{q}_2$ form an orthonormal basis of it, and the first one is an eigenvector. These are exactly the first two columns of a Schur factorization (Section 11.7): $A\vec{q}_1 = \lambda_1\vec{q}_1$ and $A\vec{q}_2 \in \operatorname{span}(\vec{q}_1, \vec{q}_2)$. So T is upper triangular in its first two columns.

12.3.2 Extracting the Eigenvalues

Orthogonal iteration gives an orthonormal basis of $\operatorname{span}(\vec{x}_1, \vec{x}_2)$, but $\vec{q}_2$ is not an eigenvector. So how do we find the eigenvalues? We put the iterates into a tall matrix with orthonormal columns, and project A onto their span:


$$Q_k \coloneqq \begin{bmatrix} \vec{q}_1^{(k)} & \vec{q}_2^{(k)} \end{bmatrix} \in \mathbb{K}^{m \times 2}, \qquad Q_k^* A Q_k \in \mathbb{K}^{2 \times 2}. \tag{12.15}$$

Proposition 85 (Projected Matrix)

As $k \to \infty$,

$$Q_k^* A Q_k = \begin{bmatrix} \vec{q}_1^* A \vec{q}_1 & \vec{q}_1^* A \vec{q}_2 \\ \vec{q}_2^* A \vec{q}_1 & \vec{q}_2^* A \vec{q}_2 \end{bmatrix} \longrightarrow \begin{bmatrix} \lambda_1 & * \\ 0 & \lambda_2 \end{bmatrix}. \tag{12.16}$$

In terms of shapes: a $2 \times m$ matrix times an $m \times m$ matrix times an $m \times 2$ matrix gives a 2×2 upper triangular matrix. Its diagonal holds the two dominant eigenvalues.



Proof sketch. Write $Q = Q_k$. Assume k is large, so that $\vec{q}_1 \approx \vec{x}_1$ and $\operatorname{span}(\vec{q}_1, \vec{q}_2) \approx \operatorname{span}(\vec{x}_1, \vec{x}_2)$.

The $(2,1)$ entry tends to zero. We use $\vec{q}_1 \approx \vec{x}_1$ twice:

$$\vec{q}_2^* A \vec{q}_1 \approx \vec{q}_2^* A \vec{x}_1 = \lambda_1 \vec{q}_2^* \vec{x}_1 \approx \lambda_1 \vec{q}_2^* \vec{q}_1 = 0. \tag{12.17}$$

The eigenvalues are λ_1, λ_2. For $i = 1, 2$, the eigenvector $\vec{x}_i$ lies (approximately) in $\mathcal{R}(Q)$. So there is $\vec{v}_i \in \mathbb{K}^2$ with $Q\vec{v}_i = \vec{x}_i$. Using $Q^*Q = I_2$, we get

$$Q^* A Q \vec{v}_i = Q^* A \vec{x}_i = \lambda_i Q^* \vec{x}_i \approx \lambda_i Q^* Q \vec{v}_i = \lambda_i \vec{v}_i. \tag{12.18}$$

So $(\lambda_i, \vec{v}_i)$ are the eigenpairs of the 2×2 matrix Q^*AQ.

Order on the diagonal. The $(1,1)$ entry $\vec{q}_1^* A \vec{q}_1$ is the Rayleigh quotient of $\vec{q}_1 \approx \vec{x}_1$, so it tends to λ_1. A triangular matrix has its eigenvalues on the diagonal. So the $(2,2)$ entry tends to λ_2. □

The proof also shows how to get the eigenvectors back. We solve the small eigenvalue problem $Q_k^* A Q_k \vec{v}_i = \lambda_i \vec{v}_i$, and then lift: $\vec{x}_i \approx Q_k \vec{v}_i$.

The $(1,2)$ entry $*$ is in general not zero. We can compute it from $A\vec{q}_2 = \lambda_2 \vec{q}_2 + \beta \vec{q}_1$. In the limit it equals $\beta = (\lambda_2 - \lambda_1)(\vec{x}_1^* \vec{x}_2) / \|P\vec{x}_2\|$. This is zero exactly when $\vec{x}_1 \perp \vec{x}_2$.

12.4 Orthogonal Iteration in General

To compute $r > 2$ eigenpairs at once, we start with r random vectors, the columns of $Q \in \mathbb{K}^{m \times r}$. Multiplying by A is then one matrix-matrix product. The Gram–Schmidt step of Algorithm 18 becomes a reduced QR factorization, which orthonormalizes all r columns at once.

Algorithm 19 Orthogonal iteration.

Input: $A \in \mathbb{K}^{m\times m}$, starting $Q \in \mathbb{K}^{m\times r}$ (e.g., random)

1: **for** $k = 1, \ldots,$ iters **do**
2: $\quad Z \leftarrow AQ$ ▷ Level 3: A is read once for all r columns
3: $\quad QR \leftarrow Z$ ▷ reduced QR, $Q \in \mathbb{K}^{m\times r}$

In the QR factorization, column j is $A\vec{q}_j$ with its components along $\vec{q}_1, \ldots, \vec{q}_{j-1}$ removed. So the first two columns do exactly what Algorithm 18 does. In general, column j does power iteration, projected onto the orthogonal complement of the first $j-1$ columns.

Remark 86 (Julia)**.** The simple loop body `Q = A*Q; Q, R = qr(Q)` does not work in Julia. The reason: `qr` returns Q in the implicit form of Lecture 7, which acts as the full $m \times m$ matrix. So from the second iteration on, `A*Q` is $m \times m$. We must ask for the thin factor explicitly:

```
for k = 1:iters
    Q = Matrix(qr(A*Q).Q)    # m x r
end
```

12.5 Convergence of Eigenvector Estimates

We run Algorithm 19 with r columns. We write $Q_k = [\vec{q}_1^{(k)} \cdots \vec{q}_r^{(k)}]$ for the iterate after k steps. The first i columns of a QR factorization depend only on the first i columns of the matrix we factor. So the first i columns of Q_k are exactly what orthogonal iteration with i vectors would give. In other words, running Algorithm 19 with r columns runs it with $1, 2, \ldots, r$ columns at the same time. So the argument of Section 12.3 generalizes to

$$\operatorname{span}(\vec{q}_1^{(k)}, \ldots, \vec{q}_i^{(k)}) \longrightarrow \operatorname{span}(\vec{x}_1, \ldots, \vec{x}_i), \qquad i = 1, \ldots, r. \tag{12.19}$$

We give a proof using $\operatorname{span}(Q_k) = \operatorname{span}(A^k Q_0)$ in Section 13.3.

Eigenvalues. We use the same argument as in the proof of Proposition 85. For $i \le r$, the eigenvector $\vec{x}_i$ lies (approximately) in $\mathcal{R}(Q_k)$. So $Q_k\vec{v}_i \approx \vec{x}_i$ for some $\vec{v}_i \in \mathbb{K}^r$, and

$$Q_k^* A Q_k \vec{v}_i \approx Q_k^* A \vec{x}_i = \lambda_i Q_k^* \vec{x}_i \approx \lambda_i Q_k^* Q_k \vec{v}_i = \lambda_i \vec{v}_i. \tag{12.20}$$

So the eigenvalues of the $r \times r$ matrix $Q_k^* A Q_k$ converge to $\lambda_1, \ldots, \lambda_r$, and $Q_k\vec{v}_i$ approximates $\vec{x}_i$. What about the columns of Q_k themselves?

The symmetric (normal) case. If A is symmetric, or more generally normal, then as before $\vec{q}_1^{(k)} \to \vec{x}_1$ and $\vec{q}_2^{(k)} \to \vec{x}_2$, up to sign. The same argument works for every column. Suppose $\vec{q}_1, \ldots, \vec{q}_{i-1}$ have converged to the orthonormal eigenvectors $\vec{x}_1, \ldots, \vec{x}_{i-1}$. Then column i does power iteration with

$$P_{i-1}A, \qquad P_{i-1} := I - \sum_{j<i} \vec{x}_j \vec{x}_j^*. \tag{12.21}$$

For normal $A = \sum_j \lambda_j \vec{x}_j \vec{x}_j^*$, this matrix is $\sum_{j\ge i} \lambda_j \vec{x}_j \vec{x}_j^*$. Its dominant eigenpair is $(\lambda_i, \vec{x}_i)$. So

$$\vec{q}_i^{(k)} \longrightarrow \pm\vec{x}_i, \qquad 1 \le i \le r, \tag{12.22}$$

at the rate $|\lambda_{i+1}/\lambda_i|$ once $\vec{q}_1, \ldots, \vec{q}_{i-1}$ have converged. Overall, the speed is limited by the slowest ratio $|\lambda_{j+1}/\lambda_j|$ with $j \le i$. For normal A, orthogonal iteration gives the eigenvectors directly.

The non-normal case. Now let $A = X\Lambda X^{-1}$ be diagonalizable, but not normal. Then the eigenvectors are no longer orthogonal. The columns of Q_k are orthogonal, so they cannot converge to the $\vec{x}_i$. To find their limit, we take a reduced QR factorization of the first r eigenvectors:

$$[\vec{x}_1 \cdots \vec{x}_r] =: Q^X R^X, \qquad Q^X = [\vec{q}_1^X \cdots \vec{q}_r^X] \in \mathbb{K}^{m\times r}. \tag{12.23}$$

Here $R^X \in \mathbb{K}^{r\times r}$ is upper triangular. The $\vec{x}_i$ are linearly independent, so R^X is invertible. As in Equation (13.6),

$$\operatorname{span}(\vec{q}_1^X, \ldots, \vec{q}_i^X) = \operatorname{span}(\vec{x}_1, \ldots, \vec{x}_i), \qquad i = 1, \ldots, r. \tag{12.24}$$

In other words, Q^X is the result of Gram–Schmidt applied to $\vec{x}_1, \ldots, \vec{x}_r$. For example, $\vec{q}_2^X$ is $(I - \vec{q}_1^X \vec{q}_1^{X*})\vec{x}_2$ normalized. This is exactly the limit of $\vec{q}_2^{(k)}$ that we found in Section 12.3.

Proposition 87 (Limit of the Orthogonal Iterates)

Let $A \in \mathbb{R}^{m\times m}$ be diagonalizable with $|\lambda_1| > \cdots > |\lambda_m|$, and assume that Equation (12.19) holds. Then $Q_k \to Q^X$ up to the signs of the columns. That is, $\vec{q}_i^{(k)} \mp \vec{q}_i^X \to \vec{0}$ for $i = 1, \ldots, r$, where the sign is chosen at each step k.

Proof sketch. We check one column at a time. We write $\vec{q}_i$ for the limit of $\vec{q}_i^{(k)}$ (up to sign). The first column $\vec{q}_1^{(k)}$ does power iteration, so $\vec{q}_1 = \pm\vec{x}_1 / \|\vec{x}_1\| = \pm\vec{q}_1^X$.

For the second column, Equation (12.19) gives

$$\operatorname{span}(\vec{x}_1, \vec{x}_2) = \operatorname{span}(\vec{q}_1, \vec{q}_2) = \operatorname{span}(\vec{q}_1^X, \vec{q}_2^X) =: S_2. \tag{12.25}$$

Now $\vec{q}_2$ is a unit vector in the orthogonal complement of $\vec{q}_1$ in S_2. Also, $\vec{q}_2^X$ is a unit vector in the orthogonal complement of $\vec{q}_1^X = \pm\vec{q}_1$ in S_2. Since S_2 is two-dimensional, this complement is one-dimensional. A real one-dimensional space contains exactly two unit vectors. So $\vec{q}_2 = \pm\vec{q}_2^X$.

For column i we repeat the argument with

$$\operatorname{span}(\vec{x}_1, \ldots, \vec{x}_i) = \operatorname{span}(\vec{q}_1, \ldots, \vec{q}_i) = \operatorname{span}(\vec{q}_1^X, \ldots, \vec{q}_i^X) =: S_i. \tag{12.26}$$

Both $\vec{q}_i$ and $\vec{q}_i^X$ are unit vectors in the orthogonal complement of $\operatorname{span}(\vec{q}_1, \ldots, \vec{q}_{i-1}) = \operatorname{span}(\vec{q}_1^X, \ldots, \vec{q}_{i-1}^X)$ in S_i, and this complement is one-dimensional. So $\vec{q}_i = \pm\vec{q}_i^X$ for all $i \le r$. □

Over $\mathbb{C}$, the same argument fixes each column only up to a factor $e^{\mathrm{i}\theta}$, instead of a sign. For normal A, the normalized eigenvectors are orthonormal. So Q^X consists of the normalized $\vec{x}_i$ themselves (up to sign), and Proposition 87 becomes $\vec{q}_i^{(k)} \to \pm\vec{x}_i$, as above.

12.6 Convergence to Schur Factorization

Now we take $r = m$, so that Q_k and Q^X are square and orthogonal. We look at the projected matrix

$$T_k := Q_k^* A Q_k \longrightarrow Q^{X*} A Q^X, \tag{12.27}$$

where the limit holds up to signs (see below). To compute the limit, we use Equation (12.23), now with $X = [\vec{x}_1 \cdots \vec{x}_m]$:

$$Q^X R^X = X \quad \Longleftrightarrow \quad Q^X = X(R^X)^{-1} \quad \Longleftrightarrow \quad Q^{X*} = (Q^X)^{-1} = R^X X^{-1}. \tag{12.28}$$

So

$$Q^{X*}AQ^X = R^X X^{-1} A X (R^X)^{-1} = R^X X^{-1} X \Lambda (R^X)^{-1} = \underbrace{R^X \Lambda (R^X)^{-1}}_{\text{upper triangular}}. \tag{12.29}$$

A product of upper triangular matrices is upper triangular, and its diagonal is $r_{ii}\lambda_i r_{ii}^{-1} = \lambda_i$. So $A = Q^X (R^X \Lambda (R^X)^{-1}) Q^{X*}$ is a Schur factorization of A (Theorem 78). The eigenvalues are on the diagonal, in decreasing order of modulus. The eigenpairs of the triangular factor are

$$(\lambda_i, R^X \vec{e}_i), \qquad \text{since} \quad R^X \Lambda (R^X)^{-1} R^X \vec{e}_i = R^X \Lambda \vec{e}_i = \lambda_i R^X \vec{e}_i. \tag{12.30}$$

If we lift them back, we get the eigenvectors of A: $Q^X R^X \vec{e}_i = X \vec{e}_i = \vec{x}_i$. (Note: the eigenvector is $R^X \vec{e}_i$, the ith column of R^X, and not $(R^X)^{-1}\vec{e}_i$. Only the first one is mapped to $\vec{x}_i$ by Q^X.)

The signs do not change this picture. If $Q_k \approx Q^X D$ with $D = \operatorname{diag}(\pm 1)$, then $T_k \approx D\, R^X \Lambda (R^X)^{-1} D$. This is still upper triangular with the same diagonal. Only some entries above the diagonal change sign.

In summary: for every k, $A = Q_k T_k Q_k^*$ holds exactly, with Q_k orthogonal. And the factor $T_k = Q_k^* A Q_k$ converges to an upper triangular matrix. So

$$Q_k, \quad Q_k^* A Q_k, \quad Q_k^* \tag{12.31}$$

converge to the Schur factorization of A. We can read off the eigenvalues from the diagonal of $Q_k^* A Q_k$. In Theorem 89 we give the exact conditions on the starting matrix, and a proof that does not assume Equation (12.19).

To compute T_{k+1} this way, we need Q_k and a matrix-matrix product with A in every step. In Section 13.5 we derive an iteration $T_k \to T_{k+1}$ that uses only T_k: the QR algorithm. In Lecture 14 we make each of its steps cheap, by first reducing A to Hessenberg form.

12.7 (SUPPLEMENT) Further Topics

12.7.1 More Consequences of Normality

Let $A = Q\Lambda Q^*$ be normal, and write $\lambda_i = |\lambda_i|\, e^{\mathrm{i}\theta_i}$. Then $A = (Q \operatorname{diag}(e^{\mathrm{i}\theta_i}))\, |\Lambda|\, Q^*$ is an SVD, up to the ordering of the entries. So the singular values of a normal matrix are the moduli of its eigenvalues [1, Lecture 24]. In particular, $\|A\|_2 = \max_i |\lambda_i|$ and $\left\|A^k\right\|_2 = \max_i |\lambda_i|^k$. So powers of a normal matrix show no transient (temporary) growth, unlike powers of a Jordan block (Section 11.6).

The Schur form also measures how far a general A is from being normal. Write the triangular factor as $T = \Lambda + N$, with N strictly upper triangular. The Frobenius norm is unitarily invariant, so

$$\|A\|_F^2 = \|T\|_F^2 = \sum_{i=1}^{m} |\lambda_i|^2 + \|N\|_F^2 . \tag{12.32}$$

So $\sum_i |\lambda_i|^2 \le \|A\|_F^2$, with equality if and only if $N = 0$, i.e., if and only if A is normal (Theorem 84). The quantity $\|N\|_F$ is called the *departure from normality*. It does not depend on which Schur factorization we choose, since $\|A\|_F$ and the eigenvalues do not.

12.7.2 The Rayleigh–Ritz Procedure

Take any $Q \in \mathbb{K}^{m \times r}$ with orthonormal columns, and form the projected matrix $H := Q^* A Q$, as in Proposition 85. This matrix generalizes the Rayleigh quotient from one vector to a subspace. Its eigenvalues θ_i are called *Ritz values*, and the lifted eigenvectors $Q\vec{v}_i$ are called *Ritz vectors*.

Among all $R \in \mathbb{K}^{r\times r}$, the choice $R = H$ gives the smallest residual $\|AQ - QR\|_2$. To see this, let $E := AQ - QH$. Then $Q^*E = 0$, so for every $\vec{y}$,

$$\|(AQ - QR)\vec{y}\|^2 = \|E\vec{y} + Q(H - R)\vec{y}\|^2 = \|E\vec{y}\|^2 + \|(H - R)\vec{y}\|^2 \geq \|E\vec{y}\|^2 . \tag{12.33}$$

For Hermitian A, there are r eigenvalues of A, each within $\|E\|_2$ of a different Ritz value. These are not necessarily the largest eigenvalues [2, §7.2].

So in subspace iteration for Hermitian matrices, we often replace Q_k by its Ritz vectors Q_kV, where $Q_k^*AQ_k = V\Theta V^*$. The Ritz vectors depend only on $\operatorname{span}(Q_k) = \operatorname{span}(A^kQ_0)$. In this subspace, the components along $\vec{x}_j$, $j > r$, are smaller than the component along $\vec{x}_i$ by roughly a factor $|\lambda_j/\lambda_i|^k \leq |\lambda_{r+1}/\lambda_i|^k$. So the ith Ritz vector converges like $|\lambda_{r+1}/\lambda_i|^k$, not like $|\lambda_{i+1}/\lambda_i|^k$. This means that iterating with a few more columns than the number of eigenpairs we want makes convergence faster. Without the Rayleigh–Ritz step, extra columns do not help the first columns, since the leading columns of Algorithm 19 never see the later ones.

12.7.3 Convergence of the Individual Entries

The leading p columns span a subspace that converges at the rate $|\lambda_{p+1}/\lambda_p|$. So the block $(T_k)_{p+1:m,\,1:p}$ decays like $|\lambda_{p+1}/\lambda_p|^k$. The diagonal entry $(T_k)_{pp}$ depends on the blocks on both sides of it. It converges to λ_p like

$$\max\left(\left|\frac{\lambda_{p+1}}{\lambda_p}\right|, \left|\frac{\lambda_p}{\lambda_{p-1}}\right|\right)^k . \tag{12.34}$$

For $p = 1$ only the first ratio is present, and for $p = m$ only the second [2, §4.4.3]. So a pair of eigenvalues with nearly equal modulus slows down only the part of T_k near their position. The rest of the Schur form still converges quickly. For symmetric A, all columns of Q_k converge to eigenvectors linearly, with constant $\max_{1\leq j\leq r} |\lambda_{j+1}/\lambda_j|$ [1, Thm. 28.1].

12.7.4 Why Orthonormalize at Every Step

Mathematically, $\operatorname{span}(Q_k) = \operatorname{span}(A^kQ_0)$. So we could compute $V_k = A^kQ_0$ and orthonormalize only once at the end ("unnormalized simultaneous iteration"). In floating point this fails. Every column of A^kQ_0 converges to a multiple of $\vec{x}_1$, so the columns become nearly parallel. The information about $\vec{x}_2, \ldots, \vec{x}_r$ is in components that are smaller by factors $|\lambda_i/\lambda_1|^k$, and rounding errors destroy it [1, Lecture 28]. If we orthonormalize after every multiplication, we keep a well-conditioned basis of the same subspace.

With $Q_0 = I$ and $r = m$, the product of all the triangular factors gives the QR factorization of A^k itself. This identity is the starting point of the QR algorithm in Lecture 13.

12.7.5 Beyond Subspace Iteration

Each step of orthogonal iteration costs one product AQ and one QR factorization. The product costs about $2m^2r$ flops for dense A, and far less for sparse A. The QR factorization costs $O(mr^2)$. So orthogonal iteration is a good choice when we need only a few dominant eigenpairs of a large sparse matrix. But its convergence is still linear, with the ratio $|\lambda_{r+1}/\lambda_r|$.

Krylov methods do better with the same products. They do not keep only A^kQ_0. Instead, they apply Rayleigh–Ritz to the whole Krylov subspace $\operatorname{span}(\vec{q}, A\vec{q}, \ldots, A^{k-1}\vec{q})$, which contains all earlier power iterates. This leads to the Arnoldi iteration and, for Hermitian A, the Lanczos iteration [1, Lectures 33, 34, and 36], [2, Chapter 7].

Lecture 13

Convergence of Orthogonal Iteration and the Basic QR Iteration


Supplementary reading:

- [1] Lecture 28.
- [2] Sections 4.4.3 and 4.4.4.

13.1 Recap

Last lecture (Lecture 12) we found the matrices whose Schur factorization is a diagonalization. We also began to compute several eigenpairs at once.

- *From Schur factorization to diagonalization* (Section 12.2). A is normal, $A^*A = AA^*$, if and only if $A = Q\Lambda Q^*$ with Q unitary (Theorem 84). Take the Schur factorization $A = QTQ^*$. If A is normal, then $T^*T = TT^*$. Comparing the $(1,1)$ entries gives $|t_{11}|^2 = \|T_{1,:}\|^2$. So the first row of T is zero off the diagonal. By induction, T is diagonal. In terms of shapes, $T^*T = TT^*$ says that lower times upper equals upper times lower:



  The $(1,1)$ entry on the left sees only the first column of T. The $(1,1)$ entry on the right sees the whole first row.

- *Orthogonal iteration for two vectors* (Section 12.3). We run power iteration on $\vec{q}_1, \vec{q}_2$. After each step, we project $\vec{q}_1$ out of $\vec{q}_2$ (Algorithm 18). Then $\vec{q}_1 \to \vec{x}_1$. Once $\vec{q}_1$ has converged, we have, up to normalization,

$$\vec{q}_2^{(k)} = (I - \vec{x}_1\vec{x}_1^*)A(I - \vec{x}_1\vec{x}_1^*)\vec{q}_2^{(k-1)}. \tag{13.1}$$

  This matrix has the eigenpairs $(\lambda_i, (I - \vec{x}_1\vec{x}_1^*)\vec{x}_i)$ for $i \geq 2$:

$$(I - \vec{x}_1\vec{x}_1^*)A(I - \vec{x}_1\vec{x}_1^*)\vec{x}_i = \lambda_i(I - \vec{x}_1\vec{x}_1^*)\vec{x}_i. \tag{13.2}$$

  So $\vec{q}_2^{(k)} \to (I - \vec{x}_1\vec{x}_1^*)\vec{x}_2$. This is different from $\vec{x}_2$, unless $\vec{x}_1 \perp \vec{x}_2$ (as for normal A). What converges is the subspace: $\operatorname{span}(\vec{q}_1, \vec{q}_2) \to \operatorname{span}(\vec{x}_1, \vec{x}_2)$.

- *Orthogonal iteration for a general number of vectors* (Section 12.4). With r columns, the projection step becomes a reduced QR factorization (Algorithm 19):

```
for k = 1:iterations
    Q = A*Q
    Q, R = qr(Q)
end
```

13.2 Schur and Eigendecomposition

In this lecture we always assume

$$A = X\Lambda X^{-1}, \qquad |\lambda_1| > |\lambda_2| > \cdots > |\lambda_m|\,. \tag{13.3}$$

In words: A is diagonalizable, and its eigenvalues have distinct moduli.

This assumption is *generic*. It holds for all matrices except a small set, and an arbitrarily small perturbation moves a matrix out of this set. Here is why. Every A has a Schur factorization $A = QTQ^*$, with the eigenvalues on the diagonal of T (Theorem 78). We add a small perturbation to the diagonal of T, and so also to A:

$$\tilde{A} \coloneqq Q(T + \operatorname{diag}(\varepsilon_1, \ldots, \varepsilon_m))Q^* = A + Q\operatorname{diag}(\varepsilon_1, \ldots, \varepsilon_m)Q^*. \tag{13.4}$$

We can choose the ε_i so that $\tilde{A}$ has eigenvalues of distinct magnitude:

- $\tilde{A} = Q(T + \operatorname{diag}(\varepsilon_i))Q^*$ is a Schur factorization of $\tilde{A}$, with diagonal entries $t_{ii} + \varepsilon_i$. So these are its eigenvalues, and we can choose the ε_i to make their moduli distinct.
- The perturbation is small: $\left\|\tilde{A} - A\right\| = \max_i |\varepsilon_i|$, since Q is unitary.
- Distinct eigenvalues imply diagonalizability. So $\tilde{A}$ satisfies Equation (13.3).

Also, eigenvalues depend continuously on the matrix. So the strict inequalities still hold after small perturbations of $\tilde{A}$.

Remark 88 (Real Matrices (SUPPLEMENT))**.** The perturbation above is complex in general. For real A, non-real eigenvalues come in conjugate pairs $\lambda, \overline{\lambda}$ with $|\lambda| = \left|\overline{\lambda}\right|$. A small real perturbation keeps them a conjugate pair. So among real matrices, Equation (13.3) is not generic. It fails on an open set, for example near every rotation of $\mathbb{R}^2$ by an angle $\theta \notin \{0, \pi\}$.

Now, given $A = X\Lambda X^{-1}$, how do we compute its Schur form? We take a QR factorization of the eigenvector matrix, $X = QR$. Since X is invertible, R is invertible too, and

$$X = QR \quad \Longrightarrow \quad A = QR\Lambda R^{-1}Q^* = Q\underbrace{(R\Lambda R^{-1})}_{\text{upper triangular}}Q^*. \tag{13.5}$$

Why is $R\Lambda R^{-1}$ upper triangular? R^{-1} is upper triangular, and products of upper triangular matrices are upper triangular. Its diagonal is $r_{ii}\lambda_i r_{ii}^{-1} = \lambda_i$. So Equation (13.5) is a Schur factorization of A. The eigenvalues are on the diagonal of $T = R\Lambda R^{-1}$, in the same order as in Λ.

Since R is upper triangular, $\vec{x}_j = \sum_{i \le j} r_{ij}\vec{q}_i$ only uses the first j columns of Q. Since R is invertible, the other direction also holds. So

$$\operatorname{span}(\vec{q}_1, \ldots, \vec{q}_j) = \operatorname{span}(\vec{x}_1, \ldots, \vec{x}_j), \qquad j = 1, \ldots, m. \tag{13.6}$$

In words: the Schur vectors span the same nested invariant subspaces as the eigenvectors. This is not an algorithm, since it needs X. But it shows us the target. In the rest of this chapter we show that orthogonal iteration converges to these subspaces without knowing X.

13.3 Convergence of Orthogonal Iteration

How should we think about the convergence of orthogonal iteration (Algorithm 19)? Let $Q_k \in \mathbb{K}^{m\times p}$ be the iterate after k steps. We start from Q_0 with p orthonormal columns. Let $A = X\Lambda X^{-1}$ be diagonalizable, with its eigenvalues ordered by modulus, $|\lambda_1| \geq |\lambda_2| \geq \cdots \geq |\lambda_m|$. In this section we need less than Equation (13.3). It is enough that there is a gap after the pth eigenvalue:

$$|\lambda_p| > |\lambda_{p+1}|\,. \tag{13.7}$$

As in Lecture 12, we track the subspace $\operatorname{span}(Q_k)$ (the span of the columns), not the individual columns.

The QR step does not change the column space, so $\operatorname{span}(Q_{k+1}) = \operatorname{span}(AQ_k)$. Unrolling the recursion gives

$$\operatorname{span}(Q_{k+1}) = \operatorname{span}(AQ_k) = \cdots = \operatorname{span}(A^{k+1}Q_0) = \operatorname{span}(X\Lambda^{k+1}X^{-1}Q_0). \tag{13.8}$$

We pull out the factor λ_p^{k+1}. This does not change the span. (Note that $\lambda_p \neq 0$, since $|\lambda_p| > |\lambda_{p+1}| \geq 0$ by Equation (13.7).) We get

$$\operatorname{span}(Q_{k+1}) = \operatorname{span}\left(\lambda_p^{k+1} X \begin{bmatrix} (\lambda_1/\lambda_p)^{k+1} & & & & \\ & \ddots & & & \\ & & 1 & & \\ & & & \ddots & \\ & & & & (\lambda_m/\lambda_p)^{k+1} \end{bmatrix} X^{-1}Q_0\right), \tag{13.9}$$

with the 1 in position (p,p). By the ordering of the eigenvalues and the gap Equation (13.7),

$$|\lambda_i/\lambda_p| \geq 1 \quad \text{for } i \leq p, \qquad |\lambda_i/\lambda_p| < 1 \quad \text{for } i > p. \tag{13.10}$$

So the first p rows of $(\Lambda/\lambda_p)^{k+1}X^{-1}Q_0$ do not decay. The last $m-p$ rows are scaled by factors that tend to zero. We split the matrix in the same way:

$$(\Lambda/\lambda_p)^{k+1}\, X^{-1}Q_0 = \begin{bmatrix} V_k \\ W_k \end{bmatrix}, \qquad V_k \in \mathbb{K}^{p\times p}, \quad W_k \in \mathbb{K}^{(m-p)\times p}, \qquad W_k \xrightarrow{k\to\infty} 0. \tag{13.11}$$

Here W_k decays like $|\lambda_{p+1}/\lambda_p|^{k+1}$. We split the columns of X in the same way: $X = [X_p\ \tilde{X}_p]$, where $X_p = [\vec{x}_1 \cdots \vec{x}_p]$ holds the first p eigenvectors. This gives

$$\operatorname{span}(Q_{k+1}) = \operatorname{span}(X\Lambda^{k+1}X^{-1}Q_0) = \operatorname{span}(X_pV_k + \tilde{X}_pW_k) \xrightarrow{k\to\infty} \operatorname{span}(X_p). \tag{13.12}$$

So $\operatorname{span}(Q_k)$ converges to $\operatorname{span}(\vec{x}_1,\ldots,\vec{x}_p)$, the invariant subspace of the p dominant eigenvectors.

To make the limit precise (SUPPLEMENT), we write $B \coloneqq X^{-1}Q_0 = \left[\begin{smallmatrix} B_1 \\ B_2 \end{smallmatrix}\right]$ for the coordinates of Q_0 in the eigenvector basis, with $B_1 \in \mathbb{K}^{p\times p}$. We also write $\Lambda = \operatorname{diag}(\Lambda_1, \Lambda_2)$ with $\Lambda_1 = \operatorname{diag}(\lambda_1,\ldots,\lambda_p)$. Then

$$V_k = (\Lambda_1/\lambda_p)^{k+1}B_1, \qquad W_k = (\Lambda_2/\lambda_p)^{k+1}B_2. \tag{13.13}$$

Assume B_1 is invertible. This means that Q_0 has a nonzero component in every direction $\vec{x}_1, \dots, \vec{x}_p$. It is the analogue of $\vec{x}_1^* \vec{q}^{(0)} \neq 0$ for power iteration, and it holds with probability one for random Q_0. Then V_k is invertible. Multiplying by V_k^{-1} on the right does not change the span:

$$\operatorname{span}(X_p V_k + \tilde{X}_p W_k) = \operatorname{span}(X_p + \tilde{X}_p W_k V_k^{-1}). \tag{13.14}$$

We have $V_k^{-1} = B_1^{-1}(\lambda_p \Lambda_1^{-1})^{k+1}$ and $|\lambda_p/\lambda_i| \leq 1$ for $i \leq p$. So $\left\|V_k^{-1}\right\| \leq \left\|B_1^{-1}\right\|$, and

$$\left\|\tilde{X}_p W_k V_k^{-1}\right\| \leq \left\|\tilde{X}_p\right\| \|B_2\| \left\|B_1^{-1}\right\| \left|\frac{\lambda_{p+1}}{\lambda_p}\right|^{k+1} \longrightarrow 0. \tag{13.15}$$

So the subspace converges at the rate $|\lambda_{p+1}/\lambda_p|$. This rate is the gap between the last eigenvalue we keep and the first one we drop.

13.4 Convergence to Schur Factorization

Now we run orthogonal iteration with $p = m$, that is, on a full orthonormal basis. We start from $Q_0 = I$.

Theorem 89 (Orthogonal Iteration Converges to Schur Form)

Consider orthogonal iteration (Algorithm 19) on $A = X\Lambda X^{-1}$ with $p = m$ and $Q_0 = I$. Assume that all eigenvalues of A have distinct absolute values, $|\lambda_1| > \cdots > |\lambda_m|$, and that the leading blocks $(X^{-1})_{1:j,1:j}$ have full rank for $j = 1, \dots, m$. Then

$$T_k \coloneqq Q_k^* A Q_k \tag{13.16}$$

converges to the Schur form of A: its strictly lower triangular part tends to zero, and its diagonal tends to $(\lambda_1, \dots, \lambda_m)$.

Note that the condition is on the leading blocks of X^{-1}, not of X. The two conditions are not equivalent. Some books state the condition with $X_{1:j,1:j}$ (e.g., [2, Thm. 4.8]).

Why X^{-1}? The argument of Section 13.3 needs a condition on $X^{-1}Q_0 = X^{-1}$. For j columns, the block B_1 consists of the first j rows and columns, $(X^{-1})_{1:j,1:j}$. If we only ask that $X_{1:j,1:j}$ have full rank for all j, this is not enough. For example, take

$$X = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}, \qquad X^{-1} = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix}. \tag{13.17}$$

Here $X_{1:1,1:1} = 1$ and X are nonsingular, but $(X^{-1})_{11} = 0$. The starting vector $\vec{e}_1 = \vec{x}_2$ is itself an eigenvector. It has no component along $\vec{x}_1$, and in exact arithmetic $\vec{q}_1^{(k)} = \vec{x}_2$ forever.

We can state the correct condition in two other ways.

- By Jacobi's identity for complementary minors, $\det(X^{-1})_{1:j,1:j} = \det X_{j+1:m,j+1:m} / \det X$. So the condition says: the trailing blocks $X_{j+1:m,j+1:m}$ are nonsingular.
- Equivalently, X^{-1} has an LU factorization without pivoting.

For normal $A = Q\Lambda Q^*$ the difference disappears. Then $X = Q$ is unitary, $X^{-1} = Q^*$, and $(Q^*)_{1:j,1:j} = (Q_{1:j,1:j})^*$. So the leading blocks of X^{-1} are nonsingular exactly when those of X are. Trefethen and

Bau treat real symmetric $A = Q\Lambda Q^\top$. This is why they can state the condition as "all leading principal submatrices of Q are nonsingular" [1, Thm. 28.4]. For non-normal A, the example above shows that only the condition on X^{-1} is correct.

Proof sketch. Fix $p \in \{1, \dots, m-1\}$. We split the columns of Q_k after the pth column:

$$Q_k = [\bar{Q}_k \, \tilde{Q}_k], \qquad \bar{Q}_k \in \mathbb{K}^{m \times p}. \tag{13.18}$$

(Here p is only the place where we split. Q_k still has all m columns.) Then

$$T_k = Q_k^* A Q_k = \begin{bmatrix} \bar{Q}_k^* A \bar{Q}_k & \bar{Q}_k^* A \tilde{Q}_k \\ \tilde{Q}_k^* A \bar{Q}_k & \tilde{Q}_k^* A \tilde{Q}_k \end{bmatrix}. \tag{13.19}$$

Column j of a QR factorization depends only on the first j columns of the factored matrix. So $\bar{Q}_k$ is exactly the iterate of orthogonal iteration with p columns, started from $\bar{Q}_0 = I_{:,1:p}$. By Section 13.3,

$$\operatorname{span}(\bar{Q}_k) \longrightarrow \operatorname{span}(\vec{x}_1, \dots, \vec{x}_p). \tag{13.20}$$

We apply A:

$$\operatorname{span}(A\bar{Q}_k) \longrightarrow \operatorname{span}(A[\vec{x}_1 \cdots \vec{x}_p]) = \operatorname{span}([\vec{x}_1 \cdots \vec{x}_p]). \tag{13.21}$$

The equality holds because $A[\vec{x}_1 \cdots \vec{x}_p] = [\vec{x}_1 \cdots \vec{x}_p] \operatorname{diag}(\lambda_1, \dots, \lambda_p)$ and $\lambda_1, \dots, \lambda_p \neq 0$ (as $|\lambda_p| > |\lambda_{p+1}| \geq 0$). In words: the span of the dominant eigenvectors is invariant under A. So the columns of $A\bar{Q}_k$ lie (asymptotically) in $\operatorname{span}(\bar{Q}_k)$. The columns of $\tilde{Q}_k$ are orthogonal to $\operatorname{span}(\bar{Q}_k)$. Hence

$$\tilde{Q}_k^* A \bar{Q}_k \longrightarrow 0. \tag{13.22}$$

More precisely, $\tilde{Q}_k^* \bar{Q}_k = 0$ gives

$$\tilde{Q}_k^* A \bar{Q}_k = \tilde{Q}_k^* (I - \bar{Q}_k \bar{Q}_k^*) A \bar{Q}_k.$$

Here $(I - \bar{Q}_k \bar{Q}_k^*) A \bar{Q}_k$ is the part of $A\bar{Q}_k$ outside $\operatorname{span}(\bar{Q}_k)$, and it tends to zero.

The lower left block of the split after column p contains the entries (i, j) with $i > p \geq j$. Every strictly lower triangular entry (i, j), $i > j$, lies in this block for $p = j$. So the argument covers all subdiagonal blocks. Letting p range over $1, \dots, m-1$ shows that

$$T_k = Q_k^* A Q_k \longrightarrow \text{upper triangular.} \tag{13.23}$$

For the diagonal (SUPPLEMENT): the block $\bar{Q}_k^* A \bar{Q}_k$ is the matrix of A restricted to $\operatorname{span}(\bar{Q}_k)$, and $\operatorname{span}(\bar{Q}_k) \to \operatorname{span}(\vec{x}_1, \dots, \vec{x}_p)$. So its eigenvalues tend to $\lambda_1, \dots, \lambda_p$, as in Proposition 85. This block becomes upper triangular, so its diagonal holds these eigenvalues. Comparing p with $p-1$ shows that the pth diagonal entry of T_k tends to λ_p. □

13.5 Basic QR Iteration

With orthogonal iteration we can compute

$$T_k \coloneqq Q_k^* A Q_k. \tag{13.24}$$

By Theorem 89, T_k converges to the triangular factor of a Schur factorization.

To compute T_{k+1} this way, we need Q_k and a matrix-matrix product with A. Can we instead find an iteration $T_k \to T_{k+1}$ that uses only the T's? We have

$$T_k = Q_k^* A Q_k, \qquad T_{k+1} = Q_{k+1}^* A Q_{k+1}. \tag{13.25}$$

One step of orthogonal iteration is the QR factorization $Q_{k+1}R_{k+1} = AQ_k$. We multiply by Q_{k+1}^* on the left and by Q_k^* on the right. (Here $p = m$, so Q_k is square and $Q_k Q_k^* = I$.) This gives

$$Q_{k+1}R_{k+1} = AQ_k \quad \Longrightarrow \quad R_{k+1}Q_k^* = Q_{k+1}^* A. \tag{13.26}$$

We put the first form into T_k and the second into T_{k+1}:

$$T_k = Q_k^*(Q_{k+1}R_{k+1}), \qquad T_{k+1} = (R_{k+1}Q_k^*)Q_{k+1}. \tag{13.27}$$

Both expressions contain the same product $Q_k^* Q_{k+1}$. We give it a name:

$$U_{k+1} \coloneqq Q_k^* Q_{k+1} : \qquad T_k = U_{k+1}R_{k+1}, \qquad T_{k+1} = R_{k+1}U_{k+1}. \tag{13.28}$$

Here U_{k+1} is unitary, since $U_{k+1}^* U_{k+1} = Q_{k+1}^* Q_k Q_k^* Q_{k+1} = I$, and R_{k+1} is upper triangular. So $T_k = U_{k+1}R_{k+1}$ is a QR factorization of T_k. We get T_{k+1} by multiplying its factors in reverse order:

$$T_k = U_{k+1}R_{k+1} \quad \longrightarrow \quad T_{k+1} = R_{k+1}U_{k+1}. \tag{13.29}$$

In short: factor as QR, multiply back as RQ. Neither A nor Q_k appears. Also, T_{k+1} is unitarily similar to T_k:

$$T_{k+1} = R_{k+1}U_{k+1} = U_{k+1}^*(U_{k+1}R_{k+1})U_{k+1} = U_{k+1}^* T_k U_{k+1}.$$

So every T_k has the eigenvalues of A.

We start orthogonal iteration from $Q_0 = I$. This gives $T_0 = Q_0^* A Q_0 = A$, so the iteration starts from A itself.

Algorithm 20 Basic QR iteration (QR algorithm without shifts).

Input: $A \in \mathbb{K}^{m \times m}$

1: $T \leftarrow A$
2: **for** $k = 1, \ldots,$ maxiter **do**
3: $\quad UR \leftarrow T$ ▷ QR factorization
4: $\quad T \leftarrow RU$
5: **return** $\operatorname{diag}(T)$ ▷ approximate eigenvalues $\lambda_1, \ldots, \lambda_m$

By Theorem 89, T converges to upper triangular form. The eigenvalues are on its diagonal, ordered by decreasing modulus.

A library QR routine may not return the same factors as orthogonal iteration. For invertible T, QR factors are unique only up to signs (phases) of the columns of U and the rows of R (Lecture 7). Every choice gives $T_{k+1} = U^* T_k U$. Different choices give iterates that differ by a diagonal unitary similarity $D^* T_k D$. This does not change the diagonal or the moduli of the entries.

How do we get Q? Algorithm 20 returns only the eigenvalues. The Schur vectors Q_k of orthogonal iteration never appear. We can recover them from the factors U, since by definition

$$U_{k+1} = Q_k^* Q_{k+1}. \tag{13.30}$$

The product of the U's telescopes. Since $Q_k Q_k^* = I$ and $Q_0 = I$,

$$U_1 U_2 \cdots U_{k+1} = (Q_0^* Q_1)(Q_1^* Q_2) \cdots (Q_k^* Q_{k+1}) = Q_0^* Q_{k+1} = Q_{k+1}. \tag{13.31}$$

So to get the Schur vectors, we start from $Q \leftarrow I$ and update $Q \leftarrow QU$ after each QR factorization in Algorithm 20. At the end, $A \approx QTQ^*$.

The factors U_k have a simple meaning.

- Near convergence, $Q_{k+1} \approx Q_k$ up to the phases of the columns. So $U_{k+1} = Q_k^* Q_{k+1}$ is a small rotation: it is close to a diagonal unitary matrix D.
- We can also see this from $T_k = U_{k+1} R_{k+1}$. Once T_k is nearly upper triangular, $U_{k+1} = T_k R_{k+1}^{-1}$ is nearly upper triangular. It is also unitary, so it is nearly diagonal.
- With the normalization $r_{ii} > 0$ we get $D \approx \operatorname{diag}(\lambda_1/|\lambda_1|, \ldots, \lambda_m/|\lambda_m|)$. So $D = I$ only when all eigenvalues are positive. A negative eigenvalue flips the sign of its column of Q_k in every step, exactly as in power iteration.

Up to these phases, the U_k are a sequence of rotation corrections. Together they turn $Q_0 = I$ into the Schur vectors Q. The corrections are large in the first steps, and they get smaller and smaller as T_k approaches triangular form.

Remark 90 (Julia)**.** For square T, we can multiply directly by the implicit Q returned by `qr` (Lecture 7). So the pitfall of Lecture 12 does not come up:

```
T = copy(A); Q = Matrix(1.0I, m, m)
for k = 1:maxiter
    F = qr(T)
    T = F.R * F.Q
    Q = Q * F.Q
end
```

13.6 (SUPPLEMENT) Further Topics

13.6.1 The QR Algorithm Factors the Powers of A

Let $\underline{R}_k := R_k R_{k-1} \cdots R_1$. This is the product of the triangular factors of orthogonal iteration started from $Q_0 = I$. Then

$$A^k = Q_k \underline{R}_k, \qquad T_k = Q_k^* A Q_k, \qquad Q_k = U_1 U_2 \cdots U_k. \tag{13.32}$$

The first identity follows by induction:

$$A^k = A\, Q_{k-1} \underline{R}_{k-1} = Q_k R_k \underline{R}_{k-1} = Q_k \underline{R}_k.$$

Also, $\underline{R}_k$ is upper triangular, as a product of upper triangular matrices. So the QR algorithm computes a QR factorization of A^k without ever forming A^k. (The columns of A^k would all be nearly parallel to $\vec{x}_1$.) The two identities in Equation (13.32) explain the whole algorithm [1, Lecture 28].

- The first says that the leading j columns of Q_k are an orthonormal basis of the span of the leading j columns of A^k. This is simultaneous power iteration on $\vec{e}_1, \ldots, \vec{e}_m$.
- The second says that the diagonal entries $(T_k)_{jj} = \vec{q}_j^{(k)*} A \vec{q}_j^{(k)}$ are Rayleigh quotients. So for Hermitian A, they converge twice as fast as the columns of Q_k (Lecture 10).

13.6.2 Simultaneous Inverse Iteration

Let A be invertible. We invert $A^k = Q_k \underline{R}_k$ and take the adjoint:

$$(A^*)^{-k} = (\underline{R}_k^{-1} Q_k^*)^* = Q_k \underline{R}_k^{-*}. \tag{13.33}$$

Here $\underline{R}_k^{-*}$ is lower triangular. Let P be the permutation that reverses the order of the rows, $P^2 = I$. If we reverse both the rows and the columns of a lower triangular matrix, it becomes upper triangular. So

$$(A^*)^{-k} P = (Q_k P)(P \underline{R}_k^{-*} P) \tag{13.34}$$

is a QR factorization of $(A^*)^{-k} P$. Hence $Q_k P$ (that is, Q_k with its columns in reverse order) is what orthogonal iteration with $(A^*)^{-1}$ produces from $Q_0 = P$.

In particular, the last column of Q_k is the result of k steps of inverse iteration (Algorithm 16) with A^*, started from $\vec{e}_m$. It converges at the rate $|\lambda_m / \lambda_{m-1}|$ to an eigenvector $\vec{y}_m$ of A^* for $\overline{\lambda_m}$. This $\vec{y}_m$ is a left eigenvector of A, $\vec{y}_m^* A = \lambda_m \vec{y}_m^*$. It is orthogonal to $\vec{x}_1, \ldots, \vec{x}_{m-1}$, which agrees with Equation (13.6). So the QR algorithm runs two methods at the same time: power iteration in its first column, and inverse iteration in its last column. For real symmetric A, $A^* = A$; see [1, Lecture 29].

13.6.3 Convergence Rates of Individual Entries

By Section 13.3, the block $(T_k)_{p+1:m,1:p}$ decays like $|\lambda_{p+1}/\lambda_p|^k$. This is the block below and to the left of the split after column p. In particular:

- the subdiagonal entry $(T_k)_{j+1,j}$ decays like $|\lambda_{j+1}/\lambda_j|^k$;
- the diagonal entry $(T_k)_{pp}$ converges to λ_p like $\max(|\lambda_{p+1}/\lambda_p|^k, |\lambda_p/\lambda_{p-1}|^k)$ [2, §4.4.3].

The overall rate is the worst ratio $\max_j |\lambda_{j+1}/\lambda_j|$ [1, Thm. 28.4]. So if two eigenvalues have nearly equal modulus, the unshifted algorithm becomes arbitrarily slow.

For example, take $\Lambda = \mathrm{diag}(30, 6, 2, 1)$ and a random X with $\kappa(X) \approx 20$. After 19 steps, the subdiagonal entries are about $7 \cdot 10^{-13}$, $1 \cdot 10^{-9}$, and $5 \cdot 10^{-6}$ [2, Example 4.5]. This agrees with $(1/5)^{19} \approx 5 \cdot 10^{-14}$, $(1/3)^{19} \approx 9 \cdot 10^{-10}$, and $(1/2)^{19} \approx 2 \cdot 10^{-6}$, up to constants.

The most important block is the last row, $(T_k)_{m,1:m-1}$. It decays like $|\lambda_m/\lambda_{m-1}|^k$. Once it is below a tolerance such as $\varepsilon_{\text{mach}} \|A\|$, we accept $(T_k)_{mm}$ as an eigenvalue. Then we continue on the leading $(m-1) \times (m-1)$ block. This is called deflation.

13.6.4 Eigenvalues of Equal Modulus

If there is no gap $|\lambda_p| > |\lambda_{p+1}|$, the block at the split after column p has no reason to decay. The extreme case is a unitary A: all its eigenvalues have modulus 1. Its QR factorization is $A = A \cdot I$, so $T_1 = I \cdot A = A$, and the iteration does not move at all [2, Example 4.6]. An example is $A = \left[\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right]$, with eigenvalues ± 1.

For real matrices, the generic case is a pair of complex conjugate eigenvalues $\lambda, \overline{\lambda}$ (see the remark in Section 13.2). For real A, all U_k, R_k, and T_k are real. So T_k cannot converge to a triangular matrix with non-real diagonal entries. But the splits between eigenvalues of different moduli still have a gap, so those blocks still decay (under the analogous rank conditions). Suppose the only ties in modulus come from conjugate pairs. Then T_k converges to a block upper triangular matrix with 1×1 and 2×2 diagonal blocks, that is, to the real Schur form (Theorem 80). The entries of each 2×2 block keep changing from step to step, but its eigenvalues converge to $\lambda, \overline{\lambda}$.

13.6.5 Why Shifts Help

The shifted matrix $A - \sigma I = Q(T - \sigma I)Q^*$ has the same Schur vectors as A, and its eigenvalues are $\lambda_i - \sigma$. So the QR algorithm on $A - \sigma I$ converges at the rates $|\lambda_{j+1} - \sigma| \,/\, |\lambda_j - \sigma|$. Now the eigenvalues are ordered by their distance from σ. If σ is close to an eigenvalue λ, the last row decays at the rate $|\lambda - \sigma| \,/\, \min_{\lambda_j \neq \lambda} |\lambda_j - \sigma|$, which is nearly zero.

The shift may change from step to step, and we do not need to undo it:

$$T_k - \sigma_k I = U_{k+1} R_{k+1}, \qquad T_{k+1} = R_{k+1} U_{k+1} + \sigma_k I = U_{k+1}^* T_k U_{k+1}. \tag{13.35}$$

So T_{k+1} is still unitarily similar to A [2, §4.4.4]. By the previous subsections, the last column of Q_k then does shifted inverse iteration. The natural choice is $\sigma_k = (T_k)_{mm}$, the Rayleigh quotient of that column. This choice gives a version of Rayleigh quotient iteration (Algorithm 17), with cubic convergence for real symmetric A [1, Lecture 29].

Shifts also break ties in modulus. For $A = \left[\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right]$ and real $\sigma \neq 0$, we have $|1 - \sigma| \neq |-1 - \sigma|$. But here $(T_k)_{mm} = 0$ gives no shift at all. This is why practical codes use a more robust shift.

Fast convergence alone is not enough, since each step of Algorithm 20 still costs $O(m^3)$. In Lecture 14 we reduce this cost with a Hessenberg reduction, and we add shifts.

Lecture 14

Hessenberg QR

Supplementary reading:

- [1] Lecture 26.
- [2] Sections 4.4.5, 4.4.6, and 4.4.8.

14.1 Recap: From Orthogonal Iteration to the QR Algorithm

We start with a recap of Lecture 13.

- *Schur vectors and eigenvectors.* If $A = X\Lambda X^{-1}$ and $X = QR$, then $A = Q(R\Lambda R^{-1})Q^*$, and $R\Lambda R^{-1}$ is upper triangular. So the Schur vectors span the same nested subspaces as the eigenvectors (Section 13.2).
- *Orthogonal iteration.* It finds these subspaces without knowing X. Since $\operatorname{span}(Q_k) = \operatorname{span}(A^k Q_0)$, the span of the first p columns converges to $\operatorname{span}(\vec{x}_1, \ldots, \vec{x}_p)$. The rate is $|\lambda_{p+1}/\lambda_p|$ (Section 13.3).
- *Convergence to Schur form.* We split $Q_k = [\bar{Q}_k \; \tilde{Q}_k]$ after each column p. This shows that $\tilde{Q}_k^* A \bar{Q}_k \to 0$. So $T_k \coloneqq Q_k^* A Q_k$ converges to the triangular factor of a Schur factorization (Theorem 89).
- *The QR iteration.* Let $U_{k+1} \coloneqq Q_k^* Q_{k+1}$. Then we can write the iteration using only T_k:

$$T_k = U_{k+1} R_{k+1} \quad \longrightarrow \quad T_{k+1} = R_{k+1} U_{k+1}, \tag{14.1}$$

  starting from $T_0 = A$. This is the basic QR iteration (Algorithm 20).

In this lecture we ask: *how expensive is the QR iteration, and how can we make it cheaper?*

14.2 The Cost of the QR Iteration

Each step of Algorithm 20 computes the QR factorization of a dense $m \times m$ matrix. This costs $O(m^3)$, and we may need many steps. Suppose every eigenvector converged in $O(1)$ iterations. There are still m of them, so we would need $O(m)$ factorizations, for a total cost of $O(m^4)$. So faster convergence alone is not enough. We must also make each step cheaper.

The idea is to *preprocess* A *once*, so that every later QR factorization is simple. The preprocessing has to satisfy some conditions.

- It must not change the eigenvalues. So it must be a similarity transformation, $A \mapsto Q^* A Q$, and not a one-sided product as in the QR factorization.
- We take Q unitary. Then the conditioning does not get worse (Theorem 110).
- We cannot reduce A all the way to triangular form. That would compute a Schur factorization in finitely many steps, which is impossible (Section 10.3).

The best we can get is triangular form plus one extra diagonal.

Definition 91 (Hessenberg Matrix)

A matrix $H \in \mathbb{K}^{m \times m}$ is *upper Hessenberg* if $h_{ij} = 0$ for $i > j + 1$, that is, it is upper triangular plus one extra subdiagonal:

$$H = \begin{bmatrix} \times & \times & \times & \times \\ \times & \times & \times & \times \\ 0 & \times & \times & \times \\ 0 & 0 & \times & \times \end{bmatrix}. \tag{14.2}$$

14.3 Reduction to Hessenberg Form

We compute Q with Householder reflections (Lecture 6). Recall that a Householder reflector is both Hermitian and unitary. So it is its own inverse and its own adjoint.

Let $P_1 \in \mathbb{K}^{(m-1)\times(m-1)}$ be the Householder reflector that maps

$$\vec{v} \coloneqq A_{2:m,1} \quad \text{to} \quad \|\vec{v}\|\,\vec{e}_1, \qquad \text{and set} \qquad Q_1 \coloneqq \begin{bmatrix} 1 & 0 \\ 0 & P_1 \end{bmatrix}. \tag{14.3}$$

Multiplying by Q_1 on the left does not change the first row of A. It zeros out the first column below the subdiagonal:

$$\underbrace{\begin{bmatrix} \times & \times & \times & \times \\ \times & \times & \times & \times \\ \times & \times & \times & \times \\ \times & \times & \times & \times \end{bmatrix}}_{A} \xrightarrow{Q_1 \cdot} \underbrace{\begin{bmatrix} \times & \times & \times & \times \\ \times & \times & \times & \times \\ 0 & \times & \times & \times \\ 0 & \times & \times & \times \end{bmatrix}}_{Q_1 A}. \tag{14.4}$$

To complete the similarity, we multiply by $Q_1^* = Q_1$ on the right. This acts only on columns $2, \ldots, m$. So the first column of $Q_1 A$ does not change, and its new zeros stay:

$$\underbrace{\begin{bmatrix} \times & \times & \times & \times \\ \times & \times & \times & \times \\ 0 & \times & \times & \times \\ 0 & \times & \times & \times \end{bmatrix}}_{Q_1 A} \xrightarrow{\cdot\, Q_1^*} \underbrace{\begin{bmatrix} \times & \times & \times & \times \\ \times & \times & \times & \times \\ 0 & \times & \times & \times \\ 0 & \times & \times & \times \end{bmatrix}}_{Q_1 A Q_1^*}. \tag{14.5}$$

This is why the reflector starts at row 2 and leaves a_{21} in place. Suppose instead we used a reflector on all rows, as in the first step of Householder QR. It would zero out a_{21} too. But then multiplying by its adjoint on the right would mix all columns, and fill the first column back in.

Now we repeat the same step on the trailing $(m-1) \times (m-1)$ block. In step k, the reflector P_k maps the part of column k below the diagonal, $\vec{v} = (\cdot)_{k+1:m,k}$, to $\|\vec{v}\|\,\vec{e}_1$. We embed it as $Q_k = \operatorname{diag}(I_k, P_k)$.

- Left multiplication by Q_k changes only rows $k+1, \dots, m$.
- Right multiplication by Q_k changes only columns $k+1, \dots, m$.

So the zeros we created in columns $1, \dots, k-1$ stay. We need $m-2$ steps, since the last two columns have nothing below the subdiagonal. Then we collect the reflectors into

$$Q := Q_1 Q_2 \cdots Q_{m-2}, \qquad Q^* A Q = Q_{m-2} \cdots Q_1 \, A \, Q_1 \cdots Q_{m-2} = H. \tag{14.6}$$

The order of the factors in Q matters. Each Q_k is Hermitian, but $Q^* = Q_{m-2} \cdots Q_1$ has them in reverse order. (On the board the reflectors are collected as "$Q_m \cdots Q_1 = Q$". Since every Q_k is Hermitian, that product is Q^* in our notation. Also, we need only $m-2$ reflectors, because the last two columns have nothing below the subdiagonal.)

As in Householder QR, we never form the Q_k or Q. We store the reflection vectors and apply them from both sides.

Algorithm 21 Householder reduction to Hessenberg form [1, Algorithm 26.1].

Input: $A \in \mathbb{K}^{m \times m}$, overwritten by $H = Q^* A Q$

for $k = 1, \dots, m-2$ **do**
  $\vec{x} \leftarrow A_{k+1:m,k}$
  $\vec{v}_k \leftarrow \operatorname{sign}(x_1)\,\|\vec{x}\|\,\vec{e}_1 + \vec{x}, \quad \vec{v}_k \leftarrow \vec{v}_k / \|\vec{v}_k\|$
  $A_{k+1:m,k:m} \leftarrow A_{k+1:m,k:m} - 2\vec{v}_k(\vec{v}_k^* A_{k+1:m,k:m})$ ▷ from the left: rows $k+1, \dots, m$
  $A_{1:m,k+1:m} \leftarrow A_{1:m,k+1:m} - 2(A_{1:m,k+1:m}\vec{v}_k)\vec{v}_k^*$ ▷ from the right: columns $k+1, \dots, m$

Here the reflector maps $\vec{x}$ to $-\operatorname{sign}(x_1)\,\|\vec{x}\|\,\vec{e}_1$, not to $\|\vec{x}\|\,\vec{e}_1$. This avoids cancellation, exactly as in Lecture 6. Either choice gives a Hessenberg matrix. The cost has two parts.

- The left updates cost about $\frac{4}{3}m^3$ flops.
- The right updates act on full columns. They cost about $2m^3$ flops.

The total is

$$\text{work for Hessenberg reduction} \sim \tfrac{10}{3}m^3 \text{ flops } [1, (26.1)]. \tag{14.7}$$

We pay this once, before the QR iteration starts.

14.4 The QR Iteration on Hessenberg Matrices

We now run Algorithm 20 on $T_0 = H$ instead of A. The first half of each step is a QR factorization $U_k R_k = T_{k-1}$. Suppose T_{k-1} is upper Hessenberg. Then $O(m)$ Givens rotations are enough, as we saw in Section 7.3. We only need to eliminate the $m-1$ subdiagonal entries, and we can remove them one at a time from the top. Multiplying by G_1^* on the left acts on rows 1 and 2, and zeros out the $(2,1)$ entry. Then G_2^* acts on rows 2 and 3, and zeros out the $(3,2)$ entry. And so on:

$$\begin{bmatrix} \times & \times & \times & \times \\ \times & \times & \times & \times \\ 0 & \times & \times & \times \\ 0 & 0 & \times & \times \end{bmatrix} \xrightarrow{G_1^*} \begin{bmatrix} \times & \times & \times & \times \\ 0 & \times & \times & \times \\ 0 & \times & \times & \times \\ 0 & 0 & \times & \times \end{bmatrix} \xrightarrow{G_2^*} \begin{bmatrix} \times & \times & \times & \times \\ 0 & \times & \times & \times \\ 0 & 0 & \times & \times \\ 0 & 0 & \times & \times \end{bmatrix} \xrightarrow{G_3^*} \begin{bmatrix} \times & \times & \times & \times \\ 0 & \times & \times & \times \\ 0 & 0 & \times & \times \\ 0 & 0 & 0 & \times \end{bmatrix}. \tag{14.8}$$

Each rotation changes only two rows, at a cost of $O(m)$. So the whole factorization

$$R_k = G_{m-1}^* \cdots G_1^* T_{k-1}, \qquad U_k = G_1 G_2 \cdots G_{m-1}, \tag{14.9}$$

costs $O(m^2)$ instead of $O(m^3)$. We never form U_k. We only store the $m-1$ rotations.

The second half of each step computes $T_k = R_k U_k$. We keep U_k in factored form:

$$T_k = R_k G_1 G_2 \cdots G_{m-1}, \tag{14.10}$$

Multiplying by G_j on the right changes only columns j and $j+1$, again at a cost of $O(m)$. So the second half also costs $O(m^2)$.

For this to repeat, T_k must again be upper Hessenberg. It is. Right multiplication by G_j replaces columns j and $j+1$ by linear combinations of these two columns. Column $j+1$ of $R_k G_1 \cdots G_{j-1}$ has nonzeros down to row $j+1$. So mixing it into column j creates one new nonzero, at position $(j+1, j)$. This position is on the subdiagonal, and nothing appears below it:

$$\underbrace{\begin{bmatrix} \times & \times & \times & \times \\ 0 & \times & \times & \times \\ 0 & 0 & \times & \times \\ 0 & 0 & 0 & \times \end{bmatrix}}_{R_k} \xrightarrow{\cdot\, G_1} \begin{bmatrix} \times & \times & \times & \times \\ \times & \times & \times & \times \\ 0 & 0 & \times & \times \\ 0 & 0 & 0 & \times \end{bmatrix} \xrightarrow{\cdot\, G_2} \begin{bmatrix} \times & \times & \times & \times \\ \times & \times & \times & \times \\ 0 & \times & \times & \times \\ 0 & 0 & 0 & \times \end{bmatrix} \xrightarrow{\cdot\, G_3} \underbrace{\begin{bmatrix} \times & \times & \times & \times \\ \times & \times & \times & \times \\ 0 & \times & \times & \times \\ 0 & 0 & \times & \times \end{bmatrix}}_{T_k}. \tag{14.11}$$

So $T_k = R_k G_1 \cdots G_{m-1}$ is upper Hessenberg, and every step of the QR iteration costs $O(m^2)$. Then m iterations cost $O(m^3)$ in total. This is the same order as the one-time reduction to Hessenberg form. On the dense matrix A, the QR iteration would cost $O(m^4)$.

14.5 Bulge Chasing

So far, one step of the QR iteration on a Hessenberg matrix has two sweeps.

- The first sweep applies $G_1^*, \ldots, G_{m-1}^*$ from the left. It turns T_{k-1} into the upper triangular R_k.
- The second sweep applies $G_1, \ldots, G_{m-1}$ from the right. It turns R_k back into the Hessenberg matrix T_k.

In one formula,

$$T_k = \left(G_{m-1}^* \cdots G_1^* T_{k-1}\right) G_1 \cdots G_{m-1}. \tag{14.12}$$

Matrix multiplication is associative, so we do not have to use this order. *Can we apply the left and right rotations at the same time*? That is, can we compute

$$T_k = G_{m-1}^* \Big(\cdots G_2^* (G_1^* T_{k-1} G_1) G_2 \cdots \Big) G_{m-1}, \tag{14.13}$$

and finish each similarity $G_j^*(\cdot)G_j$ before we move on to the next rotation?

The difficulty is that we do not know the rotations in advance. We define G_2 by one requirement: G_2^* must zero out the $(3,2)$ entry of $G_1^* T_{k-1}$. So to compute G_2, we need the matrix $G_1^* T_{k-1}$. If we apply G_1 on the right right away, we only have $G_1^* T_{k-1} G_1$. In this matrix, the second column has been mixed with the first. *How can we compute G_2 from $G_1^* T_{k-1} G_1$?*

To answer this, we look at what happens when we compute $G_1^* T_{k-1} G_1$.

- Left multiplication by G_1^* only combines rows 1 and 2. It keeps the Hessenberg form (it zeros out the $(2,1)$ entry).
- Right multiplication by G_1 then combines columns 1 and 2. Column 2 has a nonzero in row 3.

So we get a nonzero at position $(3,1)$, one below the subdiagonal:

$$\underbrace{\begin{bmatrix} \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times \\ 0 & \times & \times & \times & \times \\ 0 & 0 & \times & \times & \times \\ 0 & 0 & 0 & \times & \times \end{bmatrix}}_{T_{k-1}} \xrightarrow{G_1^* \cdot G_1} \begin{bmatrix} \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times \\ + & \times & \times & \times & \times \\ 0 & 0 & \times & \times & \times \\ 0 & 0 & 0 & \times & \times \end{bmatrix}. \tag{14.14}$$

This entry, marked $+$, is called the *bulge*. The idea is to *choose G_2 so that it removes the bulge*. So G_2^* acts on rows 2 and 3, and zeros out the $(3,1)$ entry. Then right multiplication by G_2 combines columns 2 and 3. This creates a new bulge at $(4,2)$. So the bulge moves one step further down, along the subdiagonal:

$$\begin{bmatrix} \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times \\ + & \times & \times & \times & \times \\ 0 & 0 & \times & \times & \times \\ 0 & 0 & 0 & \times & \times \end{bmatrix} \xrightarrow{G_2^* \cdot G_2} \begin{bmatrix} \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times \\ 0 & \times & \times & \times & \times \\ 0 & + & \times & \times & \times \\ 0 & 0 & 0 & \times & \times \end{bmatrix} \xrightarrow{G_3^* \cdot G_3} \begin{bmatrix} \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times \\ 0 & \times & \times & \times & \times \\ 0 & 0 & \times & \times & \times \\ 0 & 0 & + & \times & \times \end{bmatrix} \xrightarrow{G_4^* \cdot G_4} \text{Hessenberg.} \tag{14.15}$$

The last rotation chases the bulge off the bottom of the matrix. What is left is an upper Hessenberg matrix. We apply each rotation from both sides as soon as we know it, so the step is a single sweep through the matrix. This is *bulge chasing*. We still need to show that this gives the same T_k as the two-sweep algorithm (up to signs). This follows from the implicit Q theorem. We state it in Section 14.6 and prove it at the start of the next lecture (Section 15.2).

Algorithm 22 One step of the QR iteration on an upper Hessenberg matrix by bulge chasing.

Input: upper Hessenberg $T \in \mathbb{K}^{m \times m}$, overwritten by the next iterate

1: choose G_1 acting on rows $1, 2$ such that $(G_1^* T)_{21} = 0$
2: $T \leftarrow G_1^* T G_1$ ▷ creates the bulge at $(3,1)$
3: **for** $j = 2, \ldots, m-1$ **do**
4: choose G_j acting on rows $j, j+1$ such that $(G_j^* T)_{j+1,j-1} = 0$ ▷ remove the bulge
5: $T \leftarrow G_j^* T G_j$ ▷ bulge moves to $(j+2, j)$, or leaves the matrix if $j = m-1$

Each step changes only two rows and two columns. So one QR step still costs $O(m^2)$, but now in a single sweep through the matrix.

14.6 The Implicit Q Theorem

We end with the statement of the theorem that justifies bulge chasing. We give its proof in Section 15.2.

Definition 92 (Unreduced Hessenberg Matrix)

An upper Hessenberg matrix H is *unreduced* if all its subdiagonal entries are nonzero: $h_{j+1,j} \neq 0$ for $j = 1, \dots, m-1$. Equivalently, H is not block upper triangular.

Suppose some $h_{j+1,j} = 0$. Then H is block upper triangular, and its eigenvalue problem splits into the problems for the two diagonal blocks. So we can always assume that H is unreduced.

Theorem 93 (Implicit Q Theorem)

Assume that Q is orthogonal and $Q^\top AQ = H$ is unreduced upper Hessenberg. Then the first column of Q determines columns $2, \dots, m$ of Q uniquely, up to signs.

The name comes from how we use the theorem. To know the whole transformation Q, it is enough to know two things: its first column, and that the result is upper Hessenberg. We never have to compute Q explicitly.

For bulge chasing, this works as follows.

- Both the two-sweep algorithm and the one-sweep algorithm give an upper Hessenberg $T_k = U_k^* T_{k-1} U_k$.
- In both, $U_k \vec{e}_1 = G_1 \vec{e}_1$, with the same G_1.

So by the theorem, the two U_k agree up to signs, and so do the two T_k (Section 15.3).

14.7 (SUPPLEMENT) Further Topics

14.7.1 The Hermitian Case: Tridiagonal Form

If A is Hermitian, then so is $H = Q^* AQ$. A Hermitian Hessenberg matrix is tridiagonal. We can use the zeros and the symmetry to reduce the cost from $\frac{10}{3}m^3$ to $\sim \frac{4}{3}m^3$ flops. After that, each QR step on a tridiagonal matrix costs only $O(m)$ [1, Lecture 26], [2, Section 4.4.7].

14.7.2 Stability

Like Householder QR, Algorithm 21 is backward stable. Let $\tilde{H}$ be the computed result, and let $\tilde{Q}$ be the exactly unitary matrix defined by the computed reflection vectors. Then $\tilde{Q}\tilde{H}\tilde{Q}^* = A + \delta A$ with $\|\delta A\| / \|A\| = O(\varepsilon)$ [1, Theorem 26.1]. So the eigenvalues of $\tilde{H}$ are the exact eigenvalues of a nearby matrix.

14.7.3 The "Bad Idea" Revisited

Suppose we apply a full Householder reflector from both sides, as in the first step of Householder QR. This destroys the zeros it creates. But it usually does make the entries below the diagonal smaller. This is exactly what one step of the QR iteration does. In this sense, the QR algorithm "triangularizes in the limit" [1, Lecture 26].

14.7.4 In Practice

LAPACK computes the Hessenberg reduction with blocked Householder updates (`xGEHRD`). It runs the QR iteration with multishift bulge chasing and aggressive early deflation (`xHSEQR`). In Julia, `hessenberg(A)` returns the reduction, and `eigvals(A)` runs all the steps.

Lecture 15

Shifted QR

Supplementary reading:

- [2] Section 4.4.8 (Theorem 4.9, the implicit Q theorem).
- [2] Sections 4.4.4 and 4.4.5 (QR iteration with a shift, making it practical).
- [2] Section 5.3.1 (Wilkinson's shift).
- [1] Lecture 33 (Arnoldi iteration).

15.1 Recap: QR Iteration on Hessenberg Matrices

We begin with a recap of Lecture 14. First we reduce A once to upper Hessenberg form $H = Q^\top A Q$ (Section 14.3). Then we run the QR iteration on $T_0 = H$. Each step does a QR factorization by Givens rotations, and then multiplies the factors in reverse order (Section 14.4):

$$R_k = G_{m-1}^\top \cdots G_1^\top T_{k-1}, \qquad T_k = R_k G_1 \cdots G_{m-1}. \tag{15.1}$$

The new T_k is again upper Hessenberg. So each step costs $O(m^2)$.

In class we compared two orders of applying the same rotations.

- In the *two-sweep* order, we do all left multiplications first, and then all right multiplications:

$$G_1^\top T_{k-1}, \quad G_2^\top G_1^\top T_{k-1}, \quad \ldots, \quad R_k, \quad R_k G_1, \quad R_k G_1 G_2, \quad \ldots, \quad T_k. \tag{15.2}$$

- In *bulge chasing* (Section 14.5), we apply each rotation from both sides as soon as we know it:

$$G_1^\top T_{k-1} G_1, \quad G_2^\top G_1^\top T_{k-1} G_1 G_2, \quad \ldots, \quad T_k. \tag{15.3}$$

By associativity, both orders give the same product *if they use the same rotations*. But the rotations are chosen differently. In bulge chasing, we choose $G_2, \ldots, G_{m-1}$ to remove the bulge at $(3,1)$ that $G_1^\top T_{k-1} G_1$ creates. In the two-sweep order, we choose them to remove the subdiagonal entries of $G_1^\top T_{k-1}$. Why do we still get the same T_k? The answer is the implicit Q theorem, which we stated at the end of Lecture 14. In this lecture we first prove it, and then we study shifts.

15.2 Proof of the Implicit Q Theorem

Recall the statement (Theorem 93). Let Q be orthogonal, and let $Q^\top AQ = H$ be unreduced upper Hessenberg. Then the first column of Q fixes columns $2, \dots, m$ of Q, up to signs.

Proof. We follow [2, Theorem 4.9]. Let both $Q^\top AQ = H$ and $V^\top AV = G$ be as in the theorem. Assume they have the same first column, $\vec{q}_1 = \vec{v}_1$. We want to show that

$$W \coloneqq V^\top Q = \operatorname{diag}(\pm 1, \dots, \pm 1). \tag{15.4}$$

Then $Q = VW$, i.e., $\vec{q}_i = \pm \vec{v}_i$ for every i.

W is a product of orthogonal matrices, so it is orthogonal. Its first column is $W\vec{e}_1 = V^\top \vec{q}_1 = V^\top \vec{v}_1 = \vec{e}_1$. We also have $GV^\top = V^\top AVV^\top = V^\top A$ and $AQ = QH$. With these,

$$GW = GV^\top Q = V^\top AQ = V^\top QH = WH. \tag{15.5}$$

Write $\vec{w}_i$ for the ith column of W. We take column i of both sides. Since H is upper Hessenberg, column i of H is zero below row $i+1$. So

$$G\vec{w}_i = (GW)_{:,i} = (WH)_{:,i} = \sum_{j=1}^{i+1} \vec{w}_j H_{ji}. \tag{15.6}$$

Only the term $j = i+1$ contains $\vec{w}_{i+1}$. We move the other terms to the right-hand side:

$$H_{i+1,i}\, \vec{w}_{i+1} = G\vec{w}_i - \sum_{j=1}^{i} H_{ji}\vec{w}_j. \tag{15.7}$$

We show by induction that each $\vec{w}_i$ is nonzero in entries $1, \dots, i$ only:

$$\vec{w}_i = \begin{bmatrix} \times \\ \vdots \\ \times \\ 0 \\ \vdots \\ 0 \end{bmatrix} \leftarrow i \tag{15.8}$$

This holds for $\vec{w}_1 = \vec{e}_1$. Suppose it holds for $\vec{w}_1, \dots, \vec{w}_i$. We look at the right-hand side of (15.7).

- $G\vec{w}_i$ is a combination of the first i columns of G. Since G is upper Hessenberg, these columns are zero below row $i+1$.
- The sum is zero below row i, by the induction hypothesis.

So the right-hand side is zero below row $i+1$. Since H is unreduced, $H_{i+1,i} \neq 0$. We divide by it, and get that $\vec{w}_{i+1}$ is zero below row $i+1$.

So W is upper triangular. W is also orthogonal, so $W^{-1} = W^\top$. This matrix is upper triangular, as the inverse of an upper triangular matrix. It is also lower triangular, as the transpose of one. So W is diagonal. A diagonal orthogonal matrix has $w_{ii}^2 = 1$, so $W = \operatorname{diag}(\pm 1, \dots, \pm 1)$. □

This proof works like the proof that the QR factorization is unique (Theorem 33).

- We take two solutions, and look at the matrix W that maps one to the other.
- We show that W is triangular and orthogonal. Then W must be diagonal, with entries of modulus one.
- These entries are exactly the sign choices that we cannot fix.

The recursion (15.7) also tells us how to build Q from $\vec{q}_1$, one column at a time. We use $AQ = QH$. Orthogonality gives $h_{1i}, \dots, h_{ii}$. Then normalization gives $\vec{q}_{i+1}$. This is the Arnoldi iteration [1, Lecture 33]. In the step where we divide by $h_{i+1,i}$, we need H to be unreduced.

For $\mathbb{K} = \mathbb{C}$, the same proof works with Q and V unitary, and with $\top$ replaced by $*$. Then W is a unitary diagonal matrix. So each column of Q is fixed up to a factor of modulus one.

15.3 Application to Bulge Chasing

Both the two-sweep algorithm and bulge chasing compute $T_k = U_k^\top T_{k-1} U_k$. In both, U_k is a product of rotations $G_1 G_2 \cdots G_{m-1}$, where G_j acts on rows j and $j+1$. We check the conditions of Theorem 93.

- *Same first column.* Only G_1 touches the first row, so $U_k \vec{e}_1 = G_1 \vec{e}_1$ in both cases. Also, G_1 is the same in both, because only the first column of T_{k-1} decides it.
- *Hessenberg form.* Both results are upper Hessenberg.

Assume T_k is unreduced (if not, we deflate). Then Theorem 93 says that the two U_k agree up to the signs of their columns. So the two T_k agree up to a similarity $D^\top T_k D$ with $D = \operatorname{diag}(\pm 1, \dots, \pm 1)$. This similarity only flips the signs of some rows and columns. As in Section 13.5, it does not change the diagonal or the moduli of the entries.

15.4 The Shifted QR Iteration

The QR iteration of Algorithm 20 can converge slowly. Its rate is $|\lambda_{p+1}/\lambda_p|$ (Section 13.3). As for inverse iteration (Algorithm 16), the fix is a shift. Here are the two iterations side by side:

$$\text{normal QR:} \quad \begin{aligned} T_{k-1} &= U_k R_k, \\ T_k &= R_k U_k, \end{aligned} \qquad \text{shifted QR:} \quad \begin{aligned} T_{k-1} - \mu_k I &= U_k R_k, \\ T_k &= R_k U_k + \mu_k I. \end{aligned} \tag{15.9}$$

We subtract the shift μ_k before the factorization, and add it back afterwards. We check the two properties that made the unshifted iteration work.

The upper Hessenberg form is kept. Subtracting and adding $\mu_k I$ changes only the diagonal. So:

- $T_{k-1} - \mu_k I$ is upper Hessenberg, and its QR factorization takes $m - 1$ Givens rotations;
- $R_k U_k$ is upper Hessenberg, by Section 14.4.

Each step still costs $O(m^2)$.

The eigenvalues are kept. The first equation gives $R_k = U_k^*(T_{k-1} - \mu_k I)$. So

$$T_k = R_k U_k + \mu_k I = U_k^*(T_{k-1} - \mu_k I)U_k + \mu_k I = U_k^* T_{k-1} U_k, \tag{15.10}$$

since $U_k^*(\mu_k I)U_k = \mu_k I$. So T_k is again a unitary similarity transformation of T_{k-1}. If we did not add back $\mu_k I$, every step would shift the eigenvalues by $-\mu_k$.

Tracking the eigenvectors. The relation $T_k = U_k^* T_{k-1} U_k$ has exactly the same form as in the unshifted iteration. So we can track Q_k as before (Section 13.5). After each step we update $Q \leftarrow QU_k$. Then $T_k = (U_1 \cdots U_k)^* H (U_1 \cdots U_k)$. Together with the Hessenberg reduction, this gives

$$A \approx (QU_1 \cdots U_k)\, T_k\, (QU_1 \cdots U_k)^*$$

once T_k is nearly triangular.

15.5 Choosing the Shift

How does the shift change convergence? Which μ_k should we choose? We first look at the ideal case.

Exact shift. Assume μ_k is an eigenvalue of T_{k-1}. Then $T_{k-1} - \mu_k I$ is singular. Assume also that all off-diagonal entries $t_{j+1,j}$ are nonzero, i.e., T_{k-1} is unreduced. (If not, we split the problem; see below.)

First, the first $m-1$ columns of $T_{k-1} - \mu_k I$ are linearly independent. The reason: rows $2, \ldots, m$ of these columns form an upper triangular $(m-1) \times (m-1)$ matrix. Its diagonal entries are the subdiagonal entries of T_{k-1}, which are nonzero.

Now look at the QR factorization $T_{k-1} - \mu_k I = U_k R_k$. Column j is a combination of $\vec{u}_1, \ldots, \vec{u}_j$, and the coefficient on $\vec{u}_j$ is r_{jj}. If $r_{jj} = 0$ for some $j \le m-1$, then column j would lie in the span of the columns before it. So $r_{11}, \ldots, r_{m-1,m-1} \neq 0$. But R_k is singular. So *the last entry must be zero*: $r_{mm} = 0$. Since R_k is upper triangular, its whole last row is zero:

$$\underbrace{T_{k-1} - \mu_k I}_{\text{unreduced Hessenberg}} = U_k \underbrace{\begin{bmatrix} \times & \times & \times & \times \\ 0 & \times & \times & \times \\ 0 & 0 & \times & \times \\ 0 & 0 & 0 & 0 \end{bmatrix}}_{R_k}. \tag{15.11}$$

What happens to T_k? The last row of $R_k U_k$ is the last row of R_k times U_k. So it is zero. Adding $\mu_k I$ gives

$$T_k = R_k U_k + \mu_k I = \begin{bmatrix} & \hat{T} & & \vec{t} \\ 0 & \cdots & 0 & \mu_k \end{bmatrix}, \tag{15.12}$$

with $\hat{T} \in \mathbb{K}^{(m-1)\times(m-1)}$ upper Hessenberg. So $t_{m,m-1} = 0$ and $t_{mm} = \mu_k$. With an exact shift, *one step* of the QR iteration splits off the eigenvalue μ_k. As we said in class, we don't have to worry about the last entry any more.

Then we continue on $\hat{T}$ alone. A unitary Q that acts on $\hat{T}$ acts on the full matrix as $\mathrm{diag}(Q, 1)$:

$$\begin{bmatrix} Q^* & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \hat{T} & \vec{t} \\ 0 & \mu_k \end{bmatrix} \begin{bmatrix} Q & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} Q^*\hat{T}Q & Q^*\vec{t} \\ 0 & \mu_k \end{bmatrix}, \tag{15.13}$$

So the eigenvalue we already found does not change, and the problem is one size smaller. We need to update the last column $Q^*\vec{t}$ only if we want the full Schur form, not just the eigenvalues.

Splitting. In practice μ_k is only close to an eigenvalue. So the subdiagonal entries become small, but not zero. Whenever a subdiagonal entry gets close to 0, we set it to zero. Then T_k is (approximately) block upper triangular:

$$T_k \approx \begin{bmatrix} (T_k)_{11} & (T_k)_{12} \\ 0 & (T_k)_{22} \end{bmatrix}, \tag{15.14}$$

Its eigenvalues are the eigenvalues of the two diagonal blocks. Both blocks are again upper Hessenberg. We recurse on $(T_k)_{11}$ and $(T_k)_{22}$ separately.

Algorithm 23 Practical QR iteration.

Input: $A \in \mathbb{K}^{m\times m}$

compute $H = Q^* A Q$ ▷ Algorithm 21, once
$T_0 \leftarrow H$, $k \leftarrow 0$
while not converged **do**
    $k \leftarrow k+1$, choose μ_k
    $T_{k-1} - \mu_k I = U_k R_k$ ▷ Givens QR, $O(m^2)$
    $T_k \leftarrow R_k U_k + \mu_k I$
    if some subdiagonal entry of T_k is ≈ 0 **then**
        split T_k into $(T_k)_{11}$ and $(T_k)_{22}$ and recurse on both

Which eigenvalue to aim for? We want to choose μ_k close to an eigenvalue, and split the problem wherever a subdiagonal entry is ≈ 0. But close to which eigenvalue? In class we chose $\mu_k \approx \lambda_m$, for two reasons.

1. The shift helps the eigenvalue λ_i to separate if $|\lambda_i - \mu| \ll |\lambda_j - \mu|$ for all $j \neq i$. Without a shift ($\mu = 0$), this is easiest when λ_i itself is small. That is the eigenvalue of smallest modulus.
2. The QR iteration sorts the eigenvalues by size along the diagonal of T_k (Theorem 89). The smallest one, λ_m, is at the bottom right. A shift close to another eigenvalue would have to move that eigenvalue to the bottom. And *reordering the eigenvalues slows down convergence*.

So the shift should be close to the eigenvalue that is already converging in the bottom right corner. There we then check whether $t_{m,m-1} \approx 0$.

15.6 Relationship with (Inverse) Power Iteration

We have seen that the QR iteration is the same as orthogonal iteration with $Q_0 = I$ (Section 13.5). In the same way, the shifted QR iteration is the same as a *shifted* orthogonal iteration. Start with $T_0 = H$. The first two steps are

$$H - \mu_1 I = U_1 R_1, \qquad T_1 = R_1 U_1 + \mu_1 I, \qquad T_1 - \mu_2 I = U_2 R_2.$$

So

$$\begin{aligned} U_1 U_2 R_2 R_1 &= U_1 (T_1 - \mu_2 I) R_1 = U_1 (R_1 U_1 + \mu_1 I - \mu_2 I) R_1 \\ &= (H - \mu_1 I)\big((H - \mu_1 I) + \mu_1 I - \mu_2 I\big) \\ &= (H - \mu_1 I)(H - \mu_2 I), \end{aligned} \tag{15.15}$$

In the second line we used $U_1 R_1 = H - \mu_1 I$ twice:

$$U_1 R_1 U_1 R_1 = (H - \mu_1 I)^2, \qquad U_1 (\mu_1 - \mu_2) R_1 = (\mu_1 - \mu_2)(H - \mu_1 I).$$

Let $\bar{Q}_k \coloneqq U_1 \cdots U_k$ and $\bar{R}_k \coloneqq R_k \cdots R_1$. In the same way we get

$$(H - \mu_k I) \cdots (H - \mu_1 I) = \bar{Q}_k \bar{R}_k. \tag{15.16}$$

We prove this by induction. By the similarity relation of Section 15.4, $T_{k-1} = \bar{Q}_{k-1}^* H \bar{Q}_{k-1}$. So

$$\bar{Q}_k \bar{R}_k = \bar{Q}_{k-1}(T_{k-1} - \mu_k I)\bar{R}_{k-1} = (H - \mu_k I)\bar{Q}_{k-1}\bar{R}_{k-1}. \tag{15.17}$$

$\bar{Q}_k$ is unitary. $\bar{R}_k$ is upper triangular, as a product of upper triangular matrices. So (15.16) is a QR factorization of the matrix polynomial

$$p_k(H) = (H - \mu_k I) \cdots (H - \mu_1 I).$$

The factors commute, so their order does not matter.

Compare this with the unshifted case. There, $\bar{Q}_k$ is the Q factor of H^k. Now the power H^k is replaced by $p_k(H)$.

- H^k multiplies the eigenvector component for λ_j by λ_j^k.
- $p_k(H)$ multiplies it by $p_k(\lambda_j) = \prod_{i=1}^k (\lambda_j - \mu_i)$.

If the shifts are close to an eigenvalue, p_k is tiny there. This is what makes convergence faster. Also, $T_k = \bar{Q}_k^* H \bar{Q}_k$ shows how to track the eigenvectors. $\bar{Q}_k$ collects all the transformations. Once T_k has converged to upper triangular form, $Q\bar{Q}_k$ contains the Schur vectors of A.

What about inverse iteration? We invert (15.16) and get $p_k(H)^{-1} = \bar{R}_k^{-1}\bar{Q}_k^*$. Then we take the adjoint:

$$\bar{Q}_k \bar{R}_k^{-*} = (H - \mu_k I)^{-*} \cdots (H - \mu_1 I)^{-*}, \tag{15.18}$$

where $X^{-*} \coloneqq (X^{-1})^*$. Inversion and the adjoint both reverse the order of a product. But the factors on the right commute, so the order does not matter. The right-hand side is exactly the operator of shifted inverse iteration (Algorithm 16). It is applied to H^* with shifts $\bar{\mu}_i$. (H^* has the eigenvalues $\bar{\lambda}_j$.)

The difference from (15.16) is the triangular factor. $\bar{R}_k^{-1}$ is upper triangular, so $\bar{R}_k^{-*}$ is *lower* triangular.

- With an upper triangular factor, the first columns of the product use only the first columns of $\bar{Q}_k$.
- With a lower triangular factor, the last columns of the product use only the last columns of $\bar{Q}_k$.

In particular, the last column of $\bar{R}_k^{-*}$ is $(\bar{R}_k^{-*})_{mm}\vec{e}_m$, with $(\bar{R}_k^{-*})_{mm} = 1/\overline{(\bar{R}_k)_{mm}}$. So multiplying (15.18) by $\vec{e}_m$ gives

$$\bar{\vec{q}}_m = \overline{(\bar{R}_k)_{mm}}\,(H - \mu_k I)^{-*} \cdots (H - \mu_1 I)^{-*}\vec{e}_m, \tag{15.19}$$

where $\bar{\vec{q}}_m$ is the last column of $\bar{Q}_k$. So the same matrix $\bar{Q}_k$ does two iterations at once:

factorization	triangular factor	columns of $\bar{Q}_k$
$p_k(H) = \bar{Q}_k \bar{R}_k$	upper	first columns: orthogonal (power) iteration
$p_k(H)^{-*} = \bar{Q}_k \bar{R}_k^{-*}$	lower	last columns: shifted inverse iteration

The last column is shifted inverse iteration started from $\vec{e}_m$. For a single step, this is the argument of [2, Section 4.4.4]. In short, *the QR iteration is both power and inverse power iteration*. This explains two things.

- The bottom right corner of T_k converges first.
- The shift should be close to λ_m (Section 15.5). A shift close to an eigenvalue makes inverse iteration converge fast (Theorem 62).

15.7 (SUPPLEMENT) Further Topics

15.7.1 The Rayleigh Quotient Shift

In class we did not fix a concrete choice of $\mu_k \approx \lambda_m$. The natural estimate is the bottom right entry,

$$\mu_k = t_{mm}^{(k-1)} = \vec{e}_m^* T_{k-1} \vec{e}_m.$$

This is the Rayleigh quotient of T_{k-1} at $\vec{e}_m$. So we get a link to Rayleigh quotient iteration (Algorithm 17). By the inverse iteration view above, the last column of $\bar{Q}_k$ is inverse iteration, with the shift taken from the current Rayleigh quotient. This gives local *quadratic* convergence. Demmel [2, Section 4.4.4] explains why.

- Let $\eta = \|T_{k-1}(m, 1{:}m-1)\| \,/\, \|A\| \ll 1$.
- Setting the off-diagonal part of the last row to zero changes an eigenvalue by $O(\eta \,\|A\|)$, if the eigenvalue is well separated. So $|\lambda_m - t_{mm}| = O(\eta \,\|A\|)$.
- The next step shrinks the last row by the factor $|\lambda_m - \mu| \,/\, \min_{j \neq m} |\lambda_j - \mu| = O(\eta)$. This gives $O(\eta^2)$.

15.7.2 The Wilkinson Shift

The Rayleigh quotient shift can get stuck. Take $T_0 = \left[\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right]$. Then $\mu_1 = t_{22} = 0$. T_0 is itself orthogonal, so its QR factorization is $U_1 = T_0$, $R_1 = I$. So $T_1 = R_1 U_1 = T_0$. Nothing changes, because the eigenvalues ± 1 are equally far from the shift.

The *Wilkinson shift* uses a different μ_k. Look at the last 2×2 block of T_{k-1}. Of its two eigenvalues, we take the one closer to t_{mm}. In the example this is ± 1, an exact eigenvalue, so one step is enough. For symmetric tridiagonal matrices, QR iteration with the Wilkinson shift has these properties [2, Theorem 5.8]:

- it converges globally, and at least linearly;
- for almost all matrices, it converges cubically in the limit.

15.7.3 When Is a Subdiagonal Entry ≈ 0?

We get T_k from A by unitary similarity transformations in floating point. So T_k already has rounding errors of size $O(\varepsilon_{\text{mach}}) \,\|A\|$. A subdiagonal entry smaller than this might as well be zero. Setting it to zero is a backward stable perturbation [2, Section 4.4.5]. The iteration stops when all diagonal blocks are 1×1. (For a real matrix with complex conjugate pairs of eigenvalues, some blocks are 2×2.)

15.7.4 Why the Implicit Q Theorem Matters for Shifts

With a shift μ, one QR step factors $T_{k-1} - \mu I = U_k R_k$. Then it sets $T_k = R_k U_k + \mu I = U_k^* T_{k-1} U_k$. By Theorem 93, the first column of U_k fixes U_k. This first column is parallel to the first column of $T_{k-1} - \mu I$, which is $(t_{11} - \mu, t_{21}, 0, \ldots, 0)^\top$. So:

- we compute G_1 from these two numbers;
- the rest of the step is the bulge chasing of Algorithm 22 on T_{k-1} itself;
- we never form $T_{k-1} - \mu I$.

This is the *implicitly shifted* QR algorithm [2, Section 4.4.8]. A real matrix can have complex conjugate eigenvalues. Then we can combine two such steps, with shifts μ and $\bar{\mu}$, into one *double-shift* step that uses only real arithmetic (the Francis step) [2, Section 4.4.8].

15.7.5 The Krylov View

Why should the first column fix everything? Column i of $AQ = QH$ gives

$$h_{i+1,i}\vec{q}_{i+1} = A\vec{q}_i - \sum_{j \le i} h_{ji}\vec{q}_j.$$

By induction, using $h_{i+1,i} \neq 0$, we get

$$\operatorname{span}(\vec{q}_1, \ldots, \vec{q}_k) = \operatorname{span}(\vec{q}_1, A\vec{q}_1, \ldots, A^{k-1}\vec{q}_1) \qquad \text{for } k = 1, \ldots, m. \tag{15.20}$$

The right-hand side is the Krylov subspace generated by A and $\vec{q}_1$. It depends on nothing else. We can say the same with matrices. The Krylov matrix

$$K = [\vec{q}_1, A\vec{q}_1, \ldots, A^{m-1}\vec{q}_1]$$

satisfies $K = QR$, with $R = Q^\top K$ upper triangular and invertible. So Q is the orthogonal factor of the QR factorization of K. The QR factorization is unique up to signs, and this gives the implicit Q theorem [1, Lecture 33]. The proof above is the same argument, written out column by column.

15.7.6 A Direct Check for the First Bulge

For the first rotation after G_1, we can also check directly that the two algorithms agree. Write $G_1 = \left[\begin{smallmatrix} c & -s \\ s & c \end{smallmatrix}\right]$ in rows and columns $1, 2$. Let $X = G_1^\top T_{k-1}$.

- *Two-sweep.* The first column of X is zero in rows $2, \ldots, m$. The algorithm chooses G_2 from (x_{22}, x_{32}).
- *Bulge chasing.* Right multiplication by G_1 replaces column 1 by $c\,(\text{column } 1) + s\,(\text{column } 2)$. So in XG_1, the bulge column has entries $s\,(x_{22}, x_{32})$ in rows $2, 3$.

This vector is parallel to (x_{22}, x_{32}). So bulge chasing chooses the same G_2, up to sign. (If $s = 0$, then $(T_{k-1})_{21} = 0$, and T_{k-1} was reduced.) We could continue this argument down the matrix, but it is tedious. The implicit Q theorem handles all rotations at once.

Appendix A

Linear Algebra Basics

This appendix collects the first-course linear algebra that is used without comment throughout the course: matrix products, transposes, inverses, determinants, subspaces, eigenvalues, orthogonality, and symmetric matrices. Proofs are omitted; results proved in the main text are cross-referenced. Throughout, A and B are matrices of compatible sizes, c is a scalar, and $\vec{x}, \vec{y}$ are column vectors. Supplementary reading:

- [3] Chapters 2–7.
- [1] Lectures 1 and 2.
- [4] Section 7A.

A.1 Matrix Multiplication

Write $\vec{a}_j$ for the columns of $A \in \mathbb{K}^{m\times n}$ and $\vec{b}_i^{\top}$ for the rows of $B \in \mathbb{K}^{n\times p}$. The product AB can be read in four equivalent ways:

- *entries*: $(AB)_{ij} = \sum_k A_{ik}B_{kj}$;
- *columns*: $A\vec{x} = \sum_j x_j \vec{a}_j$ is a linear combination of the columns of A, and the j-th column of AB is A times the j-th column of B;
- *rows*: the i-th row of AB is the i-th row of A times B;
- *outer products*: $AB = \sum_{k=1}^{n} \vec{a}_k \vec{b}_k^{\top}$, a sum of rank-one matrices.

Multiplication is associative and distributive but not commutative. Multiplying by a diagonal matrix D on the left scales the rows (DA), on the right the columns (AD). Block matrices multiply like ordinary matrices as long as the block sizes are compatible.

A.2 Transpose and Conjugate Transpose

Definition 94 (Transpose and Conjugate Transpose)

For $A \in \mathbb{K}^{m\times n}$, the *transpose* $A^\top \in \mathbb{K}^{n\times m}$ and the *conjugate transpose* (or *adjoint*) $A^* \in \mathbb{K}^{n\times m}$ are

$$(A^\top)_{ij} = A_{ji}, \qquad (A^*)_{ij} = \overline{A_{ji}}. \tag{A.1}$$

For real matrices the two coincide.

Proposition 95 (Rules for the Transpose)

1. $(A^\top)^\top = A$.
2. $(A+B)^\top = A^\top + B^\top$ and $(cA)^\top = cA^\top$.
3. $(AB)^\top = B^\top A^\top$, and more generally $(A_1 A_2 \cdots A_k)^\top = A_k^\top \cdots A_2^\top A_1^\top$.
4. If A is invertible, so is $A^\top$, and $(A^\top)^{-1} = (A^{-1})^\top$, written $A^{-\top}$.
5. $\det A^\top = \det A$, $\operatorname{tr} A^\top = \operatorname{tr} A$, and $\operatorname{rank} A^\top = \operatorname{rank} A$.
6. $A^\top$ and A have the same characteristic polynomial, hence the same eigenvalues, but in general different eigenvectors.

The order reversal in (3) is the same as in $(AB)^{-1} = B^{-1}A^{-1}$: the factor applied last is undone first. Rule (4) follows from (3) by transposing $AA^{-1} = I$.

A.2.1 Vectors and Inner Products

Since a column vector is an $n \times 1$ matrix, the rules above apply to it directly.

- $(A\vec{x})^\top = \vec{x}^\top A^\top$.
- $\vec{x}^\top \vec{y} = \vec{y}^\top \vec{x}$, because $\vec{x}^\top \vec{y}$ is a 1×1 matrix, and a scalar is its own transpose.
- $(A\vec{x})^\top \vec{y} = \vec{x}^\top (A^\top \vec{y})$, i.e., $\langle A\vec{x}, \vec{y}\rangle = \left\langle \vec{x}, A^\top \vec{y}\right\rangle$. This *adjoint relation* moves a matrix from one side of an inner product to the other. It is the key step in deriving the normal equations (Lemma 34).

A.2.2 Block Matrices

A block matrix is transposed by transposing each block *and* the arrangement of the blocks:

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix}^\top = \begin{bmatrix} A^\top & C^\top \\ B^\top & D^\top \end{bmatrix}. \tag{A.2}$$

The off-diagonal blocks B and C swap places.

A.2.3 The Complex Case

For complex matrices the conjugate transpose $A^* = \overline{A}^\top$ plays the role of the transpose. All rules of Proposition 95 carry over, with complex conjugates wherever a scalar appears:

- $(cA)^* = \bar{c}A^*$ and $(AB)^* = B^*A^*$;
- $\det A^* = \overline{\det A}$, and the eigenvalues of A^* are the complex conjugates of those of A;
- $\vec{x}^*\vec{y} = \overline{\vec{y}^*\vec{x}}$, which in general is *not* equal to $\vec{y}^*\vec{x}$;
- $\langle A\vec{x}, \vec{y}\rangle = \langle \vec{x}, A^*\vec{y}\rangle$ for the standard inner product $\langle \vec{x}, \vec{y}\rangle = \vec{x}^*\vec{y}$ (Lecture 11).

Hermitian versus symmetric. A matrix is *Hermitian* if $A^* = A$ and *symmetric* if $A^\top = A$. For real matrices $A^* = A^\top$, so the two notions coincide, and every result about Hermitian matrices applies to real symmetric ones. For complex matrices they differ: a complex symmetric matrix is in general not Hermitian. For example,

$$A = \begin{bmatrix} 1 & i \\ i & 1 \end{bmatrix} \tag{A.3}$$

satisfies $A^\top = A$, but $A^* = \left[\begin{smallmatrix} 1 & -i \\ -i & 1 \end{smallmatrix}\right] \neq A$, and its eigenvalues $1 \pm i$ are not real. Complex symmetric matrices therefore enjoy none of the special properties of Hermitian matrices (real eigenvalues, unitary diagonalization, the Rayleigh quotient characterization). In these notes $\mathbb{K}$ is $\mathbb{R}$ or $\mathbb{C}$, and "Hermitian" means "symmetric" when $\mathbb{K} = \mathbb{R}$.

A.2.4 Useful Consequences

- A^*A and AA^* are Hermitian and positive semidefinite, since $\vec{x}^*A^*A\vec{x} = \|A\vec{x}\|_2^2 \geq 0$.
- Q is unitary if and only if $Q^*Q = I$, i.e., $Q^* = Q^{-1}$ (Section B.2).
- A^* and A have the same singular values, so $\|A^*\|_2 = \|A\|_2$ and $\kappa_2(A^*) = \kappa_2(A)$.
- $\|A^*\|_1 = \|A\|_\infty$, since transposing turns column sums into row sums (Proposition 103).

Remark 96 (Common Mistakes)**.**

- Writing $(AB)^\top = A^\top B^\top$. The order must be reversed.
- Assuming that the product of two symmetric matrices is symmetric. In fact $(AB)^\top = B^\top A^\top = BA$, which in general differs from AB.
- Using $^\top$ where * is meant for complex matrices, e.g., writing $A^\top = A$ for "Hermitian".

A.3 Inverses, Elementary and Triangular Matrices

A square matrix A is invertible if $AB = BA = I$ for some $B = A^{-1}$; for square matrices a one-sided inverse is automatically two-sided. Then $(AB)^{-1} = B^{-1}A^{-1}$ and $(A^{-1})^{-1} = A$.

Theorem 97 (Invertible Matrix Theorem)

For $A \in \mathbb{K}^{n\times n}$ the following are equivalent:

1. A is invertible;
2. $A\vec{x} = \vec{b}$ has a unique solution for every $\vec{b}$;
3. $A\vec{x} = \vec{0}$ has only the solution $\vec{x} = \vec{0}$;
4. the columns (equivalently, the rows) of A are linearly independent;
5. $\operatorname{rank} A = n$;
6. $\det A \neq 0$;
7. 0 is not an eigenvalue of A;
8. $\sigma_{\min}(A) > 0$.

Elementary matrices. Each elementary row operation (swap two rows, scale a row, add a multiple of one row to another) is left multiplication by the matrix obtained by applying that operation to I; the corresponding column operation is right multiplication. Gaussian elimination is therefore a sequence of left multiplications, which is the starting point of LU factorization (Lecture 3). A *permutation matrix* P, a row-permuted identity, satisfies $P^{-1} = P^{\top}$.

Triangular matrices. Products and inverses of lower (upper) triangular matrices are lower (upper) triangular, and if the diagonal entries are all 1 they stay 1. A triangular matrix is invertible if and only if all diagonal entries are nonzero, and its eigenvalues are its diagonal entries.

A.4 Determinants

- For triangular A, $\det A = \prod_i a_{ii}$.
- Row operations: swapping two rows flips the sign, scaling a row by c scales $\det A$ by c, and adding a multiple of one row to another leaves $\det A$ unchanged. Hence if $PA = LU$ with unit lower triangular L, then $\det A = \pm \prod_i u_{ii}$, which is how determinants are computed in practice.
- $\det(AB) = \det A \det B$, $\det A^{-1} = 1/\det A$, $\det(cA) = c^n \det A$ for $A \in \mathbb{K}^{n\times n}$, and $|\det Q| = 1$ for unitary Q. In general $\det(A + B) \neq \det A + \det B$.
- For block triangular matrices, $\det \left[\begin{smallmatrix} A_{11} & A_{12} \\ 0 & A_{22} \end{smallmatrix}\right] = \det A_{11} \det A_{22}$.
- $\det A = \prod_i \lambda_i$ and $\operatorname{tr} A = \sum_i \lambda_i$, with eigenvalues counted with algebraic multiplicity.

A.5 Subspaces and Rank

For $A \in \mathbb{K}^{m\times n}$, the *range* (column space) $\mathcal{R}(A) = \{A\vec{x}\} \subseteq \mathbb{K}^m$ and the *null space* $\mathcal{N}(A) = \{\vec{x} : A\vec{x} = \vec{0}\} \subseteq \mathbb{K}^n$ are subspaces. A *basis* of a subspace is a linearly independent spanning set; all bases have the same size, the *dimension*.

- $\operatorname{rank} A = \dim \mathcal{R}(A)$, and column rank equals row rank, so $\operatorname{rank} A \le \min(m, n)$.
- *Rank–nullity*: $\operatorname{rank} A + \dim \mathcal{N}(A) = n$.
- $\mathcal{R}(A)^\perp = \mathcal{N}(A^*)$, so $\mathbb{K}^m = \mathcal{R}(A) \oplus \mathcal{N}(A^*)$ as an orthogonal direct sum.
- $\mathcal{N}(A^*A) = \mathcal{N}(A)$, hence $\operatorname{rank} A^*A = \operatorname{rank} A$, and A^*A is invertible if and only if A has linearly independent columns (Lecture 8).
- $\operatorname{rank}(AB) \le \min(\operatorname{rank} A, \operatorname{rank} B)$, and multiplying by an invertible matrix does not change the rank.

A.6 Eigenvalues and Diagonalization

λ is an *eigenvalue* of $A \in \mathbb{K}^{n\times n}$ with *eigenvector* $\vec{x} \neq \vec{0}$ if $A\vec{x} = \lambda\vec{x}$. The eigenvalues are the roots of the characteristic polynomial $\det(\lambda I - A)$, so there are exactly n of them in $\mathbb{C}$, counted with multiplicity.

- A real matrix may have complex eigenvalues; they come in conjugate pairs $\lambda, \bar{\lambda}$, with conjugate eigenvectors.
- If $A\vec{x} = \lambda\vec{x}$, then $A^k\vec{x} = \lambda^k\vec{x}$, $(A - \sigma I)\vec{x} = (\lambda - \sigma)\vec{x}$, and $A^{-1}\vec{x} = \lambda^{-1}\vec{x}$ for invertible A. These are the facts behind the power iteration, shifts, and inverse iteration (Lecture 10).
- *Similar* matrices $B = S^{-1}AS$ have the same characteristic polynomial, hence the same eigenvalues; eigenvectors transform as $\vec{x} \mapsto S^{-1}\vec{x}$. Row operations alone are *not* similarity transformations and change the eigenvalues, which is why eigenvalue algorithms transform $A \mapsto Q^*AQ$ from both sides.
- Eigenvectors belonging to distinct eigenvalues are linearly independent.
- The *algebraic multiplicity* of λ is its multiplicity as a root of the characteristic polynomial, and the *geometric multiplicity* is $\dim \mathcal{N}(A - \lambda I)$; always $1 \le$ geometric $\le$ algebraic.
- A is *diagonalizable*, $A = X\Lambda X^{-1}$, if and only if it has n linearly independent eigenvectors, if and only if the two multiplicities agree for every eigenvalue. Then $A^k = X\Lambda^k X^{-1}$. A matrix with n distinct eigenvalues is diagonalizable; a matrix that is not is called *defective* (Lecture 11).

A.7 Orthogonality and Projections

- A set of nonzero, mutually orthogonal vectors is linearly independent. In an orthonormal basis $\vec{q}_1, \ldots, \vec{q}_n$ the coordinates are inner products: $\vec{y} = \sum_j (\vec{q}_j^*\vec{y})\vec{q}_j$.
- If $Q \in \mathbb{K}^{m\times n}$ has orthonormal columns, then $Q^*Q = I_n$ and $\|Q\vec{x}\|_2 = \|\vec{x}\|_2$. For $m > n$, however, $QQ^* \neq I_m$: it is the orthogonal projector onto $\mathcal{R}(Q)$.
- *Projection onto a line*: the projection of $\vec{y}$ onto $\operatorname{span}\{\vec{a}\}$ is $\frac{\vec{a}^*\vec{y}}{\vec{a}^*\vec{a}}\vec{a}$.
- *Orthogonal decomposition and best approximation*: for a subspace W, every $\vec{y}$ splits uniquely as $\vec{y} = \hat{\vec{y}} + \vec{z}$ with $\hat{\vec{y}} \in W$ and $\vec{z} \perp W$, and $\hat{\vec{y}}$ is the point of W closest to $\vec{y}$. If the columns of Q form an orthonormal basis of W, then $\hat{\vec{y}} = QQ^*\vec{y}$.
- *Gram–Schmidt* turns linearly independent columns into orthonormal ones and yields $A = QR$ (Lectures 5 and 7).

- *Least squares*: $\min_{\vec{x}} \left\| A\vec{x} - \vec{b} \right\|_2$ is attained exactly when $A^* A\vec{x} = A^*\vec{b}$; with $A = QR$ this becomes $R\vec{x} = Q^*\vec{b}$ (Lectures 5 and 8).

A.8 Hermitian Matrices, Quadratic Forms, and the SVD

- *Spectral theorem* (Theorem 69): a Hermitian A has real eigenvalues and an orthonormal basis of eigenvectors, $A = Q\Lambda Q^*$ with Q unitary and Λ real diagonal. Equivalently $A = \sum_i \lambda_i \vec{q}_i \vec{q}_i^*$ (spectral decomposition).
- A Hermitian A is *positive definite* if $\vec{x}^* A\vec{x} > 0$ for all $\vec{x} \neq \vec{0}$, equivalently if all its eigenvalues are positive; *semidefinite* versions use $\geq$. *Negative* (semi)definite is defined in the same way. A is *indefinite* if $\vec{x}^* A\vec{x}$ takes both signs, equivalently if it has both positive and negative eigenvalues. For example, $\left[\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right]$ is indefinite: for real $\vec{x}$, $\vec{x}^\top A\vec{x} = 2x_1x_2$ is positive along $(1, 1)$ and negative along $(1, -1)$. Changing variables $\vec{x} = Q\vec{y}$ turns the quadratic form into $\sum_i \lambda_i |y_i|^2$ (principal axes theorem).
- On the unit sphere, $\vec{x}^* A\vec{x}$ ranges over $[\lambda_{\min}, \lambda_{\max}]$, attained at the corresponding eigenvectors (Definition 54).
- *SVD*: every $A \in \mathbb{K}^{m\times n}$ factors as $A = U\Sigma V^*$ with U, V unitary and Σ diagonal with entries $\sigma_1 \geq \cdots \geq 0$. The σ_i^2 are the eigenvalues of A^*A, the columns $\vec{v}_i$ of V are its orthonormal eigenvectors, and $\vec{u}_i = A\vec{v}_i/\sigma_i$ for $\sigma_i > 0$. The number of nonzero σ_i is $\operatorname{rank} A$ (Section B.3).

Appendix B

Norms and Unitary Matrices

This appendix collects facts about unitary matrices and matrix norms that are used throughout the course, starting from Lecture 6. We begin with vector and matrix norms in general, following [1, Lecture 3], and then specialize to the 2-norm and the Frobenius norm. Supplementary reading:

- [1] Lectures 2, 3, and 5.
- [2] Sections 3.2.3 and 3.4.

B.1 Vector and Matrix Norms

Norms are the yardsticks of numerical linear algebra: every statement about the size of an error, the accuracy of an approximation, or the convergence of an iteration is made in some norm.

B.1.1 Vector Norms

Definition 98 (Vector Norm)

A *norm* on $\mathbb{K}^m$ is a function $\|\cdot\| : \mathbb{K}^m \to \mathbb{R}$ such that for all $\vec{x}, \vec{y} \in \mathbb{K}^m$ and all $\alpha \in \mathbb{K}$:

1. $\|\vec{x}\| \geq 0$, and $\|\vec{x}\| = 0$ only if $\vec{x} = \vec{0}$;
2. $\|\vec{x} + \vec{y}\| \leq \|\vec{x}\| + \|\vec{y}\|$ (the *triangle inequality*);
3. $\|\alpha\vec{x}\| = |\alpha| \, \|\vec{x}\|$.

In words, a nonzero vector has positive length, the length of a sum is at most the sum of the lengths, and scaling a vector scales its length by the same amount. The most important norms are the p-norms.


Definition 99 (p-Norms)

For $\vec{x} \in \mathbb{K}^m$,

$$\|\vec{x}\|_1 = \sum_{i=1}^{m} |x_i|\,, \qquad \|\vec{x}\|_2 = \left(\sum_{i=1}^{m} |x_i|^2\right)^{1/2} = \sqrt{\vec{x}^*\vec{x}}, \qquad \|\vec{x}\|_\infty = \max_{1\le i\le m} |x_i|\,, \tag{B.1}$$

and more generally $\|\vec{x}\|_p = (\sum_{i=1}^m |x_i|^p)^{1/p}$ for $1 \le p < \infty$.

A norm is determined by its closed unit ball $\{\vec{x} : \|\vec{x}\| \le 1\}$. For $m = 2$, this gives the following picture.



Unit balls of $\mathbb{R}^2$ in the p-norms. As p grows, the ball grows from the diamond ($p = 1$) to the square ($p = \infty$).

The 2-norm is the Euclidean length, and its unit ball is round. Since the balls are nested, the p-norms decrease as p increases; Corollary 107 below makes this precise, with constants depending on the dimension.

Aside from the p-norms, the most useful norms are the *weighted norms*. Given any norm $\|\cdot\|$ and a diagonal matrix $W = \operatorname{diag}(w_1, \ldots, w_m)$ with all $w_i \ne 0$, we set

$$\|\vec{x}\|_W := \|W\vec{x}\|\,. \tag{B.2}$$

For example, the weighted 2-norm is $\|\vec{x}\|_W = \left(\sum_i |w_i x_i|^2\right)^{1/2}$, and its unit ball is an axis-aligned ellipse with semiaxes $1/|w_i|$. More generally, $\|W\vec{x}\|$ is a norm for *any* nonsingular W: properties 2 and 3 are inherited from $\|\cdot\|$ because W is linear, and $\|W\vec{x}\| = 0$ forces $W\vec{x} = \vec{0}$, hence $\vec{x} = \vec{0}$. Weighted norms are natural when the entries of $\vec{x}$ are measured in different units.

B.1.2 Induced Matrix Norms

An $m \times n$ matrix can be viewed as a vector in an mn-dimensional space, so any vector norm on $\mathbb{K}^{mn}$ measures the size of a matrix. More useful, however, are norms that measure what A *does* as a map from $\mathbb{K}^n$ to $\mathbb{K}^m$.

Definition 100 (Induced Matrix Norm)

Let $\|\cdot\|_{(n)}$ and $\|\cdot\|_{(m)}$ be norms on $\mathbb{K}^n$ and $\mathbb{K}^m$. The *induced matrix norm* of $A \in \mathbb{K}^{m\times n}$ is the smallest C such that $\|A\vec{x}\|_{(m)} \le C\,\|\vec{x}\|_{(n)}$ for all $\vec{x} \in \mathbb{K}^n$, that is,

$$\|A\|_{(m,n)} := \sup_{\vec{x}\ne\vec{0}} \frac{\|A\vec{x}\|_{(m)}}{\|\vec{x}\|_{(n)}} = \sup_{\|\vec{x}\|_{(n)}=1} \|A\vec{x}\|_{(m)}\,. \tag{B.3}$$


When both norms are p-norms, we write $\|A\|_p$.

So $\|A\|_{(m,n)}$ is the largest factor by which A can stretch a vector. The two expressions agree by property 3 of Definition 98: scaling $\vec{x}$ does not change the ratio, so it suffices to look at unit vectors. In particular, $\|A\|$ is determined by the image of the unit ball under A, which makes induced norms easy to visualize.

Example 101 (A 2×2 Matrix in Three Norms)**.** Take

$$A = \begin{bmatrix} 1 & 2 \\ 0 & 2 \end{bmatrix}. \tag{B.4}$$

Since A is real, we may draw it as a map of $\mathbb{R}^2$. It sends $\vec{e}_1$ to its first column $(1,0)^\top$ and $\vec{e}_2$ to its second column $(2,2)^\top$. Mapping the unit balls of the 1-, 2-, and ∞-norms gives the following picture.



Left: unit balls of $\mathbb{R}^2$. Right: their images under A. The dashed vectors are the ones that A stretches the most in each norm.

In the 1-norm, the most amplified unit vector is $\vec{e}_2$, which is mapped to $(2,2)^\top$ with 1-norm 4. In the ∞-norm, it is $(1,1)^\top$, which is mapped to $(3,2)^\top$ with ∞-norm 3. In the 2-norm, the image of the unit circle is an ellipse, and $\|A\|_2$ is the length of its longest semiaxis, about 2.9208. This must be at least $\|A\vec{e}_2\|_2 = \sqrt{8} \approx 2.8284$; the exact value is $\sigma_{\max}(A)$ by Theorem 113.

Example 102 (The p-Norm of a Diagonal Matrix)**.** Let $D = \operatorname{diag}(d_1, \ldots, d_m)$. For $1 \le p < \infty$,

$$\|D\vec{x}\|_p^p = \sum_{i=1}^{m} |d_i x_i|^p \le \left(\max_i |d_i|\right)^p \|\vec{x}\|_p^p, \tag{B.5}$$

with equality for $\vec{x} = \vec{e}_i$ where i maximizes $|d_i|$; the case $p = \infty$ is the same argument with sums replaced by maxima. Hence $\|D\|_p = \max_i |d_i|$ for every p. Geometrically, D maps the 2-norm unit sphere to an ellipsoid with semiaxes $|d_i|$, and $\|D\|_2$ is the longest semiaxis. The SVD shows that *every* matrix maps the unit sphere to an ellipsoid; only the axes may be rotated.

For $p = 1$ and $p = \infty$, the induced norm has a closed form in terms of the entries.

Proposition 103 (The 1- and ∞-Norms of a Matrix)

Let $A \in \mathbb{K}^{m\times n}$ have columns $\vec{a}_1, \dots, \vec{a}_n$ and rows $\vec{\alpha}_1^*, \dots, \vec{\alpha}_m^*$. Then

$$\|A\|_1 = \max_{1\le j\le n} \|\vec{a}_j\|_1 \quad \text{(maximum column sum)}, \qquad \|A\|_\infty = \max_{1\le i\le m} \|\vec{\alpha}_i\|_1 \quad \text{(maximum row sum)}. \tag{B.6}$$

Proof. For the 1-norm, let $\|\vec{x}\|_1 = 1$. By the triangle inequality,

$$\|A\vec{x}\|_1 = \left\| \sum_{j=1}^n x_j \vec{a}_j \right\|_1 \le \sum_{j=1}^n |x_j| \, \|\vec{a}_j\|_1 \le \max_j \|\vec{a}_j\|_1 , \tag{B.7}$$

since the weights $|x_j|$ sum to 1. Taking $\vec{x} = \vec{e}_j$ for the column j with the largest 1-norm attains the bound.

For the ∞-norm, let $\|\vec{x}\|_\infty = 1$. Each entry of $A\vec{x}$ satisfies $|(A\vec{x})_i| \le \sum_j |A_{ij}| \, |x_j| \le \|\vec{\alpha}_i\|_1$, so $\|A\vec{x}\|_\infty \le \max_i \|\vec{\alpha}_i\|_1$. To attain it, pick the row i with the largest 1-norm and set $x_j = \overline{A_{ij}} / |A_{ij}|$ (and $x_j = 1$ if $A_{ij} = 0$). Then $\|\vec{x}\|_\infty = 1$ and $(A\vec{x})_i = \sum_j |A_{ij}| = \|\vec{\alpha}_i\|_1$. □

In the example above, the column sums are 1 and 4 and the row sums are 3 and 2, matching the picture. Both norms cost only $O(mn)$ operations, whereas $\|A\|_2$ requires the largest singular value. Computing $\|A\|_p$ for other p is harder still.

B.1.3 Cauchy–Schwarz and Hölder Inequalities

To bound matrix norms when $p \neq 1, \infty$, we bound inner products by p-norms.

Theorem 104 (Hölder and Cauchy–Schwarz Inequalities)

Let $1 \le p, q \le \infty$ with $1/p + 1/q = 1$. Then for all $\vec{x}, \vec{y} \in \mathbb{K}^m$,

$$|\vec{x}^* \vec{y}| \le \|\vec{x}\|_p \, \|\vec{y}\|_q . \tag{B.8}$$

The case $p = q = 2$ is the *Cauchy–Schwarz inequality* $|\vec{x}^*\vec{y}| \le \|\vec{x}\|_2 \, \|\vec{y}\|_2$.

Both bounds are tight: for every $\vec{x}$ there is a $\vec{y} \neq \vec{0}$ attaining equality. For Cauchy–Schwarz, $\vec{y} = \vec{x}$ works. Proofs can be found in any linear algebra text. These inequalities immediately give the 2-norm of two simple kinds of matrices.

Example 105 (The 2-Norm of a Row Vector)**.** Let $A = \vec{a}^*$ be a single row. By Cauchy–Schwarz, $\|A\vec{x}\|_2 = |\vec{a}^*\vec{x}| \le \|\vec{a}\|_2 \, \|\vec{x}\|_2$, with equality for $\vec{x} = \vec{a}$. Hence $\|\vec{a}^*\|_2 = \|\vec{a}\|_2$.

Example 106 (The 2-Norm of an Outer Product)**.** Let $A = \vec{u}\vec{v}^*$ with $\vec{u} \in \mathbb{K}^m$ and $\vec{v} \in \mathbb{K}^n$. For every $\vec{x} \in \mathbb{K}^n$,

$$\|A\vec{x}\|_2 = \|\vec{u}(\vec{v}^*\vec{x})\|_2 = \|\vec{u}\|_2 \, |\vec{v}^*\vec{x}| \le \|\vec{u}\|_2 \, \|\vec{v}\|_2 \, \|\vec{x}\|_2 , \tag{B.9}$$

with equality for $\vec{x} = \vec{v}$. Hence $\|\vec{u}\vec{v}^*\|_2 = \|\vec{u}\|_2 \, \|\vec{v}\|_2$. The Frobenius norm gives the same value: $\|\vec{u}\vec{v}^*\|_F^2 = \sum_{i,j} |u_i|^2 \, |v_j|^2 = \|\vec{u}\|_2^2 \, \|\vec{v}\|_2^2$, consistent with item 3 of Proposition 115 for $r = 1$.

Cauchy–Schwarz also lets us compare the p-norms in a fixed dimension.

Corollary 107 (Comparing p-Norms)**.** *For $\vec{x} \in \mathbb{K}^m$,*

$$\|\vec{x}\|_\infty \le \|\vec{x}\|_2 \le \|\vec{x}\|_1 \le \sqrt{m}\, \|\vec{x}\|_2 \le m\, \|\vec{x}\|_\infty , \tag{B.10}$$

and for $A \in \mathbb{K}^{m\times n}$,

$$\frac{1}{\sqrt{n}} \|A\|_\infty \le \|A\|_2 \le \sqrt{m}\, \|A\|_\infty , \qquad \frac{1}{\sqrt{m}} \|A\|_1 \le \|A\|_2 \le \sqrt{n}\, \|A\|_1 . \tag{B.11}$$

Proof. The first two inequalities hold because $\max_i |x_i|^2 \le \sum_i |x_i|^2 \le (\sum_i |x_i|)^2$. The third is Cauchy–Schwarz applied to the vector of the $|x_i|$ and the all-ones vector, and the fourth is $\sum_i |x_i|^2 \le m \max_i |x_i|^2$.

For the matrix bounds, chain the vector bounds through the definition. For instance,

$$\|A\vec{x}\|_2 \le \sqrt{m}\, \|A\vec{x}\|_\infty \le \sqrt{m}\, \|A\|_\infty \|\vec{x}\|_\infty \le \sqrt{m}\, \|A\|_\infty \|\vec{x}\|_2 \tag{B.12}$$

gives $\|A\|_2 \le \sqrt{m}\, \|A\|_\infty$, and $\|A\vec{x}\|_\infty \le \|A\vec{x}\|_2 \le \|A\|_2 \|\vec{x}\|_2 \le \sqrt{n}\, \|A\|_2 \|\vec{x}\|_\infty$ gives the other bound. The 1-norm bounds follow in the same way. □

In particular, all p-norms agree up to factors that depend only on the dimension. Such factors are harmless in $O(\varepsilon_{\text{mach}})$ statements, so a method that is stable in one p-norm is stable in all of them.

B.1.4 Bounding $\|AB\|$

Proposition 108 (Submultiplicativity of Induced Norms)

Let $\|\cdot\|_{(\ell)}$, $\|\cdot\|_{(m)}$, $\|\cdot\|_{(n)}$ be norms on $\mathbb{K}^\ell$, $\mathbb{K}^m$, $\mathbb{K}^n$, and let $A \in \mathbb{K}^{\ell\times m}$ and $B \in \mathbb{K}^{m\times n}$. Then

$$\|AB\|_{(\ell,n)} \le \|A\|_{(\ell,m)}\, \|B\|_{(m,n)} . \tag{B.13}$$

Proof. Apply the definition twice: $\|AB\vec{x}\|_{(\ell)} \le \|A\|_{(\ell,m)}\, \|B\vec{x}\|_{(m)} \le \|A\|_{(\ell,m)}\, \|B\|_{(m,n)}\, \|\vec{x}\|_{(n)}$. □

In particular, $\left\|A^k\right\| \le \|A\|^k$ for a square matrix in any induced norm. This is generally not an equality. For example, $A = \left[\begin{smallmatrix} 0 & 1 \\ 0 & 0 \end{smallmatrix}\right]$ has $\|A\|_p = 1$ for every p, but $A^2 = 0$.

A related bound concerns eigenvalues. If $A\vec{x} = \lambda\vec{x}$ with $\vec{x} \ne \vec{0}$, then $|\lambda|\, \|\vec{x}\| = \|A\vec{x}\| \le \|A\|\, \|\vec{x}\|$. So the *spectral radius* $\rho(A) \coloneqq \max\{|\lambda| : \lambda \text{ an eigenvalue of } A\}$ satisfies

$$\rho(A) \le \|A\| \tag{B.14}$$

in every induced norm. The same example shows that the inequality can be strict: $\rho(A) = 0$ but $\|A\| = 1$.

B.1.5 General Matrix Norms

Matrix norms do not have to be induced by vector norms. In general, a *matrix norm* is any function on $\mathbb{K}^{m\times n}$ that satisfies the three conditions of Definition 98 on this mn-dimensional vector space:

$$\|A\| \ge 0 \text{ with equality only if } A = 0, \qquad \|A + B\| \le \|A\| + \|B\| , \qquad \|\alpha A\| = |\alpha|\, \|A\| . \tag{B.15}$$

The most important matrix norm that is not induced by a vector norm is the Frobenius norm, the subject of Section B.4. Finally, the 2-norm and the Frobenius norm are both invariant under multiplication by unitary matrices; this is Theorem 110 below.

B.2 Unitary Transformations

The error analysis of Lecture 6 uses one property of unitary matrices: they do not change the length of a vector. In this section we collect this and a few related facts, which we also use in Lectures 5 and 7.

Definition 109 (Unitary and Orthogonal Matrices)

We say that a square matrix $Q \in \mathbb{K}^{m\times m}$ is *unitary* if $Q^*Q = I$. When $\mathbb{K} = \mathbb{R}$, this condition reads $Q^\top Q = I$, and we call Q *orthogonal*. We say that a tall matrix $\hat{Q} \in \mathbb{K}^{m\times n}$ has *orthonormal columns* if $\hat{Q}^*\hat{Q} = I_n$.

The condition $Q^*Q = I$ just says that the columns of Q are orthonormal. Since Q is square, it also gives $Q^{-1} = Q^*$ and $QQ^* = I$. In particular, inverting a unitary matrix costs nothing: the solution of $Q\vec{x} = \vec{b}$ is $\vec{x} = Q^*\vec{b}$.

Theorem 110 (Unitary Matrices Preserve the 2-Norm)

Let $Q \in \mathbb{K}^{m\times m}$ be unitary. Then for all $\vec{x}, \vec{y} \in \mathbb{K}^m$,

$$\langle Q\vec{x}, Q\vec{y}\rangle = \langle \vec{x}, \vec{y}\rangle\,, \qquad \|Q\vec{x}\|_2 = \|\vec{x}\|_2\,. \tag{B.16}$$

As a consequence, for any $A \in \mathbb{K}^{m\times n}$ and unitary $U \in \mathbb{K}^{m\times m}$, $V \in \mathbb{K}^{n\times n}$:

1. $\|UAV\|_2 = \|A\|_2$ and $\|UAV\|_F = \|A\|_F$.
2. UAV has the same singular values as A. In particular, $\kappa_2(UAV) = \kappa_2(A)$.
3. $\|Q\|_2 = \|Q^{-1}\|_2 = 1$ and $\kappa_2(Q) = 1$. All singular values of Q are 1, every eigenvalue of Q satisfies $|\lambda| = 1$, and $|\det Q| = 1$.

Proof. We have $\langle Q\vec{x}, Q\vec{y}\rangle = \vec{x}^*Q^*Q\vec{y} = \vec{x}^*\vec{y}$. Taking $\vec{y} = \vec{x}$ gives $\|Q\vec{x}\|_2 = \|\vec{x}\|_2$.

For item 1, the norm identity gives $\|UA\vec{x}\|_2 = \|A\vec{x}\|_2$ for every $\vec{x}$, so $\|UA\|_2 = \|A\|_2$. For the right factor, note that $\vec{y} = V\vec{x}$ runs over all unit vectors as $\vec{x}$ does, so

$$\|AV\|_2 = \sup_{\|\vec{x}\|_2=1} \|AV\vec{x}\|_2 = \sup_{\|\vec{y}\|_2=1} \|A\vec{y}\|_2 = \|A\|_2\,. \tag{B.17}$$

The statement for the Frobenius norm is item 1 of Proposition 115.

For item 2, let $A = W\Sigma Z^*$ be an SVD. Then $UAV = (UW)\Sigma(V^*Z)^*$. Products of unitary matrices are unitary, so this is an SVD of UAV with the same Σ.

For item 3, $Q = Q \cdot I \cdot I^*$ is an SVD with $\Sigma = I$. If $Q\vec{x} = \lambda\vec{x}$ with $\vec{x} \neq \vec{0}$, then $\|\vec{x}\|_2 = \|Q\vec{x}\|_2 = |\lambda|\,\|\vec{x}\|_2$, so $|\lambda| = 1$. Finally, $|\det Q|^2 = \det(Q^*Q) = \det I = 1$. □

The converse is also true: if $\|Q\vec{x}\|_2 = \|\vec{x}\|_2$ for every $\vec{x}$, then Q is unitary. (One recovers inner products from norms by the polarization identity.) In other words, unitary matrices are exactly the linear maps that preserve lengths. Geometrically, they are rotations and reflections.

B.2.1 Examples

Several matrices we have already seen are unitary.

- *Householder reflectors* $F = I - 2\vec{v}\vec{v}^*/(\vec{v}^*\vec{v})$. They are Hermitian, satisfy $F^2 = I$, and have $\det F = -1$.
- *Givens rotations* (Lecture 7). They have $\det = 1$ and change only two coordinates.
- *Permutation matrices*. So the row swaps of partial pivoting (Lecture 4) do not change any norms, and all of the instability of LU comes from the triangular factors.
- *Diagonal matrices* $D = \operatorname{diag}(d_1, \dots, d_m)$ with $|d_i| = 1$. These describe exactly the freedom in the QR factorization, since $A = (QD)(D^*R)$. The condition $r_{jj} > 0$ removes this freedom (Theorem 33).

B.2.2 Tall Matrices with Orthonormal Columns

Now let $\hat{Q} \in \mathbb{K}^{m\times n}$ with $m > n$ have orthonormal columns. The same one-line proof shows that $\left\|\hat{Q}\vec{x}\right\|_2 = \|\vec{x}\|_2$, and hence, as in item 1 of Theorem 110, $\left\|\hat{Q}B\right\|_2 = \|B\|_2$ and $\left\|\hat{Q}B\right\|_F = \|B\|_F$ for every $B \in \mathbb{K}^{n\times p}$. Taking conjugate transposes, the same holds for multiplication on the right by a matrix with orthonormal rows. However, $\hat{Q}^*$ does *not* preserve lengths. The matrix $\hat{Q}\hat{Q}^*$ is not the identity; it is the orthogonal projector onto $\mathcal{R}(\hat{Q})$. So $\left\|\hat{Q}^*\vec{y}\right\|_2 \le \|\vec{y}\|_2$, with equality only if $\vec{y} \in \mathcal{R}(\hat{Q})$.

This is exactly what makes QR useful for least squares. Take a full QR factorization $A = QR = \begin{bmatrix}\hat{Q} & \hat{Q}_\perp\end{bmatrix}\begin{bmatrix}\hat{R}\\ 0\end{bmatrix}$. Since Q^* preserves lengths, we can apply it inside the norm:

$$\left\|A\vec{x} - \vec{b}\right\|_2^2 = \left\|Q^*(A\vec{x} - \vec{b})\right\|_2^2 = \left\|\hat{R}\vec{x} - \hat{Q}^*\vec{b}\right\|_2^2 + \left\|\hat{Q}_\perp^*\vec{b}\right\|_2^2. \tag{B.18}$$

The second term does not depend on $\vec{x}$. So the minimizer solves $\hat{R}\vec{x} = \hat{Q}^*\vec{b}$, and the minimal residual is $\left\|\hat{Q}_\perp^*\vec{b}\right\|_2$ (compare Lecture 5).

B.2.3 Why Unitary Transformations Are Good for Computation

1. *One step does not amplify errors.* In Theorem 26 we have $\kappa(Q_k) = 1$, so each step has relative backward error $O(\varepsilon_{\text{mach}})$ with no extra factor.
2. *Many steps do not amplify errors.* A product of unitary matrices is unitary. So when we pull an error back through all the previous steps, its size does not change. The backward errors of n steps add up to $O(n\varepsilon_{\text{mach}})$ instead of multiplying (see Section 6.7).
3. *No element growth.* Every intermediate matrix has the same 2-norm as A, so there is nothing like the growth factor of LU. Compare this with Example 27: a triangular matrix with entries bounded by 1 can still have an exponentially large inverse, while $\left\|Q^{-1}\right\|_2 = \|Q\|_2 = 1$ always.
4. *Errors look the same before and after.* An error of size δ in $\vec{x}$ is an error of size δ in $Q\vec{x}$. So we can move errors between the input and the output without losing anything.

Note that this invariance holds for the 2-norm and the Frobenius norm, but not for other norms. For example, rotating $\vec{e}_1$ by $45°$ gives $(1/\sqrt{2}, 1/\sqrt{2})^\top$, whose ∞-norm is $1/\sqrt{2}$ and whose 1-norm is $\sqrt{2}$. This is why stability results for orthogonal methods are stated in the 2-norm. In other norms they still hold, but with extra constants that depend on the dimension.

As a final note, everything above assumes that the transformation is *exactly* unitary. A matrix that we compute column by column, such as the $\hat{Q}$ of modified Gram–Schmidt, is only unitary up to $O(\varepsilon_{\text{mach}}\kappa(A))$, and it

loses these advantages to the same extent. Householder's Q is different. Each reflector built from a computed vector $\tilde{\vec{v}}_k$ is exactly unitary, so rounding errors only change *which* unitary matrix we get, not *whether* it is unitary (Lecture 7).

B.3 The 2-Norm, Singular Values, and Condition Numbers

Most error bounds in this course are stated in the 2-norm. Here we record how it is expressed through the SVD; these facts are proved in Homework 1, Problem 3.

Definition 111 (Full and Reduced SVD)

Let $A \in \mathbb{K}^{m\times n}$ and $p = \min(m,n)$. A *full SVD* is a factorization

$$A = U\Sigma V^*, \qquad U \in \mathbb{K}^{m\times m}, \quad \Sigma \in \mathbb{R}^{m\times n}, \quad V \in \mathbb{K}^{n\times n}, \tag{B.19}$$

with U and V unitary and Σ "diagonal" of the same shape as A: only $\Sigma_{11}, \dots, \Sigma_{pp}$ can be nonzero, and these are the *singular values* $\sigma_1 \geq \cdots \geq \sigma_p \geq 0$. Dropping the zero blocks of Σ together with the columns of U or V they multiply gives the *reduced SVD*

$$A = \hat{U}\hat{\Sigma}\hat{V}^*, \qquad \hat{U} \in \mathbb{K}^{m\times p}, \quad \hat{\Sigma} = \operatorname{diag}(\sigma_1, \dots, \sigma_p) \in \mathbb{R}^{p\times p}, \quad \hat{V} \in \mathbb{K}^{n\times p}, \tag{B.20}$$

where $\hat{U}$ and $\hat{V}$ have orthonormal columns.

In the full SVD, Σ has a zero block below or beside its square diagonal part $D = \operatorname{diag}(\sigma_1, \dots, \sigma_p)$:

$$\Sigma = \begin{bmatrix} D \\ 0 \end{bmatrix} \ (m > n), \qquad \Sigma = \begin{bmatrix} D & 0 \end{bmatrix} \ (m < n). \tag{B.21}$$

The reduced form is more economical for computation, while the full form is more convenient in proofs, because square unitary U and V preserve norms in both directions (Theorem 110). Every matrix has an SVD, and the singular values are unique [1, Theorem 4.1].

Definition 112 (Induced 2-Norm)

The 2-*norm* of $A \in \mathbb{K}^{m\times n}$ is the matrix norm induced by the vector 2-norm (Definition 100),

$$\|A\|_2 \coloneqq \sup_{\vec{x}\neq\vec{0}} \frac{\|A\vec{x}\|_2}{\|\vec{x}\|_2}. \tag{B.22}$$

Theorem 113 (The 2-Norm Through the SVD)

Let $A \in \mathbb{K}^{m\times n}$ have singular values $\sigma_{\max} = \sigma_1 \geq \cdots \geq \sigma_{\min(m,n)} = \sigma_{\min}$. Then:

1. $\|A\|_2 = \sigma_{\max}$.
2. If $m \geq n$, then $\|A\vec{x}\|_2 \geq \sigma_{\min} \|\vec{x}\|_2$ for all $\vec{x}$, with equality for the last right singular vector.
3. If A is square and invertible, then $\left\|A^{-1}\right\|_2 = 1/\sigma_{\min}$, and hence

$$\kappa_2(A) \coloneqq \|A\|_2 \left\|A^{-1}\right\|_2 = \frac{\sigma_{\max}}{\sigma_{\min}}. \tag{B.23}$$

Proof. Take an SVD $A = U\Sigma V^*$ and put $\vec{y} = V^*\vec{x}$. Since U and V are unitary (Theorem 110), $\|A\vec{x}\|_2 = \|\Sigma\vec{y}\|_2$ and $\|\vec{y}\|_2 = \|\vec{x}\|_2$. Now

$$\|\Sigma\vec{y}\|_2^2 = \sum_{i=1}^{\min(m,n)} \sigma_i^2 |y_i|^2 . \tag{B.24}$$

The sum stops at $\min(m,n)$, not at n: for $m < n$ the last $n-m$ entries of $\vec{y}$ are multiplied by the zero block of Σ and do not contribute. In particular a wide matrix has a nontrivial null space, which is why item 2 needs $m \geq n$. This is at most $\sigma_{\max}^2 \|\vec{y}\|_2^2$, with equality for $\vec{y} = \vec{e}_1$, which gives item 1. If $m \geq n$, the sum runs over all n entries of $\vec{y}$, so it is at least $\sigma_{\min}^2 \|\vec{y}\|_2^2$, with equality for $\vec{y} = \vec{e}_n$; this gives item 2. For item 3, $A^{-1} = V\Sigma^{-1}U^*$ is an SVD up to reordering, so the singular values of A^{-1} are the $1/\sigma_i$, and the largest of them is $1/\sigma_{\min}$. □

For a tall matrix with full column rank we *define* $\kappa_2(A) \coloneqq \sigma_{\max}/\sigma_{\min}$; this is the condition number that governs least squares (Lecture 5). In both cases, $\kappa_2(A)$ is the worst-case condition number of the map $\vec{x} \mapsto A\vec{x}$ from Lecture 1:

$$\sup_{\vec{x}\neq\vec{0}} \frac{\|A\|_2 \|\vec{x}\|_2}{\|A\vec{x}\|_2} = \frac{\sigma_{\max}}{\sigma_{\min}}, \tag{B.25}$$

since $\|A\vec{x}\|_2 \,/\, \|\vec{x}\|_2$ is smallest, equal to $\sigma_{\min}$, for the last right singular vector. In words, $\kappa_2(A)$ is the ratio of the largest to the smallest stretching factor of A.

B.4 The Frobenius Norm

In Lecture 6 we bounded $\|L\|_2$ through the Frobenius norm. It is often the easiest matrix norm to compute and to estimate.

Definition 114 (Frobenius Norm)

The *Frobenius norm* of $A \in \mathbb{K}^{m\times n}$ is

$$\|A\|_F \coloneqq \left(\sum_{i=1}^{m} \sum_{j=1}^{n} |A_{ij}|^2 \right)^{1/2}. \tag{B.26}$$

In other words, $\|A\|_F$ is the Euclidean norm of A if we stack its columns into one long vector of length mn. Equivalently, $\|A\|_F^2 = \sum_j \|\vec{a}_j\|_2^2$ is the sum of the squared column norms. These squared column norms are

exactly the diagonal entries of A^*A, and the squared row norms are the diagonal entries of AA^*, so

$$\|A\|_F^2 = \operatorname{trace}(A^*A) = \operatorname{trace}(AA^*). \tag{B.27}$$

Proposition 115 (Properties of the Frobenius Norm)

Let $A \in \mathbb{K}^{m\times n}$ have rank r and singular values $\sigma_1 \geq \cdots \geq \sigma_r > 0$. Then:

1. *Unitary invariance.* $\|UAV\|_F = \|A\|_F$ for all unitary $U \in \mathbb{K}^{m\times m}$ and $V \in \mathbb{K}^{n\times n}$.
2. *Singular values.* $\|A\|_F = (\sigma_1^2 + \cdots + \sigma_r^2)^{1/2}$.
3. *Comparison with the 2-norm.* $\|A\|_2 \leq \|A\|_F \leq \sqrt{r}\,\|A\|_2 \leq \sqrt{\min(m,n)}\,\|A\|_2$.
4. *Compatibility and submultiplicativity.* $\|A\vec{x}\|_2 \leq \|A\|_F\,\|\vec{x}\|_2$, and $\|AB\|_F \leq \|A\|_2\,\|B\|_F \leq \|A\|_F\,\|B\|_F$.
5. *Only absolute values matter.* $\|A\|_F = \||A|\|_F$, and if $|A| \leq |B|$ entrywise, then $\|A\|_F \leq \|B\|_F$.

Proof. For item 1, using $U^*U = I$ and the cyclic property of the trace,

$$\|UAV\|_F^2 = \operatorname{trace}(V^*A^*U^*UAV) = \operatorname{trace}(V^*A^*AV) = \operatorname{trace}(A^*A) = \|A\|_F^2\,. \tag{B.28}$$

For item 2, take an SVD $A = U\Sigma V^*$. By item 1, $\|A\|_F = \|\Sigma\|_F = (\sum_i \sigma_i^2)^{1/2}$.

For item 3, use item 2 and $\|A\|_2 = \sigma_1$: we have $\sigma_1^2 \leq \sum_i \sigma_i^2 \leq r\sigma_1^2$.

For item 4, let $\vec{\alpha}_i^*$ be the i-th row of A. By the Cauchy–Schwarz inequality, $|\vec{\alpha}_i^*\vec{x}|^2 \leq \|\vec{\alpha}_i\|_2^2\,\|\vec{x}\|_2^2$. Summing over i gives $\|A\vec{x}\|_2^2 \leq \|A\|_F^2\,\|\vec{x}\|_2^2$. For the product, apply $\left\|A\vec{b}_j\right\|_2 \leq \|A\|_2\left\|\vec{b}_j\right\|_2$ to each column $\vec{b}_j$ of B and sum the squares. The last inequality then follows from $\|A\|_2 \leq \|A\|_F$.

Item 5 follows directly from the definition. □

Note that the Frobenius norm is *not* induced by any vector norm. Every induced norm satisfies $\|I\| = \sup_{\vec{x}\neq\vec{0}} \|\vec{x}\| / \|\vec{x}\| = 1$, but $\|I_n\|_F = \sqrt{n}$. Still, by item 4 it is compatible with the Euclidean norm on vectors.

B.4.1 Why It Is Useful in Error Analysis

- *It is cheap.* Computing $\|A\|_F$ takes $O(mn)$ operations, while $\|A\|_2 = \sigma_1$ requires the SVD (or an estimate of it). By item 3, the two norms differ by at most a factor of $\sqrt{\min(m,n)}$, and such dimension-dependent constants do not matter in $O(\varepsilon_{\text{mach}})$ statements.
- *It turns entrywise bounds into norm bounds.* The backward error of LU from Lecture 4 satisfies $|E| \leq m\varepsilon_{\text{mach}}\,|L|\,|U|$. By items 5 and 4,

$$\|E\|_F \leq m\varepsilon_{\text{mach}}\,\||L|\,|U|\|_F \leq m\varepsilon_{\text{mach}}\,\|L\|_F\,\|U\|_F\,. \tag{B.29}$$

  With partial pivoting, $\|L\|_F \leq m$ (Lecture 6), so everything depends on $\|U\|_F$, i.e., on the growth factor.
- *It is unitarily invariant.* Just like the 2-norm, the Frobenius norm of an error does not change under orthogonal transformations such as Householder reflectors.

B.4.2 Low-Rank Approximation

Since $\|A-B\|_F^2 = \sum_{i,j} |A_{ij} - B_{ij}|^2$ is a sum of squared entrywise errors, approximating a matrix in the Frobenius norm is a least squares problem. The Eckart–Young theorem also holds in this norm: the truncated SVD $A_k = \sum_{i \le k} \sigma_i \vec{u}_i \vec{v}_i^*$ satisfies

$$\min_{\operatorname{rank}(B) \le k} \|A - B\|_F = \|A - A_k\|_F = \left(\sigma_{k+1}^2 + \cdots + \sigma_r^2\right)^{1/2}. \tag{B.30}$$

In the 2-norm, the corresponding error is $\|A - A_k\|_2 = \sigma_{k+1}$ [2, §3.2.3]. We will come back to this when the course covers low-rank approximation.

Appendix C

The Floating-Point Model

This appendix collects background on floating-point arithmetic that is used throughout the course. Supplementary reading:

- [1] Lectures 13 and 14.
- [2] Section 1.5.
- [5] Section 2.4.

C.1 IEEE 754 Formats

The formats in common use are listed below; b_p counts the implicit bit, so a significand field with $b_p - 1$ stored bits provides b_p bits of precision.

Format	Total bits	Exponent bits	Stored significand bits	b_p	Bias	Decimal digits
Float64	64	11	52	53	1023	≈ 15.9
Float32	32	8	23	24	127	≈ 7.2
Float16	16	5	10	11	15	≈ 3.3
BFloat16	16	8	7	8	127	≈ 2.4

The exponent field uses a biased encoding, and its two extreme values are reserved:

- An all-zeros exponent encodes ± 0 (if the significand is also zero) or a *subnormal* number, which drops the implicit leading 1 and uses $0.f \times 2^{E_{\min}}$. Subnormals provide *gradual underflow*, which guarantees $x \neq y \implies x - y \neq 0$.
- An all-ones exponent encodes $\pm\infty$ (zero significand) or `NaN` (nonzero significand).

The normal range of Float64 is therefore roughly 2.2×10^{-308} to 1.8×10^{308}. IEEE defines five exceptions (invalid, division by zero, overflow, underflow, inexact), and specifies that ∞ and `NaN` propagate through arithmetic instead of halting it. Demmel [2, §1.5] stresses the value of this: an algorithm can run to completion and check its result at the end, instead of branching at every step. Note that `NaN != NaN` is always true, so one must test with `isnan`.

C.2 Machine Epsilon and the Model of Arithmetic

C.2.1 Two Conventions for Machine Epsilon

Two different quantities are both called "machine epsilon" in the literature, and they differ by a factor of 2.

Definition 116 (Machine Epsilon and Unit Roundoff)

Let b_p be the precision of a floating-point format.

- The *machine epsilon* $\varepsilon = \varepsilon_{\text{mach}} := 2^{-(b_p-1)}$ is the distance from 1 to the next larger floating-point number. For Float64, $\varepsilon = 2^{-52} \approx 2.22 \times 10^{-16}$. This is the convention we use, as do Julia's `eps(Float64)`, and of C's `DBL_EPSILON`.
- The *unit roundoff* $u := 2^{-b_p} = \varepsilon/2$ is the largest relative rounding error, i.e., half a spacing. For Float64, $u = 2^{-53} \approx 1.11 \times 10^{-16}$. Trefethen and Bau [1] and Demmel [2] call this quantity *machine epsilon* ($\epsilon_{\text{machine}}$, resp. ε), and Golub and Van Loan [5] call it the unit roundoff. So when copying a bound from these books, their "machine epsilon" is our $u = \varepsilon_{\text{mach}}/2$.

The two quantities answer different questions: ε describes how dense the floating-point grid is, while u describes how much error rounding to that grid causes. They differ by a factor of 2 because rounding goes to the *nearest* floating-point number. On $[2^e, 2^{e+1})$ the spacing is $2^e\varepsilon$, a real number x in this interval is at most half a spacing from its nearest neighbour, and therefore

$$\frac{|\mathrm{fl}(x) - x|}{|x|} \le \frac{2^e\varepsilon/2}{2^e} = \frac{\varepsilon}{2} = u. \tag{C.1}$$

Both conventions survive for historical and practical reasons. Many early machines rounded by *chopping*, i.e., discarding the extra bits, and then the largest relative error is a whole spacing, so the two quantities coincided. They separated only once IEEE 754 made round to nearest the default. Since then, authors on error analysis (Trefethen–Bau, Higham, Golub–Van Loan) use u, because what they need is the bound on δ in $\mathrm{fl}(x) = x(1+\delta)$, whereas programming languages report ε, because the gap from 1 to `nextfloat(1.0)` is what can be observed directly.

Any statement of the form $O(\varepsilon)$ is unaffected by the choice, but constants copied from different textbooks, such as $|\delta| \le u$ or $\gamma_n = nu/(1-nu)$, must be read with care.

The IEEE default rounding mode is *round to nearest, ties to even*, under which both orders in Example 1 (Lecture 1) give 4. Non-associativity does not depend on the rounding mode, however. With ties to even,

$$(1+u)+u = 1 \qquad \text{but} \qquad 1+(u+u) = 1+\varepsilon, \tag{C.2}$$

since on the left each intermediate result is a tie that rounds back to 1, while on the right $u+u = \varepsilon$ and $1+\varepsilon$ are both exact.

C.2.2 The Fundamental Axiom

Error analysis needs rounding to be captured by a rule we can reason with. Let $\mathrm{fl}(x)$ denote the floating-point number nearest to x. Because the relative spacing is bounded, every x in the normal range satisfies

$$\mathrm{fl}(x) = x(1+\delta), \qquad |\delta| \le u. \tag{C.3}$$

IEEE requires $+$, $-$, $\times$, $\div$, and $\sqrt{\cdot}$ to be *correctly rounded*: the exact result is computed in infinite precision and then rounded once. This gives the following model, stated in the same form in [1, Lecture 13], [5, §2.4], and [2, §1.5].

Theorem 117 (Fundamental Axiom of Floating-Point Arithmetic)

For all floating-point numbers x, y and $\circledast \in \{+, -, \times, \div\}$,

$$\mathrm{fl}(x \circledast y) = (x \circledast y)(1+\delta), \qquad |\delta| \le u. \tag{C.4}$$

The axiom has an immediate *backward* reading. For addition,

$$\mathrm{fl}(x+y) = (x+y)(1+\delta) = x(1+\delta) + y(1+\delta), \tag{C.5}$$

so the computed sum is *exactly* the sum of two slightly perturbed inputs. Pushing errors back onto the inputs in this way is the main thread of error analysis in this course; see Section C.3.

C.2.3 Accumulation of Errors

A single operation incurs an error of size $O(u)$. For n operations, Golub and Van Loan write

$$\gamma_n \coloneqq \frac{nu}{1-nu} \approx nu \qquad (nu \ll 1). \tag{C.6}$$

Proposition 118 (Error in Inner Products)

For $\vec{x}, \vec{y} \in \mathbb{R}^n$,

$$\left|\mathrm{fl}(\vec{x}^\top \vec{y}) - \vec{x}^\top \vec{y}\right| \le \gamma_n \sum_{i=1}^{n} |x_i|\,|y_i| = \gamma_n \left|\vec{x}\right|^\top |\vec{y}|\,. \tag{C.7}$$

Similarly, for matrices, $|\mathrm{fl}(AB) - AB| \le \gamma_n |A|\,|B|$ entrywise.

The *shape* of this bound matters much more than its constant. The right-hand side involves $|\vec{x}|^\top |\vec{y}|$, not $\left|\vec{x}^\top \vec{y}\right|$. If all terms have the same sign, the two agree and the relative error is $O(nu)$, which is fine. If instead $\vec{x}^\top \vec{y}$ is a small number obtained by cancelling many terms of both signs, then $|\vec{x}|^\top |\vec{y}| \gg \left|\vec{x}^\top \vec{y}\right|$ and the relative error can be arbitrarily bad. The linear factor n is almost never the real issue (it is a very pessimistic worst case, and random errors grow more like $\sqrt{n}$). The real issue is always *cancellation*.

C.3 Backward Stability

The textbooks [1, Lecture 14] mostly use a stronger notion than the stability defined in Section 1.7 of Lecture 1.

Definition 119 (Backward Stability)

An algorithm $\hat{f}$ for f is *backward stable* if for every x there exists $\tilde{x}$ with

$$\hat{f}(x) = f(\tilde{x}) \qquad \text{and} \qquad \frac{|\tilde{x} - x|}{|x|} = O(u), \tag{C.8}$$

where $\tilde{x}$ is a hypothetical input constructed in the analysis, never computed.

In words: *the computed answer is the exact answer to a nearby problem*. Both halves of the definition matter.

- The two sides of $\hat{f}(x) = f(\tilde{x})$ are computed differently. The left side runs the algorithm on the computer, with rounding, on the true input x. The right side evaluates the exact mathematical function f, with no rounding at all, on the perturbed input $\tilde{x}$.
- The perturbation must be small, $|\tilde{x} - x| \,/\, |x| = O(u)$. Without this condition the definition would be empty, since for most f some $\tilde{x}$ reproduces any output whatsoever.

Backward stability does not make the rounding errors disappear; it is a way of *accounting* for them. All errors committed inside the algorithm are pushed back to the input and reinterpreted as a perturbation $\tilde{x} - x$ of the data. The backward reading of Theorem 117 is the simplest example: $\mathrm{fl}(x + y)$ is the exact sum of $\tilde{x} = x(1 + \delta)$ and $\tilde{y} = y(1 + \delta)$, which are within relative distance u of x and y.

This accounting is useful because the input carries errors of size $O(u)$ anyway, from rounding when it is stored or from measurement. A backward stable algorithm gives the exact answer for some input that is indistinguishable from the one we have, which is as much as can be asked of any algorithm. Whatever error remains is due to the problem, not the algorithm. Backward stability is the gold standard of numerical linear algebra because it cleanly separates the problem from the algorithm, and it yields the classical rule of thumb

$$\text{forward relative error} \lesssim \kappa_f(x) \cdot u, \tag{C.9}$$

or, in a form that is easier to remember,

$$\text{accuracy} \approx \text{condition number (problem)} \times \text{backward error (algorithm)}. \tag{C.10}$$

A backward stable algorithm still returns inaccurate answers for ill-conditioned problems, and this is *not the algorithm's fault*; it cannot be improved upon. Conversely, when a result is inaccurate, the first step should be to estimate the condition number, not to suspect the code.

Concept	Belongs to	Question it answers	Measure
Conditioning	problem f and input x	how much input perturbations are intrinsically amplified	$\kappa_f(x)$
Stability	algorithm $\hat{f}$	whether the implementation amplifies errors further	backward error $O(u)$
Accuracy	final result	whether the computed number is right	$\lesssim \kappa \cdot u$

Some common condition numbers:

Problem	κ	Remark
$f(x) = \sqrt{x}$	$1/2$	always well-conditioned; even shrinks errors
$f(x) = x^n$	n	grows linearly with the power
$f(x) = e^x$	$\lvert x\rvert$	ill-conditioned for large $\lvert x\rvert$
$f(x_1, x_2) = x_1 - x_2$	$\dfrac{\lvert x_1\rvert + \lvert x_2\rvert}{\lvert x_1 - x_2\rvert}$	cancellation (componentwise perturbations)
$\vec{x} \mapsto A\vec{x}$	$\dfrac{\lVert A\rVert \, \lVert\vec{x}\rVert}{\lVert A\vec{x}\rVert} \le \kappa(A)$	depends on the direction of $\vec{x}$
solve $A\vec{x} = \vec{b}$	$\kappa(A) = \lVert A\rVert \, \lVert A^{-1}\rVert$	
polynomial roots	can be enormous	Wilkinson's polynomial

For $f : \mathbb{R}^n \to \mathbb{R}^m$ one replaces f' by the Jacobian $J(\vec{x})$: the absolute condition number is $\hat{\kappa} = \lVert J(\vec{x})\rVert$ and the relative condition number is $\kappa = \lVert J(\vec{x})\rVert \, \lVert\vec{x}\rVert \, / \, \lVert f(\vec{x})\rVert$ [1, Lecture 12].

Bibliography

[1] L. N. Trefethen and D. Bau III. *Numerical Linear Algebra*. SIAM, 1997.

[2] J. W. Demmel. *Applied Numerical Linear Algebra*. SIAM, 1997.

[3] D. C. Lay, S. R. Lay, and J. J. McDonald. *Linear Algebra and Its Applications*, 6th edition. Pearson, 2021.

[4] S. Axler. *Linear Algebra Done Right*, 4th edition. Springer, 2024.

[5] G. H. Golub and C. F. Van Loan. *Matrix Computations*, 3rd edition. Johns Hopkins University Press, 1996.

[6] M. L. Overton. *Numerical Computing with IEEE Floating Point Arithmetic*. SIAM, 2001.

[7] N. J. Higham. *Accuracy and Stability of Numerical Algorithms*, 2nd edition. SIAM, 2002.

[8] N. J. Higham. *Functions of Matrices: Theory and Computation*. SIAM, 2008.