

Tuesday, March 27 * Solutions * Introduction to multiple integrals


1. Evaluate the following integral by reversing the order of integration:

$$\int_0^1 \int_{\sqrt{y}}^1 \sqrt{x^3 + 1} \, dx \, dy.$$

(Hint: When you change to $dx \, dy$, be sure to also change the bounds of integration.)

SOLUTION:

We are integrating over the region below:



Changing the order of integration we get


$$\int_0^1 \int_{\sqrt{y}}^1 \sqrt{x^3 + 1} \, dx \, dy = \int_0^1 \int_0^{x^2} \sqrt{x^3 + 1} \, dy \, dx$$

$$\int_0^1 \int_0^{x^2} \sqrt{x^3 + 1} \, dy \, dx = \int_0^1 x^2 \sqrt{x^3 + 1} \, dx = 2/9[(x^3 + 1)^{3/2}]_0^1 = 2/9(2^{3/2} - 1).$$

2. Consider the region bounded by the curves determined by $-2x + y^2 = 6$ and $-x + y = -1$.

(a) Sketch the region R in the plane.

SOLUTION:



(b) Set up and evaluate an integral of the form $\iint_R dA$ that calculates the area of R .


SOLUTION:

$$\int_{-2}^4 \int_{\frac{y^2-6}{2}}^{y+1} dx dy = \int_{-2}^4 y + 1 - \frac{y^2-6}{2} dy = \left[-1/6 y^3 + 1/2 y^2 + 4y \right]_{-2}^4 = 18$$

3. Consider the region R in the first quadrant which lies above the x -axis and between the circles of radius 1 and 2 centered at $(0,0)$. *Without using polar coordinates*, evaluate

$$\iint_R y \, dA.$$

SOLUTION: Notice that both the function y and the region R are symmetric about the y -axis, so we can integrate y over the half of R which lies in the first quadrant (Call this R') and double our answer. R' is shown below.



Break up R' into two parts A and B as above. Integrating, we obtain

$$\begin{aligned} \iint_R y \, dA &= \iint_A y \, dA + \iint_B y \, dA = \int_0^1 \int_{\sqrt{1-x^2}}^{\sqrt{4-x^2}} y \, dy \, dx + \int_1^2 \int_0^{\sqrt{4-x^2}} y \, dy \, dx \\ &= \int_0^1 [y^2/2]_{\sqrt{1-x^2}}^{\sqrt{4-x^2}} dx + \int_1^2 [y^2/2]_0^{\sqrt{4-x^2}} dx = \int_0^1 3/2 \, dx + \int_1^2 1/2(4-x^2) \, dx \\ &= 7/3 \end{aligned}$$

Now double this value to get $14/3$, which is the integral over the entire region R .


4. Evaluate

$$\int_{-2}^0 \int_0^{\sqrt{4-x^2}} (x^2 + y^2) dy dx.$$

Hint: don't do it directly.

SOLUTION:

The region over which we are integrating is:




Converting to polar we get

$$\int_{-2}^0 \int_0^{\sqrt{4-x^2}} (x^2 + y^2) dy dx = \int_{\pi/2}^{\pi} \int_0^2 (r^2) r dr d\theta = 2\pi$$

5. The function $P(x) = e^{-x^2}$ is fundamental in probability.

(a) Sketch the graph of $P(x)$. Explain why it is called a “bell curve.”

SOLUTION:



(b) Compute $I = \int_{-\infty}^{\infty} e^{-x^2} dx$ using the following brilliant strategy of Gauss.

i. Instead of computing I , compute $I^2 = \left(\int_{-\infty}^{\infty} e^{-x^2} dx \right) \left(\int_{-\infty}^{\infty} e^{-y^2} dy \right)$.

ii. Rewrite I^2 as an integral of the form $\iint_R f(x, y) dA$ where R is the entire Cartesian plane.

SOLUTION:

$$I^2 = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-x^2-y^2} dy dx$$

iii. Convert that integral to polar coordinates.

SOLUTION:

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-x^2-y^2} dy dx = \int_0^{2\pi} \int_0^{\infty} r e^{-r^2} dr d\theta$$

iv. Evaluate to find I^2 . Deduce the value of I .

SOLUTION:

$$\begin{aligned} \int_0^{2\pi} \int_0^{\infty} r e^{-r^2} dr d\theta &= 2\pi \int_0^{\infty} r e^{-r^2} dr = 2\pi \lim_{t \rightarrow \infty} \int_0^t r e^{-r^2} dr = 2\pi \lim_{t \rightarrow \infty} \left[-1/2 e^{-r^2} \right]_0^t \\ &= \pi \lim_{t \rightarrow \infty} (-e^{-t^2} + 1) = \pi \end{aligned}$$

So $I = \sqrt{\pi}$.

6. Compute $\int_0^\infty \int_0^\infty \frac{1}{(1+x^2+y^2)^2} dx dy$.

SOLUTION:

As in the previous problem, let's convert to polar coordinates.

$$\int_0^\infty \int_0^\infty \frac{1}{(1+x^2+y^2)^2} dx dy = \int_0^{\pi/2} \int_0^\infty \frac{r}{(1+r^2)^2} dr d\theta = \pi/2 \int_0^\infty \frac{r}{(1+r^2)^2} dr$$

This is an improper integral, so

$$\pi/2 \int_0^\infty \frac{r}{(1+r^2)^2} dr = \pi/2 \lim_{t \rightarrow \infty} \int_0^t \frac{r}{(1+r^2)^2} dr = \pi/4 \lim_{t \rightarrow \infty} \left[-\frac{1}{1+r^2} \right]_0^t$$

$$= \pi/4 \lim_{t \rightarrow \infty} \left(-\frac{1}{1+t^2} + 1 \right) = \pi/4$$