

1. (a) Find the gradient vector field of  $f(x, y, z) = xye^yz + x^2 + yz$ .  
 (b) State the Fundamental Theorem of Line Integrals.  
 (c)  $\vec{F}(x, y, z) = \langle yz + 2(y + z), xz + 2(x + z), xy + 2(x + y) \rangle$  is conservative. Let  $C$  be the curve parameterized by  $\vec{r}(t) = \langle t, t^2, t^3 \rangle$  for  $0 \leq t \leq 1$ . Evaluate  $\int_C \vec{F} \cdot d\vec{r}$ .  
 (d) Let  $\vec{F}(x, y, z)$  be a vector field, with  $\int_C \vec{F} \cdot d\vec{r} = \pi$  for the curve  $C$  parameterized by  $\vec{r}(t) = \langle 6, \cos 2t, \sin 2t \rangle$  for  $0 \leq t \leq \pi$ . TRUE or FALSE:  $\vec{F}$  is conservative.  
 (e) Which of the three vector fields:
  - i.  $\vec{F}(x, y) = \langle (1 + xy)e^{xy}, x^2e^{xy} \rangle$
  - ii.  $\vec{G}(x, y) = \langle x^3y^2 + 2, x^2y^3 + 2 \rangle$
  - iii.  $\vec{H}(x, y) = \langle ye^x + \sin y, e^x + x \cos y \rangle$
 are conservative? Find a potential function for those which are conservative.

2. Set up (but do **NOT** evaluate) a double integral to calculate the volume of the solids  $E$  and  $S$ , where:

- (a)  $E$  is the solid under the surface  $z = x^2 + 2y^4$  and above the region  $D = \{(x, y, 0) \mid -1 \leq x \leq 1, x^2 \leq y \leq 2x^2\}$ .
- (b)  $S$  is the solid under the plane  $3x + 2y - z = 0$  and above the region enclosed by  $y = x^2$  and  $x = y^2$ .

3. Evaluate the following integrals by reversing the order of integration.

(a)

$$\int_0^1 \int_{3y}^3 e^{x^2} dx dy$$

(b)

$$\int_0^2 \int_{\frac{y}{2}}^1 y \cos(x^3 - 1) dx dy$$

4. Find the mass of the lamina the occupies the region  $R$  bounded by the semicircle  $x = -\sqrt{4 - y^2}$  and the  $y$ -axis, whose density at any point in  $R$  is  $\rho(x, y) = e^{-x^2 - y^2}$ .

5. Use polar coordinates to combine the sum

$$\int_0^{\sqrt{2}} \int_0^x xy \, dy \, dx + \int_{\sqrt{2}}^2 \int_0^{\sqrt{4-x^2}} xy \, dy \, dx$$

into one double integral. Then evaluate the double integral.

6. Set up (but do **NOT** evaluate) the triple integral

$$\iiint_E 6xy \, dV,$$

as an iterated integral, where  $E$  lies under the plane  $z = 1 + x + y$  and above the region in the  $xy$ -plane bounded by the curves  $y = \sqrt{x}$ ,  $y = 0$ , and  $x = 1$ .

7. Evaluate  $\iiint_E (x - y) \, dV$ , where  $E$  is the solid that lies between the cylinders  $x^2 + y^2 = 1$  and  $x^2 + y^2 = 16$ , above the  $xy$ -plane, and below the plane  $z = y + 4$ .
8. Let  $R = \{(x, y, z) \mid 2 \leq z \leq 2 + \sqrt{4 - x^2 - y^2}\}$ . Set up the iterated integrals to find the volume of region  $R$  in rectangular, cylindrical, and spherical coordinates, then calculate the volume of  $R$ .

## TRIGONOMETRIC IDENTITIES

$$\sin(\theta + \phi) = \sin\theta \cos\phi + \cos\theta \sin\phi$$

$$\cos(\theta + \phi) = \cos\theta \cos\phi - \sin\theta \sin\phi$$

$$\tan(\theta + \phi) = \frac{\tan(\theta) + \tan(\phi)}{1 - \tan\theta \tan\phi}$$

$$\sin(\theta - \phi) = \sin\theta \cos\phi - \cos\theta \sin\phi$$

$$\cos(\theta - \phi) = \cos\theta \cos\phi + \sin\theta \sin\phi$$

$$\tan(\theta - \phi) = \frac{\tan(\theta) - \tan(\phi)}{1 + \tan\theta \tan\phi}$$

$$\sin(\theta) \sin(\phi) = \frac{\cos(\theta - \phi) - \cos(\theta + \phi)}{2}$$

$$\cos(\theta) \cos(\phi) = \frac{\cos(\theta - \phi) + \cos(\theta + \phi)}{2}$$

$$\sin(\theta) \cos(\phi) = \frac{\sin(\phi + \theta) - \sin(\phi - \theta)}{2}$$

$$\sin(2\theta) = 2 \sin\theta \cos\theta$$

$$\cos(2\theta) = \cos^2\theta - \sin^2\theta$$

$$\tan(2\theta) = \frac{2 \tan\theta}{1 - \tan^2\theta}$$

$$\sin\theta = \pm \sqrt{\frac{1 - \cos(2\theta)}{2}}$$

$$\cos\theta = \pm \sqrt{\frac{1 + \cos(2\theta)}{2}}$$

$$\tan\theta = \frac{\sin(2\theta)}{1 + \cos(2\theta)}$$