

Higher condensation theory

Liang Kong ^{*a,b}, Zhi-Hao Zhang ^{†c,d,a}, Jiaheng Zhao ^{‡e,f}, and Hao Zheng ^{§g,c}

^aInternational Quantum Academy, Shenzhen 518048, China

^bShenzhen Institute for Quantum Science and Engineering,

Southern University of Science and Technology, Shenzhen, 518055, China

^cBeijing Institute of Mathematical Sciences and Applications, Beijing 101408, China

^dWu Wen-Tsun Key Laboratory of Mathematics of Chinese Academy of Sciences,

School of Mathematical Sciences,

University of Science and Technology of China, Hefei, 230026, China

^eAcademy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China

^fUniversity of Chinese Academy of Sciences, Beijing 100049, China

^gInstitute for Applied Mathematics, Tsinghua University, Beijing, 100084, China

Abstract

We develop a unified mathematical theory of defect condensations for topological orders in all dimensions based on higher categories, higher algebras and higher representations. A k -codimensional topological defect A in an $n+1$ D (potentially anomalous) topological order C^{n+1} is condensable if it is equipped with the structure of a condensable E_k -algebra. Condensing such a defect A amounts to a k -step process. In the first step, we condense the defect A along one of its transversal directions, thus obtaining a $(k-1)$ -codimensional defect ΣA , which is naturally equipped with the structure of a condensable E_{k-1} -algebra. In the second step, we condense the defect ΣA along one of the remaining transversal directions, thus obtaining a $(k-2)$ -codimensional defect $\Sigma^2 A$, so on and so forth. In the k -th step, we condense the 1-codimensional defect $\Sigma^{k-1} A$ along the only transversal direction, thus defining a phase transition from C^{n+1} to a new $n+1$ D topological order D^{n+1} . We give precise mathematical descriptions of each step in above process, including the precise mathematical characterization of i -codimensional topological defects in the condensed phase D^{n+1} for $1 \leq i \leq k$ in terms of E_i -modules over the E_i -algebra $\Sigma^{k-i} A$. When C^{n+1} is anomaly-free, the same phase transition, as a k -step process, can be alternatively defined by replacing the last two steps by a single step of condensing the E_2 -algebra $\Sigma^{k-2} A$ directly along the remaining two transversal directions. When $n = 2$, this modified last step is precisely a usual anyon condensation in a 2+1D topological order. In addition to above algebraic approach, we also provide a geometric theory of defect condensations. We derive many new mathematical results physically along the way, and develop some powerful tools for the construction and classification of condensable E_1 -algebras and Lagrangian E_2 -algebras from the precise characterization of the non-surjectivity and non-fully-faithfulness of the center functors. We also establish the equivalences among various notions of ‘gauging’ symmetries. At the end of this work, we briefly discuss questions, generalizations and applications that naturally arise from our theory, including higher Morita theory, a theory of integrals and the condensations of liquid-like gapless defects in topological orders.

*Email: kongustc@outlook.com

†Email: zhangzhihao@bimsa.cn

‡Email: zhaojiaheng171@mails.ucas.ac.cn

§Email: haozheng@mail.tsinghua.edu.cn

Contents

1	Introduction	4
1.1	Motivations and historical remarks	4
1.2	Main results	5
1.3	Layout, conventions and notations	12
2	Categories of topological defects	15
2.1	Categories of topological defects	15
2.1.1	2+1D topological orders	15
2.1.2	Remote detectable principle and ‘intrinsic’ topological defects	20
2.1.3	Boundary-bulk relation	24
2.2	Condensation completion	27
2.2.1	Category $n\text{Vec}$	27
2.2.2	Separable n -categories	31
2.2.3	Composite topological orders	33
2.2.4	Morita equivalence between topological orders	35
2.2.5	Multi-fusion n -categories	36
2.3	Higher algebras	42
2.3.1	Monoidal 1-categories as E_1 -algebras	42
2.3.2	Braided monoidal 1-categories as E_2 -algebras	43
2.3.3	E_m -multi-fusion n -categories	46
2.3.4	Condensable higher algebras	47
2.3.5	E_m -center and E_m -centralizer	50
2.4	Center functor	52
2.4.1	Complete boundary-bulk relation as a center functor	52
2.4.2	Measure non-surjectivity and non-fully-faithfulness	55
2.4.3	Generalized center functors	57
3	Condensations of 1-codimensional topological defects	60
3.1	Particle condensations in 1+1D: algebraic approach	60
3.1.1	Algebraic bootstrap	60
3.1.2	Internal homs and structure theorem	62
3.1.3	Examples of particle condensations in 1+1D	64
3.2	Particle condensations in 1+1D: geometric approach	66
3.2.1	Physical intuitions	66
3.2.2	Precise geometric theory	68
3.3	Condensations of 1-codimensional defects in $n+1D$	70
3.3.1	General theory	70
3.3.2	General constructions	76
3.3.3	Examples in 2+1D: anomaly-free cases	83
3.3.4	Examples in 2+1D: anomalous cases	86
3.3.5	Examples in 3+1D	89
3.3.6	Examples in higher dimensions	92
4	Condensations of 2-codimensional topological defects	95
4.1	Anyon condensations in 2+1D	95
4.1.1	Algebraic approach	95
4.1.2	Examples	101
4.1.3	Geometric approach	105
4.2	Condensations of 2-codimensional defects in $n+1D$	108
4.2.1	General theory	108
4.2.2	General construction	113
4.2.3	Witt equivalence	117
4.2.4	Examples in 3+1D	118
4.2.5	Examples in higher dimensions	121

5	Condensations of higher codimensional topological defects	123
5.1	Particle condensations in 2+1D via a two-step process	123
5.1.1	General theory	123
5.1.2	Examples	127
5.2	Condensation of higher codimensional topological defects	129
5.2.1	General theory	129
5.2.2	General constructions	135
5.2.3	Concrete examples	138
5.3	Gauging a G -symmetry	140
5.3.1	Gauging a G -symmetry in a 2+1D SET order	140
5.3.2	General constructions and examples	143
5.3.3	Generalizations, examples and remarks	146
6	Generalizations and Applications	149
6.1	Higher Morita theory	149
6.1.1	Higher Morita equivalences	149
6.1.2	Enveloping algebras	151
6.2	Theory of integrals	153
6.3	Condensations of gapless liquid-like defects	157
6.3.1	General theory	157
6.3.2	An illustrating example in 2+1D	159
6.4	Generalized lattice models with categorified local degrees of freedom	162
A	Appendix	166
A.1	Proof of Theorem 4.1.19	166
A.2	Module Eilenberg-Watts calculus	169
A.3	E_k -center of an E_k -algebra	174
A.4	Geometric condensation via topological Wick rotation	175
	References	176

1 Introduction

In this work, we develop a unified mathematical theory of defect condensations for topological orders in all dimensions based on higher categories, higher algebras and higher representations, and explore its consequences and applications. In this section, we explain the motivations, summarize the main results, and fix some conventions and notations. The symbol “ nD ” represents the spacetime dimension.

1.1 Motivations and historical remarks

One of fundamental themes in modern theoretical physics is to establish a new paradigm of phases and phase transitions that can unify traditional spontaneous symmetry-breaking (SSB) orders with exotic new phases discovered since 1980’s, including topological orders [Wen90], symmetry protected/enriched topological (SPT/SET) orders [GW09, CGW10] and gapless quantum liquids [KZ22b]. This yet-to-be-established new paradigm should be a theoretical framework that can catch the universal and model independent features of all these quantum phases and phase transitions among them. It was known that many phase transitions among topological orders are driven by the condensations of topological defects. In 2+1D, a condensation of 0+1D defects was known as an anyon condensation [MS89c, BSS02, BSS03, BS09, BSH09, BSS11, KS11a, Lev13, BJQ13, Kon14]. A condensation of 1+1D topological defects was known as a string condensation [ZLZH⁺23]. A defect condensation can also be viewed as the (higher) gauging of generalized or categorical symmetries (see for example [FFRS10, GKSW15, BT18, TW24, RSS23, CH25, HPRW25])¹. Defect condensations can also be used to control or define phase transitions among gapless quantum liquids [KZ20, KZ21, CJKY20, CW23, LY23, CJW25, CAW24]. Therefore, the mathematical theory of defect condensations, which is indeed universal and model independent, is an indispensable ingredient of the new paradigm of phase transitions.

Since both the experimental discovery of topological orders [TSG82, Wen90] (see [Wen17, Wen19] for reviews) and the theoretical discovery of its macroscopic descriptions occurred in 2+1D [MS89a, Wit89, FRS89, FG90, Reh90, MR91, Wen91, Kit06], the first thoroughly studied defect condensation is the boson condensation (somehow misnamed as anyon condensation) in 2+1D topological orders. It was originated in the study of the extensions of chiral algebras [MS89b] and was first studied by Moore and Seiberg in a special case in [MS89c]. It was later systematically developed and named as ‘boson condensation’ by Bais, Schroers, Slingerland in [BSS02, BSS03] as a generalization of Landau’s symmetry-breaking theory, and was further developed in [BS09, BSH09, BSS11]. The mathematical description of an anyon condensation was partially settled for 2+1D finite abelian gauge theories in [KS11b, Lev13, BJQ13], and was fully established in categorical language for all 2+1D topological orders in [Kon14] based on many earlier mathematical works on algebras in braided fusion categories or subfactors [Par95, BEK99, BEK00, BEK01, KO02, FFRS06, DMNO13]. More works on this categorical description followed [ERB14, HW15, NHvKSB16a, NHvKSB16b, Bur18]. We provide more references on recent developments in Remark 4.1.6.

However, the anyon condensation theory do not provide the complete model-independent story of a phase transition between two 2+1D topological orders. A manifestation of this fact is that an anyon condensation from one 2+1D topological order to another is not reversible. On the other hand, an experimental or lattice model realization of a phase transition is always reversible. Another manifestation of this fact is that two 2+1D topological orders connected by a gapped domain wall, in general, cannot be obtained from each other via anyon condensations. The reason behind both phenomena is that anyons do not cover all topological defects in a 2+1D topological order. Indeed, anyons are topological defects of codimension 2 (or 2-codimensional topological defects). There are non-trivial 1-codimensional topological defects in a 2+1D topological order [KS11a, KK12, FSV13, CR16, Kon14]. These topological defects form a monoidal 2-category [KK12, KW14]. It turns out that all of these 1-codimensional topological defects can be obtained from anyons via condensations [FSV13, CR16, Kon14], thus are called condensation descendants or condensed defects [KW14]. Mathematically, the process of finding all condensation descendants from a subset of defects (such as anyons) is called condensation completion. It was first introduced by Carqueville and Runkel in [CR16] for 2-categories under the name of “orbifold completion”, and was proposed as a fundamental question for higher dimensional topological orders in [KW14]. It was later thoroughly developed by

¹Within the framework of TQFT, there is a program, which is parallel to defect condensations, of ‘orbifold TQFTs’ developed by Carqueville, Runkel, Schaumann and Mulevičius in a series of works [CR16, CRS18, CRS19, CRS20, CMRSS21, Mul24, MR23, CM23, CMRSS24] (see [Car23] for a survey, see also Remark 4.1.7) based on state-sum constructions. It was preceded by some earlier works on QFTs with defects [FFRS07, FFRS10, DKR11]. See Remark 1.2.10 and Remark 3.3.28 for further comments.

Douglas and Reutter in [DR18] under the name of “idempotent completion”, and was used there to give a mathematical definition of multi-fusion 2-category and a state-sum construction of 3+1D TQFT’s. Combining it with other known mathematical results, it becomes clear that the monoidal 2-category of topological defects of codimension 1 or higher in a 2+1D topological order [KK12, Kon14] is actually a fusion 2-category (see [KLWZZ20b] for a detailed explanation). The theory of condensation completion was generalized to higher categories by Gaiotto and Johnson-Freyd in [GJF19] under the name of “Karoubi completion” or “condensation completion” (see also [CRS19] and a rigorous treatment for 3-categories in [CM23]). This theory leads Johnson-Freyd to a mathematical theory of a (multi-)fusion n -category in [JF22]. The compatibility between condensation completion and boundary-bulk relation [KWZ15, KWZ17] was first shown in [KTZ20b] for 3+1D Dijkgraaf-Witten theories, and was proved in general context by Johnson-Freyd in [JF22] (see [KZ22b] for a slightly different approach). This compatibility was also proposed in [KLWZZ20b] as general physical principle beyond topological orders (see also [KZ22b]). The theory of condensation completion was also generalized to gapless quantum liquids [KZ22b, KZ22c].

Although the theory of defect condensations was significantly advanced by that of condensation completion, it is still largely incomplete. The condensation of a 1-codimensional topological defect was studied for 1+1D anomalous topological orders in [KS11a, FSV13, CR16, Kon14, DR18] and for higher dimensional cases in the condensation completion theory [GJF19]. The condensation theory of k -codimensional topological defects was known for $k = 2$ in 2+1D [Kon14] and 3+1D [ZLZH⁺23, DX24, Xu24a], and is widely open for $k > 2$ in higher dimension theories. Most of these results are too brief and formal, and are lack of physical explanations and systematic constructions of examples. The relation between the condensations of the defects of different codimensions have not been studied. Moreover, there are serious gaps in turning the theory into a computable one, i.e., expressing the condensation data and results in explicit coordinate systems (see Theorem^{ph} 1.2.6 (2),(4),(5)). Recently, the category theory of anyon condensations has inspired a lot of works on topological phase transitions in 1+1D (see for example [CJKYZ20, CW23, LY23, CJW25, CAW24] and references therein). As we are entering a new era of systematically studying topological phase transitions, there are increasing demands to have a rather complete theory of defect condensations in all dimensions.

The main goal of this work is to develop such a theory, i.e., a theory of the condensations of k -codimensional topological defects in a (potentially anomalous) $n+1$ D topological order for $1 \leq k \leq n$. We show that this theory naturally leads us to the mathematical theory of higher algebras in higher monoidal categories and their higher representations. We also provide some systematic constructions of examples. This theory can be naturally generalized to the condensations of liquid-like gapless defects in topological orders or in gapped/gapless quantum liquids. We provide a glimpse of this generalized theory at the end of this work, together with other questions or generalizations naturally arising from our theory.

1.2 Main results

Since invertible topological orders do not play any role in our defect condensation theory, throughout this work, a topological order is defined up to invertible ones.


In order to state the main results, we need first introduce some mathematical notions. We want to keep it brief because a detailed review of these notions is given in later sections.

Consider a potentially anomalous $n+1$ D topological order C^{n+1} (see Definition 2.1.9). We want to define the condensation of a k -codimensional topological defect A in C^{n+1} . This condensation produces a condensed phase denoted by D^{n+1} and a gapped domain wall denoted by M^n . We denote the category of topological defects of codimension 1 and higher in C^{n+1} (resp. D^{n+1}) by $\mathcal{C}$ (resp. $\mathcal{D}$), which is a multi-fusion n -category [JF22] with a tensor unit $\mathbb{1}_{\mathcal{C}}$ (resp. $\mathbb{1}_{\mathcal{D}}$). If C^{n+1} is simple (see Definition 2.2.15), then $\mathcal{C}$ is a fusion n -category. The gapped domain wall M^n can be described by a pair $(\mathcal{M}, m)$, where $\mathcal{M}$ is the category of all wall conditions (as objects) and topological defects (as morphisms) and m is a distinguished wall condition (i.e., $m \in \mathcal{M}$). We set $\Omega\mathcal{C} := \text{hom}_{\mathcal{C}}(\mathbb{1}_{\mathcal{C}}, \mathbb{1}_{\mathcal{C}})$, $\Omega_m\mathcal{M} := \text{hom}_{\mathcal{M}}(m, m)$ and $\Omega^k\mathcal{C} = \Omega(\Omega^{k-1}\mathcal{C})$. It is known that $\Omega^{k-1}\mathcal{C}$ is an E_k -multi-fusion $(n-k+1)$ -category [KZ22b], which is a direct sum of indecomposable separable $(n-k+1)$ -categories. We denote the indecomposable summand of $\Omega^{k-1}\mathcal{C}$ consisting of the tensor unit by $\Sigma\Omega^k\mathcal{C}$, which is called the delooping of $\Omega^k\mathcal{C}$. Mathematically, the delooping of an E_1 -multi-fusion l -category $\mathcal{A}$, denoted by $\Sigma\mathcal{A}$, can be defined by a one-point delooping of $\mathcal{A}$ followed by condensation completion [DR18, GJF19] (see a physical review in Section 2.2), and can be mathematically identified with the category $\text{RMod}_{\mathcal{A}}((l+1)\text{Vec})$ of right separable $\mathcal{A}$ -modules, or equivalently, right $\mathcal{A}$ -modules in $(l+1)\text{Vec}$ [DR18, GJF19], where $n\text{Vec} := \Sigma^n\mathbb{C}$ [GJF19] and Vec is the 1-category of finite dimensional vector spaces over $\mathbb{C}$.

Now we summarize the main results of this work.

Theorem^{ph} 1.2.1. Consider a (potentially anomalous) $n+1$ D simple topological order C^{n+1} , i.e., $\mathcal{C}$ is a fusion n -category. A k -codimensional topological defect A in C^{n+1} is fully condensable if A is a condensable E_k -algebra in $\Omega^{k-1}\mathcal{C}$ (see Definition 2.3.13). By condensing A , we mean a k -step process explained below.

1. In the first step, we condense A along one of the transversal directions $x^1, \dots, x^k$ (see Convention 2.1.16, 2.3.2) of A , thus obtaining a $(k-1)$ -codimensional defect ΣA in $\Sigma\Omega^{k-1}\mathcal{C} \hookrightarrow \Omega^{k-2}\mathcal{C}$.



It turns out that $\Sigma\Omega^{k-1}\mathcal{C}$ can be identified with $\text{RMod}_{\Omega^{k-1}\mathcal{C}}((n-k+2)\text{Vec})$, which can be viewed as a coordinate system of $\Sigma\Omega^{k-1}\mathcal{C}$. In this coordinate system, ΣA can be identified with the category $\text{RMod}_A(\Omega^{k-1}\mathcal{C})$ of right A -modules in $\Omega^{k-1}\mathcal{C}$, i.e., $\Sigma A = \text{RMod}_A(\Omega^{k-1}\mathcal{C}) \in \Sigma\Omega^{k-1}\mathcal{C} \hookrightarrow \Omega^{k-2}\mathcal{C}$.

2. It turns out that ΣA is automatically a condensable E_{k-1} -algebra in $\Sigma\Omega^{k-1}\mathcal{C}$ and, therefore, a condensable E_{k-1} -algebra in $\Omega^{k-2}\mathcal{C}$, which can be further condensed along one of the remaining transversal directions as the second step. So on and so forth. In the $(k-1)$ -th step, we obtain a 1-codimensional topological defect $\Sigma^{k-1}A$, which is automatically a condensable E_1 -algebra in $\mathcal{C}$.
3. In the k -th step, we condense $\Sigma^{k-1}A$ in $\mathcal{C}$ along the unique remaining transversal direction. We obtain a new topological order D^{n+1} and a gapped domain wall M^n such that

$$\begin{array}{c}
 \Omega^i \mathcal{C} \xrightarrow{L_{i+1}} \Omega_m^i \mathcal{M} \quad \Omega_m^i \mathcal{M} \xleftarrow{R_{i+1}} \Omega^i \mathcal{D} \\
 \begin{array}{|c|c|}
 \hline
 \mathcal{C}^{n+1} & \mathcal{D}^{n+1} \\
 \hline
 \end{array} \\
 \begin{array}{|c|}
 \hline
 M^n \\
 \hline
 \end{array} \\
 M^n = (\mathcal{M}, m)
 \end{array}
 \quad
 \begin{array}{l}
 \mathcal{D} \simeq \text{Mod}_{\Sigma^{k-1}A}^{E_1}(\mathcal{C}); \\
 \mathcal{M} \simeq \text{RMod}_{\Sigma^{k-1}A}(\mathcal{C}) = \Sigma^k A, \\
 m = \Sigma^{k-1}A \in \mathcal{M},
 \end{array}
 \quad (1.2.1)$$

where $\text{Mod}_{\Sigma^{k-1}A}^{E_1}(\mathcal{C})$ denotes the category of E_1 -modules (i.e., bimodules) over $\Sigma^{k-1}A$ in $\mathcal{C}$ and is an indecomposable multi-fusion n -category; and $\text{RMod}_{\Sigma^{k-1}A}(\mathcal{C})$ denotes the category of right $\Sigma^{k-1}A$ -modules in $\mathcal{C}$ and is a separable n -category. Moreover, $\mathcal{D}$ is Morita equivalent to $\mathcal{C}$ with the Morita equivalence defined by the invertible $\mathcal{C}$ - $\mathcal{D}$ -bimodule $\mathcal{M}$. When the algebra A is indecomposable, $\mathcal{D}$ is a fusion n -category and $\mathcal{M}$ is an indecomposable left $\mathcal{C}$ -module.

4. For $1 \leq i \leq k$, the category of topological defects of codimension i and higher in D^{n+1} is determined by the following canonical E_i -monoidal equivalence:

$$\Omega^{i-1}\mathcal{D} = \Omega^{i-1}\text{Mod}_{\Sigma^{k-1}A}^{E_1}(\mathcal{C}) \simeq \text{Mod}_{\Sigma^{k-i}A}^{E_i}(\Omega^{i-1}\mathcal{C}),$$

where $\text{Mod}_{\Sigma^{k-i}A}^{E_i}(\Omega^{i-1}\mathcal{C})$ denotes the category of E_i -modules over $\Sigma^{k-i}A$ in $\Omega^{i-1}\mathcal{C}$. Roughly speaking, an E_i -module over $\Sigma^{k-i}A$ in $\Omega^{i-1}\mathcal{C}$ is an object in $\Omega^{i-1}\mathcal{C}$ equipped with an i -dimensional A -action.


For $1 \leq i \leq k$, the category of topological defects of codimension $(i-1)$ and higher in M^n is determined by the following canonical equivalences:

$$\Omega_m^{i-1}\mathcal{M} = \Omega_m^{i-1}\Sigma^k A = \Omega_m^{i-1}\text{RMod}_{\Sigma^{k-1}A}(\mathcal{C}) \simeq \text{RMod}_{\Sigma^{k-i}A}(\Omega^{i-1}\mathcal{C}).$$

5. For $1 \leq i \leq k$, i -codimensional topological defects in $\Omega^{i-1}\mathcal{C}$ and those in $\Omega^{i-1}\mathcal{D}$ move onto the wall according to the following functors, respectively, as illustrated in (1.2.1),

$$L_i := - \otimes \Sigma^{k-i} A: \Omega^{i-1}\mathcal{C} \rightarrow \Omega_m^{i-1}\mathcal{M}; \quad R_i := \Sigma^{k-i} A \otimes_{\Sigma^{k-i}A} -: \Omega^{i-1}\mathcal{D} \rightarrow \Omega_m^{i-1}\mathcal{M}.$$

6. C^{n+1} and D^{n+1} share the same gravitational anomaly. In other words, they are gapped boundaries of the same anomaly-free $n+2$ D topological order as illustrated below,



where $Z(C)^{n+2}$ and $Z(D)^{n+2}$ denote the bulk of C^{n+1} and D^{n+1} , respectively. Mathematically, it implies the following results.

$$\mathfrak{Z}_1(\mathcal{C}) \simeq \mathfrak{Z}_1(\text{Mod}_{\Sigma^{k-1}A}^{E_1}(\mathcal{C})), \quad (1.2.2)$$

$$\mathfrak{Z}_0(\text{RMod}_{\Sigma^{k-1}A}(\mathcal{C})) \simeq \text{Mod}_{\Sigma^{k-1}A}^{E_1}(\mathcal{C})^{\text{op}} \boxtimes_{\mathfrak{Z}_1(\mathcal{C})} \mathcal{C}, \quad (1.2.3)$$

$$\mathcal{D}^{\text{op}} \simeq \text{Mod}_{\Sigma^{k-1}A}^{E_1}(\mathcal{C})^{\text{op}} \simeq \mathfrak{Z}_1(\mathcal{C}, \mathfrak{Z}_0(\text{RMod}_{\Sigma^{k-1}A}(\mathcal{C}))), \quad (1.2.4)$$


where $\mathfrak{Z}_0(-)$, $\mathfrak{Z}_1(-)$, $\mathfrak{Z}_1(-, -)$ denote the E_0 -center, the E_1 -center and E_1 -centralizer (see Definition 2.3.17) and the superscript $^{\text{op}}$ is defined in Definition 2.3.5. Our convention of left and right is explained below Figure 7. $\square$

Remark 1.2.2. In most physical literature, physicists study condensations in an anomaly-free topological order (see Definition 2.1.9). In this work, we provide a condensation theory for potentially anomalous topological orders (e.g., the boundary phase C^{n+1} of a non-trivial bulk phase $Z(C)^{n+2}$). It is not only because our theory naturally covers all the anomalous cases, but also because a defect condensation in an anomalous $n+1$ D topological order, according to the sandwich construction or the topological Wick rotation [KZ18b, KZ20, KZ21, KLWZZ20b, KLWZZ20a, KZ22b], automatically defines a phase transition between two anomaly-free $n+1$ D gapped quantum liquids (including SPT/SET/SSB orders) with a finite internal symmetry, and automatically defines the gauging of categorical symmetries in a gapless QFT. In other words, our main results automatically give a mathematical theory of condensations in SPT/SET/SSB orders (see some concrete examples in 1+1D in [KWZ22, XZ24]), and a theory of the gauging of categorical symmetries in gapless QFT's (see Section 5.3.3 for more discussions on the gauging of symmetries in QFT's). $\diamond$

Remark 1.2.3. We state the main results for simple topological orders. In the later sections, we slightly generalize them to composite topological orders (see Definition 2.2.15) for the sake of completeness. Physics oriented readers can ignore the composite cases at least in their first reading. $\diamond$

In the special case when C^{n+1} is anomaly-free, we can define the same condensation alternatively by combining the last two-steps into one as stated below (see Section 4.2.1 for more related results).

Theorem^{ph} 1.2.4. When C^{n+1} is anomaly-free and simple, if A is a simple condensable E_k -algebra in $\Omega^{k-1}\mathcal{C}$ for $k \geq 2$, then we can modify the k -step process of condensing A by combining the last two steps into a single step of condensing $\Sigma^{k-2}A$ in $\Omega\mathcal{C}$ directly in the remaining two transversal directions. Note that $\Sigma^{k-2}A$ is automatically a condensable E_2 -algebra in $\Omega\mathcal{C}$. This condensation reproduces the same condensed phase D^{n+1} and the same gapped domain wall M^n as explained below.



1. We have the following monoidal equivalences:

$$\Omega\mathcal{D} \simeq \text{Mod}_{\Sigma^{k-2}A}^{E_2}(\Omega\mathcal{C}) \quad \text{and} \quad \Omega_m\mathcal{M} \simeq \text{RMod}_{\Sigma^{k-2}A}(\Omega\mathcal{C}), \quad (1.2.5)$$

where the first one is a braided equivalence or an E_2 -monoidal equivalence.

2. The 2-codimensional topological defects in C^{n+1} and D^{n+1} move onto the wall according to the following functors:

$$L_2 := - \otimes \Sigma^{k-2} A : \Omega \mathcal{C} \rightarrow \Omega_m \mathcal{M}, \quad R_2 := \Sigma^{k-2} A \otimes_{\Sigma^{k-2} A} - : \Omega \mathcal{D} \rightarrow \Omega_m \mathcal{M}.$$

both of which are central functors that can be lifted to an E_2 -monoidal (or braided) equivalence:

$$\Omega \mathcal{C} \boxtimes \text{Mod}_{\Sigma^{k-2} A}^{E_2}(\Omega \mathcal{C})^{\text{op}} \simeq \mathfrak{Z}_1(\text{RMod}_{\Sigma^{k-2} A}(\Omega \mathcal{C})),$$

or equivalently, an E_2 -monoidal (or braided) equivalence:

$$\text{Mod}_{\Sigma^{k-2} A}^{E_2}(\Omega \mathcal{C})^{\text{op}} \simeq \mathfrak{Z}_2(\Omega \mathcal{C}, \mathfrak{Z}_1(\text{RMod}_{\Sigma^{k-2} A}(\Omega \mathcal{C}))).$$

3. Its compatibility with the k -step condensation leads to the following equivalences:

$$\mathcal{D} \simeq \text{Mod}_{\Sigma^{k-1} A}^{E_1}(\mathcal{C}) \simeq \Sigma \text{Mod}_{\Sigma^{k-2} A}^{E_2}(\Omega \mathcal{C}), \quad \text{RMod}_{\Sigma^{k-1} A}(\mathcal{C}) \simeq \Sigma \text{RMod}_{\Sigma^{k-2} A}(\Omega \mathcal{C}). \quad \square$$

Some other related results are better summarized as pure mathematical results.

Theorem^o 1.2.5. Let $\mathcal{B}$ be an E_k -fusion n -category and A a condensable E_k -algebra in $\mathcal{B}$. We set $\Sigma^0 A = A$ and $\Sigma^i A = \text{RMod}_{\Sigma^{i-1} A}(\Sigma^{i-1} \mathcal{B})$ for $0 \leq i \leq k$. We have a natural E_{k-i} -monoidal equivalence:

$$\text{Mod}_{\Sigma^i A}^{E_{k-i}}(\Sigma^i \mathcal{B}) \simeq \Omega^{k-i} \text{Mod}_{\Sigma^k A}^{E_0}(\Sigma^k \mathcal{B}), \quad (1.2.6)$$

a natural E_{k-1-i} -monoidal equivalence $\Sigma^i \text{RMod}_A(\mathcal{B}) \simeq \text{RMod}_{\Sigma^i A}(\Sigma^i \mathcal{B})$ and an E_{k-1} -monoidal equivalence $\text{RMod}_A(\mathcal{B}) \simeq \Omega^i \text{RMod}_{\Sigma^i A}(\Sigma^i \mathcal{B})$ (see Theorem 5.2.9). We have an E_k -monoidal equivalence:

$$\text{Mod}_A^{E_k}(\mathcal{B})^{\text{op}} \simeq \mathfrak{Z}_k(\mathcal{B}, \mathfrak{Z}_{k-1}(\text{RMod}_A(\mathcal{B}))), \quad (1.2.7)$$

where $\mathfrak{Z}_k(-, -)$ is the E_k -centralizer and $\mathfrak{Z}_{k-1}(-)$ is the E_{k-1} -center (see Definition 2.3.17). When $\mathcal{B} = n\text{Vec}$, we obtain $\text{Mod}_A^{E_k}(n\text{Vec})^{\text{op}} \simeq \mathfrak{Z}_{k-1}(\Sigma A)$ (see Theorem 5.2.11).

In addition to the general theory summarized above, we also provide some general constructions of k -codimensional defect condensations. These results are summarized below.

Theorem^{ph} 1.2.6. When $\mathcal{C}$ is equipped with an explicit coordinate system, we can give a classification or a mathematical characterization of all possible condensations.

- (1) When C^{n+1} is the trivial phase, i.e., $C^{n+1} = \mathbf{1}^{n+1}$, we have $\mathcal{C} = n\text{Vec}$. An indecomposable condensable E_k -algebra $\mathcal{A}$ in $\Omega^{k-1} \mathcal{C} = (n-k+1)\text{Vec}$ is precisely an indecomposable E_k -multi-fusion $(n-k)$ -categories. By condensing $\mathcal{A}$, we obtain a new phase D^{n+1} and a gapped domain wall M^n such that (see Section 5.2.2)

$$\begin{aligned} \mathcal{D} &\simeq \text{Mod}_{\Sigma^{k-1} \mathcal{A}}^{E_1}(n\text{Vec}) \simeq \mathfrak{Z}_0(\Sigma^k \mathcal{A})^{\text{op}}, & (\mathcal{M}, m) &\simeq (\text{RMod}_{\Sigma^{k-1} \mathcal{A}}(n\text{Vec}), \Sigma^{k-1} \mathcal{A}), \\ \Omega^{k-1} \mathcal{D} &\simeq \text{Mod}_{\mathcal{A}}^{E_k}((n-k+1)\text{Vec}) \simeq \mathfrak{Z}_{k-1}(\Sigma \mathcal{A})^{\text{op}}, & \Omega_m^{k-1} \mathcal{M} &\simeq \text{RMod}_{\mathcal{A}}((n-k+1)\text{Vec}). \end{aligned}$$

When $k = 1$, we can obtain all non-chiral $n+1$ D topological orders D^{n+1} in this way by choosing a proper $\mathcal{A}$, and $\mathcal{A}$ is ‘Lagrangian’ (in the sense that $D^{n+1} = \mathbf{1}^{n+1}$) if and only if $\mathcal{A}$ is a non-degenerate indecomposable multi-fusion $(n-1)$ -category (see Definition 2.3.31).

When $k = 2$, $\mathcal{D} \simeq \Sigma \text{Mod}_{\mathcal{A}}^{E_2}((n-1)\text{Vec})$, and $\mathcal{A}$ is ‘Lagrangian’ (in the sense that $D^{n+1} = \mathbf{1}^{n+1}$) if and only if $\mathcal{A}$ is a non-degenerate braided fusion $(n-2)$ -category (see Definition 2.3.31).

- (2) When C^{n+1} is anomaly-free and simple, we have $\mathcal{C} = \Sigma \Omega \mathcal{C} = \text{RMod}_{\Omega \mathcal{C}}(n\text{Vec})$ as fusion n -categories. An indecomposable condensable E_1 -algebra in $\mathcal{C}$ is precisely an indecomposable multi-fusion $(n-1)$ -category $\mathcal{A}$ equipped with a braided monoidal functor $\phi : \Omega \mathcal{C} \rightarrow \mathfrak{Z}_1(\mathcal{A})$. By condensing $\mathcal{A}$ in $\mathcal{C}$, we obtain a condensed phase D^{n+1} and a gapped domain wall M^n such that (see Theorem^{ph} 3.3.26)

$$\begin{aligned} \mathcal{D} &\simeq \text{Mod}_{\mathcal{A}}^{E_1}(\mathcal{C}) \simeq \Sigma \mathfrak{Z}_2(\Omega \mathcal{C}, \mathfrak{Z}_1(\mathcal{A}))^{\text{op}}, & \Omega \mathcal{D} &\simeq \mathfrak{Z}_2(\Omega \mathcal{C}, \mathfrak{Z}_1(\mathcal{A})); \\ \mathcal{M} &\simeq \text{RMod}_{\mathcal{A}}(\mathcal{C}) \simeq \text{RMod}_{\mathcal{A}}(n\text{Vec}), & m &= \mathcal{A} \in \text{RMod}_{\mathcal{A}}(\mathcal{C}). \end{aligned}$$

Moreover, $\mathcal{A}$ is Lagrangian in the sense that $D^{n+1} = \mathbf{1}^{n+1}$ if and only if ϕ is an equivalence.

- (3) Given two Morita equivalent anomaly-free simple topological orders C^{n+1} and D^{n+1} connected by a gapped domain wall K^n as illustrated in (3.3.20), there is a one-to-one correspondence between

- $(\mathcal{K} \boxtimes \mathcal{T})$ -modules modulo the equivalence relation: $\mathcal{X} \sim \mathcal{Y}$ if $\mathcal{Y} \simeq \mathcal{X} \boxtimes \mathcal{V}$ for $\mathcal{V} \in (n+1)\text{Vec}^\times$;
- indecomposable condensable E_1 -algebras in $\mathcal{C}$ that define 1-codimensional defect condensations from C^{n+2} to D^{n+2} in the Morita class $[K^{n+1} \boxtimes T^{n+1}]$,


defined by $\mathcal{X} \mapsto A_{\mathcal{X}} = \text{Fun}_{\mathcal{K} \boxtimes \mathcal{T}}(\mathcal{X}, \mathcal{X})^{\text{op}}$ (in the coordinate $\mathcal{C} = \text{RMod}_{\Omega \mathcal{C}}(n\text{Vec})$). The condensable E_1 -algebra $A_{\mathcal{X}}$ is simple if and only if the module $\mathcal{X}$ is indecomposable (see Theorem^{ph} 3.3.29).

- (4) When C^{n+1} is anomaly-free, simple and non-chiral, i.e., C^{n+1} admitting a gapped boundary B^n , in this case, we have a new coordinate system $\mathcal{C} = \text{BMod}_{\mathcal{B}|\mathcal{B}}(n\text{Vec})^{\text{op}}$ for an indecomposable multi-fusion $(n-1)$ -category $\mathcal{B}$. In this case, an indecomposable condensable E_1 -algebra in $\mathcal{C}$ is precisely an indecomposable multi-fusion $(n-1)$ -category $\mathcal{A}^{\text{op}}$ equipped with a monoidal functor $\psi : \mathcal{B} \rightarrow \mathcal{A}$. By condensing $\mathcal{A}^{\text{op}}$ in $\mathcal{C}$, we obtain a condensed phase D^{n+1} and a gapped domain wall M^n such that (see Theorem^{ph} 3.3.32)

$$\begin{aligned} \mathcal{D} &\simeq \text{Mod}_{\mathcal{A}^{\text{op}}}^{E_1}(\mathcal{C}) \simeq \text{Mod}_{\mathcal{A}^{\text{op}}}^{E_1}(n\text{Vec}) \simeq \text{Mod}_{\mathcal{A}}^{E_1}(n\text{Vec})^{\text{op}} \simeq \Sigma \mathfrak{Z}_1(\mathcal{A}), \\ (\mathcal{M}, m) &= (\text{LMod}_{\mathcal{A}}(\mathcal{C}), \mathcal{A}) = (\text{BMod}_{\mathcal{A}|\mathcal{B}}(n\text{Vec}), \mathcal{A}). \end{aligned}$$

Moreover, $\mathcal{A}^{\text{op}}$ is Lagrangian if and only if $\mathcal{A}$ is a non-degenerate multi-fusion $(n-1)$ -category.


- (5) When C^{n+1} is anomaly-free, simple and non-chiral, there are two coordinate systems, i.e., $\mathcal{C} = \text{RMod}_{\Omega \mathcal{C}}(n\text{Vec})$ and $\mathcal{C} = \text{BMod}_{\mathcal{B}|\mathcal{B}}(n\text{Vec})^{\text{op}}$. The same condensable E_1 -algebra in $\mathcal{C}$ can be expressed in two coordinates. The coordinate transformation for condensable E_1 -algebras in $\mathcal{C}$ is illustrated below (see Theorem^{ph} 3.3.33).



Two definitions of the same algebra in $\mathcal{C}$:

- (2) the central functor $\mathfrak{Z}_1(\mathcal{B}) \rightarrow \text{Fun}_{\mathcal{A}|\mathcal{B}}(\mathcal{A}, \mathcal{A})$ defines $\text{Fun}_{\mathcal{A}|\mathcal{B}}(\mathcal{A}, \mathcal{A}) \in \text{Alg}_{E_1}^c(\text{RMod}_{\mathfrak{Z}_1(\mathcal{B})}(n\text{Vec}))$;
- (1) the monoidal functor $\mathcal{B} \rightarrow \mathcal{A}$ defines $\mathcal{A}^{\text{op}} \in \text{Alg}_{E_1}^c(\text{BMod}_{\mathcal{B}|\mathcal{B}}(n\text{Vec})^{\text{op}})$.

- (6) Let D^{n+1} be the anomaly-free, simple and non-chiral phase obtained in the case (1) by condensing an E_k -fusion $(n-k)$ -category $\mathcal{A}$ as a k -codimensional defect in 1^{n+1} . A simple condensable E_k -algebra B in $\Omega^{k-1}\mathcal{D} \simeq \text{Mod}_{\mathcal{A}}^{E_k}((n-k+1)\text{Vec}) \simeq \mathfrak{Z}_{k-1}(\Sigma \mathcal{A})^{\text{op}}$ is precisely defined by an E_k -monoidal functor $\mathcal{A} \rightarrow \mathcal{B}$ for an E_k -fusion $(n-k)$ -category $\mathcal{B}$. By condensing $B \in \Omega^{k-1}\mathcal{D}$ in D^{n+1} in k steps, we obtain a new condensed phase E^{n+1} such that



$$\mathcal{E} = \text{Mod}_{\Sigma^{k-1}\mathcal{B}}^{E_1}(\mathcal{D}) \simeq \text{Mod}_{\Sigma^{k-1}\mathcal{B}}^{E_1}(n\text{Vec}),$$

$$\Omega^{k-1}\mathcal{E} \simeq \text{Mod}_{\mathcal{B}}^{E_k}((n-k+1)\text{Vec}) \simeq \mathfrak{Z}_{k-1}(\Sigma \mathcal{B})^{\text{op}},$$

where $\Sigma^{k-1}\mathcal{B}$ is equipped with an E_1 -monoidal functor $\Sigma^{k-1}\mathcal{A} \rightarrow \Sigma^{k-1}\mathcal{B}$ and should be viewed as a condensable E_1 -algebra in $\mathcal{D}$. Moreover, the gapped domain wall $N^n = (\mathcal{N}, n)$ produced by this condensation is given by


$$(\mathcal{N}, n) = (\text{BMod}_{\Sigma^{k-1}\mathcal{A}|\Sigma^{k-1}\mathcal{B}}(\Sigma^{k-1}\mathcal{B}, \Sigma^{k-1}\mathcal{B}), \Sigma^{k-1}\mathcal{B}),$$

$$\Omega_n^{k-1}\mathcal{N} = \Omega^{k-2}\text{Fun}_{\Sigma^{k-1}\mathcal{A}|\Sigma^{k-1}\mathcal{B}}(\Sigma^{k-1}\mathcal{B}, \Sigma^{k-1}\mathcal{B}) \simeq \text{RMod}_{\mathcal{B}}(\text{Mod}_{\mathcal{A}}^{E_k}((m+1)\text{Vec})).$$

Note that this result is an E_k -analogue of (4) and is explained in details in Theorem^{ph} 5.2.20, and the $k=2$ case is explained in Theorem^{ph} 4.2.25. $\square$

Using (2) in above theorem and Theorem^{ph} 2.4.12 and 2.4.18, we obtain an efficient way to construct and classify all 1-codimensional defect condensations between any two $n+2$ D topological orders (see Corollary^{ph} 3.3.37), and a one-to-one correspondence between Lagrangian E_1 -algebras (or Lagrangian E_2 -algebras) in a non-chiral $n+2$ D topological order and certain module n -categories (see Theorem^{ph} 4.2.23, 4.2.25 and Corollary^{ph} 2.4.15). We summarize the key results below.

Theorem^{ph} 1.2.7. For a non-chiral simple topological order $Z(A)^{n+2}$ with a simple gapped boundary A^{n+1} (i.e., $\mathcal{A}$ is a fusion n -category) and a non-degenerate fusion n -category $\mathcal{T}$, there are one-to-one correspondences among the following sets: (the idea is illustrated below)



- (1) left indecomposable $\mathcal{A} \boxtimes \mathcal{T}$ -modules modulo the equivalence relation: $\mathcal{X} \sim \mathcal{Y}$ if $\mathcal{Y} \simeq \mathcal{X} \boxtimes \mathcal{V}$ for an invertible separable n -category $\mathcal{V}$;
- (2) simple gapped boundaries K^{n+1} of $Z(A)^{n+2}$ in the Morita class $[A^{n+1} \boxtimes T^{n+1}]$ (see Definition 2.2.19);
- (3) Lagrangian E_1 -algebras in $\Sigma\mathfrak{Z}_1(\mathcal{A})$ in the Morita class $[\mathcal{A} \boxtimes \mathcal{T}]$ (see Remark 3.3.27);
- (4) Lagrangian E_2 -algebras in $\mathfrak{Z}_1(\mathcal{A})$ in the 2-Morita class $[\mathcal{A} \boxtimes \mathcal{T}]$ (see Theorem^{ph} 4.2.6).

By allowing $\mathcal{T}$ to run through all Morita classes, we obtain a classification of all gapped boundaries of $Z(A)^{n+2}$, all Lagrangian E_1 -algebras in $\Sigma\mathfrak{Z}_1(\mathcal{A})$ and all Lagrangian E_2 -algebra in $\mathfrak{Z}_1(\mathcal{A})$. When $n = 1$, we recover an old result in 1-categories [KR08, DMNO13].

Moreover, when the fusion n -category $\mathcal{A}$ is indecomposable as a separable n -category, i.e., $\mathcal{A} \simeq \Sigma\Omega\mathcal{A}$, there is a one-to-one correspondence between the set of braided monoidal functors $\Omega\mathcal{A} \rightarrow \mathcal{O}$ (up to isomorphisms) for a non-degenerate braided fusion $(n-1)$ -category $\mathcal{O}$ and that of Lagrangian E_2 -algebras in $\mathfrak{Z}_1(\mathcal{A})$. This result generalizes an earlier result for the $n = 2$ case [JFR23, DX24]. $\square$

We also obtain many other interesting new results along the way as we develop our theory. We list a few important examples and highlight other important contents of this work.

- (1) The theory of the categories of topological defects in topological orders in all dimensions was initiated in [KW14, KWZ15], and was fully established in [GJF19, JF22] (see also [KLWZZ20b]), and was further developed in [KZ22c, KZ24, KZ22b]. We give a comprehensive review of the historical developments of this theory with an emphasis on the physical intuitions and physical principles instead of mathematical rigor. Although the main results of this theory have already appeared in literature [KWZ15, JF22, KLWZZ20b, KZ22b], the complete physical reasoning behind this theory, including its limitations, is told for the first time in Section 2.
- (2) We review the complete mathematical formulation of boundary-bulk relation as the center functor in Section 2.4. In Section 2.4.2, we point out that the proof of Proposition* 3.13 in [KZ24] actually provide a precise mathematical characterization of the non-surjectivity and the non-fully-faithfulness of the center functor, which is stated explicitly in Theorem 2.4.7 or 2.4.8. This further leads to interesting result such as Theorem^{ph} 2.4.12, Theorem^{ph} 2.4.18 and Corollary^{ph} 2.4.15, Theorem^{ph} 2.2.26, Theorem^{ph} 3.3.14 and results summarized in Theorem^{ph} 1.2.7.
- (3) In addition to the general constructions of condensable E_k -algebras (see Section 3.3.2, 4.2.2, 5.2.2), we also provide some concrete examples of 1-codimensional, 2-codimensional and k -codimensional condensations in Section 3.3.3, 3.3.5, 4.1.2, 4.2.4, 5.1.2 and 5.2.3. Even for the well-known theory of anyon condensations in 2+1D, we retell the story from a new perspective, emphasizing the role of higher algebras and higher representations in the condensation theory. This perspective makes the later generalizations to condensations in higher (co)-dimensions very natural. We also provide many examples of anyon condensations in Section 4.1.2. Readers can use it as a guide to literature.

- (4) We develop a geometric approach towards defect condensations. In particular, we develop the geometric approach towards the particle condensations in 1+1D in Section 3.2.2 and the anyon condensations in 2+1D in Section 4.1.3 (see the key result Theorem 4.1.19). In this geometric approach, we emphasize the construction of condensable algebras via internal homs, a mathematical notion which is reviewed in Section 3.1.2. The complete geometric theory is quite involved. In order to control the complexity of mathematics, we move some mathematically technical results to Appendix A.1 and A.2. In particular, in Appendix A.2, we develop a relative version of classical Eilenberg-Watts calculus that has independent values. These geometric theories can be generalized to higher dimensions and lead to a theory of integrals, which is briefly outlined in Section 6.2. It seems that it includes but generalizes the so-called α -version of factorization homology [Lur17, AFT17, AFT16, AF20] and should be related (or even equivalent) to the so-called β -version of factorization homology [AFR18, AF24] (see Remark 6.2.2).
- (5) In Section 5.3, we establish the connections or equivalences among various notions of gauging 0-form G -symmetries in a 2+1D topological order, including G -crossed extension approach [BBCW19], minimal nondegenerate extension approach [LKW16a, KIWZZ20b] and the 1-codimensional defect condensation approach (or via a monoidal functor [KZ24, LYW24]). In Remark 5.3.16, we largely generalize the notion of a symmetry as a morphism between two ‘higher quiches’. The condensation theory in this work automatically provides a theory of gauging such a generalized symmetry.
- (6) Although we mainly focus on the condensations of topological defects in topological orders, our theory naturally generalizes to situations beyond gapped defects. In Section 6.3, we briefly discuss how the theory generalizes to the condensations of liquid-like gapless defects in topological orders and the condensations of quantum liquids by focusing only on the topological data of gapless defects or phases. We also provide an illustrating example in 2+1D (see the key result in Theorem^{ph} 6.3.4). More detailed study of the condensations of gapless defects will appear elsewhere.
- (7) The higher representation theory of higher algebras is much richer than that of an E_1 -algebra. In addition to results summarized in this section, we obtain some other interesting results along the way. In Section 6.1, we briefly discuss some basic ingredients in the higher Morita theory for condensable E_k -algebras, including introducing the notions of various higher Morita equivalences and proving some basic results. We also introduce the notion of the enveloping algebra of a condensable E_k -algebra, which is an important tool in the study of higher representation theory.

In addition to many important algebraic constructions, we emphasize a geometric way to understand higher algebras and higher representations (see Section 2.3.1, 2.3.2 and 4.1.1) and almost all mathematical results appeared in this work. Moreover, we emphasize a geometric way to construct these condensable E_k -algebras as a theory of integrals (i.e., integrating local observables to a global ones). We show that some computations can be carried out by purely geometrical and physical intuitions (see Section 3.2, 3.2, 4.1.3, 4.2.1, 6.2 and 6.3.2). One of the main goals of this work is to convince readers that almost all ingredients in our theory of topological orders and defect condensations, based on higher categories, higher algebras and higher representations, have geometric meanings. Moreover, many deep and new algebraic results can be derived from the geometric intuitions with the help from only a handful physical principles.

Remark 1.2.8. Our theory shows that defect condensations is essentially a theory of higher algebras (e.g., E_k -algebras) and higher representations (e.g., E_k -modules). Although we try to explain many of these mathematical notions in details, a mathematically rigorous treatment of this subject is far beyond this paper, which originally aimed at physics oriented readers. A more mathematical treatment of this subject will appear in a mathematical companion of this paper [KZZZ25]. In this work, we provide an almost self-contained story, in which most mathematical complexity is replaced by physical principles and self-evident geometric or physical intuitions. We show that these geometric and physical intuitions, together with necessary physical principles, are powerful enough for us to build a self consistent and rather complete theoretical framework. This framework allows us to establish a precise correspondence between geometric or physical manipulations and algebraic computations. Once this correspondence is established, we are able to do precise computations based on purely geometric intuitions. This leads us to many new results that are not available in mathematical literature. In order to make this framework to work, we make a technical assumption that we always work within separable higher categories [KZZ2b], and (higher) algebras are always separable (also called condensable in this work). Their higher centers or higher centralizers, internal homs and their categories

of E_k -modules are automatically separable as higher algebras or higher categories [KZ22b, KZ24] (see also Remark 2.3.27). Some parts of this statement were known in mathematical literature for 1-categories and 2-categories (see for example [Ost03a, Déc25, Déc23b, Déc23a, DX24, Xu24a]). $\diamond$

Remark 1.2.9. All the condensations studied in this work are bosonic in nature. We leave a systematic study of fermionic condensations to the future. See [ALW19, WW17, LSCH21, GK21, ZWG22] and references therein for existing works on this topics. Although we restrict our study in the setting of E_k -multi-fusion higher categories, we believe that all types of condensation phenomena, including those in QFT's with continuous symmetries, can be formulated as higher algebras and higher representations in more general higher-monoidal higher categories (see some examples in Section 6.3 and see also [Sto25]). $\diamond$

Remark 1.2.10. On the one hand, for high energy theorists or the mathematical physicists following the program of functorial quantum field theories (FQFT), the notion of a QFT is often defined via path integrals over all spacetime manifolds. In this setup, gauging a q -form symmetry in an n D QFT was defined roughly by a process of “summing over insertions of topological defects of codimension $(q + 1)$ in an n -dimensional spacetime manifold” [RSS23]. It turns out that, before the rise of higher form symmetries, there was a precisely and rigorous mathematical theory of this gauging process under the name of ‘orbifold TQFT’ [FFRS07, FFRS10, CR16, CRS18, CRS19, CRS20, CMRSS21, Mul24, MR23, CM23, CMRSS24]. On the other hand, for condensed matter theorists, the notion of a topological order (or a SPT/SET/SSB order) is naturally defined on a spatial open disk (often by a lattice model). Therefore, it is natural to view a topological order as a ‘fully extended TQFT defined on a spatial open disk’ or an algebraic quantum field theory (AQFT) defined by local observables (which often form a higher categorical structure [KW14, KWZ15, JF22, KZ22b]). The relation between a FQFT and an AQFT was known in principle. At least for fully dualizable QFT's, in principle or as a folklore, the partition function on a non-trivial spacetime manifold M in a FQFT can be obtained from an AQFT by integrating observables on an open disk over M . For TQFT's, the mathematical foundation of this integration is an on-going program of the so-called factorization homology [Lur17, AF20, AFT17, AFT16, AFR18, AF24] (see Remark 6.2.2).

The manifest difference between the mathematical formulation of ‘gaugings’ in FQFT's and that of ‘condensations’ in AQFT's is due to the fact that an integration over a spacetime manifold changes the mathematical structure of the observables. For example, integrating a system of anyons (i.e., a modular tensor category) over a closed spatial surface M produces a pair (Vec, u_M) , where u_M is the space of ground states on M [AKZ17]. Note that the data u_M is the only global observable on M . All the E_2 -monoidal structures (i.e., the fusions and braidings of anyons) that are well-defined on an open 2-disk are lost after the integration. In other words, anyons, together with their fusions and braidings, are not global observables (see [AKZ17, KZ22a] for more discussion). Therefore, the explicitly structured mathematical theory of defect condensations based on higher categories, higher algebras and higher representations is a distinct feature of an AQFT². It remains an important open problem to compare the partition functions obtained by integrating our condensation results with those obtained by gauging q -form symmetries and those in orbifold TQFT's. $\diamond$

1.3 Layout, conventions and notations

The discouraging length of this paper is an avoidable consequence of the incredible richness of the theory of defect condensations and our goal of providing a comprehensive story of it. For readers convenience, we summarize the layout of this work, the conventions and the notations used in this work.

Layout: In Section 2, we review the mathematical theory of topological defects in an $n+1$ D topological orders based on the remote detectable principle, boundary-bulk relation and condensation complete principle. We also review the notions of E_m -multi-fusion n -categories and explain that an E_m -monoidal n -category can be viewed as an n -category equipped with a tensor product in each of n independent dimensions. We also introduce other useful notations, new notions and new results along the way. Importantly, in Section 2.4, we derive some new results from some earlier results on the center functor in [KZ24]. These results are very important for later constructions and classifications of the condensations of defects of codimension 1 or 2. In Section 3, we show that a phase transition from a topological order C^{n+1} to another D^{n+1} can be defined by the condensation of a 1-codimensional topological defects. In Section 4, we show that, when

²In orbifold TQFT's [CRS20], it is unavoidable to introduce something similar or equivalent to a higher algebra as the initial data in their construction (see Remark 3.3.28).

C^{n+1} is anomaly-free, the same condensation can be defined by condensing a 2-codimensional topological defect directly. When $n = 2$, this condensation is precisely the usual anyon condensation in 2+1D topological orders. We start from a review of anyon condensation in 2+1D but from a new perspective in Section 4.1.1 and provide a large amount of examples in Section 4.1.2. We also develop a geometric anyon condensation theory in Section 4.1.3. Then in Section 4.2, we generalize it to higher dimension. In Section 5, we study the condensation of k -codimensional topological defects and give the main results of this work as summarized in this Section. In Section 6, we outline a few directions that this theory should be further developed or generalized, including a under-developed higher Morita theory, a under-developed theory of integrals (which might be related to the theory of factorization homology), a condensation theory of (potentially gapless) liquid-like defects or quantum liquids and some applications.

Conventions: Throughout this work, a topological order is defined up to invertible ones. In order to highlight different types of results, we introduce the following notations.

1. We use ‘Theorem^{ph}’ to represent a physical result, which usually contains physical notions such as topological orders, quantum liquids, boundaries, walls, defects and condensations.
2. We use ‘Theorem^o’ to represent a result, which makes sense mathematically without referring to its physical meaning but lacks of a complete mathematical proof. Therefore, it should be viewed as a mathematical conjecture. In this work, it is either derived from our physical theory of condensations or based on the ‘mathematical theory’ of separable or E_k -multi-fusion higher categories [GJF19, JF22, KZ22b]. This ‘mathematical theory’ lacks of a complete mathematical foundation.
3. We use ‘Theorem’ to represent a mathematically rigorous result.

We use $n+1D$ to denote the spacetime dimension. We mainly use spacetime dimension in this work because the $n+1D$ spacetime dimension matches with the $n+1$ -levels of hierarchical structure of an n -category. However, almost all the illustrating pictures are drawn in the spatial dimensions because E_k -monoidal structure matches with the spatial dimension. For example, particles on a line in the spatial dimension can be fused in one spatial dimension, thus forming an E_1 -fusion 1-category or simply a fusion 1-category.

Notations: In this work, we systematically introduce a lot of formal notations. For example, we denote a simple statement like: “ $\mathcal{A}$ is an indecomposable multi-fusion n -category” by “ $\mathcal{A} \in \text{Alg}_{E_1}^c((n+1)\text{Vec})$ ”. Although we are aware of the danger of turning physics oriented readers away, we do it for an important reason. We slowly introduce the formal language when we review the more familiar situations in lower dimensions (but often from a new perspective). Once the readers get used to the new formal language, they will start to appreciate the power of categorical language when we move on to higher dimensional theories. For example, a simple notation $A \in \text{Alg}_{E_1}^c(\text{Mod}_A^{E_k}(\mathcal{C}))$ means a thousand words. Indeed, it summarizes incredible amount of structures, which were completely out of reach before the advent of this categorical language. Without such a powerful language, the higher condensation theory is simply impossible.

Most of the notations are introduced carefully when they first appear. Here, we highlight a few important conventions on notations.

1. We denote $n+1D$ (potentially anomalous) topological orders by $A^{n+1}, B^{n+1}, C^{n+1}, D^{n+1}, \dots$, and denote their categories of (‘intrinsic’ topological defects see Section 2.1.2):
 - topological defects of codimension 1 and higher by $\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}, \dots$, respectively;
 - topological defects of codimension 2 and higher by $\Omega\mathcal{A}, \Omega\mathcal{B}, \Omega\mathcal{C}, \Omega\mathcal{D}, \dots$, respectively;
 - topological defects of codimension $(k-1)$ and higher by $\Omega^k\mathcal{A}, \Omega^k\mathcal{B}, \Omega^k\mathcal{C}, \Omega^k\mathcal{D}, \dots$, respectively.

Note that we carefully distinguish A^{n+1} with $\mathcal{A}$ simply because they have different meanings and play different roles in the category of topological orders.

2. Given an n -category $\mathcal{S}$, we denote the category obtained from flipping all k -morphisms in $\mathcal{S}$ by $\mathcal{S}^{\text{op}k}$. For convenience, we abbreviate $\mathcal{S}^{\text{op}1}$ to $\mathcal{S}^{\text{op}}$.
3. If a topological order A^{n+1} is anomalous, it has a unique $n+2D$ bulk phase denoted by $Z(A)^{n+2}$ such that A^{n+1} becomes a gapped boundary of $Z(A)^{n+2}$ [KW14]. Recall that $Z(A)^{n+2}$ is also called *gravitational anomaly* of A^{n+1} in [KW14].

4. We use TC^3 and Is^3 to denote 2+1D toric code phase and 2+1D Ising topological order, respectively. For a finite group G and $\omega \in Z^{n+1}(G, U(1))$, we use $\text{GT}_{(G,\omega)}^{n+1}$ to denote the $n+1$ D twisted finite gauge theory (or Dijkgraaf-Witten theory). When ω is trivial, it is called a finite gauge theory and is denoted by GT_G^{n+1} . Note that $\text{TC}^3 = \text{GT}_{\mathbb{Z}_2}^3$. Following the first convention, we denote the categories of all topological defects of codimension 1 and higher in TC^3 , Is^3 , $\text{GT}_{(G,\omega)}^{n+1}$ and GT_G^{n+1} by $\mathcal{TC}$, $\mathcal{Is}$, $\mathcal{GT}_{(G,\omega)}^{n+1}$ and $\mathcal{GT}_G^{n+1}$, respectively. According to our first notation convention, the usual Ising modular tensor category, as a category of anyons (or defects of codimension 2 and higher), is denoted by $\Omega\mathcal{Is}$. Restricted in Example 5.3.10, the notations $\text{Is}_\pm$ are used to denote two Ising type fusion 1-categories, respectively.
5. We use Vec to denote the category of finite dimensional vector spaces over the complex number field $\mathbb{C}$. It can be treated as the definition of the delooping of $\mathbb{C}$, i.e., $\Sigma\mathbb{C} := \text{Vec}$ [GJF19].
6. We use $(n+1)\text{Vec}$, which is defined by iterated deloopings $\Sigma^{n+1}\mathbb{C}$ [GJF19] (see Section 2.2 for more details), to denote the category of separable n -categories. Intuitively, one can view a separable n -category $\mathcal{S}$ as a “finite dimensional n -vector space”. We set $\mathcal{S}^{\text{op}} := \mathcal{S}^{\text{op}1}$, i.e., flipping all 1-morphisms.
7. We denote the full subcategory of $(n+1)\text{Vec}$ consisting of invertible objects by $(n+1)\text{Vec}^\times$.
8. $\mathcal{B} \in \text{Alg}_{E_k}^c((n+1)\text{Vec})$ means that $\mathcal{B}$ is an E_k -multi-fusion n -category, and, at the same time, a condensable E_k -algebra in $(n+1)\text{Vec}$.
9. $A, B \in \text{Alg}_{E_k}^c(\mathcal{B})$ means that A and B are condensable E_k -algebras in $\mathcal{B}$.
10. $\text{LMod}_A(\mathcal{B})$, $\text{RMod}_A(\mathcal{B})$, $\text{BMod}_{A|B}(\mathcal{B})$ represent the categories of left A -modules, right A -modules, A - B -bimodules in $\mathcal{B}$, respectively.
11. $\text{Mod}_A^{E_k}(\mathcal{B})$ denotes the categories of E_k - A -modules (or E_k -modules over A) in $\mathcal{B}$. See Section 4.1.1 and Section 5.2.1 for more explanation of the notion of an E_k - A -module.
12. $B \in \text{Alg}_{E_k}(\text{Mod}_A^{E_k}(\mathcal{B}))$ means that B is an E_k - A -module in $\mathcal{B}$ and, at the same time, an E_k -algebra in $\mathcal{B}$ such that the defining data of the E_k -algebra structure on B are all (higher) morphisms in $\text{Mod}_A^{E_k}(\mathcal{B})$.
13. For $\mathcal{A}, \mathcal{B} \in \text{Alg}_{E_1}^c((n+1)\text{Vec})$ and $\mathcal{M}, \mathcal{N} \in \text{BMod}_{\mathcal{A}|B}((n+1)\text{Vec})$, we use $\text{Fun}_{\mathcal{A}|B}(\mathcal{M}, \mathcal{N})$ to denote the category of bimodule functors from $\mathcal{M}$ to $\mathcal{N}$.
14. In this work, $[x, y]$ denotes the internal hom which is briefly reviewed in Section 3.1.2.
15. For an E_k -monoidal functor $F : \mathcal{A} \rightarrow \mathcal{B}$, we use $\mathfrak{Z}_k(F)$ or $\mathfrak{Z}_k(\mathcal{A}, \mathcal{B})$ to denote the E_k -centralizer of F (see Definition 2.3.17), and use $\mathfrak{Z}_k(\mathcal{A}) = \mathfrak{Z}_k(\text{id}_{\mathcal{A}})$ to denote E_k -center of $\mathcal{A}$.
16. For a monoidal n -category $\mathcal{C}$, the meaning of the notation $\mathcal{C}^{\text{op}}$ is explained in Convention 2.1.16 and Definition 2.3.5.

Acknowledgements: We would like to thank David Ayala, Ansi Bai, Nils Carqueville, John Francis, Ling-Yan Hung, Tian Lan, Gregory Moore, Ingo Runkel, Nathan Seiberg, Rui Wen, Xiao-Gang Wen and Hao Xu for useful discussion and comments. LK and ZHZ are supported by NSFC (Grant No. 11971219 & 12574175) and by Guangdong Provincial Key Laboratory (Grant No. 2019B121203002) and by Guangdong Basic and Applied Basic Research Foundation (Grant No. 2020B1515120100). ZHZ is also supported by Wu Wen-Tsun Key Laboratory of Mathematics at USTC of Chinese Academy of Sciences. JHZ is partially supported by NSFC (Grant No.11571343). HZ is supported by NSFC under Grant No. 11871078 and by Startup Grant of Tsinghua University and BIMS.

2 Categories of topological defects

Based on three important guiding principles: Remote Detectable Principle [Lev13, KW14], Boundary-Bulk Relation [KK12, KWZ15, KWZ17] and Condensation Completion Principle [KS11a, KK12, FSV13, CR16, Kon14, KW14, DR18, KTZ20b, GJF19, JF22, KLWZZ20b], the mathematical theory of topological defects in topological orders was initiated and developed in some earlier works [KW14, KWZ15, LKW18, LW19, KTZ20b], and was fully established in [GJF19, JF22] (also in [KLWZZ20b]), and was further developed and generalized to SPT/SET orders and gapped/gapless quantum liquids in [KLWZZ20b, KLWZZ20a, KZ22b, KZ24, KZ22c]. It forms the theoretical foundation of our higher condensation theory and are used throughout this work. In this section, we review this theory in details and introduce some important notions and notations along the way. Although the main results of this theory have already appeared in literature [KWZ15, JF22, KLWZZ20b, KZ22b], we believe that the complete physical reasoning behind this theory, including its limitations, is told here for the first time. In Section 2.4, we use center functor to derive some new results that become important to the classification of 1-codimensional defect condensations and that of Lagrangian E_2 -algebras in later sections.

2.1 Categories of topological defects

2.1.1 2+1D topological orders

The systematic study or classification of topological defects in TQFT's or topological orders was initiated from that of the Wilson lines in 2+1D TQFT's [Wit89, RT91] and, around the same time, that of the anyons in 2+1D anomaly-free topological orders [MS89c, FFK89, FRS89, FG90, Reh90, MR91, Wen91] (preceded by many earlier works on fractional statistics of fields and anyons [Frö76, LM77, GMS80, GMS81, Wil82, Wu84, ASW84, Fro88]). We start from a review of the theory of anyons in a 2+1D anomaly-free topological order before we move onto the higher dimensional theories.

Consider a 2+1D anomaly-free topological order C^3 (see Remark 2.1.1). It can have 2-codimensional topological defects, which are particle-like topological excitations in spatial dimension picture and are also called anyons or particles. In the spacetime picture, an anyon is also called a Wilson line, which is the world line of the anyon. It is possible to have 0D defects connecting two potentially different Wilson lines. Such a 0D defect, or a 3-codimensional defect, is also called an instanton. Anyons can be fused. More precisely, for two anyons a and b , their fusion is denoted by $a \otimes b$. Moreover, anyons can be braided. This amounts to a braiding isomorphism

$$c_{a,b} : a \otimes b \xrightarrow{\sim} b \otimes a.$$

The composed isomorphism $(a \otimes b \xrightarrow{c_{a,b}} b \otimes a \xrightarrow{c_{b,a}} a \otimes b)$ is called the double braiding, which amounts to an adiabatic move of the a -particle along a circle around the b -particle as illustrated in Figure 1. The complete set of anyons and instantons, together with the fusions and braidings among anyons, form a mathematical structure called a braided fusion 1-category³ [Wit89, MS89a, FRS89, ZHK89, FG90, Reh90, MR91, Wen91, Kit06] (see [KZ22a] for a recent review). We denoted this braided fusion 1-category by $\Omega\mathcal{C}$, a notation which represents the looping of $\mathcal{C}$ and is explained later in (2.1.1). Moreover, the braidings of $\Omega\mathcal{C}$ are required to be *non-degenerate* in the sense that the S -matrix, whose entries are the trace of double braidings of two simple objects, is non-degenerate [MS89a, Tur20] (see also [BK01, Kit06] for reviews). The simplest non-degenerate braided fusion 1-category is the category Vec of finite dimensional vector spaces over $\mathbb{C}$. It is precisely the category of anyons in the 2+1D trivial topological order, which is denoted by $\mathbf{1}^3$.

Concrete lattice models realizing non-chiral 2+1D topological orders (i.e. those admitting gapped boundaries), such as Kitaev's quantum double model [Kit03] and Levin-Wen models [LW05], were constructed. Anyons in these models were intensively studied and were shown to form a non-degenerate braided fusion 1-category [Kit03, KK12]. We want to emphasize that not only simple anyons (i.e., simple objects in $\Omega\mathcal{C}$) are physical, but also all the composite anyons (i.e., direct sums of simple objects in $\Omega\mathcal{C}$), which can be realized in lattice models (see [KZ22a] for a more detailed explanation). It was proposed in [Kit06] that a 2+1D topological order C^3 can be completely characterized by the data $\Omega\mathcal{C}$ (up to invertible topological orders).

³In physics, $\Omega\mathcal{C}$ is necessarily a modular tensor category, which is a non-degenerate braided fusion 1-category equipped with a ribbon structure [MS89a, Tur20]. Moreover, unitarity is often assumed. We prefer to ignore both structures for the reason that the roles played by them in condensation theory are not essential. But we secretly assume the unitarity in all pictures.




Figure 1: This picture illustrate some defects in a 2+1D topological order C^3 in spatial dimensions: two 1-codimensional defect $x, y \in \mathcal{C}$, a domain wall between $\mathbb{1}_e$ and x , a domain wall between $\mathbb{1}_e$ and y and two anyons $a, b \in \Omega\mathcal{C}$.

For example, a non-chiral 2+1D topological order (see Definition 2.1.9) can be described by the Drinfeld center $\mathfrak{Z}_1(\mathcal{A})$ [LW05, KK12] of a fusion 1-category $\mathcal{A}$, where $\mathfrak{Z}_1(\mathcal{A})$ is automatically a non-degenerate braided fusion 1-category [Müg03b]. This proposal was soon accepted by the community.

Remark 2.1.1. The 2+1D anomaly-free topological orders in above discussion are assumed to satisfy a *simpleness* (or a stability) condition: the ground state degeneracy of this topological order defined on a 2-sphere is trivial. This condition is automatically assumed in most physical literature. We proceed our story with this simpleness assumption until Section 2.2, where we introduce a slightly more general notion of a composite topological order (see Definition 2.2.15). It is natural and indispensable in the study of the fusions among topological orders and condensations [KWZ15, KZ22b]. $\diamond$

However, this proposal became puzzling and questionable after a new discovery. In [KS11a, KK12, FSV13, CR16, Kon14], it was shown that a 2+1D topological order C^3 can have non-trivial 1-codimensional topological defects as depicted in Figure 1, and these defects, together with all higher codimensional topological defects form a monoidal 2-category $\mathcal{C}$. We explain the ingredients of this 2-category $\mathcal{C}$ below.

- 0-morphisms (or objects) are 1-codimensional topological defects (i.e., string-like defects in the spatial dimension, or strings for simplicity);
- 1-morphisms are 2-codimensional topological defects, including those 2-codimensional topological defects that are domain walls between two (potentially non-trivial) 1-codimensional topological defects.
- 2-morphisms are 3-codimensional topological defects (i.e., instantons or 0D operators in spacetime).

The tensor product of $\mathcal{C}$ is given by the horizontal fusion of two 1-codimensional topological defects (e.g., $x \otimes y$ in Figure 1), and the tensor unit of $\mathcal{C}$ is the trivial 1-codimensional topological defect, denoted by $\mathbb{1}_e$. Since a particle can be viewed as a domain wall between two trivial 1-codimensional defects, the category of particles in C^3 is precisely given by the following 1-category:

$$\Omega\mathcal{C} := \text{hom}_{\mathcal{C}}(\mathbb{1}_e, \mathbb{1}_e), \quad (2.1.1)$$

which is called the looping of $\mathcal{C}$. We denote the full sub-2-category of $\mathcal{C}$ consisting of a single object $\mathbb{1}_e$ by $B\Omega\mathcal{C}$, which is called one-point delooping of $\Omega\mathcal{C}$, i.e.

$$B\Omega\mathcal{C} := \begin{array}{c} \Omega\mathcal{C} \\ \curvearrowright \\ \mathbb{1}_e \end{array}. \quad (2.1.2)$$

Notice that $B\Omega\mathcal{C}$ does not contain more information than $\Omega\mathcal{C}$.

Remark 2.1.2. We do not give a precise definition of a topological defect. Instead, we give a brief clarification. In this work, by a k -codimensional topological defect in C^{n+1} , we mean an anomalous $(n - k + 1)$ D topological order (i.e., a gapped quantum liquid without symmetry). In other words, we assume a liquid-like property [KZ22b, KZ22c]. Physically, by ‘liquid-like’ we mean that the defect can be bent freely without any change⁴, and the fusions among such defects are well-defined. Mathematically, it means that the category of

⁴This condition amounts to say that a liquid-like defect commutes with the energy-momentum tensor, and, therefore, can be viewed as a generalized symmetry [FFRS10, GKS15] (see [SN23] for a recent review).

all such gapped liquid-like defects are well-defined and fully dualizable [Lur09]. Note that it was known that there are non-liquid quantum phases [Cha05, Haa11], which can be viewed as non-liquid-like defects in the higher dimensional trivial phase. In general, if we stack a k D topological defect with a gapped k D non-liquid, we obtain a gapped non-liquid defect. Non-liquid-like defects are not dualizable in general. In this work, we ignore non-liquid gapped defects or domain walls completely. In particular, by a ‘gapped defect’, we always mean a topological defect unless we specified otherwise. $\diamond$

When C^3 admits gapped boundaries, we have $\Omega C = \mathfrak{Z}_1(\mathcal{A})$ for a fusion 1-category $\mathcal{A}$ [KK12, FSV13, Kon14]. In this case, it was shown in [KK12] that a 1-codimensional topological defect in C^3 can be constructed in a generalized Levin-Wen models [LW05] based on the defining data of a finite semisimple $\mathcal{A}$ - $\mathcal{A}$ -bimodule $\mathcal{M}$. By “finite semisimple”, we mean a 1-category that is equivalent to a finite direct sum of Vec . We denote the 2-category of finite semisimple 1-categories by 2Vec . Then a finite semisimple $\mathcal{A}$ - $\mathcal{A}$ -bimodule simply means an $\mathcal{A}$ - $\mathcal{A}$ -bimodule in 2Vec . We denote by $\text{BMod}_{\mathcal{A}|\mathcal{A}}(2\text{Vec})$ the 2-category of $\mathcal{A}$ - $\mathcal{A}$ -bimodules in 2Vec (as objects), bimodule functors (as 1-morphisms) and bimodule natural transformations (as 2-morphisms). Then the main results in [KK12] says that

$$\mathcal{C} = \text{BMod}_{\mathcal{A}|\mathcal{A}}(2\text{Vec}). \quad (2.1.3)$$

The 2-category $\text{BMod}_{\mathcal{A}|\mathcal{A}}(2\text{Vec})$ has a natural monoidal structure defined by the relative tensor product $\boxtimes_{\mathcal{A}}$ (see [EGNO15] for a review) and the tensor unit $\mathcal{A}$. In this case, we have

$$\Omega C = \Omega \text{BMod}_{\mathcal{A}|\mathcal{A}}(2\text{Vec}) = \text{Fun}_{\mathcal{A}|\mathcal{A}}(\mathcal{A}, \mathcal{A}) \simeq \mathfrak{Z}_1(\mathcal{A}), \quad (2.1.4)$$

which is exactly our original setup.

Remark 2.1.3. There is another way to look at the 1-codimensional topological defects in C^3 . By the folding trick, a 1-codimensional topological defect in C^3 can be viewed as a gapped boundary of the double-layered system $C^3 \boxtimes \overline{C^3}$ (see Example 2.1.10). As a consequence, there are one-to-one correspondences among simple 1-codimensional topological defects in C^3 , indecomposable ΩC -modules (up to equivalences) and Lagrangian algebras in $\Omega C \boxtimes \Omega C^{\text{op}}$ [KR09, DMNO13, KK12]. $\diamond$

We give some examples of $\mathcal{C}$ that are frequently used in this work.

Example 2.1.4. 2+1D trivial topological order 1^3 : The 1-category of anyons is given by Vec . Since 1^3 can be realized by Levin-Wen model defined by the trivial fusion 1-category Vec . The monoidal 2-category of all defects in 1^3 can be identified with $\text{BMod}_{\text{Vec}|\text{Vec}}(2\text{Vec}) \simeq 2\text{Vec}$. The objects in 2Vec are precisely the labels of 1+1D anomaly-free topological orders.

1. the tensor unit $\mathbb{1} = \text{Vec}$ in 2Vec labels the trivial 1-codimensional topological defects in 1^3 , which is, at the same time, the trivial 1+1D topological order 1^2 . Note that $\text{hom}_{2\text{Vec}}(\mathbb{1}, \mathbb{1}) \simeq \text{Vec}$ is precisely the category of particles in 1^2 .
2. What about a generic object $\mathbb{1}^{\oplus k}$ in 2Vec ? It is a composite 1-codimensional topological defect in 1^3 and, at the same time, a composite 1+1D topological order $(1^2)^{\oplus k}$, which naturally occurs in the process of dimensional reduction. For example, if we define a toric code model on a narrow strap, viewed as a quasi-1+1D system, with both boundaries chosen to be the smooth boundary, we obtain the composite 1+1D topological order $(1^2)^{\oplus 2}$ (see [KZ22a] for a more detailed explanation). Note that $\text{hom}_{2\text{Vec}}(\mathbb{1}^{\oplus k}, \mathbb{1}^{\oplus k})$ is precisely the multi-fusion 1-category of particles living on the composite 1+1D topological order $(1^2)^{\oplus k}$ [KW15, AKZ17].

Example 2.1.5. 2+1D $\mathbb{Z}_2$ topological order TC^3 : Since it is realizable by the 2+1D toric code model [Kit03], we denote it by TC^3 . We denote the 2-category of topological defects in TC^3 by $\mathcal{TC}$. In this case, $\Omega \mathcal{TC}$ has four simple objects $1, e, m, f$ and the following fusion rules:

$$e \otimes e \simeq m \otimes m \simeq f \otimes f \simeq 1, \quad f \simeq e \otimes m \simeq m \otimes e.$$

The double braiding between e and m is -1 . The self-double-braidings of e and m are both trivial, and the self-double-braiding of f is -1 . Mathematically, $\Omega \mathcal{TC}$ can be identified with the Drinfeld center $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$

of $\text{Rep}(\mathbb{Z}_2)$, where $\text{Rep}(\mathbb{Z}_2)$ denote the category of finite dimensional representations of the $\mathbb{Z}_2$ group. The monoidal 2-category $\mathcal{TC}$ can be obtained either by the following monoidal equivalence [KK12]:

$$\mathcal{TC} \simeq \text{BMod}_{\text{Rep}(\mathbb{Z}_2)|\text{Rep}(\mathbb{Z}_2)}(2\text{Vec})$$

or by explicit constructions in the toric code model [KZ22a].

1. There are six simple 1-codimensional topological defects $\mathbb{1}, \vartheta, ss, sr, rs, rr$ in $\mathcal{TC}^3$, where $\mathbb{1}$ denotes the trivial 1-codimensional defect; ϑ denotes the invertible defect that exchanges the e -particles with the m -particles thus defines the e - m -duality [Bom10, KK12]; the remaining four defects ss, sr, rr, rs denote the defects obtained by stacking of two smooth/rough gapped boundaries of the toric code model as illustrated below:



where the letter ‘s’ represents the smooth boundary and the letter ‘r’ represents the rough boundary. The fusion rules given in the following table [KZ22a, Table 1]:

$\otimes$	$\mathbb{1}$	ϑ	ss	sr	rs	rr
$\mathbb{1}$	$\mathbb{1}$	ϑ	ss	sr	rs	rr
ϑ	ϑ	$\mathbb{1}$	rs	rr	ss	sr
ss	ss	sr	$2ss$	$2sr$	ss	sr
sr	sr	ss	ss	sr	$2ss$	$2sr$
rs	rs	rr	$2rs$	$2rr$	rs	rr
rr	rr	rs	rs	rr	$2rs$	$2rr$

(2.1.5)

where ‘ $2ss$ ’ represents $ss \oplus ss$ (see [KZ22a] for more explanation of this fusion rule). All six defects $\mathbb{1}, \vartheta, ss, sr, rs, rr$ can be obtained by condensing certain (potentially composite) anyons as we show explicitly in Example 5.1.8 (see also the second table in this example).

2. There are six simple $\text{Rep}(\mathbb{Z}_2)$ - $\text{Rep}(\mathbb{Z}_2)$ -bimodules: $\text{Rep}(\mathbb{Z}_2), \text{Vec}_-, \text{Rep}(\mathbb{Z}_2) \boxtimes \text{Rep}(\mathbb{Z}_2), \text{Rep}(\mathbb{Z}_2) \boxtimes \text{Vec}, \text{Vec} \boxtimes \text{Rep}(\mathbb{Z}_2), \text{Vec} \boxtimes \text{Vec}$, where Vec_- is an invertible bimodule defined by the associator $(a \odot x) \odot a \xrightarrow{-1} a \odot (x \odot a)$ when $a \in \text{Rep}(\mathbb{Z}_2)$ is the non-trivial simple object and $x \in \text{Vec}_-$. They correspond to six simple 1-codimensional topological defects $\mathbb{1}, \vartheta, ss, sr, rs, rr$ in $\mathcal{TC}^3$, respectively.

Above two ways of representing $\mathcal{TC}^3$ provide two coordinate systems of $\mathcal{TC}^3$. Similar to differential geometry, computations are often carried out in a concrete coordinate systems of a chart, and the coordinate transformations allow us to relate different charts and play an important role in the theory. It turns out that $\mathcal{TC}^3$ has a few more natural coordinate systems.

1. The monoidal equivalence $\mathcal{TC}^3 \simeq \text{RMod}_{\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))}(2\text{Vec})$ [KLWZZ20b], explained in later subsections (see Eq. (2.1.14) and Theorem 2.2.32), provides another important coordinate system.
2. Since $\text{Vec}_{\mathbb{Z}_2}$ and $\text{Rep}(\mathbb{Z}_2)$ are Morita equivalent, we have $\mathcal{TC}^3 \simeq \text{BMod}_{\text{Vec}_{\mathbb{Z}_2}|\text{Vec}_{\mathbb{Z}_2}}(2\text{Vec})$.

The coordinate transformations among different coordinate systems are summarized in the following table.

Coordinates of $\mathcal{TC}^3$	$\mathbb{1}$	ϑ	ss	sr	rs	rr
$\text{RMod}_{\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))}(2\text{Vec})$	$\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$	$\text{Vec} \oplus \text{Vec}$	$\text{Rep}(\mathbb{Z}_2)$	Vec	Vec	$\text{Vec}_{\mathbb{Z}_2}$
$\text{BMod}_{\text{Rep}(\mathbb{Z}_2) \text{Rep}(\mathbb{Z}_2)}(2\text{Vec})$	$\text{Rep}(\mathbb{Z}_2)$	Vec_-	$\text{Rep}(\mathbb{Z}_2) \boxtimes \text{Rep}(\mathbb{Z}_2)$	$\text{Rep}(\mathbb{Z}_2) \boxtimes \text{Vec}$	$\text{Vec} \boxtimes \text{Rep}(\mathbb{Z}_2)$	$\text{Vec} \boxtimes \text{Vec}$
$\text{BMod}_{\text{Vec}_{\mathbb{Z}_2} \text{Vec}_{\mathbb{Z}_2}}(2\text{Vec})$	$\text{Vec}_{\mathbb{Z}_2}$	Vec_-	$\text{Vec} \boxtimes \text{Vec}$	$\text{Vec} \boxtimes \text{Vec}_{\mathbb{Z}_2}$	$\text{Vec}_{\mathbb{Z}_2} \boxtimes \text{Vec}$	$\text{Vec}_{\mathbb{Z}_2} \boxtimes \text{Vec}_{\mathbb{Z}_2}$
(H, α)	$(\{1\}, 0)$	$(\{1, f\}, 0)$	$(\{1, m\}, 0)$	$(\{1, e, m, f\}, \alpha_0)$	$(\{1, e, m, f\}, \alpha_1)$	$(\{1, e\}, 0)$
ΣA for $A \in \text{Alg}_{E_1}^c(\Omega\mathcal{TC})$	1	$A_f = 1 \oplus f$	$A_m = 1 \oplus m$	$A_e = A_m \otimes A_e$	$A_4^{\text{op}} = A_e \otimes A_m$	$A_e = 1 \oplus e$

Table 1: Coordinates in $\mathcal{TC}^3$ and coordinate transformations

In the second row, two simple objects in $\text{Vec} \oplus \text{Vec}$ are precisely the two twist defects χ_+, χ_- constructed in [Bom10, KK12] and the right $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$ -module structure is defined by $\chi_{\pm} \otimes e = \chi_{\pm} \otimes m = \chi_{\mp}$; the fifth row is explained in Example 2.1.6 (see (2.1.6)); the sixth row is explained in Example 5.1.8. $\heartsuit$

Example 2.1.6. 2+1D finite gauge theories GT_G^3 for a finite gauge group G : Note that $\text{GT}_{\mathbb{Z}_2}^3 = \text{TC}^3$. The category $\Omega\mathcal{GT}_G^3$ of topological defects of codimension 2 (or anyons) and higher in GT_G^3 form the non-degenerate braided fusion 1-category $\mathfrak{Z}_1(\text{Rep}(G))$, which is the Drinfeld center of $\text{Rep}(G)$. The category $\mathcal{GT}_G^3$ of all topological defects form the monoidal 2-category:

$$\mathcal{GT}_G^3 \simeq \text{BMod}_{\text{Rep}(G)|\text{Rep}(G)}(2\text{Vec}).$$

A simple object in $\mathcal{GT}_G^3$ is an indecomposable $\text{Rep}(G)$ - $\text{Rep}(G)$ -bimodules in 2Vec . Since $\text{Rep}(G) \boxtimes \text{Rep}(G)^{\text{op}} \simeq \text{Rep}(G) \boxtimes \text{Rep}(G) \simeq \text{Rep}(G \times G)$, it is also an indecomposable left $\text{Rep}(G \times G)$ -module in 2Vec , which had been classified in [BO04, Ost03a]. More precisely, an indecomposable left $\text{Rep}(G \times G)$ -module in 2Vec is given by $\text{Rep}'(\tilde{H})$, where $\tilde{H}$ is the central extension of a subgroup H of $G \times G$ determined by a 2-cohomology class of $\alpha \in Z^2(H, \mathbb{C}^\times)$:

$$1 \rightarrow \mathbb{C}^\times \rightarrow \tilde{H} \rightarrow H \rightarrow 1$$

and $\text{Rep}'(\tilde{H})$ is the category of $\tilde{H}$ -representations such that $\mathbb{C}^\times$ acts as identity. The category $\text{Rep}'(\tilde{H})$ is also denoted by $\text{Rep}(H, \alpha)$, in which an object is a vector space V equipped with a map $\rho: H \rightarrow \text{GL}(V)$ such that $\rho(g)\rho(h) = \alpha(g, h)\rho(gh)$ for all $g, h \in H$. The $\text{Rep}(G \times G)$ -action on $\text{Rep}'(\tilde{H})$ or $\text{Rep}(H, \alpha)$ is defined by the usual vector space tensor product.

The fusion product in $\mathcal{GT}_G^3$ is defined by $\boxtimes_{\text{Rep}(G)}$, where the two-side $\text{Rep}(G)$ -actions on $\text{Rep}'(\tilde{H})$ are induced from the two natural monoidal functors $-\boxtimes \mathbb{1}_{\text{Rep}(G)}, \mathbb{1}_{\text{Rep}(G)} \boxtimes - : \text{Rep}(G) \rightarrow \text{Rep}(G) \boxtimes \text{Rep}(G)$. It is possible to work out the fusion rules, but we do not need them in this work. We give a complete classification of condensable E_1 -algebras in $\mathcal{GT}_G^3$ in later sections without using the fusion rules.

We give only one example, when $G = \mathbb{Z}_2$, by abusing the notation, we denote the elements in $\mathbb{Z}_2 \times \mathbb{Z}_2$ by $1, e, m, f$ with the obvious multiplication rules. The six simple $\text{Rep}(\mathbb{Z}_2)$ - $\text{Rep}(\mathbb{Z}_2)$ -bimodules given above are precisely $\text{Rep}(H, \alpha)$ for the following six pairs (H, α) :

$$(\{1\}, 0), (\{1, f\}, 0), (\{1, m\}, 0), (\{1, e, m, f\}, \alpha_0), (\{1, e, m, f\}, \alpha_1), (\{1, e\}, 0), \quad (2.1.6)$$

where $0 \in H^3(\mathbb{Z}_2, U(1)) = 0$ and $\alpha_0, \alpha_1 \in H^3(\mathbb{Z}_2 \times \mathbb{Z}_2, U(1)) \simeq \mathbb{Z}_2$, respectively (see Table 1 for a more complete dictionary). The fusion rules are given in the table (2.1.5). $\heartsuit$

Example 2.1.7. 2+1D Ising topological order Is^3 : We denote the 2-category of topological defects in Is^3 by $\mathcal{Is}$. In this case, the 1-category $\Omega\mathcal{Is}$ of anyons is the well known Ising modular tensor category. It consists of three simple anyons $1, \psi, \sigma$ with the commutative fusion rules $\psi \otimes \psi = 1, \psi \otimes \sigma \simeq \sigma, \sigma \otimes \sigma \simeq 1 \oplus \psi$. All its non-trivial braidings are given as follows:

$$\begin{aligned} \psi \otimes \psi &= 1 \xrightarrow{c_{\psi, \psi} = -1} 1 = \psi \otimes \psi; & \psi \otimes \sigma &= \sigma \xrightarrow{c_{\psi, \sigma} = e^{-\frac{\pi i}{2}}} \sigma = \sigma \otimes \psi; \\ \sigma \otimes \psi &= \sigma \xrightarrow{c_{\sigma, \psi} = e^{-\frac{\pi i}{2}}} \sigma = \psi \otimes \sigma; & \sigma \otimes \sigma &= 1 \oplus \psi \xrightarrow{c_{\sigma, \sigma} = e^{-\frac{\pi i}{8}} \oplus e^{\frac{3\pi i}{8}}} 1 \oplus \psi = \sigma \otimes \sigma. \end{aligned}$$

According to [FRS02], all condensable E_1 -algebras in $\Omega\mathcal{Is}$ are Morita equivalent to 1. This result implies that Is^3 has no non-trivial 1-codimensional topological defect. Therefore, we obtain that $\mathcal{Is} \simeq \text{B}\Omega\mathcal{Is}$ as monoidal 2-categories. $\heartsuit$

Remark 2.1.8. While we are preparing the paper, a few more examples of $\mathcal{C}$ in 2+1D have been worked out explicitly in [YWL25]. $\diamond$

The construction of 1-codimensional topological defects in [KS11a, KK12, FSV13, CR16, Kon14] makes the proposal in [Kit06] questionable because the characterization by the pair $(\Omega\mathcal{C}, c)$ does not contain any 1-codimensional topological defects except the trivial one.

$$\text{Does it mean that the characterization } (\Omega\mathcal{C}, c) \text{ is incomplete?} \quad (2.1.7)$$

The work [KW14] was devoted to answer this question. In order to proceed, one needs guidances from new physical principles.

2.1.2 Remote detectable principle and ‘intrinsic’ topological defects

Since these new physics principles apply to topological orders in all dimensions, we explain them in an arbitrary dimension. We start from recalling some basic notions that are used throughout this work.

Definition 2.1.9. Let C^{n+1} be an $n+1$ D topological order.

1. C^{n+1} is called *anomaly-free* if it can be realized by an n D gapped lattice model; it is called *anomalous* otherwise [KW14].
2. An anomaly-free C^{n+1} is called *non-chiral* if it admits a gapped boundary; and is called *chiral* otherwise (i.e., all its boundaries are gapless).

We denote the trivial $n+1$ D topological order by $\mathbf{1}^{n+1}$. □

Since an anomalous topological order C^{n+1} should be always realizable as a topological defect in a higher (but still finite) dimensional lattice model, by a dimensional reduction argument [KW14], C^{n+1} is always realizable as a gapped boundary of an $n+2$ D anomaly-free topological order. Moreover, it was shown that this $n+2$ D anomaly-free topological order is necessarily unique [KW14], and is called the *gravitational anomaly* or the *bulk* of C^{n+1} . We denoted it by $Z(C)^{n+2}$. Then the condition that C^{n+1} is anomaly-free can be mathematically expressed as

$$Z(C)^{n+2} = \mathbf{1}^{n+2}. \quad (2.1.8)$$

For an anomalous topological order C^{n+1} , we also have an interesting identity⁵ [KW14]:

$$Z(Z(C))^{n+3} = \mathbf{1}^{n+3}. \quad (2.1.9)$$

Example 2.1.10. When C^{n+1} is anomaly-free, then the trivial 1-codimensional topological defect in C^{n+1} can be viewed as an anomalous n D topological order, denoted by C^n , and is precisely the trivial domain wall between C^{n+1} and C^{n+1} . We apply the folding trick such that C^n becomes a gapped boundary of a double-layered system as illustrated below:

$$\xrightarrow{x^1} \quad \xrightarrow{C^{n+1}} \xrightarrow{C^n} \xrightarrow{C^{n+1}} \quad \xrightarrow{\text{folding trick}} \quad \begin{array}{c} \overline{C^{n+1}} \\ \searrow \\ C^n \\ \nearrow \\ C^{n+1} \end{array} \quad (2.1.10)$$

where $\overline{C^{n+1}}$ represents the $n+1$ D topological order obtained from C^{n+1} by flipping the orientation (i.e., $x^1 \mapsto -x^1$) and $\boxtimes$ represents the stacking of two $n+1$ D topological orders (without introducing interactions between them). We obtain the following identity:

$$Z(C^n)^{n+1} = C^{n+1} \boxtimes \overline{C^{n+1}}. \quad (2.1.11)$$

In this work, we assume that a folding is done in the x^1 -th spatial dimension, which is transversal to the domain wall and is depicted as the horizontal direction in pictures (see Convention 2.1.16). ♡

Since a topological order is a macroscopic notion, it is reasonable that it can be characterized by the complete set of macroscopic observables. Although the completeness is hard to define, it should contain topological defects⁶ in all codimensions. Therefore, the anomaly-free condition of a topological order might be translated to a condition on its macroscopic observables or topological defects.

Now we are ready to state a guiding principle: Remote Detectable Principle, which was proposed in [Lev13, KW14]. By a condensation descendant, we mean a defect obtained from an initial defect via condensation either as the condensed defect (of the same dimension) or as a gapped domain wall (of 1-dimension lower) between the initial defect and the condensed defect.

Remote Detectable Principle: In an anomaly-free topological order, a topological defect should be detectable up to condensation descendants by some topological defects via double braidings.

(2.1.12)

⁵This identity says that the bulk of a bulk is trivial. It is somewhat dual to the well-known statement in topology: the boundary of a boundary is empty, which leads to a homology theory. Therefore, we expect that the identity (2.1.9) should lead us to a non-trivial but yet-unknown cohomology theory.

⁶Instead of providing a microscopic definition of a topological defect, we recommend a working definition, i.e., an m D topological defect is an m D anomalous topological order.

This principle immediately provides an answer to the question (2.1.7). Consider an anomaly-free 2+1D topological order C^3 . All topological defects in C^3 form a fusion 2-category $\mathcal{C}$.

- (1) In this case, only defects of codimension 2 (i.e., anyons) can be braided. If a simple anyon x has trivial double braidings with all anyons, by the Remote Detectable Principle, x must be the trivial particle, i.e. $x = 1_{\mathbb{1}_{\mathcal{C}}}$. We can restate this result mathematically. For a braided fusion 1-category $\mathcal{B}$, we define its E_2 -center $\mathfrak{Z}_2(\mathcal{B})$ (also called Müger center) by the full subcategory consisting of objects with trivial double braidings with all objects in $\mathcal{B}$, i.e.,

$$\mathfrak{Z}_2(\mathcal{B}) := \{z \in \mathcal{B} | c_{z,b} \circ c_{b,z} = 1_{z \otimes b}, \forall b \in \mathcal{B}\}.$$

Using this language, the anomaly-free condition of C^3 can be translated into the following condition:

$$\mathfrak{Z}_2(\Omega\mathcal{C}) \simeq \text{Vec}. \quad (2.1.13)$$

It turns out that this equivalence (2.1.13), viewed as an anomaly-free condition, is equivalent to the non-degeneracy condition of the S -matrix [Müg03b] (see also [Lev13] for a physical discussion). The compatibility between this condition (2.1.13) and Definition 2.1.9 has not yet completely established⁷.

- (2) Note that 1-codimensional defects in C^3 cannot be braided at all. In this case, the Remote Detectable Principle implies that all 1-codimensional topological defects in $\mathcal{C}$ are necessarily the condensation descendants (or condensed defects) of anyons or the trivial 1-codimensional topological defect $1_{\mathcal{C}}$. Indeed, it was shown in [Kon14] that all 1-codimensional topological defects and those higher codimensional topological defects in C^3 that are not in $\Omega\mathcal{C}$ are the condensation descendants of those in $\Omega\mathcal{C}$. This result implies that all 1-codimensional defects can be obtained from the trivial 1-codimensional defect 1 or $B\Omega\mathcal{C}$ via condensations. The process of including all condensation descendants of $B\Omega\mathcal{C}$ is called the condensation completion (or Karoubi completion) of $B\Omega\mathcal{C}$, and is denoted by $\text{Kar}(B\Omega\mathcal{C})$ [GJF19], and can be mathematically defined [DR18, GJF19] (see Section 2.2). Following [DR18, GJF19], we set

$$\Sigma(-) := \text{Kar}(B(-)).$$

In this new notation, we have $\mathcal{C} = \Sigma\Omega\mathcal{C} := \text{Kar}(B\Omega\mathcal{C})$. Note that all objects in $\mathcal{C}$ are connected by 2-codimensional topological defects because they are all condensation descendants of the trivial 1-codimensional topological defect $1_{\mathcal{C}}$.

In summary, we have answered the question (2.1.7). Namely, the pair $(\Omega\mathcal{C}, c)$ indeed provides a complete mathematical characterization of the anomaly-free 2+1D topological order C^3 , and all the missing topological defects can be recovered from those in $\Omega\mathcal{C}$ via condensations, i.e., $\mathcal{C} = \Sigma\Omega\mathcal{C}$.

Remark 2.1.11. When C^3 is non-chiral, we have $\mathcal{C} \simeq \text{BMod}_{\mathcal{A}|\mathcal{A}}(2\text{Vec})$ for a fusion 1-category $\mathcal{A}$ (recall (2.1.3)) [KK12]. In this case, Remote Detectable Principle implies the following monoidal equivalence:

$$\Sigma\mathfrak{Z}_1(\mathcal{A}) \simeq \text{BMod}_{\mathcal{A}|\mathcal{A}}(2\text{Vec}). \quad (2.1.14)$$

In Section 3.3.2, we rederive this result as a consequence of a 1-codimensional defect condensation in 1^3 . $\diamond$

Before we discuss how Remote Detectable Principle applies to $n+1$ D anomaly-free topological orders for $n \geq 3$, we first review some useful notions and notations in higher category theory. An n -category $\mathcal{S}$ consists of objects, which are also called 0-morphisms, and k -morphisms between two $(k-1)$ -morphisms for $1 \leq k \leq n$. For a k -morphism f in $\mathcal{S}$, we denote the identity morphism on f by 1_f . We simplify the notation 1_{1_f} to 1_f^2 and define $1_f^k := 1_{1_f^{k-1}}$ inductively. We define $\Omega^0\mathcal{S} := \mathcal{S}$ and $\Omega_f\mathcal{S} := \text{hom}_{\mathcal{S}}(f, f)$. If an n -category $\mathcal{A}$ is monoidal, it is equipped with a canonical object the tensor unit $1_{\mathcal{A}}$. In this case, we abbreviate $\Omega_{1_{\mathcal{A}}}\mathcal{A}$ to $\Omega\mathcal{A}$. For example, the category $\Omega_f\mathcal{S} := \text{hom}_{\mathcal{S}}(f, f)$ is monoidal. Therefore, we can define:

$$\Omega_f^k\mathcal{S} := \Omega(\Omega_f^{k-1}\mathcal{S}) = \Omega^{k-1}(\Omega_f\mathcal{S}) = \text{hom}_{\mathcal{S}}(1_f^{k-1}, 1_f^{k-1}).$$

⁷It was established only recently that the condition (2.1.13) suffices a 2+1D lattice model realization [LW05, IO24]. As far as we know, it is not yet known if the converse statement is true.

In particular, $\Omega^{n-k}\mathcal{S}$ is a k -category for $k \leq n$. For a monoidal n -category $\mathcal{A}$, we define the looping of $\mathcal{A}$, denoted by $\Omega\mathcal{A}$, by $\text{hom}_{\mathcal{A}}(\mathbb{1}_{\mathcal{A}}, \mathbb{1}_{\mathcal{A}})$; and define the one-point delooping of $\mathcal{A}$, denoted by $B\mathcal{A}$, to be the $(n+1)$ -category consisting of a single object $\bullet$ such that $\Omega_{\bullet}B\mathcal{A} := \text{hom}_{B\mathcal{A}}(\bullet, \bullet) = \mathcal{A}$ as illustrated below.

$$B\mathcal{A} := \begin{array}{c} \mathcal{A} = \text{hom}_{B\mathcal{A}}(\bullet, \bullet) \\ \curvearrowright \\ \bullet \end{array} \quad (2.1.15)$$

Note that the monoidal structure of $\mathcal{A}$ is necessary for $B\mathcal{A}$ to be well-defined because $\text{hom}_{B\mathcal{A}}(\bullet, \bullet)$ has a monoidal structure defined by the composition of 1-morphisms.


Then we state a result which is a consequence of a stronger result: Theorem^{ph} 3.3.9. We simply take it for granted in this section (see Remark 2.1.13 for a discussion on the limitation of this result).

Theorem^{ph} 2.1.12 ([KW14, GJF19]). Given two k D topological defects A^k and B^k , the defect B^k can be obtained from A^k via a condensation of higher codimensional topological defects in A^k if and only if they can be connected by a $(k-1)$ D gapped domain wall⁸ $\square$

Now we consider an anomaly-free $n+1$ D topological order C^{n+1} for $n \geq 3$. It can have topological defects of all codimensions. All possible topological defects in C^{n+1} form a monoidal n -category $\tilde{\mathcal{C}}$, in which k -morphisms are $(k+1)$ -codimensional topological defects for $0 \leq k \leq n$; when $k = n$, they are instantons (i.e., 0D defects or $(n+1)$ -codimensional defects). We reserve the notation $\mathcal{C}$ for a subcategory of $\tilde{\mathcal{C}}$ described later. The monoidal structure on $\tilde{\mathcal{C}}$, i.e., a fusion product $\otimes$ in $\tilde{\mathcal{C}}$, is defined by the fusion of two 1-codimensional topological defects. The tensor unit is the trivial 1-codimensional topological defect, denoted by $\mathbb{1}$. The category $\tilde{\mathcal{C}}^{n+1}$ is very large. In particular, for $k \leq n$, by stacking a k D anomaly-free topological order to an $(n-k)$ -morphism in $\tilde{\mathcal{C}}$, we obtain a new $(n-k)$ -morphism in $\tilde{\mathcal{C}}$ [KW14, KLWZZ20a]. When $C^{n+1} = \mathbf{1}^{n+1}$, an anomaly-free k D topological order for $k \leq n$ should be viewed as $(n-k+1)$ -codimensional topological defects in $\mathbf{1}^{n+1}$, thus defines an $(n-k)$ -morphism in $\tilde{\mathcal{C}}$ [KLWZZ20a]. Since we have excluded gapless defects, a k D chiral topological order is not connected to the trivial k D topological order $\mathbf{1}^k$ in $C^{n+1} = \mathbf{1}^{n+1}$ by any $(n-k+1)$ -morphisms. Moreover, for $n \geq 3$, in general, two 1-codimensional topological defects are not connected by gapped domain walls. When $C^{n+1} = \mathbf{1}^{n+1}$, we denote the $\tilde{\mathcal{C}}$ in this case by $n\widehat{\text{Vec}}$.

Now we discuss how Remote Detectable Principle affects the categorical description of the anomaly-free $n+1$ D topological order C^{n+1} .

- (1) Topological defects of codimension 2 are braidable and should be detectable via double braidings. As a consequence, if a simple k -codimensional defect for $k \geq 2$ has trivial double braidings with all topological defects of codimension 2, it must be the trivial k -codimensional topological defect $\mathbf{1}_{\mathbb{1}}^k$ or its condensation descendants. Note that the double braiding should be insensitive to condensation as illustrated below.



During the adiabatic move of a double braiding, a condensation in b , which produces a new defect b' , should not create any physically detectable difference in the result of the double braiding.

In higher dimensional cases, the ‘trivial double braiding’ does not require the double braiding to be trivial on the nose, instead, it only requires the double braiding to be trivial up to higher isomorphisms. More explicitly, for $a, b \in \Omega\mathcal{C}$, the double braiding being trivial means that there is a higher isomorphism $c_{b,a} \circ c_{a,b} \rightarrow 1_{a \otimes b}$ satisfying natural physical conditions. When $n = 3$, these conditions have been explicitly spelled out in [KTZ20b, Definition 3.10] (preceded by a definition in the semistrict⁹ cases in [Gra98]) as the defining properties of the so-called sylleptic center, which is an E_2 -center¹⁰ and a 2-categorical generalization of the Mger center.

⁸The condition is also the defining axiom of the notion of Morita equivalence introduced in Definition 2.2.19.

⁹Physical applications demand us to consider non-semistrict monoidal 2-categories [KTZ20b].

¹⁰The notion of E_2 -center is a well-defined mathematical notion [Lur17] (see Definition 2.3.17). Actually, all types of centers, such as usual center of an algebra, Drinfeld center, Mger center and left/right or full centers in 2D conformal field theories, are all universally defined by the same universal property but in different (higher) categories [Lur17, Ost03a, Dav10a, KYZ21].

- (2) The category $\widetilde{n\text{Vec}}$ of topological defects in $\mathbf{1}^{n+1}$ is not compatible with Remote Detectable Principle.
- (a) Lower dimensional anomaly-free kD topological orders for $k \leq n$ can not be detected by double braidings because braidings among lower dimensional anomaly-free kD topological orders for $k \leq n$ are completely symmetric. Mathematically, it means that the category $\widetilde{n\text{Vec}}$ is symmetric monoidal. As a consequence, Remote Detectable Principle requires that we either modify the categorical description $\widetilde{n\text{Vec}}$ or demands that all lower dimensional anomaly-free kD topological orders are condensation descendants of the trivial mD topological orders for $k \leq m \leq n$.
 - (b) However, by Theorem^{ph} 2.1.12, nD chiral topological orders are not condensation descendants of the trivial nD topological order¹¹. As pointed out in [KW14], if the domain wall between two topological defects are gapless and not gappable, it is still possible to relate these two topological defects via a ‘generalized condensation’. However, such a ‘generalized condensation’ is necessarily beyond the condensation theory discussed in this work, which focus only on a finite type of condensations in the sense that neither infinite direct sums nor infinite dimensional vector spaces are allowed in the definition of a condensable algebra or in that of a (multi-)fusion n -category. It was suggested in [KW14] that we should exclude all nD chiral topological orders in the categorical description of $\mathbf{1}^{n+1}$ (or C^{n+1} in general). See Remark 2.1.13 for more discussion.

Therefore, by restricting to only the finite type of condensations studied in this work, Remote Detectable Principle demands us to replace $\widetilde{n\text{Vec}}$ by a subcategory $n\text{Vec}$, which consists of only trivial $(k-1)$ -codimensional topological defects $1_{\mathbb{1}}^k$ for $k = 0, 1, \dots, n$ and their condensation descendants. As a consequence, we obtain $n\text{Vec} = \text{Kar}(\mathbf{B}(n-1)\text{Vec}) = \Sigma(n-1)\text{Vec} = \Sigma^{n-1}\text{Vec}$. By setting $\Sigma\mathbb{C} := \text{Vec}$, we obtain $n\text{Vec} = \Sigma^n\mathbb{C}$. We define $n\text{Vec}$ more carefully in Section 2.2.1.

- (3) Similarly, in the categorical description of C^{n+1} , we would like to replace $\tilde{\mathcal{C}}$ by a subcategory $\mathcal{C}$, which contains only a set of ‘elementary’ (or ‘intrinsic’) topological defects [KW14]. In spite of many attempts to make the meaning of ‘intrinsic’ (or ‘elementary’) precise in [KW14, Section XI], it was still not very successful because a full fledged mathematical theory of condensations in higher categories is missing as it was explicitly pointed out in [KW14, Conjecture 10 & 11]. This missing theory was partially developed later in [GJF19] and is reviewed in Section 2.2. Since $\tilde{\mathcal{C}}$ is never used in this work, we assume that the category $\mathcal{C}$ of all ‘intrinsic’ topological defects can be properly defined, and, by abusing language, we abbreviate it to the category $\mathcal{C}$ of all topological defects. In later sections, we provide a mathematical approach towards $\mathcal{C}$ based on the notion of a separable n -category and that of a multi-fusion n -category. Now we derive a few important properties of $\mathcal{C}$ via Remote Detectable Principle.
- (a) $\mathcal{C}$ consists of the trivial 1-codimensional topological defect $\mathbb{1}$ (also denoted by $\mathbb{1}_{\mathcal{C}}$) and is necessarily closed under fusion. Therefore, $\mathcal{C}$ is a monoidal n -category. Since condensation descendants of ‘intrinsic’ topological defects are necessarily ‘intrinsic’, $\mathcal{C}$ is necessarily condensation complete (or Karoubi complete) (see Section 2.2 for more discussion).
 - (b) Since topological defects of codimension 2 and higher are braidable, Remote Detectable Principle demands that only topological defects that have the trivial double braidings with all topological defects of codimension 2 and higher are the trivial topological defects $1_{\mathbb{1}}^k$ for $k = 0, 1, \dots, n$ and their condensation descendants. Mathematically, the category of all defects of codimension 2 or higher that have trivial double braidings¹² to all defects is called E_2 -center of $\Omega\mathcal{C}$, denoted by $\mathfrak{Z}_2(\Omega\mathcal{C})$. Therefore, we obtain the following anomaly-free condition, for $n \geq 2$,

$$\mathfrak{Z}_2(\Omega\mathcal{C}) \simeq (n-1)\text{Vec}. \quad (2.1.16)$$

- (c) Since all the other defects not in $\Omega\mathcal{C}$, including all non-trivial 1-codimensional defects, cannot be braided, they are necessarily the condensation descendants of $\Omega\mathcal{C}$ or better $B\Omega\mathcal{C}$ [KW14, JF22, KLWZZ20b]. Namely, those defects not in $B\Omega\mathcal{C}$ can all be obtained from those in $B\Omega\mathcal{C}$ via condensations. In other words, we have

$$\mathcal{C} = \Sigma\Omega\mathcal{C} := \text{Kar}(B\Omega\mathcal{C}),$$

¹¹We show in Section 3.3.2 that a kD chiral topological order is a condensation descendant (as a gapped domain wall) of $\mathbf{1}^{k+1}$.

¹²In higher categories, a ‘trivial double braiding’ does not mean that the double braiding is trivial on the nose. Instead, it means that it is ‘trivial’ up to higher isomorphisms, which are the defining data of the ‘trivialness’. It means that $\mathfrak{Z}_2(\Omega\mathcal{C})$ is not a subcategory of $\Omega\mathcal{C}$ in general.

where $\text{Kar}(-)$ denotes the condensation completion or Karoubi completion. As a consequence, all objects in $\mathcal{C}$ are connected by 1-morphisms (see Remark 2.1.14).

Remark 2.1.13. Strictly speaking, replacing $\widetilde{n\text{Vec}}$ by its subcategory $n\text{Vec}$ of ‘intrinsic topological defects’ is more of a mathematical convenience than a demand of a physical principle because the mathematical theory of ‘generalized condensations’ (of infinite type) is simply not available. However, a complete condensation theory should include the process of approaching the critical point of the phase transition, where the system closes the energy gap and becomes gapless. This process is not a finite process in nature. Therefore, a complete condensation theory might demand us to include certain ‘generalized condensations’ of infinite type and ‘generalized condensation descendants’ in the theory. As an example of thinking along this line, $n\text{Vec}$ was used in [KLWZZ20a]. We provides only a glimpse of such a ‘generalized condensation’ in Section 6.3, where some relevant physical spaces that define the condensation become infinitely dimensional (see Remark 6.3.8). $\diamond$

Remark 2.1.14. Above discussion applies to the case when C^{n+1} is anomaly-free. If C^{n+1} is anomalous, in general, 1-codimensional topological defects in C^{n+1} are not necessarily the condensation descendants of those in $\Omega\mathcal{C}$. It means that $\mathcal{C}$ is disconnected in general when C^{n+1} is anomalous. $\diamond$

Remark 2.1.15. Although $\Omega\mathcal{C}$ has already encoded all the data in $\mathcal{C}$ when C^{n+1} is anomaly-free, as we show later, $\Omega\mathcal{C}$ does not directly provide a complete story of defect condensations. For certain problems in condensation theory, it is necessary to consider $\mathcal{C} = \Sigma\Omega\mathcal{C}$ instead (see Section 4). $\diamond$

Convention 2.1.16. In this work, a monoidal n -category $\mathcal{S}$ is always interpreted physically as the category of topological defects. Therefore, we always endows $\mathcal{S}$ with a local spacetime coordinate system

$$(x^1, x^2, \dots, x^n, x^{n+1} = t)$$

such that the fusion product in $\Omega^{i-1}\mathcal{S}$, denoted by $\otimes^i$, is the fusion of defects along the positive x^i -th direction (see Figure 1 and (2.1.10)). In pictures, we always depicted the fusion of the lower-codimensional topological defects in the horizontal direction. According to this convention, the n -category of all topological defects of $\overline{C}^{n+1}$ is given by $\mathcal{C}^{\text{op}}$, which is the same category as $\mathcal{C}$ but equipped with a new fusion product $(\otimes^1)^{\text{op}}$, which is defined by $a(\otimes^1)^{\text{op}}b := b \otimes^1 a$ for $a, b \in \mathcal{C}$ (see also Definition 2.3.5).

Definition 2.1.17. Given an n -category $\mathcal{S}$, we denote the category obtained from flipping all k -morphisms in $\mathcal{S}$ by $\mathcal{S}^{\text{op}k}$ and that obtained from flipping both k -morphisms and l -morphisms by $\mathcal{S}^{\text{op}k, \text{op}l}$. For convenience, we abbreviate $\mathcal{S}^{\text{op}1}$ to $\mathcal{S}^{\text{op}}$. $\square$

2.1.3 Boundary-bulk relation

There is an obvious question: can we use $\mathcal{C}$ (instead of $\Omega\mathcal{C}$) to give a mathematically equivalent characterization of the anomaly-free condition on C^{n+1} ? We show below how this question naturally leads us to a new guiding principle.

This question is already non-trivial when $n = 1$. In this case, only topological defects in C^2 are particles and instantons, and they form a 1-category $\mathcal{C}$. Since particles can be fused along the 1-dimensional space, $\mathcal{C}$ is a monoidal 1-category. By Remote Detectable Principle, if C^2 is anomaly-free, then $\mathcal{C}$ can only contain the trivial particles. In other words, $\mathcal{C}$ must be trivial, i.e. $\mathcal{C} \simeq \text{Vec}$. This is compatible with the fact that there is no non-trivial 1+1D anomaly-free topological order, proved first in a microscopic approach [CGW10]. In order to formulate the anomaly-free condition on $\mathcal{C}$, we need consider the case that C^2 is anomalous, i.e., $B^3 := Z(C)^3 \neq \mathbf{1}^3$. This case can be physically realized in lattice model [LW05, KK12]. In this case, $\mathcal{C}$ have non-trivial particles. The bulk particles move onto the boundary and become a boundary particle. This process defines a functor $f : \Omega\mathcal{B} \rightarrow \mathcal{C}$, which clearly preserves the fusion products, i.e., $f(a) \otimes f(b) \simeq f(a \otimes b)$ [KK12, FSV13]. Mathematically, such a functor is called a monoidal functor. Moreover, for $a \in \Omega\mathcal{B}$, $f(a)$ can be half-braided with $x \in \mathcal{C}$ as illustrated in Figure 2. This half-braiding structure endows the functor f with a structure called a central functor [Bez04] (see [FSV13] and a recent review [KZ22a] for its physical meanings). This notion is defined by the property that the functor f can be factorized as the composition of two functors $\Omega\mathcal{B} \rightarrow \mathfrak{Z}_1(\mathcal{C}) \rightarrow \mathcal{C}$, where $\mathfrak{Z}_1(\mathcal{C})$ is the E_1 -center (or Drinfeld center) of $\mathcal{C}$ and the second functor is the so-called forgetful functor. Then the condition that particles in $\mathcal{C}$ can be detected by




Figure 2: the half-braiding of $f(a)$ with x

bulk particles via the half braidings [Lev13] can be mathematically reformulated precisely as the following condition [KK12, Kon14]:

$$\Omega\mathcal{B} \simeq \mathfrak{Z}_1(\mathcal{C}), \quad (2.1.17)$$

which is also called boundary-bulk relation. In other words, the functor $f : \Omega\mathcal{B} \rightarrow \mathcal{C}$ can be identified with the forgetful functor $f : \mathfrak{Z}_1(\mathcal{C}) \rightarrow \mathcal{C}$. As a consequence, the anomaly-free condition on $\mathcal{C}^2$ can be translated to a condition on $\mathcal{C}$ expressed as follows:

$$\mathfrak{Z}_1(\mathcal{C}) \simeq \text{Vec}. \quad (2.1.18)$$

Note that the only fusion 1-category¹³ solution of above equation is $\mathcal{C} \simeq \text{Vec}$ [EGNO15]. This mathematical fact provides a categorical proof of the physical fact that there is no non-trivial 1+1D topological order [KW14, KWZ15].

The boundary-bulk relation was generalized to formulations other than (2.1.17) (see (2.1.23)) and to all dimensions and codimensions as the center functor (see Section 2.4) and to quantum phases far beyond topological orders [KWZ15, KWZ17, KZ22b]. In summary, boundary-bulk relation, in its simplest form, can be stated as follows.

Boundary-Bulk Relation: The bulk $Z(\mathcal{C})^{n+2}$ is the center of a boundary $\mathcal{C}^{n+1}$.

(2.1.19)

This result¹⁴ was proved via an important notion of a morphism between topological orders [KWZ15, KWZ17], a notion which can be generalized to gapped/gapless quantum liquids [KWZ17, KZ22b]. We should regard (2.1.19) as a formal guiding principle, which leads to concrete predictions when we replace $Z(\mathcal{C})^{n+2}$ and $\mathcal{C}^{n+1}$ by the concrete categories of topological defects. For example, we should still have (2.1.17) for $n > 1$, i.e.,

$$\Omega\mathcal{B} \simeq \mathfrak{Z}_1(\mathcal{C}). \quad (2.1.20)$$

It is important to note that this replacement is guaranteed by the fact that the notion of a monoidal functor is compatible with that of a morphism between two topological orders [KZ18a, Theorem 3.2.3] (see also [KWZ15]) and [KZ24, Remark 3.30].

Remark 2.1.18. One should use the boundary-bulk relation with caution. Notice that both $\Omega\mathcal{B}$ and $\mathcal{C}$ give completable descriptions of defects in $\mathcal{B}^{n+2}$ and $\mathcal{C}^{n+1}$, respectively. This is a necessary condition in the applications of the boundary-bulk relation. If we do not respect this rule and apply it naively, we could have, for example, $\Omega^2\mathcal{B} \simeq \mathfrak{Z}_2(\Omega\mathcal{C})$, which is not true in general (unless $\mathcal{C} = \Sigma\Omega\mathcal{C}$). For example, when $\mathcal{C} = 2\text{Vec}_G$ for a finite group G , $\Omega^2\mathcal{B} = \text{Rep}(G)$ [LKW18, KTZ20b] but $\mathfrak{Z}_2(\Omega\mathcal{C}) = \text{Vec}$. Therefore, although the boundary-bulk relation (2.1.19) is clearly compatible with the anomaly-free condition on $\Omega\mathcal{C}$ given in (2.1.16), we should not view (2.1.16) as a direct application of the boundary-bulk relation, but instead, a consequence of the condition (2.1.21). $\diamond$

The boundary-bulk relation (2.1.20) immediately implies the following anomaly-free condition on $\mathcal{C}$ generalizing (2.1.18) [KWZ15, JF22, KLWZZ20b] (see Remark 2.1.19):

$$\mathfrak{Z}_1(\mathcal{C}) \simeq n\text{Vec}, \quad (2.1.21)$$

¹³There are more multi-fusion 1-category solutions discussed later.

¹⁴This result has a precise meaning in the symmetric monoidal higher category TO^n of potentially anomalous n D topological orders. A morphism in TO^n is defined in [KWZ15, KWZ17] (see also Figure 3), its symmetric fusion product is defined by the stacking of two topological orders, and its tensor unit is 1^n . The notion of the center in TO^n is well defined by its universal property (see Definition 2.3.17).

where $n\text{Vec}$ is the category of ‘intrinsic’ topological defects in $\mathbf{1}^{n+1}$ (recall Section 2.1.2) and its precisely definition is given in Section 2.2. Since $\Omega\mathcal{C}$ encodes all the information of $\mathcal{C}$ up to condensation descendants, we expect that the two anomaly-free condition (2.1.16) and (2.1.21) are mathematically equivalent [KWZ15, KIWZZ20b]. Based on a corrected definition of a multi-fusion n -category, this conjecture was proved (at a physical level of rigor) by Johnson-Freyd in [JF22] (see also [KZ24] for a different approach).

Remark 2.1.19. We give a remark on the history and limitation of the proposal (2.1.21). Although, in retrospect, (2.1.21) seems a natural guess, there is no mathematical evidence to support this guess beyond the known result (2.1.18) in 2014-2015¹⁵ when the program was initiated in [KW14, KWZ15]. There are some natural questions that made the conjecture (2.1.21) questionable. The bulk $Z(\mathcal{C})^{n+2}$ is needed simply because it fix the braiding non-detectability of C^{n+1} by the half-braidings between the bulk defects and the boundary defects. Therefore, one could view (2.1.21) as a reformulation of Remote Detectable Principle. On the one hand, condensation descendants do not play an essential role in braidings. It is possible that condensation descendants are not needed for boundary-bulk relation to hold. For example, replacing $\mathcal{C}$ by $B\Omega\mathcal{C}$ in (2.1.21) could be another possibility. Moreover, ‘extrinsic’ topological defects should be excluded [KW14]. But how to distinguish ‘intrinsic’ and ‘extrinsic’ defects contributed additional complication to the theory [KW14, Section XI]. On the other hand, the mathematical notion of center is universal (or maximal in some sense). It suggests that it is reasonable to include all condensation descendants for boundary-bulk relation to hold. Since $\mathcal{C}$ is the maximal possible choice (for all ‘intrinsic’ topological defects), it is called a maximal BF^{pre} -category instead of a (multi-)fusion higher category in [KW14]. The later term was reserved for the one that is compatible with the notion of center (see [KWZ15, Definition 2.21]). The key question is whether the notion of center is compatible with condensation completion. In mathematics, the notion of a fusion 2-category was officially introduced by Douglas and Reutter in [DR18] in 2018 without knowing the physically required compatibility. It was based on a detailed study of the idempotent completion of 2-categories previously introduced by Carqueville and Runkel in [CR16], and was also supported by their success in the state-sum construction of 3+1D TQFT. However, it is unclear if this mathematical notion of a fusion 2-category is compatible with the notion of center as required by physics. The first evidence of its compatibility with center appeared in 2019 in [KTZ20b]. The original goal of this work was to find the correct mathematical definition of a (modular) fusion 2-category. By explicitly computing the E_1 -center $\mathfrak{Z}_1(2\text{Vec}_G^\omega)$ of 2Vec_G^ω , it was shown that $\mathfrak{Z}_1(2\text{Vec}_G^\omega)$ contains all 2-codimensional condensation descendants in 3+1D Dijkgraaf-Witten theories. Then it becomes clear that the conjecture (2.1.21) is reasonable¹⁶. In 2020, two independent works [JF22, KIWZZ20b] on this conjecture appeared within the same week on arXiv. Based on Gaiotto and Johnson-Freyd’s work [GJF19] on condensation completion in higher categories, in the beautiful work [JF22], Johnson-Freyd introduced the mathematical definition of a (multi-)fusion n -category and established the category theory of ‘intrinsic’ topological defects, including a proof of the equivalence between two anomaly-free conditions (2.1.16) and (2.1.21) (first conjectured in [KWZ15, Definition 2.21]). Somewhat orthogonal to [JF22], the work [KIWZZ20b] was devoted to developing a categorical theory of SPT/SET orders, which automatically includes that of topological orders as a special case, and the categorical theory of topological orders and (multi-)fusion n -categories were largely taken for granted. In [KIWZZ20b], the compatibility between center and condensation completion was proposed as a general principle called *Condensation Completion Principle*¹⁷, which applies to all gapped/gapless quantum liquids and demands a generalization to include certain gapless defects [KIWZZ20a]. A more systematic approach based on separable n -categories was developed later in [KZ22b] and is reviewed in Section 2.2.2.

Although a proper mathematical characterization of a topological order is required to be compatible with boundary-bulk relation, it does not mean that the possible characterization is unique. From a physical point of view, it seems also natural to extend $\mathcal{C}$ by including certain liquid-like gapless defects (see also [KIWZZ20a] and Section 6.3) or generalizing condensations to certain infinite types (see for example [Sto25]) such that the extended theory is still consistent with boundary-bulk relation. Therefore, the challenge told in above story will remain to be a guidance for us to find correct extended theories. $\diamond$

¹⁵Although an obvious thing to do is check if it is true for $n = 2$ when $\mathcal{C} = \text{BMod}_{\mathcal{A}|\mathcal{A}}(2\text{Vec})$, how to compute the Drinfeld center of $\mathcal{C}$ was not known until years later.

¹⁶Based on the new results in [DR18, KTZ20b], the first author of this paper explained to Johnson-Freyd in the spring of 2019 the conjectured equivalence between two anomaly-free conditions (2.1.16) and (2.1.21) in its correct form instead of the incorrect one in [KWZ15] (missing condensation completion), but did not correct this mistake before [JF22, KIWZZ20b] due to an ambitious but unfinished plan of rewriting the whole paper [KWZ15].

¹⁷There is no mathematical proof of the necessity of the condensation completion of the category of boundary defects. However, it is possible to argue from the Naturality Principle that things should appear naturally unless they are forbidden by a physical law.

Notice that the boundary-bulk relation (2.1.20) only involves $\Omega\mathcal{B}$. It is natural to ask if it is possible to have a formulation of boundary-bulk relation in terms of $\mathcal{B}$? Observe that if we include all 1-codimensional topological defects in $\mathcal{B}^{n+2}$, then fusing a non-trivial 1-codimensional topological defect to the boundary $\mathcal{C}^{n+1}$ changes the boundary condition. Since $\mathcal{B}^{n+2}$ is anomaly-free, an 1-codimensional topological defects in $\mathcal{B}^{n+2}$ is necessarily a condensation descendant of the trivial 1-codimensional topological defect $\mathbb{1}_{\mathcal{B}}$. It means that a 1-codimensional topological defect in $\mathcal{B}^{n+2}$ is always connected to $\mathbb{1}_{\mathcal{B}}$ by a n D gapped domain wall. Therefore, new boundary conditions obtained by fusing 1-codimensional topological defects in $\mathcal{B}^{n+2}$ with $\mathcal{C}^{n+1}$ are necessarily obtained from $\mathcal{C}^{n+1}$ via condensations and connected to $\mathcal{C}^{n+1}$ by a n D gapped domain wall¹⁸. We denote by $\Sigma\mathcal{C}$ the $(n+1)$ -category of all gapped boundary conditions of $\mathcal{B}^{n+2}$ that can be obtained from $\mathcal{C}^{n+1}$ via condensations. The notation $\Sigma\mathcal{C}$ represents the condensation completion of $\mathcal{B}\mathcal{C}$ and is explained in Section 2.2. Applying the boundary-bulk relation again, we should expect $\mathcal{B} \simeq \mathfrak{Z}_0(\Sigma\mathcal{C})$, where $\mathfrak{Z}_0(\Sigma\mathcal{C})$ is the E_0 -center of $\Sigma\mathcal{C}$ (explained in Section 2.3.5) and can be defined explicitly by $\text{Fun}(\Sigma\mathcal{C}, \Sigma\mathcal{C})$, i.e., the category of $\mathbb{C}$ -linear functors from $\Sigma\mathcal{C}$ to $\Sigma\mathcal{C}$. We summarize boundary-bulk relation below.

Theorem^{ph} 2.1.20 (Boundary-Bulk Relation). Let $\mathcal{B}^{n+2}$ be an anomaly-free topological order with a gapped boundary $\mathcal{C}^{n+1}$. We have

$$\Omega\mathcal{B} \simeq \mathfrak{Z}_1(\mathcal{C}), \quad (2.1.22)$$

$$\mathcal{B} \simeq \mathfrak{Z}_0(\Sigma\mathcal{C}) = \text{Fun}(\Sigma\mathcal{C}, \Sigma\mathcal{C}). \quad (2.1.23)$$

Moreover, these two formulations of boundary-bulk relation are necessarily equivalent. This equivalence immediately implies the following braided equivalence:

$$\mathfrak{Z}_1(\mathcal{C}) \simeq \Omega\mathfrak{Z}_0(\Sigma\mathcal{C}). \quad (2.1.24)$$

□

Remark 2.1.21. The result (2.1.24) is a mathematical fact proved in [Fra12, JF22, KZ22b]. ◇

In summary, we have given a description of the category of topological defects in an $n+1$ D anomaly-free (simple) topological order $\mathcal{C}^{n+1}$, and we show that the anomaly-free condition on $\mathcal{C}^{n+1}$ can be translated into a condition on $\Omega\mathcal{C}$ (2.1.16) or a condition on $\mathcal{C}$ (2.1.21). It is clear that, in order to move further, we need make the $\Omega\mathcal{C}, \mathcal{C}, n\text{Vec}$ in these conditions (2.1.16) and (2.1.21) mathematically precise. This is the subject of the next subsection.

2.2 Condensation completion

For a topological order $\mathcal{C}^{n+1}$, the process of completing $\mathcal{B}\Omega\mathcal{C}$ by its condensation descendants is called the *condensation completion* (or *Karoubi completion*) of $\mathcal{B}\Omega\mathcal{C}$ [CR16, DR18, GJF19]. More generally, for an arbitrary k -category $\mathcal{A}$, it is possible to define the Karoubi completion of $\mathcal{A}$, denoted by $\text{Kar}(\mathcal{A})$. This notion was first introduced in the $k = 2$ cases by Carqueville and Runkel in [CR16], and was later thoroughly developed by Douglas and Reutter in [DR18], and was later generalized to all $k \geq 2$ by Gaiotto and Johnson-Freyd in [GJF19]. In this subsection, we briefly review the basic ideas of condensation completion [GJF19] via a physical approach along the line of [KW14, KWZ15, KLWZZ20a], and introduce the notions of separable n -categories [KZ22b] and multi-fusion n -categories [JF22, KZ22b].

2.2.1 Category $n\text{Vec}$

In the previous subsection, $\mathcal{C}$ is assumed a priori and automatically condensation complete because it contains all (‘intrinsic’) topological defects in $\mathcal{C}^{n+1}$, including all condensation descendants, by definition. We want to take a closer look at topological defects in $\mathcal{C}^{n+1}$, especially when $\mathcal{C}^{n+1} = \mathbf{1}^{n+1}$.

Recall Theorem^{ph} 2.1.12. We show in Theorem^{ph} 3.3.9 that the condensation in Theorem^{ph} 2.1.12 can be defined by the gapped domain wall directly. Now we simply take it for granted. One immediate corollary is that a k D topological defect $\mathcal{B}^k$ can be obtained from another one $\mathcal{A}^k$ via a condensation of higher

¹⁸However, it is not true that all gapped boundaries of $\mathcal{B}^{n+2}$ can be connected to $\mathcal{C}^{n+1}$ by an n D gapped domain wall. For example, when $\mathcal{B}^{n+2} = \mathbf{1}^4$, a chiral anomaly-free 2+1D topological order $\mathcal{C}^3$ cannot connect to a non-chiral anomaly-free 2+1D topological order via a gapped domain wall, or equivalently, such a domain wall is necessarily gapless.

codimensional defects living in A^k if and only if A^k can be obtained from B^k via a condensation of higher codimensional defects living in B^k . Therefore, any two k -morphisms $f, g \in \mathcal{C}$, the $(n-k)$ -category $\text{hom}_{\mathcal{C}}(f, g)$ is a direct sum of connected full sub-categories. When $\mathcal{C}^{n+1}$ is anomaly-free, the monoidal n -category $\mathcal{C}$ is connected because every 1-codimensional defect is a condensation descendant of the trivial one $\mathbb{1}_{\mathcal{C}}$ and, therefore, is connected to $\mathbb{1}_{\mathcal{C}}$ via a gapped domain wall (recall discussion in Section 2.1.2). We used the following connected diagram to give an intuitive description of $\mathcal{C}$: for $x \in \mathcal{C}$,

$$\begin{array}{ccc} \Omega\mathcal{C} & & \text{hom}_{\mathcal{C}}(x, x) \\ \downarrow \text{hom}_{\mathcal{C}}(\mathbb{1}_{\mathcal{C}}, x) & \xrightarrow{\quad} & \downarrow \text{hom}_{\mathcal{C}}(x, \mathbb{1}_{\mathcal{C}}) \\ \mathbb{1}_{\mathcal{C}} & \xrightarrow{\quad} & x \end{array}$$

Since all objects in $\mathcal{C}$ are condensation descendants of $\mathbb{1}_{\mathcal{C}}$, $\mathcal{C}$ is the condensation completion (or Karoubi completion) of $B\Omega\mathcal{C}$, i.e. $\mathcal{C} = \text{Kar}(B\Omega\mathcal{C})$. The notion of a condensation completion can be mathematically defined for any higher category [GJF19]. Recall that the one-point delooping is defined for any monoidal higher category $\mathcal{A}$. We define the delooping $\Sigma\mathcal{A}$ of a condensation complete $\mathcal{A}$ by:

$$\Sigma\mathcal{A} := \text{Kar}(B\mathcal{A}) = \text{Kar}\left(\begin{array}{c} \mathcal{A} \\ \bullet \end{array}\right) = \begin{array}{ccc} \mathcal{A} & & B\text{Mod}_{\mathcal{A}|\mathcal{A}}(\mathcal{A}) \\ \downarrow \text{LMod}_{\mathcal{A}}(\mathcal{A}) & \xrightarrow{\quad} & \downarrow \text{RMod}_{\mathcal{A}}(\mathcal{A}) \\ \bullet & \xrightarrow{\quad} & A \end{array} \quad (2.2.1)$$

where an object A in $\Sigma\mathcal{A}$ is a ‘nice’ algebra (called condensation monad) in $\mathcal{A}$ as illustrated in the last ‘=’ in (2.2.1) [CR16, DR18, GJF19]. Then we have $\Sigma\Omega\mathcal{C} = \mathcal{C}$. We set $\Sigma\mathcal{C} := \text{Vec}$.

Theorem^{ph} 2.2.1 ([GJF19]). $n\text{Vec} = \Sigma^n\mathcal{C}$. Physically, $n\text{Vec}$ is the category of ‘intrinsic’ topological defects in 1^{n+1} , consisting of ‘roughly’ all lower dimensional non-chiral topological orders and gapped domain walls among them (see Remark 2.2.11 and Theorem^{ph} 2.2.12 for more precise statement). $\square$

Proof. This follows immediately from the fact that $n\text{Vec}$ contains only the trivial k -morphisms for $k \geq 0$ and their condensation descendants, and, as we have shown earlier, this fact is a consequence of Remote Detectable Principle and the limitation of our condensation theory (see Remark 2.1.13). $\square$

We can gain a better understanding of $n\text{Vec}$ by working out $n\text{Vec}$ for $n = 2, 3$ explicitly.

Example 2.2.2. $2\text{Vec} = \Sigma\text{Vec}$ is the 2-category of topological defects in 1^3 . In Example 2.1.4, we have shown that it is precisely that 2-category of finite semisimple 1-categories or finite direct sums of Vec . Now we work it out from the point of view of condensation completion. The trivial defect in 2Vec is precisely the trivial 1+1D topological order 1^2 , particles in which form the trivial fusion 1-category Vec . Condensation completion demands us to add all the condensation descendants of the trivial 1+1D topological order. We can proceed in two slightly different but equivalent ways.

$$\begin{array}{ccc} \text{Vec} & & \text{Fun}(\mathcal{X}, \mathcal{X}) \\ \downarrow \text{hom}_{2\text{Vec}}(\text{Vec}, \mathcal{X}) & \xrightarrow{\quad} & \downarrow \text{hom}_{2\text{Vec}}(\mathcal{X}, \text{Vec}) \\ 1^2 & \xrightarrow{\quad} & \mathcal{X} \end{array} \quad \begin{array}{ccc} \text{Vec} & & B\text{Mod}_{\mathcal{A}|\mathcal{A}}(\text{Vec}) \\ \downarrow B\text{Mod}_{\mathcal{C}|\mathcal{A}}(\text{Vec}) & \xrightarrow{\quad} & \downarrow B\text{Mod}_{\mathcal{A}|\mathcal{C}}(\text{Vec}) \\ 1^2 & \xrightarrow{\quad} & A \end{array} \quad (2.2.2)$$

1. The category of particles of each of these condensation descendants is necessarily a solution to the anomaly-free condition (2.1.18). All solutions are given by the indecomposable multi-fusion 1-category $\text{Fun}(\mathcal{X}, \mathcal{X})$ for a finite semisimple 1-category $\mathcal{X}$. We see immediately that these condensation descendants form the 2-category 2Vec because objects in 2Vec are precisely finite semisimple 1-category $\mathcal{X}$ and hom 1-category in 2Vec is precisely $\text{Fun}(\mathcal{X}, \mathcal{X})$ as illustrated in the first diagram in (2.2.2)¹⁹.

Note that $\mathcal{X}$ labels a 2D anomaly-free topological order, the category of topological defects in which is precisely the indecomposable multi-fusion 1-category $\text{Fun}(\mathcal{X}, \mathcal{X})$. When $\mathcal{X} \simeq \text{Vec}^{\oplus k}$ for $k > 1$, however,

¹⁹For readers who are familiar with Levin-Wen models with gapped boundaries [KK12], the 2-category 2Vec is just a special case of the 2-category of gapped boundary conditions of a fixed 2+1D Levin-Wen model based on a fusion 1-category $\mathcal{A}$. The objects in this 2-category are labeled by finite semisimple right $\mathcal{A}$ -modules and hom 1-categories are $\text{Fun}_{\text{cop}}(\mathcal{M}, \mathcal{N})$. This 2-category is precisely the 2-category of right $\mathcal{A}$ -modules in 2Vec , denoted by $\text{RMod}_{\mathcal{A}}(2\text{Vec})$. As a special case, $2\text{Vec} = \text{RMod}_{\text{Vec}}(2\text{Vec})$.

the associated 2D topological order does not satisfy the simpleness condition (recall Remark 2.1.1) and can be identified with the direct sum of k trivial phases: $(\mathbf{1}^2)^{\oplus k} = \mathbf{1}^2 \oplus \dots \oplus \mathbf{1}^2$. Such a topological order is called a composite topological order (see Definition 2.2.15) and naturally appear in dimensional reduction processes and condensations. For example, a 2+1D toric code model defined on a narrow strap with the smooth boundary condition on two sides can be viewed as a quasi-1+1D system that realizes the composite 1+1D anomaly-free topological order $\mathbf{1}^2 \oplus \mathbf{1}^2$ [KWZ15, KZ22a]. More discussion and examples of composite 1+1D topological orders can be found in [KZ22a].

2. According to [CR16, Kon14, DR18], a condensation in $\mathbf{1}^2$ is defined by a separable algebra A in Vec . The category of particles in the condensed 1+1D phase is given by the 1-category of A - A -bimodules in Vec , denoted by $\text{BMod}_{A|A}(\text{Vec})$ as illustrated in the second diagram in (2.2.2). Then the condensation completion of the trivial 1+1D topological order can be identified with the 2-category of separable algebras in Vec (as objects), bimodules (as 1-morphisms) and bimodule maps (as 2-morphisms). Since a separable algebra A in Vec is nothing but a finite direct sum of matrix algebras, $\text{BMod}_{A|A}(\text{Vec}) \simeq \text{Fun}(\mathcal{X}, \mathcal{X})$ for some $\mathcal{X} \in 2\text{Vec}$. Moreover, $\mathcal{X}$ can be identified with $\text{RMod}_A(\text{Vec})$, which is the category of right A -modules in Vec and is invariant up to the Morita equivalence of A . For this reason, one can relabel A by its complete Morita invariant $\mathcal{X} = \text{RMod}_A(\text{Vec})$. Therefore, this 2-category is precisely 2Vec .

In both cases, the finite semisimple 1-category $\mathcal{X}$ (i.e. $\mathcal{X} \in 2\text{Vec}$) pops out. Its physical meaning is the category of particles on the domain wall between $\mathbf{1}^2$ and one of its condensation descendants (say x), i.e., $\mathcal{X} = \text{hom}_{2\text{Vec}}(\text{Vec}, x)$. It is easy to see that $2\text{Vec} = \text{RMod}_{\text{Vec}}(2\text{Vec})$, which is a special case of a more general fact (see Theorem 2.2.32). $\heartsuit$

Example 2.2.3. $3\text{Vec} = \Sigma 2\text{Vec}$ is the monoidal 2-category of topological defects in $\mathbf{1}^4$. The tensor unit $\bullet = \mathbb{1}$ is the trivial 1-codimensional topological defect, which is precisely the trivial 2+1D topological order $\mathbf{1}^3$, i.e., $\bullet = \mathbb{1} = \mathbf{1}^3$. A simple object X in 3Vec is a condensation descendant of $\mathbf{1}^3$, and, in this case only, can be conveniently²⁰ interpreted as a 2+1D non-chiral topological orders X^3 (recall Definition 2.1.9). The 3-category 3Vec is connected as shown in the following diagram.

$$\begin{array}{ccccc}
 & 2\text{Vec} & & \mathcal{X} & & \mathcal{Y} \\
 & \downarrow & & \downarrow & & \downarrow \\
 \mathbf{1}^3 & \xrightarrow{\text{hom}_{3\text{Vec}}(\mathbf{1}^3, X^3)} & & X^3 & \xrightarrow{\text{hom}_{3\text{Vec}}(X^3, Y^3)} & Y^3 \\
 & \xleftarrow{\text{hom}_{3\text{Vec}}(X^3, \mathbf{1}^3)} & & \xleftarrow{\text{hom}_{3\text{Vec}}(Y^3, X^3)} & &
 \end{array}$$

The 2-category $\text{hom}_{3\text{Vec}}(\mathbf{1}^3, X^3)$ is the 2-category of gapped boundary conditions of X^3 . Its objects $a, b, c, \dots$ are the labels for gapped boundary conditions of X^3 , and the 1-category of morphisms $\text{hom}_{3\text{Vec}}(a, a)$ is the category of particles on the boundary a and is an indecomposable multi-fusion 1-category, and $\text{hom}_{3\text{Vec}}(a, b) = \text{hom}_{3\text{Vec}}(b, a)^{\text{op}}$ is the category of particles on the 0+1D gapped domain wall connecting a and b . By Theorem^{ph} 2.1.12, the 2-category $\text{hom}_{3\text{Vec}}(\mathbf{1}^3, X^3)$ is connected by non-zero 1-morphisms as illustrated below:

$$\text{hom}_{3\text{Vec}}(\mathbf{1}^3, X^3) = \begin{array}{ccc} & \text{hom}_{3\text{Vec}}(a, a) & \text{hom}_{3\text{Vec}}(b, b) \\ & \downarrow & \downarrow \\ a & \xrightarrow{\text{hom}_{3\text{Vec}}(a, b)} & b \\ & \xleftarrow{\text{hom}_{3\text{Vec}}(b, a)} & \end{array} = \Sigma \text{hom}_{3\text{Vec}}(a, a).$$

Since $\mathcal{X} := \text{hom}_{3\text{Vec}}(X^3, X^3)$ is the precisely the monoidal 2-category of topological defects in X^3 , by (2.1.23), we obtain

$$\mathcal{X} \simeq \mathfrak{Z}_0(\Sigma \text{hom}_{3\text{Vec}}(a, a)) = \text{Fun}(\text{hom}_{3\text{Vec}}(\mathbf{1}^3, X^3), \text{hom}_{3\text{Vec}}(\mathbf{1}^3, X^3)). \quad (2.2.3)$$

The $\Omega\mathcal{X}$ is the category of particles in the non-chiral 2+1D topological order X^3 . By the boundary-bulk relation [KK12, Kon14], it must be the Drinfeld center of the category of particles on each boundary $a \in \text{hom}_{3\text{Vec}}(\mathbf{1}^3, X^3)$, i.e.,

$$\Omega\mathcal{X} \simeq \mathfrak{Z}_1(\text{hom}_{3\text{Vec}}(a, a)),$$

²⁰This convenient interpretation of an object in 3Vec needs to be slightly modified for that of an object in $n\text{Vec}$ for $n > 3$ (see Remark 2.2.11, Theorem^{ph} 2.2.12 and Example 2.2.13).




Figure 3: a domain wall f defines a functor $f \boxtimes_X -$

which is a mathematical fact following from the equivalence (2.1.24). Note that $\text{hom}_{3\text{Vec}}(a, a)$ and $\text{hom}_{3\text{Vec}}(b, b)$ are Morita equivalent. By the Morita theory of multi-fusion 1-categories [Müg03a, Ost03a], we obtain the following equivalence of 2-categories:

$$\Sigma \text{hom}_{3\text{Vec}}(a, a) \simeq \text{RMod}_{\text{hom}_{3\text{Vec}}(a, a)}(2\text{Vec}).$$

As a mathematical result, it was first obtained by Douglas and Reutter in [DR18].

The direct sum of two non-chiral topological orders $X^3 \oplus Y^3$ also lives in 3Vec and we have

$$\text{hom}_{3\text{Vec}}(1^3, X^3 \oplus Y^3) \simeq \text{hom}_{3\text{Vec}}(1^3, X^3) \oplus \text{hom}_{3\text{Vec}}(1^3, Y^3).$$

Such a generalized ‘topological order’ is called a composite topological order (see Definition 2.2.15). It is not only a mathematical convenience but also a natural physical requirement when we study the dimensional reductions and the condensation theory of topological orders. In particular, we show in Section 3.3.2 that this generalized ‘topological order’ $X^3 \oplus Y^3$ can be obtained from 1^3 by condensing the algebra $A = \text{hom}_{3\text{Vec}}(x, x) \oplus \text{hom}_{3\text{Vec}}(y, y)$ in 2Vec for $x \in \text{hom}_{3\text{Vec}}(1^3, X^3)$ and $y \in \text{hom}_{3\text{Vec}}(1^3, Y^3)$. Therefore, from now on, we use X^3 and Y^3 to denote generic objects in 3Vec which can be a direct sum.

Note that a gapped domain wall $f \in \text{hom}_{3\text{Vec}}(X^3, Y^3)$ naturally defines a functor

$$\text{hom}_{3\text{Vec}}(1^3, X^3) \xrightarrow{f \boxtimes_X -} \text{hom}_{3\text{Vec}}(1^3, Y^3).$$

The physical meaning of this functor is illustrated in Figure 3. Conversely, we claim that any functor from $\text{hom}_{3\text{Vec}}(1^3, X^3)$ to $\text{hom}_{3\text{Vec}}(1^3, Y^3)$ should be realizable by such a domain wall, i.e.,

$$\text{hom}_{3\text{Vec}}(X^3, Y^3) = \text{Fun}(\text{hom}_{3\text{Vec}}(1^3, X^3), \text{hom}_{3\text{Vec}}(1^3, Y^3)). \quad (2.2.4)$$

There are two ways to prove this claim.

1. From a physical point of view, this claim follows from the so-called *Naturality Principle*, which says that if there is no physical law to forbid a natural set of mathematical structures (functors in our case) to occur physically, then all of them should occur physically. This principle was used in the bootstrap theory of rational CFT’s [MS89a] and that of anyon condensations [Kon14].
2. From a mathematical point of view, the precise mathematical definitions of the categories of boundary conditions of X^3 and Y^3 were known [JF22, KLWZZ20b, KZ22b] (or see [KZ22a] for a review). Then this claim is simply a mathematical fact, which follows immediately from [KZ18a, Theorem 3.2.3] (see also [KWZ15]).

These two results (2.2.3) and (2.2.4) simply say that the same category 3Vec can be equivalently defined by the following replacement:

$$\begin{aligned} X^3 &\mapsto \text{hom}_{3\text{Vec}}(1^3, X^3), \\ \text{hom}_{3\text{Vec}}(X^3, Y^3) &\mapsto \text{Fun}(\text{hom}_{3\text{Vec}}(1^3, X^3), \text{hom}_{3\text{Vec}}(1^3, Y^3)). \end{aligned}$$

This fact can be reformulated in more mathematical language. Let $\text{Cat}_2^{\mathbb{C}}$ be the 3-category of $\mathbb{C}$ -linear 2-categories [DR18]. This fact means that the following functor is fully faithful.

$$\text{hom}_{3\text{Vec}}(\bullet, -) : 3\text{Vec} \rightarrow \text{Cat}_2^{\mathbb{C}} \quad (2.2.5)$$

It turns out that the fully faithfulness of the functor (2.2.5) can be generalized to $(n+1)\text{Vec}$. The generalization provides a natural way to define the notion of a separable n -category. $\heartsuit$

2.2.2 Separable n -categories

Let $\text{Cat}_n^{\mathbb{C}}$ be the $(n+1)$ -category of $\mathbb{C}$ -linear n -categories. The following result is taken from [KZ22b, Cor. 3.2] based on the works [GJF19, JF22].

Theorem° 2.2.4. The functor $\text{hom}_{(n+1)\text{Vec}}(\bullet, -) : (n+1)\text{Vec} \rightarrow \text{Cat}_n^{\mathbb{C}}$ is fully faithful.


Proof. All the arguments in Example 2.2.3 automatically generalizes to $(n+1)\text{Vec}$ to provide a physical proof of this theorem except that the reference [KZ18a, Theorem 3.2.3] should be replaced by [KZ24, Theorem 3.26, Remark 3.27]. $\square$

Definition 2.2.5. A *separable n -category* is a $\mathbb{C}$ -linear n -category that lies in the essential image of $\text{hom}_{(n+1)\text{Vec}}(\bullet, -)$. A separable n -category is *condensation complete* by definition. If it is not a direct sum of two non-zero ones, it is called *indecomposable* (or *connected*). We call an indecomposable summand of a separable n -category $\mathcal{A}$ as an indecomposable (or connected) component of $\mathcal{A}$. $\square$


Example 2.2.6. Separable 0-categories are finite dimensional vector spaces over $\mathbb{C}$, i.e., objects in Vec . Separable 1-categories are finite semisimple $\mathbb{C}$ -linear 1-categories. $\heartsuit$

Example 2.2.7. We gives a few examples of separable 2-categories.

1. For a finite group G , 2Vec_G is the 2-category of G -graded separable 1-categories. A generic object is a direct sum $X = \oplus_{g \in G} X_g$, where the subscript of X_g is the grading and X_g is a separable 1-category, which is equivalent to $\text{Vec}^{\oplus n_g}$ for $n_g \in \mathbb{N}$. The only simple objects are Vec_g for $g \in G$. We illustrate the hom spaces associated to simple objects by the following diagram:




2. $2\text{Rep}(\mathbb{Z}_2) = \Sigma\text{Rep}(\mathbb{Z}_2)$ has two simple objects $\mathbb{1}$ and $\mathbb{1}_c$ [DR18]. We illustrate the hom spaces associated to simple objects by the following diagram:



Note that $2\text{Rep}(\mathbb{Z}_2)$ has a unique connected component.

3. The monoidal center $\mathfrak{Z}_1(2\text{Vec}_{\mathbb{Z}_2})$ of $2\text{Vec}_{\mathbb{Z}_2}$ as a separable 2-category is equivalent to a direct sum of two $2\text{Rep}(\mathbb{Z}_2)$ [KTZ20b] as illustrated below.



Since $\mathfrak{Z}_1(2\text{Vec}_{\mathbb{Z}_2})$ is the category of topological defects of codimension 2 and higher, each objects represents a string-like defect. In particular, $\mathbb{1}$ is the trivial string; $\mathbb{1}_c$ is the condensation descendant of the $\mathbb{1}$ -string [KW14, EN17, KTZ20a] and is sometimes called the Cheshire string [EN17]; m is the m -string or magnetic string [HZW05]; and m_c is the condensation descendant of the m -string. The only non-trivial simple object in $\text{hom}(\mathbb{1}, \mathbb{1}) = \text{Rep}(\mathbb{Z}_2)$ is the e -particle [HZW05].

4. The monoidal center $\mathfrak{Z}_1(2\text{Vec}_G)$ of 2Vec_G as a separable 2-category is equivalent to a direct sum [KTZ20b]:

$$\mathfrak{Z}_1(2\text{Vec}_G) \simeq \oplus_{[h] \in \text{Cl}} 2\text{Rep}(C_G(h)),$$

where Cl is the set of conjugacy classes of G and $C_G(h)$ is the centralizer of $h \in G$. $\heartsuit$

Example 2.2.8. We give a few useful separable n -categories

1. $n\text{Rep}(G) := \Sigma^{n-1}\text{Rep}(G)$. It can also be defined by the category of functors $\text{Fun}(\text{BG}, n\text{Vec})$ [KZ24], where BG is the 1-point delooping of G and is viewed as an n -category with only identity k -morphisms for $k > 1$.
2. $n\text{Vec}_G$ is the category of G -graded separable $(n-1)$ -categories, defined by the condensation complete $\mathbb{C}$ -linear monoidal n -category $\bigoplus_{g \in G} n\text{Vec}$ [KZ24]. Its number of connected components is $|G|$. It has a monoidal structure with the tensor product induced by the multiplication of G . For $\omega \in Z^{n+2}(G, \mathbb{C}^\times)$, there is a way to ‘twist’ the monoidal structure on $n\text{Vec}_G$ by ω to give a new monoidal n -category denoted by $n\text{Vec}_G^\omega$. It turns out that $\Sigma n\text{Vec}_G^\omega$ is a separable $(n+1)$ -category (see Definition 2.2.28).
3. The monoidal center $\mathfrak{Z}_1(n\text{Vec}_G^\omega)$ is a separable n -category²¹. Its physical meaning is the category of topological defects of codimension 2 and higher in $n+2$ D twisted G -gauge theory²² $\text{GT}_{(G,\omega)}^{n+2}$, i.e., $\Omega\mathcal{G}\mathcal{T}_{(G,\omega)}^{n+2} = \mathfrak{Z}_1(n\text{Vec}_G^\omega)$. If the twist ω is trivial, then $\text{GT}_{(G,\omega)}^{n+2}$ is called a finite gauge theory and is denoted by GT_G^{n+2} for simplicity. It was conjectured in [KTZ20b] that we have the following equivalence of separable n -categories:

$$\Omega\mathcal{G}\mathcal{T}_{(G,\omega)}^{n+2} = \mathfrak{Z}_1(n\text{Vec}_G^\omega) \simeq \bigoplus_{[h] \in \text{Cl}} n\text{Rep}(C_G(h), \tau_h(\omega)),$$

where Cl is the set of conjugacy classes of G , $C_G(h)$ is the centralizer of $h \in G$, and τ_h is the transgression map $C^{k+1}(G, \mathbb{C}^\times) \rightarrow C^k(C_G(h), \mathbb{C}^\times)$, and $n\text{Rep}(C_G(h), \tau_h(\omega))$ is the category of right separable module categories over the fusion n -category $n\text{Vec}_{C_G(h)}^{\tau_h(\omega)}$. When ω is trivial, we simply have

$$\Omega\mathcal{G}\mathcal{T}_G^{n+2} = \mathfrak{Z}_1(n\text{Vec}_G) \simeq \bigoplus_{[h] \in \text{Cl}} n\text{Rep}(C_G(h)).$$

When $n = 1$, this conjecture was rigorously proved in [Wil08], and the $n = 2$ cases were rigorously proved in [KTZ20b]. In this work, we simply takes above conjectured equivalences for granted for all n . Moreover, we expect $\mathfrak{Z}_1(n\text{Vec}_G^\omega)$ to be E_2 -fusion. It turns out that $\Sigma^2 \mathfrak{Z}_1(n\text{Vec}_G^\omega)$ is a separable $(n+2)$ -category (see Definition 2.3.7).

4. Let $z \in G$ be the fermion parity symmetry and lives in the center of G . The category $\text{Rep}(G, z)$ is a symmetric fusion 1-category, then $n\text{Rep}(G, z) := \Sigma^{n-1}\text{Rep}(G, z)$ is a separable n -category (actually a symmetric fusion n -category). $\heartsuit$

As a consequence, we can identify the category $(n+1)\text{Vec}$ as the $(n+1)$ -category of separable n -categories (as objects), $\mathbb{C}$ -linear n -functors (as 1-morphisms), $\mathbb{C}$ -linear n -natural transformations (as 2-morphisms), so on and so forth. When $n = 0$, a separable 0-category is a finite dimensional vector space over $\mathbb{C}$; when $n = 1$, a separable 1-category is a finite semisimple 1-categories; when $n = 2$, a separable 1-category is a finite semisimple 2-category [DR18].

By the folding trick, a gapped domain wall between two non-chiral topological order X^n and Y^n is automatically a gapped boundary of $X^n \boxtimes \overline{Y^n}$. For $x, y \in n\text{Vec}$, this means that $\text{hom}_{n\text{Vec}}(x, y) \simeq \text{hom}_{n\text{Vec}}(\mathbb{1}, y \otimes x^R)$, where x^R is the right dual of x . By Definition 2.2.5, $\text{hom}_{n\text{Vec}}(x, y)$ is a separable $(n-1)$ -category. It further implies that every hom space in a separable higher category is automatically separable.

Definition 2.2.9. A 0-morphism f in a separable 1-category $\mathcal{A}$ is called *simple* if $\text{hom}_{\mathcal{A}}(f, f) \simeq \mathbb{C}$. By induction, a k -morphism g in a separable n -category $\mathcal{B}$ is called *simple* if 1_g is simple in $\text{hom}_{\mathcal{B}}(g, g)$. $\square$

Remark 2.2.10. An indecomposable separable n -category is precisely a simple object in $(n+1)\text{Vec}$. $\diamond$

Remark 2.2.11. In Example 2.2.3, for the sake of convenience, we have used the non-chiral 2+1D topological order X^3 to label a simple object X in 3Vec . There is nothing wrong in this case. For $n\text{Vec}$ and $n > 3$, however, we need slightly modify the physical meaning of a simple object $X \in n\text{Vec}$. Note that X as a label for an object in $n\text{Vec}$ is not physically detectable. What is physically detectable is the category $\text{hom}_{n\text{Vec}}(\bullet, X)$, which can be identified with X via the fully faithful functor $\text{hom}_{(n+1)\text{Vec}}(\bullet, -) : (n+1)\text{Vec} \rightarrow \text{Cat}_n^{\mathbb{C}}$. It suggests

²¹For $\omega_1 \neq \omega_2$, in general, $\mathfrak{Z}_1(n\text{Vec}_G^{\omega_1})$ and $\mathfrak{Z}_1(n\text{Vec}_G^{\omega_2})$ are not equivalent even as separable n -categories. Example 3.4 in [KTZ20b] provided an example of this non-equivalence for $n = 2$.

²²Twisted finite gauge theories are also called Dijkgraaf-Witten theories.

that the physical meaning of the object X should come from the category $\text{hom}_{n\text{Vec}}(\bullet, X)$. Therefore, as an indecomposable separable $(n-1)$ -category in $n\text{Vec}$, X is precisely a connected component of the 2-category of all gapped boundary conditions of a non-chiral topological order. This slight modification is necessary because the higher category of gapped boundary conditions of a higher dimensional non-chiral topological order is in general disconnected, i.e., splitting into a direct sum of connected full sub-categories (see Example 2.2.13). We summarize this discussion into the following physical theorem. $\diamond$

Theorem^{ph} 2.2.12. Physically, a simple object in $(n+1)\text{Vec}$, or equivalently, an indecomposable separable n -category $\mathcal{S}$ is a connected component of the category of all gapped boundary conditions (up to invertible $n\text{D}$ topological orders²³) of a non-chiral $n+1\text{D}$ topological order X^{n+1} , i.e., a pair (X^{n+1}, x) where x labels the connected component. An object $s \in \mathcal{S}$ is a gapped boundary condition (up to invertible $n\text{D}$ topological orders) of X^{n+1} , and $\Omega_s \mathcal{S} := \text{hom}_{\mathcal{S}}(s, s)$ is the category of all topological defects on the boundary s of X^{n+1} . A 1-morphism in $\mathcal{S}$ is a domain wall between two gapped boundaries of X^{n+1} and a k -morphism for $1 \leq k \leq n$ is a k -codimensional topological defect on the boundaries of X^{n+1} . $\square$

Example 2.2.13. We explain Remark 2.2.11 and Theorem^{ph} 2.2.12 via the example 4Vec . Note that 4Vec is the precisely the category of all topological defects in $\mathbf{1}^4$.

1. The tensor unit $\bullet = \mathbf{1}$ is precisely the trivial 4D topological order $\mathbf{1}^4$.
2. Let $\mathcal{P}^3$ be a $2+1\text{D}$ chiral topological order and $\mathcal{P}$ the category of all topological defects in $\mathcal{P}^3$. As we show in Section 3.3.2, by condensing $A = \mathcal{P}$ in 3Vec , we obtain an indecomposable separable 3-category $X = \Sigma\mathcal{P}$ (as a gapped domain wall), which is an invertible object in 4Vec [JF22, KZ22b] and is precisely a connected component of the category of all gapped boundary conditions (up to invertible 3D topological orders) of $\mathbf{1}^4$. The simple objects in $X = \Sigma\mathcal{P}$ labels all $2+1\text{D}$ chiral topological orders that are connected to $\mathcal{P}^3$ via $1+1\text{D}$ gapped domain walls.
3. A simple 1-morphism $f \in \text{hom}_{4\text{Vec}}(\bullet, X) \simeq \text{hom}_{4\text{Vec}}(3\text{Vec}, \Sigma\mathcal{P})$ can be identified with a simple object in $\Sigma\mathcal{P}$. It is precisely the label of a simple gapped boundary of $\mathbf{1}^4$, or equivalently, a $2+1\text{D}$ chiral topological order (up to invertible ones), whose category of particles is Witt equivalent to the non-degenerate braided fusion 1-category $\Omega\mathcal{P}$ [DMNO13].
4. A generic simple object $X \in 4\text{Vec}$ is a condensation descendant of $\mathbf{1}^4$. As an indecomposable separable 3-category, X is precisely a connected component of the category of all gapped boundaries of a non-chiral 4D topological order, whose category of topological defects is given by $\mathfrak{Z}_0(X) = \text{Fun}(X, X)$.

Notice that, although 3Vec consists of only $2+1\text{D}$ non-chiral topological orders, 4Vec consists of all chiral or non-chiral $2+1\text{D}$ topological orders (up to chiral central charges) as simple 1-morphisms from $\bullet$ to invertible objects. When the invertible object is also the tensor unit $\bullet$, then such simple 1-morphisms are precisely those non-chiral $2+1\text{D}$ topological orders. It is worth reminding readers that the group of all invertible objects in 4Vec are precisely the Witt group of non-degenerate braided fusion 1-categories [JF22, KZ22b]. $\heartsuit$

2.2.3 Composite topological orders

The physical meanings of the morphisms in $(n+1)\text{Vec}$ provides a way to generalize the notion of a topological order to composite ones. This generalization is not necessary if we limit ourselves to those $n\text{D}$ topological orders that are simple in the sense that they have a unique ground state on an $(n-1)$ -sphere (or stable under local perturbations) [KWZ15]. However, the generalization is necessary for a complete theory as illustrated in the following picture.

$$\begin{array}{ccc}
 \begin{array}{c} \text{Z(A)}^{n+1} = \text{Z(B)}^{n+1} \\ \xrightarrow{\quad} \\ \text{A}^n \quad \text{M}^{n-1} \quad \text{B}^n \end{array} & \xrightarrow{\text{closing the fan}} & \begin{array}{c} \text{X}^n := \overline{\text{B}}^n \boxtimes_{\text{Z(A)}} \text{A}^n \\ \downarrow \\ \text{M}^{n-1} \end{array}
 \end{array} \tag{2.2.7}$$

The topological orders A^n and B^n in (2.2.7), i.e., sharing the same bulk and connected by a gapped domain wall M^{n-1} , are called Morita equivalent (see also Definition 2.2.19). Note that, even if we choose A^n and B^n

²³Invertible $n\text{D}$ topological orders are also gapped boundaries of $\mathbf{1}^{n+1}$. However, they are invisible in the theory of separable higher categories.

to be simple, the domain wall M^{n-1} , as an $(n-1)$ D topological order, is not simple in general. Moreover, by closing the fan, we obtain an n D topological order X^n having M^{n-1} as its gapped boundary, and X^n is not simple in general. For example, when $n = 2$ and $\mathcal{A} = \mathcal{M} = \mathcal{B} = \text{Vec}_G$ for a non-trivial finite group G , A^2 is a simple topological order but M^1 and X^2 are not simple. In this case, it turns out that M^1 and X^2 are non-simple or composite in two different ways. More precisely, X^2 is of type-I and M^1 is of type-II as defined below.

Definition 2.2.14. Roughly speaking, a composite topological order C^{n+1} is a ‘direct sum’ of simple ones. Moreover, C^{n+1} is called *type-I* if all its simple direct summands are Morita equivalent, and is called *type-II* otherwise. (see Definition 2.2.15 for a more complete definition). $\square$

In order to give a precise meaning of the ‘direct sum’, we need work within a precise category. It is possible to do it for topological orders (up to invertible ones) as we show in Definition 2.2.15, which was already implicitly used in [KZ22b]. However, physically oriented readers are recommended to understand the notion of a composite topological order at above intuitive level.

Definition 2.2.15. We can define a generalized notion of an n D (potentially anomalous) topological order (up to invertible ones) C^n by a 1-morphism $x : \bullet \rightarrow \mathcal{S}$ for an object $\mathcal{S} \in (n+1)\text{Vec}$. Since $\mathcal{S}$ can be viewed as a separable n -category, a 1-morphism $x : \bullet \rightarrow \mathcal{S}$ can be identified with an object $x \in \mathcal{S}$. The category of topological defects in the topological order x is defined by the category $\Omega_x \mathcal{S} := \text{hom}_{\mathcal{S}}(x, x)$.

1. If $x = 0$, then C^n is called the zero n D topological order denoted by 0. We identify all zero n D topological orders and identify the topological order $y : \bullet \rightarrow \mathcal{T}$ with the topological order $0 \oplus y : \bullet \rightarrow \mathcal{S} \oplus \mathcal{T}$, i.e., $y = 0 \oplus y$.
2. If x lives in a single connected component of $\mathcal{S}$, then C^n is called *type-I*. In this case, if x is simple, then the topological order C^n is called *simple*; if x is not simple, then C^n is called *type-I composite*. We denote the number of simple direct summands of x by $\text{gsd}(x)$ or $\text{gsd}(C^n)$ because it coincides with the ground state degeneracy of C^n defined on an $(n-1)$ -sphere.
3. If $\mathcal{S}$ is not connected and x does not live in a single connected component of $\mathcal{S}$, then the topological order C^n is called *type-II composite* or *type-II* for simplicity. We denote the number of simple direct summands of x by $\text{gsd}(x)$. We call the minimal number of non-zero type-I direct summands of x the multiplicity of C^n and denote it by $\text{Mult}(C^n)$.

C^n is called *composite* if it is either type-I composite or type-II composite. When C^n is a boundary/wall, the boundary/wall is called *simple*, type-I/type-II composite if C^n is simple, type-I/type-II composite, respectively. $\square$

Remark 2.2.16. We provide some examples of type-I and type-II topological orders later in Example 2.2.38. Strictly speaking the definition of a mathematical object is incomplete if we do not define the (iso)morphisms between this notion. We do not want to do it here. One can consult with the definition of the slice category $\bullet/(n+1)\text{Vec}$ in [KZ22b]. In this work, we simply identify all isomorphic objects in $(n+1)\text{Vec}$, i.e., identify all equivalent separable n -categories, for all physical discussion. $\diamond$

Remark 2.2.17. Both type-I and type-II topological orders are necessary for a complete condensation theory. When C^n is anomaly-free, the simpleness of C^n defined in Definition 2.2.15 coincides with the following simpleness condition: the ground state degeneracy of C^n defined on an $(n-1)$ -sphere is trivial. This simpleness condition is essentially a stability condition. As we shown in [KWZ15], a type-I topological order is not stable unless it is simple (see Remark 2.2.35), so is a type-II topological order, which is necessarily a gapped boundary of a type-I topological order. It means that if we restricted ourselves to only stable phases, we can restricted our theory to only simple topological orders. However, this restriction is not very natural if we work with a condensation complete theory. $\diamond$

Remark 2.2.18. The notion of a composite topological order is necessary for the study of the categories of topological orders and the condensation theory (see Remark 2.2.39). However, for most physicists who are only interested in the simple condensations of anomaly-free simple topological orders, it is beneficial to ignore anomalous and composite cases and skip relevant discussion at least in the first reading. $\diamond$

2.2.4 Morita equivalence between topological orders

Definition 2.2.19. Two $n+1$ D topological orders A^{n+1}, B^{n+1} are called *Morita equivalent*²⁴ if they share the same bulk (or gravitational anomaly), i.e., $Z(A)^{n+2} = Z(B)^{n+2}$, and are connected by a gapped domain wall M^n as illustrated below. We use $[A^{n+1}]$ to denote the Morita class of A^{n+1} .

$$\begin{array}{c} \boxed{Z(A)^{n+2} = Z(B)^{n+2}} \\ \hline A^{n+1} \quad \square \quad B^{n+1} \\ \quad \quad M^n \end{array} \quad (2.2.8)$$

(This terminology comes from the fact that the associated multi-fusion n -categories $\mathcal{A}$ and $\mathcal{B}$ are Morita equivalent. We also use $[\mathcal{A}]$ to denote the Morita class of $\mathcal{A}$. If A^{n+1} and B^{n+1} share the same bulk but are not Morita equivalent, then a domain wall M^n between them is necessarily gapless.) $\square$

Example 2.2.20. Any two 2+1D anomaly-free simple topological orders A^3 and B^3 share the same trivial 3+1D bulk. In this case, A^3 and B^3 are Morita equivalent if and only if two fusion 2-categories $\mathcal{A}$ and $\mathcal{B}$ are Morita equivalent, or equivalently, $\Omega\mathcal{A}$ and $\Omega\mathcal{B}$ are Witt equivalent [DMNO13]. If A^3 and B^3 are not Morita equivalent, then a domain wall between them is necessarily gapless. $\heartsuit$

Remark 2.2.21. By the structure theorem of multi-fusion categories [KZ22b] (reviewed in Proposition 2.2.34), a type-I topological order is always Morita equivalent to a simple one. Two type-II topological orders are Morita equivalent if and only if they share the same multiplicity and there is one-to-one pairing among their type-I direct summands such that each pair are Morita equivalent. $\diamond$

Remark 2.2.22. Two (potentially anomalous) topological orders A^{n+1} and B^{n+1} are Morita equivalent if and only if they share the same bulk and $A^{n+1} \boxtimes_{Z(A)^{n+2}} \overline{B^{n+1}} \simeq Z(M)^{n+1}$. There is another equivalent way to characterize the Morita equivalence as a centralizer $Z(A, Z(M))^{n+1} \simeq \overline{B^{n+1}}$ (explained in Example 2.3.24). $\diamond$

We recall a result from [KW14, Proposition 4] on Morita equivalence (recall Definition 2.2.19) between two anomaly-free topological orders.

Proposition^{ph} 2.2.23 ([KW14]). Two anomaly-free $n+1$ D topological order C^{n+1} and D^{n+1} are Morita equivalent if and only if there exist two n D topological orders P^n and Q^n such that

$$C^{n+1} \boxtimes Z(P)^{n+1} = D^{n+1} \boxtimes Z(Q)^{n+1}. \quad (2.2.9)$$

Proof. ($\Rightarrow$). If C^{n+1} and D^{n+1} are Morita equivalent, then they can be connected by a gapped anomaly-free domain wall M^n . By the folding trick, we have $Z(M)^{n+1} = C^{n+1} \boxtimes \overline{D^{n+1}}$. Then we obtain

$$C^{n+1} \boxtimes Z(D^n)^{n+1} = C^{n+1} \boxtimes \overline{D^{n+1}} \boxtimes D^{n+1} = Z(M)^{n+1} \boxtimes D^{n+1},$$

where D^n is the anomalous topological order defined by the trivial 1-codimensional defect in D^{n+1} .

($\Leftarrow$). It is quite obvious because C^{n+1} is clearly Morita equivalent to the left side of (2.2.9), and D^{n+1} is Morita equivalent to the right side of (2.2.9), as illustrated in the following picture.

$$\begin{array}{|c|c|c|} \hline C^{n+1} & C^{n+1} \boxtimes Z(P)^{n+1} & D^{n+1} \\ \hline M_1^n & D^{n+1} \boxtimes Z(Q)^{n+1} & M_2^n \\ \hline \end{array}$$

$\square$

We denote by $MC^n(C^{n+1})$ the set of Morita classes of gapped (necessarily type-I) boundaries of a non-chiral simple topological order C^{n+1} . For simplicity, we set $MC^n := MC^n(1^{n+1})$. Let M^n be a gapped boundary of C^{n+1} . We denote the associated equivalence class by $[M^n]$. Stacking topological orders gives a well-defined map:

$$\begin{aligned} MC^n(B^{n+1}) \times MC^n(C^{n+1}) &\xrightarrow{\boxtimes} MC^n(B^{n+1} \boxtimes C^{n+1}) \\ ([X^n], [M^n]) &\mapsto [X^n \boxtimes M^n]. \end{aligned}$$

²⁴This notion was called Witt equivalence in [KW14].

When $B^{n+1} = \mathbf{1}^{n+1}$, this map defines a MC^n -action on $MC^n(C^{n+1})$. When $B^{n+1} = C^{n+1} = \mathbf{1}^{n+1}$, this map defines an associative and commutative multiplication on MC^n .


Theorem^{ph} 2.2.24 ([KW14]). $MC^n = (MC^n, \boxtimes, [\mathbf{1}^n])$ is an abelian group and $[X^n]^{-1} = [\overline{X^n}]$ for any anomaly-free X^n . $\square$

Remark 2.2.25. When $n = 3$, MC^3 is precisely the usual Witt group [DMNO13]. Therefore, MC^n for $n > 3$ can be viewed as higher Witt groups. $\diamond$

The following result is essentially a slightly refined reformulation of (1) $\Leftrightarrow$ (3) part of Proposition 3.13 in [KZ24] (see Theorem 2.4.7 and 2.4.8 and also Theorem^{ph} 3.3.14), and first appeared (but formulated slightly differently but equivalently) in Theorem^{ph} 3.2.13 in the first version of this paper, and was independently obtained a few months later in the third version of [LYW24].

Theorem^{ph} 2.2.26. For a non-chiral simple topological order C^{n+1} , the MC^n -action on $MC^n(C^{n+1})$ is free and transitive, i.e., $MC^n(C^{n+1})$ is a MC^n -torsor. (A mathematical reformulation of this result can be found in Theorem 2.2.42.) $\square$

Proof. Let M^n and N^n be two gapped boundaries of C^{n+1} . We set $X^n := \overline{M^n} \boxtimes_{C^{n+1}} N^n$. Since $M^n \boxtimes \overline{M^n}$ is connected to C^n (i.e. the trivial 1-codimensional topological defect in C^{n+1}) by the gapped domain wall M^{n-1} (recall Example 2.1.10: M^{n-1} is the trivial 1-codimensional topological defect in M^n), then



$M^{n-1} \boxtimes_{C^n} N^{n-1}$ defines a gapped domain wall between $M^n \boxtimes X^n = M^n \boxtimes (\overline{M^n} \boxtimes_{C^{n+1}} N^n) = (M^n \boxtimes \overline{M^n}) \boxtimes_{C^{n+1}} N^n$ and $C^n \boxtimes_{C^{n+1}} N^n = N^n$.

Therefore, N^n and $M^n \boxtimes X^n$ are Morita equivalent. $\square$

Remark 2.2.27. Once we give a coordinate system $C^{n+1} = Z(B)^{n+1}$, then $MC^n(C^{n+1})$ is equipped with a natural coordinate given by the bijection $MC^n \rightarrow MC^n(C^{n+1})$ defined by $[X^n] \mapsto [X^n \boxtimes B^n]$. $\diamond$

Once we adapt the new notion of a topological order, certain things discussed in Section 2.1 need slight modifications. We postpone the discussion to the next subsection after we introduce the notion of a multi-fusion n -category.

2.2.5 Multi-fusion n -categories

Definition 2.2.28. A *multi-fusion n -category*, or an E_1 -*multi-fusion n -category*, is a condensation complete monoidal n -category $\mathcal{A}$ such that $\Sigma\mathcal{A}$ is a separable $(n+1)$ -category. It is called *indecomposable* if it is not a direct sum of two multi-fusion n -categories. It is called a *fusion n -category* if $\mathbb{1}_{\mathcal{A}}$ is simple. $\square$

Remark 2.2.29. The notion of a (multi-)fusion 2-category was first introduced by Douglas and Reutter in [DR18]. That of a (multi-)fusion n -category was first introduced by Johnson-Freyd in [JF22]. Here we follow an approach given in [KZ22b]. $\diamond$

Example 2.2.30. We provide a few examples of fusion 2-categories below.

1. The separable 2-category 2Vec_G introduced in Example 2.2.7 has an obvious structure of fusion 2-category with the tensor product defined by $\text{Vec}_g \otimes \text{Vec}_h = \text{Vec}_{gh}$ for $g, h \in G$ and the tensor unit defined by Vec_1 .
2. The separable 2-category $2\text{Rep}(\mathbb{Z}_2)$ is also equipped with a structure of fusion 2-category with the (non-trivial) fusion rule given by $\mathbb{1}_c \otimes \mathbb{1}_c = \mathbb{1}_c \oplus \mathbb{1}_c$.
3. The separable 2-category $\mathfrak{Z}_1(2\text{Rep}(\mathbb{Z}_2)) \simeq 2\text{Rep}(\mathbb{Z}_2) \oplus 2\text{Rep}(\mathbb{Z}_2)$ (recall (2.2.6)) is equipped with a structure of a fusion 2-category. The fusion rules are given below.

$$\mathbb{1}_c \otimes \mathbb{1}_c = \mathbb{1}_c \oplus \mathbb{1}_c, \quad m \otimes m = \mathbb{1}, \quad m_c = m \otimes \mathbb{1}_c = \mathbb{1}_c \otimes m, \quad m_c \otimes m_c \simeq m_c \oplus m_c.$$

Example 2.2.31. The separable n -categories $n\text{Rep}(G)$, $n\text{Vec}_G$, $\mathfrak{Z}_1(n\text{Vec}_G)$ introduced in Example 2.2.8 are examples of fusion n -categories. An indecomposable multi-fusion n -category $\mathcal{A}$ can be easily constructed from two separable n -categories $\mathcal{S}_1$ and $\mathcal{S}_2$ as

$$\mathcal{A} = \text{Fun}(\mathcal{S}_1 \oplus \mathcal{S}_2, \mathcal{S}_1 \oplus \mathcal{S}_2) = \begin{pmatrix} \text{Fun}(\mathcal{S}_1, \mathcal{S}_1) & \text{Fun}(\mathcal{S}_1, \mathcal{S}_2) \\ \text{Fun}(\mathcal{S}_2, \mathcal{S}_1) & \text{Fun}(\mathcal{S}_2, \mathcal{S}_2) \end{pmatrix}.$$

A separable n -category is a finite direct sum of indecomposable ones. An indecomposable separable n -category $\mathcal{S}$ is illustrated by the following connected diagram (connected by 1-morphisms).

$$\begin{array}{ccc} \text{hom}_{\mathcal{S}}(a, a) & & \text{hom}_{\mathcal{S}}(b, b) \\ \downarrow & \text{hom}_{\mathcal{S}}(a, b) & \downarrow \\ a & \xleftrightarrow{\quad} & b \\ \uparrow & \text{hom}_{\mathcal{S}}(b, a) & \uparrow \end{array} \quad \forall a, b \in \mathcal{S}. \quad (2.2.10)$$

Note that $\text{hom}_{\mathcal{S}}(a, a)$ and $\text{hom}_{\mathcal{S}}(b, b)$ are indecomposable multi-fusion $(n-1)$ -categories, which define physically the categories of topological defects on the gapped boundaries labeled by a and b , respectively. Since a and b are connected, two multi-fusion $(n-1)$ -categories are necessarily Morita equivalent. Moreover, $\text{hom}_{\mathcal{S}}(a, b)$ and $\text{hom}_{\mathcal{S}}(b, a)$ are precisely the invertible bimodules that define the Morita equivalence. Physically, $\text{hom}_{\mathcal{S}}(a, b)$ and $\text{hom}_{\mathcal{S}}(b, a)$ are the category of wall conditions of gapped domain walls between a and b . In particular, we have the following monoidal equivalence:

$$\text{hom}_{\mathcal{S}}(b, b) \simeq \text{Fun}_{\text{hom}_{\mathcal{S}}(a, a)^{\text{op}}}(\text{hom}_{\mathcal{S}}(a, b), \text{hom}_{\mathcal{S}}(a, b)). \quad (2.2.11)$$

The physical meaning of this fact is illustrated in the picture in (2.2.8), where two gapped boundaries A^{n+1} and B^{n+1} (corresponding to the labels a and b , respectively) of the non-chiral topological order $Z(A)^{n+2}$ are connected by a gapped domain wall M^n . In this case, by the functoriality of the boundary-bulk relation (see Theorem 2.4.5), we obtain $\mathcal{B}^{\text{op}} \simeq \text{Fun}_{\mathcal{A}}(\mathcal{M}, \mathcal{M})$, which is precisely the equivalence (2.2.11) when we set $\mathcal{A} = \text{hom}_{\mathcal{S}}(a, a)^{\text{op}}$ and $\mathcal{M} = \text{hom}_{\mathcal{S}}(a, b)$.

By replacing b by $b \oplus c$ in (2.2.11), we obtain a new equivalence:

$$\text{hom}_{\mathcal{S}}(b, c) \simeq \text{Fun}_{\text{hom}_{\mathcal{S}}(a, a)^{\text{op}}}(\text{hom}_{\mathcal{S}}(a, b), \text{hom}_{\mathcal{S}}(a, c)), \quad (2.2.12)$$

which simply says that, $\forall a \in \mathcal{S}$, we have

$$\mathcal{S} = \Sigma \text{hom}_{\mathcal{S}}(a, a) \simeq \text{RMod}_{\text{hom}_{\mathcal{S}}(a, a)}(n\text{Vec}). \quad (2.2.13)$$

This result provides a convenient and useful characterization of the delooping.

Theorem 2.2.32 ([GJF19, KZ22b]). For a multi-fusion n -category $\mathcal{A}$, we have $\Sigma\mathcal{A} \simeq \text{RMod}_{\mathcal{A}}((n+1)\text{Vec})$ and the equivalence is defined by the functor $\text{hom}_{\Sigma\mathcal{A}}(\bullet, -) : x \mapsto \text{hom}_{\Sigma\mathcal{A}}(\bullet, x)$.

Remark 2.2.33. If we identify $\Sigma\mathcal{A}$ with $\text{RMod}_{\mathcal{A}}((n+1)\text{Vec})$, i.e., $\Sigma\mathcal{A} = \text{RMod}_{\mathcal{A}}((n+1)\text{Vec})$, then we can also identify $\text{hom}_{\Sigma\mathcal{A}}(\bullet, -)$ with $\text{id}_{\Sigma\mathcal{A}}$. This naive fact turns out to be very useful in the study of topological orders because all the physical observables are encoded directly in the hom spaces in $\Sigma\mathcal{A}$ instead of its objects, which are, in priori, only abstract labels of topological defects and are not directly observable. Then the functor $\text{hom}_{\Sigma\mathcal{A}}(\bullet, -) = \text{id}_{\Sigma\mathcal{A}}$ allows us to translate abstract data of the objects in $\Sigma\mathcal{A}$ into physically observable data.

One physical application of this result is when $\mathcal{A}$ is a non-degenerate braided fusion $(n-1)$ -category. In this case, $\Sigma\mathcal{A} = \text{RMod}_{\mathcal{A}}(n\text{Vec})$ can be viewed as the category of topological defects of an anomaly-free topological order as illustrated below.

$$\begin{array}{c} \Sigma\mathcal{A} = \text{RMod}_{\mathcal{A}}(n\text{Vec}) \\ \begin{array}{ccc} & \text{hom}_{\Sigma\mathcal{A}}(x, x') & \\ \uparrow & & \uparrow \\ x & \text{hom}_{\Sigma\mathcal{A}}(x, x) & y \\ \uparrow & & \uparrow \\ x = \text{hom}_{\Sigma\mathcal{A}}(\mathbb{1}, x) & & \text{hom}_{\Sigma\mathcal{A}}(\mathbb{1}, y) = y \end{array} \\ \begin{array}{ccc} & \mathbb{1} & \\ \downarrow & & \downarrow \\ & \mathcal{A} & \end{array} \end{array} \quad (2.2.14)$$

In the coordinate system $\Sigma\mathcal{A} = \text{RMod}_{\mathcal{A}}(n\text{Vec})$, an object $x \in \Sigma\mathcal{A}$ (the label of a 1-codimensional topological defect) can be identified with $\text{hom}_{\Sigma\mathcal{A}}(\mathbb{1}, x)$, i.e., the defect junction between two 1-codimensional defects $\mathbb{1} \in \Sigma\mathcal{A}$ and x . In this coordinate system, the identity $x = \text{hom}_{\Sigma\mathcal{A}}(\mathbb{1}, x)$ amounts to the canonical equivalence: $\mathcal{M} \simeq \text{Fun}_{\mathcal{A}}(\mathcal{A}, \mathcal{M})$ for $\mathcal{M} \in \text{RMod}_{\mathcal{A}}(n\text{Vec})$ defined by the functor $m \mapsto - \otimes m$ for $m \in \mathcal{M}$. The horizontal tensor product $x \otimes^1 y$ in $\Sigma\mathcal{A}$ is precisely the horizontal fusion of two defect junctions $\text{hom}_{\Sigma\mathcal{A}}(\mathbb{1}, x) \boxtimes_{\mathcal{A}} \text{hom}_{\Sigma\mathcal{A}}(\mathbb{1}, y)$. Moreover, a morphism $f : x \rightarrow x'$ induces a morphism $f \circ - : \text{hom}_{\Sigma\mathcal{A}}(\mathbb{1}, x) \rightarrow \text{hom}_{\Sigma\mathcal{A}}(\mathbb{1}, x')$. Therefore, f is faithfully and physically realized by vertically fusing the defect junction $\text{hom}_{\Sigma\mathcal{A}}(\mathbb{1}, x')$ with $\text{hom}_{\Sigma\mathcal{A}}(\mathbb{1}, x)$. We use these facts in many physical discussions in later sections. $\diamond$

In a separable n -category, an object $x \in \mathcal{S}$ is simple if and only if its identity 1-morphism 1_x is a simple object in $\text{hom}_{\mathcal{S}}(x, x)$. In this case, $\text{hom}_{\mathcal{S}}(x, x)$ is a fusion $(n-1)$ -category. When x is not simple, it represents a composite defect or a composite boundary condition. In this case, $\text{hom}_{\mathcal{S}}(x, x)$ is a multi-fusion n -category. If all direct summands in x all live in the same connected component of $\mathcal{S}$, then $\text{hom}_{\mathcal{S}}(x, x)$ is an indecomposable multi-fusion n -category; otherwise, $\text{hom}_{\mathcal{S}}(x, x)$ is a direct sum of indecomposable multi-fusion n -category. As an illustrating example, when $x = a \oplus b$ and a, b are simple and connected, then we have the following decomposition (see also (2.2.10)):

$$\text{hom}_{\mathcal{S}}(x, x) = \begin{pmatrix} \text{hom}_{\mathcal{S}}(a, a) & \text{hom}_{\mathcal{S}}(a, b) \\ \text{hom}_{\mathcal{S}}(b, a) & \text{hom}_{\mathcal{S}}(b, b) \end{pmatrix}.$$

All indecomposable multi-fusion n -category is of such a matrix-type [KZ22b] as stated in details in the following proposition.

Proposition° 2.2.34 ([KZ22b]). If $\mathcal{A}$ is an indecomposable multi-fusion n -category, then we have

1. $\mathcal{A} \simeq \bigoplus_{i,j=1}^p \mathcal{A}_{i,j}$, where $\mathcal{A}_{i,j} := e_i \otimes \mathcal{A} \otimes e_j$ and e_i are simple summands of $\mathbb{1}_{\mathcal{A}}$, i.e. $\mathbb{1}_{\mathcal{A}} = \bigoplus_{i=1}^p e_i$;
2. $\Sigma\mathcal{A} = \Sigma\mathcal{A}_{ii}$ and $\mathcal{A}_{ii}$ is a fusion n -category;
3. the canonical $\mathcal{A}_{ii}$ - $\mathcal{A}_{kk}$ -bimodule map $\mathcal{A}_{ij} \boxtimes_{\mathcal{A}_{jj}} \mathcal{A}_{jk} \rightarrow \mathcal{A}_{ik}$ induced from the tensor product functor is an equivalence;
4. the $\mathcal{A}_{ii}$ - $\mathcal{A}_{jj}$ -bimodule $\mathcal{A}_{ij}$ is the inverse of the $\mathcal{A}_{jj}$ - $\mathcal{A}_{ii}$ -bimodule $\mathcal{A}_{ji}$, i.e., $[\mathcal{A}_{ii}] = [\mathcal{A}_{jj}]$.

Remark 2.2.35. The physical meaning of an indecomposable multi-fusion n -category $\mathcal{A}$ is the category of topological defects in an $n+1$ D type-I topological order A^{n+1} [KWZ15]. The physical meaning of the number of simple direct summands of the tensor unit $\mathbb{1}_{\mathcal{A}}$ (i.e., the number p) is the ground state degeneracy of the associated topological order A^{n+1} defined on an n -sphere S^n . In other words, if $\mathbb{1}_{\mathcal{A}}$ is not simple, then A^{n+1} is unstable, which means that by adding local perturbations the phase A^{n+1} can flow to a stable one associated to $\mathcal{A}_{ii}$ for an $i = 1, \dots, p$ [KWZ15]. Some 1D and 2D examples have been explicitly constructed in lattice models (see for example [KK12, KZ22a]). A decomposable multi-fusion n -category is the category of all topological defects in an $n+1$ D type-II topological order. $\diamond$

Remark 2.2.36. This definition of a multi-fusion 1-category is equivalent to the usual definition, i.e., a $\mathbb{C}$ -linear monoidal 1-category satisfying the following properties.

- (1) *Rigidity*: Every particle x has a right dual particle x^R , together with a creation morphism $b_x^R : \mathbb{1} \rightarrow x^R \otimes x$ and an annihilation morphism $d_x : x \otimes x^R \rightarrow \mathbb{1}$, satisfying some natural conditions; and a left dual particle x^L , together with a creation morphism $b_x^L : \mathbb{1} \rightarrow x \otimes x^L$ and an annihilation morphism $d_x^L : x^L \otimes x \rightarrow \mathbb{1}$, satisfying some natural conditions (see [EGNO15] for a review).
- (2) *Separability*: $\mathcal{A}$ is a separable 1-category, i.e., a finite direct sum of Vec , or equivalently, an object in 2Vec (recall Example 2.2.2).

Both properties follow immediately from the fact $\Sigma\mathcal{A} = \text{RMod}_{\mathcal{A}}(2\text{Vec})$ is a separable 2-category and the fully-dualizability of 3Vec . Conversely, a multi-fusion 1-category $\mathcal{A}$ in the usual sense satisfies the condition that $\Sigma\mathcal{A} \simeq \text{RMod}_{\mathcal{A}}(2\text{Vec})$ is a separable 2-category [DR18]. $\diamond$

Recall that the entire Section 2.1 is based on the assumption of the simpleness of the topological order (recall Remark 2.1.1). Once we adapt the generalized notion of a topological order in Definition 2.2.15, one result in Section 2.1 needs to be modified slightly. A topological order C^n (in the sense of Definition 2.2.15) is called *anomaly-free* if $Z(C)^{n+1} = \mathbf{1}^{n+1}$. We encounter a new phenomenon.

- (1) If C^n is anomaly-free, then C^n is necessarily type-I. If, in addition, C^n is type-I composite, then $\mathcal{C} \neq \Sigma\Omega\mathcal{C}$ and $\mathfrak{Z}_2(\Omega\mathcal{C})$ is non-trivial.
- (2) Both boundary-bulk relations (2.1.22) and (2.1.23) still hold for the generalized notion of a topological order. But we encounter new phenomena worth discussion. If C^n is type-II composite, then $B^{n+1} = Z(C)^{n+1}$ is type-I composite. More explicitly, assume $C^n = C_1^n \oplus C_2^n$ is a 1-morphism in $(n+1)\text{Vec}$:

$$\bullet \xrightarrow{C_1^n \oplus C_2^n} \mathcal{S}_1 \oplus \mathcal{S}_2$$

for two indecomposable separable n -categories $\mathcal{S}_1$ and $\mathcal{S}_2$. In this case, we have

$$\begin{aligned} B^{n+1} &= Z(C)^{n+1} = Z(C_1)^{n+1} \oplus Z(C_2)^{n+1}, \\ \mathcal{B} &\simeq \mathfrak{Z}_0(\mathcal{S}_1 \oplus \mathcal{S}_2) = \begin{pmatrix} \text{Fun}(\mathcal{S}_1, \mathcal{S}_1) & \text{Fun}(\mathcal{S}_1, \mathcal{S}_2) \\ \text{Fun}(\mathcal{S}_2, \mathcal{S}_1) & \text{Fun}(\mathcal{S}_2, \mathcal{S}_2) \end{pmatrix}, \\ \Omega\mathcal{B} &\simeq \Omega\mathfrak{Z}_0(\mathcal{S}_1) \oplus \Omega\mathfrak{Z}_0(\mathcal{S}_2) \simeq \Omega\mathfrak{Z}_0(\Sigma\mathcal{C}_1) \oplus \Omega\mathfrak{Z}_0(\Sigma\mathcal{C}_2) \simeq \mathfrak{Z}_1(\mathcal{C}_1) \oplus \mathfrak{Z}_1(\mathcal{C}_2). \end{aligned}$$

Note that the key point here is that even though C_1^n and C_2^n are not connected by higher morphisms in $(n+1)\text{Vec}$, their bulks $Z(C_1)^{n+1}$ and $Z(C_2)^{n+1}$ are both non-chiral, and, therefore, as 1-morphisms in $(n+2)\text{Vec}$, are connected by higher morphisms. In this case, we still have the bulk of a bulk is trivial, i.e., $Z(\mathcal{B})^{n+2} = \mathbf{1}^{n+2}$. In particular, we have $\mathfrak{Z}_1(\mathcal{B}) \simeq n\text{Vec}$ [JF22, KZ22b]. Therefore, B^{n+1} is type-I composite and anomaly-free.

Now we summarize and update results in Section 2.1 and 2.2 in the following physical theorem, which is essentially a reformulation (with a minor refinement) of [KWZ15, Definition 2.21].

Theorem^{ph} 2.2.37. Let C^{n+1} be an $n+1$ D topological order (in the sense of Definition 2.2.15). The category $\mathcal{C}$ of topological defects in C^{n+1} is a multi-fusion n -category.

1. If C^{n+1} is type-I, then $\mathcal{C}$ is an indecomposable multi-fusion n -category, and its bulk $Z(C)^{n+2}$ is simple. If C^{n+1} is simple, then $\mathcal{C}$ is a fusion n -category.
2. If C^{n+1} is anomaly-free (i.e., $Z(C)^{n+2} = \mathbf{1}^{n+2}$), then C^{n+1} is necessarily type-I, $\mathfrak{Z}_1(\mathcal{C}) \simeq n\text{Vec}$ and $\mathcal{C}$ is an indecomposable multi-fusion n -category.
 - (a) If, in addition, C^{n+1} is simple, then $\mathcal{C}$ is a simple separable n -category, i.e. $\mathcal{C} = \Sigma\Omega\mathcal{C}$, and we have $\mathfrak{Z}_2(\Omega\mathcal{C}) \simeq (n-1)\text{Vec}$.
 - (b) If, in addition, C^{n+1} is type-I composite, then $\mathcal{C} \neq \Sigma\Omega\mathcal{C}$, and $\Omega\mathcal{C}$ is non-trivial and splits into a direct sum. □
3. If C^{n+1} is type-II, then $B^{n+2} = Z(C)^{n+2}$ is type-I composite and anomaly-free, and we have

$$\text{gsd}(Z(C)^{n+2}) = \text{Mult}(C^{n+1}).$$

In this case, $\mathcal{C} = \bigoplus_{i=1}^{\text{Mult}(C^{n+1})} \mathcal{C}_i$ is a direct sum of indecomposable multi-fusion n -categories $\mathcal{C}_i$, and $\mathcal{B} = \mathfrak{Z}_0(\Sigma\mathcal{C})$ is an indecomposable multi-fusion $(n+1)$ -category, and we have $\Omega\mathcal{B} \simeq \mathfrak{Z}_1(\mathcal{C}) \simeq \bigoplus_i \mathfrak{Z}_1(\mathcal{C}_i)$.

In the rest of this work, by a (simple/composite) topological order we always means the one defined in Definition 2.2.15.

Example 2.2.38. We provide some examples of type-I and type-II 2+1D topological orders.

1. A 2+1D anomaly-free simple topological order is the one that satisfies the simpleness condition: the ground state degeneracy on S^1 is trivial. As we have reviewed in Section 2.1.1, an anomaly-free simple topological order C^3 can be described by a pair $(\mathcal{C}, c)$ or $(\Omega\mathcal{C}, c)$, where $\Omega\mathcal{C}$ is a non-degenerate braided fusion 1-category and $\mathcal{C} = \Sigma\Omega\mathcal{C}$ is a fusion 2-category. It can be identified with the simple object $\bullet \in \Sigma\mathcal{C}$ (or equivalently, a 1-morphism $\bullet \rightarrow \Sigma\mathcal{C}$ in 4Vec), where $\Sigma\mathcal{C}$ is a separable 3-category and an object in 4Vec .

- For each simple object $x \in \Sigma\mathcal{C}$, $\Omega_x^2 \Sigma\mathcal{C}$ is a non-degenerate braided fusion 1-category that is Witt equivalent to $\Omega\mathcal{C}$. Moreover, $\Sigma\mathcal{C}$ is an invertible object in 4Vec . For example, if $\mathcal{C}^3$ is non-chiral, then $\Sigma\mathcal{C} \simeq 3\text{Vec} = \bullet \in 4\text{Vec}$. All invertible objects arise in this way and form an abelian group which is isomorphic to the usual Witt group [JF22].
- Let $x, y \in \Sigma\mathcal{C}$ be two simple objects. We denote the 2+1D anomaly-free topological orders defined by x, y by X^3 and Y^3 , respectively. Both X^3 and Y^3 are Morita equivalent to $\mathcal{C}^3$. Now we consider the direct sum $x \oplus y : \bullet \rightarrow \Sigma\mathcal{C}$, as a 1-morphism in 4Vec , defines a type-I composite anomaly-free 2+1D topological order, denoted by $(\Sigma\mathcal{C}, x \oplus y)^3$, i.e., $Z(\Sigma\mathcal{C}, x \oplus y)^4 = \mathbf{1}^4$. Its category of topological defects is given by the following indecomposable multi-fusion 2-category:

$$\text{hom}_{\Sigma\mathcal{C}}(x \oplus y, x \oplus y) = \begin{pmatrix} \text{hom}_{\Sigma\mathcal{C}}(x, x) & \text{hom}_{\Sigma\mathcal{C}}(x, y) \\ \text{hom}_{\Sigma\mathcal{C}}(y, x) & \text{hom}_{\Sigma\mathcal{C}}(y, y) \end{pmatrix}.$$

Note that $\text{hom}_s(x, x) = \mathcal{X}$ and $\text{hom}_s(y, y) = \mathcal{Y}$ are the categories of defects in X^3 and Y^3 , respectively. We have the following monoidal equivalences:


$$\mathfrak{Z}_1(\text{hom}_{\Sigma\mathcal{C}}(x \oplus y, x \oplus y)) \simeq \mathfrak{Z}_1(\mathcal{X}) \simeq \mathfrak{Z}_1(\mathcal{Y}) \simeq 2\text{Vec},$$

where the first and second $\simeq$ follows from Example 2.3.28, justifying the boundary-bulk relation. However, the category of particles in $(x \oplus y)^3$ is given by $\Omega\mathcal{X} \oplus \Omega\mathcal{Y}$, whose Müger center is $\text{Vec} \oplus \text{Vec}$ and non-trivial.

The separable 2-category $\text{hom}_{\Sigma\mathcal{C}}(x, y) = \text{hom}_{\Sigma\mathcal{C}}(y, x)^{\text{op}}$ is precisely the category of gapped walls between X^3 and Y^3 , and we have a monoidal equivalence

$$\mathcal{X}^{\text{op}} \boxtimes \mathcal{Y} \simeq \mathfrak{Z}_0(\text{hom}_{\Sigma\mathcal{C}}(x, y)).$$

Moreover, for two gapped walls $p, q \in \text{hom}_{\Sigma\mathcal{C}}(x, y)$ between X^3 and Y^3 , $\mathcal{P} = \Omega_p \text{hom}_{\Sigma\mathcal{C}}(x, y)$ and $\mathcal{Q} = \Omega_q \text{hom}_{\Sigma\mathcal{C}}(x, y)$ are the categories of topological defects on the walls p and q , respectively, as illustrated in the following picture and diagram.



They are Morita equivalent indecomposable multi-fusion 1-categories both equipped with a braided equivalence

$$\Omega\mathcal{X} \boxtimes \Omega\mathcal{Y}^{\text{op}} \simeq \mathfrak{Z}_1(\mathcal{P}), \quad \Omega\mathcal{X} \boxtimes \Omega\mathcal{Y}^{\text{op}} \simeq \mathfrak{Z}_1(\mathcal{Q}),$$

respectively. The 0+1D domain wall $\text{hom}_{\Sigma\mathcal{C}}(p, q) = \text{hom}_{\Sigma\mathcal{C}}(q, p)^{\text{op}}$ is the separable 1-category uniquely fixed by $\mathcal{P}, \mathcal{Q}$ and $\Omega\mathcal{X}, \Omega\mathcal{Y}$ (see [AKZ17]).

- Let $\mathcal{S}$ and $\mathcal{T}$ be two simple separable 3-categories. Two objects $x \in \mathcal{S}$ and $y \in \mathcal{T}$ define two 2+1D type-I topological orders denoted by $(\mathcal{S}, x)^3$ and $(\mathcal{T}, y)^3$. Their category of topological defects are $\Omega_x \mathcal{S}$ and $\Omega_y \mathcal{T}$, respectively. Now we consider the direct sum of 1-morphisms $x \oplus y \in \mathcal{S} \oplus \mathcal{T}$. It defines a 2+1D type-II topological orders, denoted by $(\mathcal{S} \oplus \mathcal{T}, x \oplus y)^3$. In this case, we have $(\mathcal{S} \oplus \mathcal{T}, x \oplus y)^3 = (\mathcal{S}, x)^3 \oplus (\mathcal{T}, y)^3$. The category of topological defects in $(\mathcal{S} \oplus \mathcal{T}, x \oplus y)^3$ is given by the following (decomposable) multi-fusion 2-category

$$\text{hom}_{\mathcal{S} \oplus \mathcal{T}}(x \oplus y, x \oplus y) \simeq \text{hom}_{\mathcal{S}}(x, x) \oplus \text{hom}_{\mathcal{T}}(y, y) = \Omega_x \mathcal{S} \oplus \Omega_y \mathcal{T}.$$

The bulk of $(\mathcal{S} \oplus \mathcal{T}, x \oplus y)^3$ is non-trivial and type-I composite, i.e.,

$$\mathcal{B}^4 = Z(\mathcal{S} \oplus \mathcal{T}, x \oplus y)^4 = Z(\mathcal{S}, x)^4 \oplus Z(\mathcal{T}, y)^4 \neq \mathbf{1}^4.$$

We have the following (braided) monoidal equivalences:

$$\begin{aligned} \mathcal{B} &\simeq \mathfrak{Z}_0(\mathcal{S} \oplus \mathcal{T}) = \begin{pmatrix} \text{Fun}(\mathcal{S}, \mathcal{S}) & \text{Fun}(\mathcal{S}, \mathcal{T}) \\ \text{Fun}(\mathcal{T}, \mathcal{S}) & \text{Fun}(\mathcal{T}, \mathcal{T}) \end{pmatrix}, \\ \Omega\mathcal{B} &= \Omega \text{Fun}(\mathcal{S}, \mathcal{S}) \oplus \Omega \text{Fun}(\mathcal{T}, \mathcal{T}) \simeq \mathfrak{Z}_1(\Omega_x \mathcal{S}) \oplus \mathfrak{Z}_1(\Omega_y \mathcal{T}), \end{aligned}$$

where we have used (2.1.24), justifying the boundary-bulk relation. Note that B^4 is anomaly-free in that $\mathfrak{Z}_1(\mathcal{B}) \simeq 3\text{Vec}$. But we have

$$\mathfrak{Z}_2(\Omega\mathcal{B}) \simeq \mathfrak{Z}_2(\mathfrak{Z}_1(\Omega_x\mathcal{S})) \oplus \mathfrak{Z}_2(\mathfrak{Z}_1(\Omega_y\mathcal{T})) \simeq 2\text{Vec} \oplus 2\text{Vec} \neq 2\text{Vec}.$$

Note that, when $\mathcal{S} = \mathcal{T} = \Sigma\mathcal{C}$, the bulk

$$B^4 = Z(\Sigma\mathcal{C} \oplus \Sigma\mathcal{C}, x \oplus y)^4 = Z(\Sigma\mathcal{C}, x)^4 \oplus Z(\Sigma\mathcal{C}, y)^4 = \mathbf{1}^4 \oplus \mathbf{1}^4 \neq \mathbf{1}^4$$

is still non-trivial and type-I composite. In this case, one can see a fundamental difference between $(\Sigma\mathcal{C}, x \oplus y)^3$ and $(\Sigma\mathcal{C} \oplus \Sigma\mathcal{C}, x \oplus y)^3$, where the first one is type-I composite and the second one is type-II composite.

All the constructions in this example generalize tautologically to higher dimensions. $\heartsuit$

Remark 2.2.39. As we show in later sections, the condensations in a simple topological order C^{n+1} can produce type-I composite topological orders D^{n+1} as the condensed phase and type-II composite topological orders M^n as the gapped domain wall between C^{n+1} and D^{n+1} . Therefore, composite topological orders should be included in order to have a complete condensation theory (recall (2.2.7)). $\diamond$

In an anomaly-free simple $n+1$ D topological order C^{n+1} , the category $\mathcal{C}$ of all topological defects is a fusion n -category. It is the condensation completion of the braided fusion $(n-1)$ -category $\Omega\mathcal{C}$ of topological defect of codimension 2 and higher, i.e., $\mathcal{C} \simeq \Sigma\Omega\mathcal{C}$. When C^{n+1} is non-chiral, $\Omega\mathcal{C} \simeq \mathfrak{Z}_1(\mathcal{A})$ for a fusion $(n-1)$ -category $\mathcal{A}$. In this case, there is another way to represent the fusion n -category $\mathcal{C}$ (recall (2.1.14)), i.e.,

$$\mathcal{C} = \Sigma\Omega\mathcal{C} \xrightarrow[\simeq]{\phi} \text{BMod}_{\mathcal{A}|\mathcal{A}}(2\text{Vec}).$$

These two representations provide two coordinate systems of $\mathcal{C}$. The coordinate transformations between them are given by the monoidal equivalence ϕ and ϕ^{-1} defined below [KLWZZ20b, Theorem^{ph} 3.28]:

$$\phi : X \mapsto X \boxtimes_{\mathfrak{Z}_1(\mathcal{A})} \mathcal{A}; \quad \phi^{-1} : Y \mapsto Y \boxtimes_{\mathcal{A} \circ \text{p} \boxtimes \mathcal{A}} \mathcal{A}.$$

The coordinate transformation between different coordinate systems are very useful to concrete computations (see Table 1, Theorem^{ph} 3.3.33 and Example 5.2.23).

Remark 2.2.40. We have reviewed the theory of condensation completion physically in the spirits of [KW14, KWZ15, KLWZZ20b] instead of the more rigorous theory by Gaiotto and Johnson-Freyd [GJF19, JF22], which is not yet completely mathematically rigorous in the sense that no concrete model of weak n -categories is chosen and certain theory of colimits in higher categories was assumed. However, Gaiotto and Johnson-Freyd's theory has a lot of advantages, which allow us to proceed to answer many questions that are essentially orthogonal to higher coherence data at least at the physical level of rigor as shown in [JF22, KZ22b, KZ24]. $\diamond$

Remark 2.2.41. A few physical and computational works on condensation completion in 2+1D appeared recently [KZ22a, XLWWC23, YWL25]. They contain some explicit calculations of the condensation completion for a few 2+1D topological orders. $\diamond$

Recall that if two topological orders are Morita equivalent, then they share the same bulk. Conversely, if they share the same bulk, then it is not necessarily true that they are Morita equivalent because they might be connected by a gapless domain wall. Mathematically, it is possible to find an equivalent characterization of the sharing-the-same-bulk condition. The following result is a mathematical reformulation of Theorem^{ph} 2.2.26, and is a direct generalization of similar results for fusion 1-categories [ENO11] and fusion 2-categories [Déc25].

Theorem^o 2.2.42. Two indecomposable multi-fusion n -categories $\mathcal{A}$ and $\mathcal{B}$ share the same E_1 -center, i.e., $\mathfrak{Z}_1(\mathcal{A}) \simeq \mathfrak{Z}_1(\mathcal{B})$, if and only if $\Sigma\mathcal{A} \simeq \Sigma\mathcal{B} \boxtimes \mathcal{S}$ for an invertible separable n -category $\mathcal{S}$.

Proof. It follows from the same proof of '(1) $\Leftrightarrow$ (3)' part of [KZ24, Proposition* 3.13]. $\square$

Remark 2.2.43. This result is essentially a special case of [KZ24, Proposition* 3.13], which is reformulated in a slightly stronger way in Theorem^{ph} 2.4.8 but followed from the same proof. It first appeared (but formulated differently but equivalently) in Theorem^{ph} 3.2.13 in the first arXiv version of this paper, which is now Theorem^{ph} 3.3.14 (see also Remark 3.3.15). The current reformulation in Theorem 2.2.42 first appeared in the third arXiv version of [LYW24], which appeared a few months later than the first arXiv version of this paper. $\diamond$




Figure 4: An anomalous topological order C^2 and its bulk $Z(C)^3$

2.3 Higher algebras

In this subsection, we explain the main idea of higher algebras in $(n+1)\text{Vec}$ based on physical or geometric intuitions.

2.3.1 Monoidal 1-categories as E_1 -algebras

A potentially anomalous 1+1D topological order C^2 can be realized as a gapped boundary of a 2+1D topological order $Z(C)^3$ as depicted in Figure 4. It can have particle-like topological defects or particles for simplicity. These particles can be fused along the unique spatial dimension. Therefore, these particles, together with their fusion, form a fusion 1-category $\mathcal{C}$ [KK12], in which 1-morphisms are instantons, i.e. 0D defect in spacetime. This result is well known. It is, however, only a part of a more complete story. We would like to sketch some key points in this more complete story (see also [KZ22a, Remark 3.4.19 & 3.4.66]) in order to introduce some important notions.

Consider a lattice model realization of the 2+1D topological order $Z(C)^3$ with the gapped boundary C^2 . Suppose two anyons $a, b \in \mathcal{C}$ are realized in the model at two different sites $r, s \in \mathbb{R}$, respectively, as depicted Figure 4. We chose the boundary line to be the real line. Recall that a particle is invariant under the action of local operators. Therefore, it represents a subspace of the total Hilbert space that are invariant under the net of local operators, and is more precisely defined as a superselection sector [HK64, Haa96]. The physical configuration depicted in Figure 4 already defines a fusion product $a \otimes_{(r,s)} b$, which can be viewed a subspace of the total Hilbert space. If we realize a, b in the same model but at different sites r', s' , then a different fusion product $a \otimes_{(r',s')} b$ is realized as a different subspace of a potentially different total Hilbert space²⁵. It means that there are infinitely many fusion products parameterized by the elements in the configuration space $\{(r, s) \in \mathbb{R}^2 \mid r \neq s\}$. An adiabatic move of (a, b) from (r, s) to (r', s') along a path γ in the configuration space defines a linear isomorphism:

$$T_{a,b}^\gamma : a \otimes_{(r,s)} b \xrightarrow{\sim} a \otimes_{(r',s')} b.$$

In mathematical language, $T^\gamma := \{T_{a,b}^\gamma\}_{a,b \in \mathcal{C}} : \otimes_{(r,s)} \rightarrow \otimes_{(r',s')}$ defines a natural isomorphism. One of the defining properties of a topological order is the following principle:

Adiabatic Principle: if two paths γ_1 and γ_2 in the configuration space $\{(r, s) \in \mathbb{R}^2 \mid r \neq s\}$ are homotopy equivalent, then $T^{\gamma_1} = T^{\gamma_2}$.

(2.3.1)

This property allows us to reduce the infinitely many fusion products $\otimes_{(r,s)}$ to two homotopically inequivalent fusion products $\otimes := \otimes_{(-1,1)}$ and $\otimes^{\text{op}} := \otimes_{(1,-1)}$. It is clear that we have $a \otimes^{\text{op}} b = b \otimes a$.

We have sketched only some key ideas of a complete story, which is essentially the same as the mathematical proof of the fact that an E_1 -algebra in the symmetric monoidal 2-category Cat_1 is precisely a monoidal 1-category [SW03, Lur17, Fre17]. We have used the notation Cat_1 to denote the category of 1-categories (as objects), functors (as 1-morphisms) and natural transformations (as 2-morphisms). Cat_1 is symmetric monoidal with the tensor product given by the Cartesian product $\times$ and the tensor unit given by the category with a single object $\bullet$ and a single 1-morphism $1_\bullet$. The E_1 -algebra (or an algebra over the E_1 -operad)

²⁵It depends on how one realize the defects. If we allow to introduce auxiliary space localized around a site r for the realization of a particle at r , then total Hilbert spaces are different for anyon at different sites but isomorphic via adiabatic moves.




Figure 5: three paths in configuration space

structure on $\mathcal{A}$ consists of the infinitely many fusion products $\otimes_{(r,s)}$ that are defined in 1-dimensional space, satisfying some natural conditions. It is different from that of a monoidal structure by definition. In Cat_1 , they are equivalent due to the fact that there is no higher morphisms in Cat_1 to catch the information of higher homotopy data in the E_1 -operad [SW03, Lur17, Fre17]. In this work, we do not distinguish the 2-category of E_1 -algebras, E_1 -algebra homomorphisms and E_1 -algebra 2-homomorphisms with that of monoidal categories, monoidal functors and monoidal natural transformations, and denote both categories by $\text{Alg}_{E_1}(\text{Cat}_1)$. Then the fact that $\mathcal{A}$ is a monoidal 1-category can be easily summarized by a compact notation $\mathcal{A} \in \text{Alg}_{E_1}(\text{Cat}_1)$. From now on, we use the physical intuition of infinitely many fusion products freely whenever we talk about a 1+1D topological order A^2 or a monoidal 1-category $\mathcal{A}$.

For $\mathcal{A} \in \text{Alg}_{E_1}(\text{Cat}_1)$, we can define the notion of a left/right $\mathcal{A}$ -module in Cat_1 , that of a module functor and that of a module natural transformation [Bén65, Ost03a] (see [EGNO15] for a review). We denote the 2-category of left, right, bi- $\mathcal{A}$ -modules in Cat_1 by $\text{LMod}_{\mathcal{A}}(\text{Cat}_1)$, $\text{RMod}_{\mathcal{A}}(\text{Cat}_1)$, $\text{BMod}_{\mathcal{A}|\mathcal{A}}(\text{Cat}_1)$, respectively.

2.3.2 Braided monoidal 1-categories as E_2 -algebras

For 2+1D topological order C^3 , the category $\mathcal{C}$ of topological defects in C^3 is monoidal with a tensor unit $1_{\mathcal{C}} = 1$. Its anyons form a braided monoidal category $\Omega\mathcal{C}$, which is usually defined by a tensor product $\otimes$, a tensor unit $1_{\mathbb{1}}$, the associators $\alpha_{a,b,c} : a \otimes (b \otimes c) \rightarrow (a \otimes b) \otimes c$, the left unitors $l_a : 1_{\mathbb{1}} \otimes a \rightarrow a$, the right unitors $r_a : a \otimes 1_{\mathbb{1}} \rightarrow a$ and the braidings isomorphisms $c_{a,b} : a \otimes b \rightarrow b \otimes a$ satisfying some natural conditions. In this subsection and the next, we reinterpret this structure in two seemingly different but equivalent ways that are ready to be generalized to higher dimensions.

In the first way, we go back to the physical intuition provided by the lattice model realizations of a 2+1D topological order. We mainly repeat the discussion in [KZ22a, Remark 3.4.19 & 3.4.66].

Suppose we have a lattice model realization of the 2+1D topological order C^3 , and suppose two anyons $a, b \in \Omega\mathcal{C}$ are realized in the model at two different sites $\xi, \eta \in \mathbb{R}^2 = \mathbb{C}$, respectively, as depicted in Figure 5. Recall that an anyon is invariant under the action of local operators. Therefore, it represents a subspace of the total Hilbert space that are invariant under the net of local operators, which can be mathematically defined as superselection sectors [HK64, Haa96]. This physical configuration already defines a fusion product $a \otimes_{(\xi,\eta)} b$, which can be viewed a subspace of the total Hilbert space. If we realize $a, b \in \Omega\mathcal{C}$ in the same model but at different sites ξ', η' , then a different fusion product $a \otimes_{(\xi',\eta')} b$ is realized as a different subspace of a potentially different total Hilbert space. Similar to 1+1D topological orders, it means that there are infinitely many fusion products parameterized by the elements in the configuration space $\{(\xi, \eta) \in \mathbb{C}^2 \mid \xi \neq \eta\}$. An adiabatic move of (a, b) from (ξ, η) to (ξ', η') along a path γ in the configuration space defines an isomorphism:

$$T_{a,b}^{\gamma} : a \otimes_{(\xi,\eta)} b \xrightarrow{\sim} a \otimes_{(\xi',\eta')} b.$$

In mathematical language, $T^{\gamma} := \{T_{a,b}^{\gamma}\}_{a,b \in \Omega\mathcal{C}} : \otimes_{(\xi,\eta)} \rightarrow \otimes_{(\xi',\eta')}$ defines a natural isomorphism. Similar to 1+1D topological order, we have the following defining property of a 2+1D topological order:

Adiabatic Principle: if two paths γ_1 and γ_2 in the configuration space $\{(\xi, \eta) \in \mathbb{C}^2 \mid \xi \neq \eta\}$ are homotopy equivalent, then $T^{\gamma_1} = T^{\gamma_2}$.

(2.3.2)

Again Adiabatic Principle allows us to reduce the infinitely many fusion products $\otimes_{(\xi,\eta)}$ to a single fusion product $\otimes := \otimes_{(-1,1)}$ and a braiding isomorphisms $c_{a,b}$ defined by

$$a \otimes b \xrightarrow{c_{a,b} := T_{a,b}^{\gamma_+}} b \otimes a, \quad \gamma_+ := \{(e^{i(1-t)\pi}, e^{i(2-t)\pi}) \in \mathbb{C}^2 | t \in [0, 1]\}.$$

A physical work on adiabatic moves and braided monoidal structures can also be found in [KL20].

The point of this discussion is that, instead of strictly following the mathematical definition of a braided monoidal category, i.e. one tensor product plus braidings, we can return freely to above physical intuition of the infinitely many fusion products $\otimes_{(\xi,\eta)}$ whenever we discuss the anyons in a 2+1D topological order. Importantly, we have only sketched some key points of a complete story, which is essentially equivalent to the proof of the well-known mathematical result that an E_2 -algebra in the 2-category Cat_1 is equivalent to a braided monoidal 1-category [SW03, Lur17, Fre17]. We denote the category of E_2 -algebras in Cat_1 (or equivalently, braided monoidal 1-categories) by $\text{Alg}_{E_2}(\text{Cat}_1)$. Its 1-morphisms are E_2 -algebra homomorphisms (or braided monoidal functors), and 2-morphisms are morphisms between two E_2 -algebra homomorphisms (or braided monoidal natural transformations). Then the fact that $\Omega\mathcal{C}$ is a braided 1-category can be simply represented by a compact notation $\Omega\mathcal{C} \in \text{Alg}_{E_2}(\text{Cat}_1)$.

In the second way, we re-derive the braiding structure from the monoidal structure of $B\Omega\mathcal{C}$. Since $B\Omega\mathcal{C}$ is monoidal, its one point delooping $B^2\Omega\mathcal{C}$ is a well defined 3-category, a fact which is equivalent to $\Omega\mathcal{C}$ being braided monoidal. The 3-category $B^2\Omega\mathcal{C}$ is sometimes called a 2-tuply monoidal 1-category (a list of proposed definitions of a weak n -category can be found in [Lei02]; see also the list in nLab). This result simply says that the braiding structure can be replaced by the fusion product in two independent directions as illustrated in the right half of Figure 6.




Figure 6: the idea of braiding and 2-tuply monoidal 1-category

- (1) Anyon $a, c \in \Omega\mathcal{C}$ can be fused along the trivial string $\mathbb{1}$ (i.e., vertically in Figure 6). We denote this fusion product in x^2 -direction by $a \otimes^2 c$. Mathematically, this fusion is precisely the composition of two 1-morphisms $a, c \in \text{hom}_{\mathcal{C}}(\mathbb{1}, \mathbb{1})$:

$$\begin{aligned} \text{hom}_{\mathcal{C}}(\mathbb{1}, \mathbb{1}) \times \text{hom}_{\mathcal{C}}(\mathbb{1}, \mathbb{1}) &\xrightarrow{\circ} \text{hom}_{\mathcal{C}}(\mathbb{1}, \mathbb{1}) \\ (c, a) &\mapsto c \circ a = a \otimes^2 c. \end{aligned}$$

$1_{\mathbb{1}}$ is the unit of this fusion product, i.e. $1_{\mathbb{1}} \otimes^2 a \simeq a \simeq a \otimes^2 1_{\mathbb{1}}$.

- (2) There is another fusion product $\otimes^1$ on $\Omega\mathcal{C}$ defined by horizontally fusion two $\mathbb{1}$ -strings in Figure 6. Mathematically, it comes from the fusion product of $B\Omega\mathcal{C}$.

$$\begin{aligned} \text{hom}_{\mathcal{C}}(\mathbb{1}, \mathbb{1}) \times \text{hom}_{\mathcal{C}}(\mathbb{1}, \mathbb{1}) &\xrightarrow{\otimes} \text{hom}_{\mathcal{C}}(\mathbb{1} \otimes \mathbb{1}, \mathbb{1} \otimes \mathbb{1}) \simeq \text{hom}_{\mathcal{C}}(\mathbb{1}, \mathbb{1}) \\ (a, b) &\mapsto a \otimes^1 b. \end{aligned}$$

It is obvious that $1_{\mathbb{1}}$ is also the unit of the fusion product $\otimes^1$.

Physically, it is obvious that these two fusion products are compatible with each other. It means that the following diagram

$$\begin{array}{ccc} \Omega\mathcal{C} \times \Omega\mathcal{C} \times \Omega\mathcal{C} \times \Omega\mathcal{C} & \xleftarrow{1 \times \tau \times 1} & \Omega\mathcal{C} \times \Omega\mathcal{C} \times \Omega\mathcal{C} \times \Omega\mathcal{C} \\ \downarrow \otimes^1 \times \otimes^1 & & \downarrow \otimes^2 \times \otimes^2 \\ \Omega\mathcal{C} \times \Omega\mathcal{C} & \xrightleftharpoons{\delta} & \Omega\mathcal{C} \times \Omega\mathcal{C} \\ \searrow \otimes^2 & & \swarrow \otimes^1 \\ & \Omega\mathcal{C} & \end{array} \quad (2.3.3)$$

where τ is defined by $(b, c) \mapsto (c, b)$ for $c, b \in \Omega\mathcal{C}$, is commutative up to a natural isomorphism δ . Equivalently, there is a natural isomorphism, for $a, b, c, d \in \Omega\mathcal{C}$,

$$\delta_{a,b,c,d} : (a \otimes^1 b) \otimes^2 (c \otimes^1 d) \xrightarrow{\sim} (a \otimes^2 c) \otimes^1 (b \otimes^1 d). \quad (2.3.4)$$

- By restricting to a special case, we obtain

$$a \otimes^2 d \simeq (a \otimes^1 1_{\mathbb{1}}) \otimes^2 (1_{\mathbb{1}} \otimes^1 d) \xrightarrow{\delta_{a,1_{\mathbb{1}},1_{\mathbb{1}},d}} (a \otimes^2 1_{\mathbb{1}}) \otimes^1 (1_{\mathbb{1}} \otimes^2 d) \simeq a \otimes^1 d. \quad (2.3.5)$$

which says that two fusion products are isomorphic. If we use the physical intuition discussed in the first way. Without loss of generality, we can set $\otimes^1 = \otimes_{(0,1)}$ and $\otimes^2 = \otimes_{(0,i)}$, then $\delta_{a,1_{\mathbb{1}},1_{\mathbb{1}},d}$ can be defined physically by the isomorphism $T_{a,d}^\gamma$, where $\gamma := \{(0, e^{i\frac{\pi}{2}(1-t)}) \in \mathbb{C}^2 | t \in [0, 1]\}$.

- By restricting to another special case, we recover the braiding isomorphism

$$\begin{aligned} c_{b,c}^1 : b \otimes^1 c &\simeq (1_{\mathbb{1}} \otimes^2 b) \otimes^1 (c \otimes^2 1_{\mathbb{1}}) \xrightarrow{\delta_{1_{\mathbb{1}},b,c,1_{\mathbb{1}}}^{-1}} (1_{\mathbb{1}} \otimes^1 c) \otimes^2 (b \otimes^1 1_{\mathbb{1}}) \\ &\simeq (c \otimes^1 1_{\mathbb{1}}) \otimes^2 (1_{\mathbb{1}} \otimes^1 b) \xrightarrow{\delta_{c,1_{\mathbb{1}},1_{\mathbb{1}},d}} (c \otimes^2 1_{\mathbb{1}}) \otimes^1 (1_{\mathbb{1}} \otimes^1 b) \simeq c \otimes^1 b, \end{aligned} \quad (2.3.6)$$

$$\begin{aligned} c_{b,c}^2 : b \otimes^2 c &\simeq (1_{\mathbb{1}} \otimes^1 b) \otimes^2 (c \otimes^1 1_{\mathbb{1}}) \xrightarrow{\delta_{1_{\mathbb{1}},b,c,1_{\mathbb{1}}}} (1_{\mathbb{1}} \otimes^2 c) \otimes^1 (b \otimes^2 1_{\mathbb{1}}) \\ &\simeq (c \otimes^2 1_{\mathbb{1}}) \otimes^1 (1_{\mathbb{1}} \otimes^2 b) \xrightarrow{\delta_{c,1_{\mathbb{1}},1_{\mathbb{1}},d}^{-1}} (c \otimes^1 1_{\mathbb{1}}) \otimes^2 (1_{\mathbb{1}} \otimes^1 b) \simeq c \otimes^2 b, \end{aligned} \quad (2.3.7)$$

where we have used the fact that $1_{\mathbb{1}}$ is the tensor unit of both $\otimes^1$ and $\otimes^2$. We leave it as an exercise to prove that the isomorphism defined in (2.3.6) and (2.3.7) satisfies the axioms of a braiding.

In summary, we have sketched a proof of the fact that a 1-category $\mathcal{A}$ being braided monoidal if and only if $B\mathcal{A}$ is well defined and monoidal, or equivalently, $B^2\mathcal{A}$ is a well-defined 3-category. In more concrete terms, we have shown that the braiding monoidal structure of $\Omega\mathcal{C}$ can be equivalently encoded by two fusions in two independent directions and their compatibilities. Now we translate this compatibility into more formal mathematical language.

Recall that the category of E_1 -algebras in Cat_1 (i.e., monoidal 1-categories) is denoted by $\text{Alg}_{E_1}(\text{Cat}_1)$. Its 1-morphisms are E_1 -algebra homomorphisms (i.e., monoidal functors), and 2-morphisms are morphisms between two E_1 -algebra homomorphisms (i.e., monoidal natural transformations). Now the fact that the pair $(\otimes^2, 1_{\mathbb{1}})$ endows $\Omega\mathcal{C}$ with a monoidal structure can be represented by the following notation:

$$(\Omega\mathcal{C}, \otimes^2, 1_{\mathbb{1}}) \in \text{Alg}_{E_1}(\text{Cat}_1) \quad \text{or} \quad \Omega\mathcal{C} \in \text{Alg}_{E_1}(\text{Cat}_1) \text{ for simplicity.}$$

Moreover, $\text{Alg}_{E_1}(\text{Cat}_1)$ is again symmetric monoidal with the same Cartesian product $\times$ and the same tensor unit $\bullet$ as in Cat_1 . In particular, we obtain $\Omega\mathcal{C} \times \Omega\mathcal{C} \in \text{Alg}_{E_1}(\text{Cat}_1)$. The monoidal structure on $\Omega\mathcal{C} \times \Omega\mathcal{C}$ is defined by the composed functor $(\otimes^2 \times \otimes^2) \circ (1 \times \tau \times 1)$. Then the compatibility condition (2.3.3) simply says that $\otimes^1 : \Omega\mathcal{C} \times \Omega\mathcal{C} \rightarrow \Omega\mathcal{C}$ is an E_1 -algebra homomorphism (or a monoidal functor). It means that $\otimes^1 : \Omega\mathcal{C} \times \Omega\mathcal{C} \rightarrow \Omega\mathcal{C}$ is a 1-morphism in $\text{Alg}_{E_1}(\text{Cat}_1)$. Therefore, $\otimes^1$ endows $\Omega\mathcal{C}$ with a structure of an E_1 -algebra in $\text{Alg}_{E_1}(\text{Cat}_1)$. In other words, we obtain

$$\Omega\mathcal{C} \in \text{Alg}_{E_1}(\text{Alg}_{E_1}(\text{Cat})). \quad (2.3.8)$$

This notation is heavily loaded. It means that $\Omega\mathcal{C}$ has two multiplication maps $\otimes^1$ and $\otimes^2$, and $\otimes^1$ is a 1-morphism in $\text{Alg}_{E_1}(\text{Cat})$. What we have explained in this second way is a key fact

$$\text{Alg}_{E_2}(\text{Cat}_1) \simeq \text{Alg}_{E_1}(\text{Alg}_{E_1}(\text{Cat}_1)).$$

It is a well-known mathematical result that holds in more general context [Dun88, Lur17].

2.3.3 E_m -multi-fusion n -categories

We briefly review the notion of an E_m -multi-fusion n -category and some basic facts associated to it from [KZ22b, Section 3] and [KZ24].

Definition 2.3.1. An E_1 -monoidal n -category $\mathcal{A}$ is a pair $(\mathcal{A}, B\mathcal{A})$, where $B\mathcal{A}$ is the one-point delooping of $\mathcal{A}$. The identity 1-morphism $1_\bullet$ is the tensor unit of $\mathcal{A}$. By induction, an E_m -monoidal n -category is a pair $(\mathcal{A}, B\mathcal{A})$ such that $B\mathcal{A}$ is an E_{m-1} -monoidal n -category. By abusing notation, we simply denote such an E_{m-1} -monoidal n -category by $\mathcal{A}$, i.e.

$$\mathcal{A} = (\mathcal{A}, B\mathcal{A}, B^2\mathcal{A}, \dots, B^m\mathcal{A}).$$

When $m = 1$, $\mathcal{A}$ is also called a *monoidal n -category*; when $m = 2$, $\mathcal{A}$ is also called a *braided monoidal n -category*. $\square$

Convention 2.3.2. We adapt the following convention that is compatible with Convention 2.1.16. Note that $B^{m-1}\mathcal{A}$ is E_1 -monoidal and is equipped with a tensor product $\otimes^1$; $B^{m-2}\mathcal{A}$ is E_2 -monoidal and is equipped with two tensor products $\otimes^1, \otimes^2$; $\mathcal{A}$ is E_m -monoidal and is equipped with m tensor products $\otimes^1, \dots, \otimes^m$. We always use a coordinate system $(x^1, x^2, \dots, x^n)$ for the n -spatial dimensions such that $\otimes^i$ is the tensor product in the positive x^i -direction for $i = 1, \dots, n$.

Example 2.3.3. A symmetric monoidal 1-category is an E_3 -algebra in Cat_1 . It turns out that an E_3 -algebra in Cat_1 is automatically E_∞ -monoidal because Cat_1 does not have higher morphisms to distinguish higher commutativity. If one consider symmetric monoidal 3-category Cat_2 (i.e., the 3-category of 2-categories), then E_3 -algebras in Cat_2 that are not E_∞ . In this case, a monoidal 2-category is an E_1 -algebra in Cat_2 ; a braided monoidal 2-category is an E_2 -algebra in Cat_2 ; a sylleptic monoidal 2-category is an E_3 -algebra in Cat_2 ; a symmetric monoidal 2-category is an E_4 -algebra in Cat_2 , which is automatically an E_∞ -algebra in Cat_2 . In general, an E_{n+2} -monoidal n -category in Cat_n is automatically an E_∞ -algebra in Cat_n . $\heartsuit$

Definition 2.3.4. An E_0 -monoidal n -category $\mathcal{A}$ is a pair $(\mathcal{A}, \mathbb{1}_{\mathcal{A}})$, where $\mathcal{A}$ is an n -category and $\mathbb{1}_{\mathcal{A}} \in \mathcal{A}$ is a distinguished object. An E_0 -monoidal functor $(\mathcal{A}, \mathbb{1}_{\mathcal{A}}) \rightarrow (\mathcal{B}, \mathbb{1}_{\mathcal{B}})$ is a functor $f : \mathcal{A} \rightarrow \mathcal{B}$ such that $f(\mathbb{1}_{\mathcal{A}}) = \mathbb{1}_{\mathcal{B}}$. An E_0 -monoidal (higher) natural transformation is a (higher) natural transformation that is trivial on the distinguished object. We use $E_0\text{Cat}_n$ to denote the $(n+1)$ -category of E_0 -monoidal categories, E_0 -monoidal functors and E_0 -monoidal (higher) natural transformations. We set

$$\text{Fun}^{E_0}((\mathcal{A}, \mathbb{1}_{\mathcal{A}}), (\mathcal{B}, \mathbb{1}_{\mathcal{B}})) := \text{hom}_{E_0\text{Cat}_n}((\mathcal{A}, \mathbb{1}_{\mathcal{A}}), (\mathcal{B}, \mathbb{1}_{\mathcal{B}})),$$

which is (not full) subcategory of $\text{Fun}(\mathcal{A}, \mathcal{B})$. We use $E_m\text{Cat}_n$ to denote the full $(n+1)$ -subcategory $E_0\text{Cat}_n$ consisting of all the iterated 1-point deloopings $B^m\mathcal{C}$. We set

$$\text{Fun}^{E_m}(\mathcal{A}, \mathcal{B}) := \text{Fun}^{E_0}(B^m\mathcal{A}, B^m\mathcal{B}).$$

Definition 2.3.5. For $k > -m$, we use $\mathcal{A}^{\text{op}k}$ to denote the E_m -monoidal n -category obtained by reversing all the k -morphisms, i.e., $B^m(\mathcal{A}^{\text{op}k}) = (B^m\mathcal{A})^{\text{op}(k+m)}$. We set $\mathcal{A}^{\text{op}} := (B^m\mathcal{A})^{\text{op}1} = \mathcal{A}^{\text{op}(1-m)}$. $\square$

Remark 2.3.6. According to Convention 2.1.16 and 2.3.2, $\mathcal{A}^{\text{op}}$ flips the fusion product $\otimes^1$ (in the x^1 -direction) to the opposite fusion product $(\otimes^1)^{\text{op}}$ defined by $a(\otimes^1)^{\text{op}}b := b \otimes^1 a$. $\diamond$

We denote by $\text{KarCat}_n^{\mathbb{C}}$ the $(n+1)$ -category of condensation complete (or Karoubi complete) $\mathbb{C}$ -linear n -categories ($\mathbb{C}$ -linear functors, etc) [JF22, KZ22b]. It is also a symmetric monoidal $(n+1)$ -category [JF22]. Its tensor product, denoted by $\boxtimes$, is defined by

$$\text{Fun}_{\mathbb{C}}(\mathcal{A} \boxtimes \mathcal{B}, -) \simeq \text{Fun}_{\mathbb{C}}(\mathcal{A}, \text{Fun}_{\mathbb{C}}(\mathcal{B}, -)),$$

where $\text{Fun}_{\mathbb{C}}(\cdot)$ denotes the category of $\mathbb{C}$ -linear n -functors. Its tensor unit is $n\text{Vec} := \Sigma^n\mathbb{C}$. We denote by $E_m\text{KarCat}_n^{\mathbb{C}}$ the $(n+1)$ -category of E_m -monoidal n -categories, E_m -monoidal functors, E_m -monoidal (higher) natural transformations, etc. The notion of an E_m -multi-fusion n -category was introduced in [JF22]. We take the definition from [KZ22b].

Definition 2.3.7. An E_m -multi-fusion n -category $\mathcal{A}$ is a condensation-complete $\mathbb{C}$ -linear E_m -monoidal n -category, i.e., $\mathcal{A} \in E_m \text{KarCat}_n^{\mathbb{C}}$, such that $\Sigma^m \mathcal{A}$ is a separable $(n+m)$ -category, i.e., $\Sigma^m \mathcal{A} \in (n+m+1)\text{Vec}$. It is called *indecomposable* if $\mathcal{A}$ is not a direct sum of two non-zero separable n -categories. If the tensor unit $\mathbb{1}_{\mathcal{A}}$ is simple, then $\mathcal{A}$ is called an E_m -fusion n -category. An E_m -fusion 0-category is just $\mathbb{C}$. $\square$

Example 2.3.8. We provide some useful examples.

1. If $\mathcal{A}$ is also E_{∞} -monoidal, it means that $\Sigma^n \mathcal{A}$ can be defined for all n . In this case, $\Sigma^n \mathcal{A}$ is also E_{∞} -monoidal. For example, symmetric fusion 1-categories are $\text{Rep}(G)$ or $\text{Rep}(G, z)$ for a finite group G . Then $(n+1)\text{Rep}(G) = \Sigma^n \text{Rep}(G)$ and $(n+1)\text{Rep}(G, z) = \Sigma^n \text{Rep}(G, z)$ are E_{∞} -monoidal for all n . One can equivalently define $n\text{Rep}(G)$ by $n\text{Rep}(G) := \text{Fun}(\text{BG}, n\text{Vec})$ and prove that $n\text{Rep}(G) \simeq \Sigma(n-1)\text{Rep}(G)$ as a theorem [KZ24]. We treat different but equivalent definitions of $n\text{Rep}(G)$ as different coordinate systems of $n\text{Rep}(G)$.
2. For $\omega \in Z^{n+2}(G, U(1))$, $n\text{Vec}_G^{\omega}$ is an E_1 -multi-fusion n -category [KZ24], and $\mathfrak{Z}_1(n\text{Vec}_G^{\omega})$ is an E_2 -multi-fusion n -category. When ω is trivial and G is abelian, $n\text{Vec}_G$ is clearly a symmetric or E_{∞} -fusion n -category.
3. An n -groupoid is an n -category such that all morphisms are invertible. An n -group is a monoidal $(n-1)$ -category $\mathcal{G}$ such that $\text{B}\mathcal{G}$ is an n -groupoid. It is called finite if all homotopy groups $\pi_i(\mathcal{G})$ are finite. For a finite n -group $\mathcal{G}$, we recall some results in [KZ24].
 - $n\text{Rep}(\mathcal{G}) := \text{Fun}(\text{B}\mathcal{G}, n\text{Vec})$ is a symmetric fusion n -category and $(n+1)\text{Rep}(\mathcal{G}) \simeq \Sigma n\text{Rep}(\mathcal{G})$.
 - $n\text{Vec}_{\mathcal{G}} := \Omega L(\text{B}\mathcal{G})$ is a fusion n -category, where L is the left adjoint functor of the forgetful functor $f : \text{KarCat}_{n+1}^{\mathbb{C}} \rightarrow \text{KarCat}_{n+1}$. $\heartsuit$

When G is a finite abelian group. Let $\chi(G) := \text{hom}(G, \mathbb{C}^{\times})$ be the character group. We have $n\text{Vec}_{\text{B}^{n-1}G} \simeq n\text{Rep}(\chi(G))$ and $n\text{Rep}(\text{B}^{n-1}G) \simeq n\text{Vec}_{\chi(G)}$ [KZ24].

4. Using the E_k -center for $k \geq 0$ (see Definition 2.3.17), we can construct an E_{k+1} -multi-fusion n -category from an E_k -multi-fusion n -category. For example, $\mathfrak{Z}_k(n\text{Rep}(G))$ is an E_{k+1} -multi-fusion n -category. In this case, unfortunately, it does not generate new examples when $k \geq 3$ (see (5.2.22)).
5. Another way to generate a new E_k -multi-fusion n -category from an old one $\mathcal{C}$ is by condensations. More precisely, the new category is the category of E_k -modules over a condensable E_k -algebra in $\mathcal{C}$ explained Section 5.2.

The following results are useful tools to determine or find new E_k -multi-fusion higher categories.

Theorem° 2.3.9 ([KZ22b]). Let $\mathcal{A}$ be a $\mathbb{C}$ -linear condensation-complete E_m -monoidal n -category for $m \geq 1$. If $\Sigma \mathcal{A}$ is a separable $(n+1)$ -category, then $\mathcal{A}$ is an E_m -multi-fusion n -category.

Corollary° 2.3.10 ([KZ22b]). If $\mathcal{A}$ is an E_m -multi-fusion n -category for $n \geq 1$, then $\Omega \mathcal{A}$ is an E_{m+1} -multi-fusion $(n-1)$ -category.

2.3.4 Condensable higher algebras

We have seen that an E_m -multi-fusion n -category is an E_m -algebra in $(n+1)\text{Vec}$. It turns out that they are also condensable (or separable) in a sense that we want to explain now. However, the mathematical theory of condensable (or separable) higher algebras in higher categories is very rich and highly non-trivial. It is beyond this work. We will develop such a theory in [KZZZ25]. In this work, we only briefly sketch some facts that are useful to later sections and provide some intuitions.

A 0-category is defined by a set. A 0-condensation is an equality between objects of a 0-category.

Definition 2.3.11 ([GJF19]). For two objects X and Y in a weak n -category $\mathcal{C}$, an n -condensation of X onto Y , denoted by $X \rightarrow Y$, is a pair of 1-morphism $f : X \rightarrow Y$ and $g : Y \rightarrow X$, together with an $(n-1)$ -condensation $f g \rightarrow \text{id}_Y$, i.e.

$$X \rightarrow Y := \{f : X \rightrightarrows Y : g, \quad f g \rightarrow \text{id}_Y\}$$

The object Y is called a condensate of X . A condensation is an n -condensation for some n . $\square$

Example 2.3.12. Let us unravel the definition in a few lower dimensional cases.

1. When $n = 1$, $\mathcal{C}$ is a 1-category. In this case, a 1-condensation is a pair of 1-morphisms $f : X \rightarrow Y$ and $g : Y \rightarrow X$ such that $fg = \text{id}_Y$. As a consequence, $p := gf$ is a idempotent, i.e. $p^2 = p$.
2. When $n = 2$, $\mathcal{C}$ is a 2-category. A 2-condensation in $\mathcal{C}$ is a pair of 1-morphisms $f : X \rightarrow Y$ and $g : Y \rightarrow X$ such that $fg \rightarrow \text{id}_Y$ is a 1-condensation, i.e. there is a pair of 2-morphism $i : fg \rightarrow \text{id}_Y$ and $j : \text{id}_Y \rightarrow fg$ such that $ij = \text{id}_{\text{id}_Y}$.
- (a) Let $\mathcal{A}$ be a multi-fusion category, and let $\text{LMod}_{\mathcal{A}}(2\text{Vec})$ be the 2-category of left finite semisimple $\mathcal{A}$ -modules. Note that

$$\Omega_{\mathcal{A}}(\text{LMod}_{\mathcal{A}}(2\text{Vec})) = \text{Fun}_{\mathcal{A}}(\mathcal{A}, \mathcal{A}) \simeq \mathcal{A}^{\text{op}} \quad (2.3.9)$$

♡

Let $A = (A, \mu_A, \eta_A)$ be an algebra in $\mathcal{A}$, where $\mu_A : A \otimes A \rightarrow A$ is the multiplication and $\eta_A : \mathbb{1}_{\mathcal{A}} \rightarrow A$ is the unit. Then $\mathcal{M} := \text{RMod}_{\mathcal{A}}(\mathcal{A}) \in \text{LMod}_{\mathcal{A}}(2\text{Vec})$. Now we would like to construct a 2-condensation $\mathcal{A} \rightarrow \mathcal{M}$ when the algebra A is separable. It means that we should construct

$$f : \mathcal{A} \rightarrow \mathcal{M}, \quad g : \mathcal{M} \rightarrow \mathcal{A}, \quad i : fg \rightarrow \text{id}_{\mathcal{M}}, \quad j : \text{id}_{\mathcal{M}} \rightarrow fg$$

- $f = - \otimes A : \mathcal{A} \rightarrow \mathcal{M}$ defines a 1-morphism in $\text{LMod}_{\mathcal{A}}(2\text{Vec})$.
- $g = [A, -] : \mathcal{M} \rightarrow \mathcal{A}$ is precisely the forgetful functor and is automatically a left $\mathcal{A}$ -module functor, and, at the same time, the right adjoint of f .
- $fg = [A, -] \otimes A$ is equipped with an evaluation 2-morphism

$$fg = [A, -] \otimes A \xrightarrow{i=\text{ev}} \text{id}_{\mathcal{M}},$$

i.e. $\text{ev}_M = \mu_M : M \otimes A \rightarrow M$, which is nothing but the right A -action on M . Note that i is a 1-morphism in $\text{Fun}_{\mathcal{A}}(\mathcal{M}, \mathcal{M})$.

- We need a natural transformation $j : \text{id}_{\mathcal{M}} \rightarrow [A, -] \otimes A$ splitting i in $\text{Fun}_{\mathcal{A}}(\mathcal{M}, \mathcal{M})$. By [KZ17, Theorem 4.10], this condition is equivalent to the condition that A is separable algebra in $\mathcal{A}$ in the usual sense, i.e., $\mu_A : A \otimes A \rightarrow A$ splits as A - A -bimodules.
- (b) Using [GJF19, Proposition 3.1.5], a separable algebra A in $\mathcal{A}^{\text{op}}$ can be defined as $A = u^R \circ u$, where $u : \mathcal{A} \rightarrow \mathcal{M} \in \text{LMod}_{\mathcal{A}}(2\text{Vec})$ extends to a 2-condensation in 2Vec and u^R is the right adjoint of u . This idea can be used to define a separable higher algebra in a higher multi-fusion categories. We will show that elsewhere.

The precise definition of a condensable E_m -algebra in an E_m -multi-fusion n -category will be given elsewhere. We only provide a working definition that provides some physical intuitions here. For $m = 0$, a condensable E_0 -algebra in an E_0 -multi-fusion n -category $(\mathcal{A}, \mathbb{1}_{\mathcal{A}})$ is a pair (A, η_A) , where A is an object in $\mathcal{A}$ equipped with a unit morphism $\mathbb{1}_{\mathcal{A}} \xrightarrow{\eta_A} A$. For $m \leq 1$, roughly speaking, there is a morphism $\mu_A^i : A \otimes A \rightarrow A$ of an E_m -algebra that defines the multiplication in the i -th independent direction for $i = 1, 2, \dots, m$. All of them are compatible in the sense that $\mu_A := \mu_A^1 \simeq \mu_A^i$ (see Section 4.1.1 for more discussion). It has a right adjoint μ_A^R , and both μ_A and μ_A^R intertwine the m -dimensional A -actions. If μ_A is an n -morphism, its right adjoint is not defined. In this case, for conveniences and physical applications, we assume that top two morphisms form a $*$ -category. In particular, the space of n -morphisms are Hilbert space and f^* is defined for each n -morphisms f .²⁶ We set $\mu_A^R = \mu_A^*$. Since A plays the role of the vacuum in the condensed phase, we need impose a stability condition of the vacuum as explained in [Kon14] for the $m = 1, 2$ and $n = 1$ cases. More explicitly, it means that $\mu_A \circ \mu_A^R = \text{id}_A$ when $m = 1, 2$ and $n = 1$, and it can be generalized to higher dimensions.

Definition 2.3.13. In an E_m -multi-fusion n -category $\mathcal{A}$, a condensable (or separable) E_m -algebra A is an E_m -algebra A equipped with an $(n - 1)$ -condensation $\mu_A \circ \mu_A^R \rightarrow \text{id}_A$ in $\text{Mod}_{\mathcal{A}}^{E_m}(\mathcal{A})$ (i.e., modules equipped with m -dimensional A -actions explained later), where μ_A^R is the right adjoint of the multiplication 1-morphism $\mu_A : A \otimes A \rightarrow A$. It is called *simple* if the morphism $\eta_A : \mathbb{1}_{\mathcal{A}} \rightarrow A$ is simple for $n > 1$ and $\text{hom}(\mathbb{1}_{\mathcal{A}}, A) \simeq \mathbb{C}$ for $n = 1$. It is called *indecomposable* if A is not a direct sum of two algebras. □

²⁶In this case, we have $g^R = g^L = g^*$ for all k -morphisms g [KWZ15, Proposition A.7].

Example 2.3.14. When $m = 1$ and $n = 2$, above definition is the same as the notion of a separable algebra in [D  c23b, DX24]. According to [D  c23b, DX24], the following statements are true.

1. Condensable E_1 -algebras in 2Vec are precisely multi-fusion 1-categories, and simple ones are fusion 1-categories, and indecomposable ones are indecomposable multi-fusion 1-categories.
2. Condensable E_2 -algebras in 2Vec are precisely E_2 -multi-fusion 1-categories or braided multi-fusion 1-categories. Indecomposable ones are automatically simple and are E_2 -fusion 1-categories (or braided fusion 1-categories).
3. Condensable E_3 -algebras in 2Vec are symmetric multi-fusion 1-categories, which is automatically E_∞ -monoidal.
4. Condensable E_2 -algebras in $\mathfrak{Z}_1(2\text{Vec}_G)$ for a finite group G are exactly G -crossed braided multi-fusion 1-categories.
5. Let $\mathcal{B}$ be an E_2 -fusion 1-category. A braided multi-fusion 1-category $\mathcal{A}$ equipped with a braided functor $\mathcal{B} \rightarrow \mathcal{A}$ is a condensable E_2 -algebra in the E_2 -fusion 2-category $\mathfrak{Z}_1(\Sigma\mathcal{B})$. $\heartsuit$

Example 2.3.15. We give a few examples of condensable E_k -algebras in higher categories.

1. Condensable E_k -algebras in $(n + 1)\text{Vec}$ are precisely E_k -multi-fusion categories.
2. When G is abelian, condensable E_k -algebras in $(n + 1)\text{Vec}_G$ are precisely G -graded E_k -multi-fusion categories.
3. Condensable E_k -algebras in $(n + 1)\text{Rep}(G)$ in the coordinate $(n + 1)\text{Rep}(G) = \text{Fun}(BG, (n + 1)\text{Vec})$ are precisely G -equivariant E_k -multi-fusion categories, a notion which means an E_k -multi-fusion category $\mathcal{A}$ equipped with a monoidal functor $G \rightarrow \text{Aut}^{E_k}(\mathcal{A})$ and $\text{Aut}^{E_k}(\mathcal{A})$ is the category of E_k -monoidal auto-equivalences of $\mathcal{A}$.
4. A condensable E_2 -algebras in $\mathfrak{Z}_1((n + 1)\text{Vec}_G)$ is precisely a G -crossed braided multi-fusion n -category (see Example 4.2.44 (1) for a physical explanation).

We will develop the theory of condensable E_m -algebras in an E_m -multi-fusion category elsewhere. In this work, we simply take it for granted that, for $m \geq 1$, condensable E_m -algebras in $(n + 1)\text{Vec}$ are precisely E_m -multi-fusion n -categories. Indecomposable condensable E_m -algebras in $(n + 1)\text{Vec}$ are E_m -fusion n -categories for $m > 1$ and are indecomposable multi-fusion n -categories for $m = 1$; simples ones are E_m -fusion n -categories for all $m \geq 1$. For an E_m -multi-fusion n -category $\mathcal{A}$, we denote the category of condensable E_m -algebras in $\mathcal{A}$ by $\text{Alg}_{E_m}^c(\mathcal{A})$. We denote the statement that A is a condensable E_m -algebra A in $\mathcal{A}$ by $A \in \text{Alg}_{E_m}^c(\mathcal{A})$. Then we also have $\mathcal{A} \in \text{Alg}_{E_m}^c((n + 1)\text{Vec})$. As we have explained in Section 2.3.2, by iterating the same argument, we obtain

$$\text{Alg}_{E_m}^c(\mathcal{A}) \simeq \text{Alg}_{E_1}^c(\text{Alg}_{E_{m-1}}^c(\mathcal{A})).$$

Moreover, from Example 2.3.12, we see that a condensable E_1 -algebra A in a multi-fusion 1-category can always be realized by an internal hom $[A, A]$ (see Section 3.1.2 for a brief introduction of this notion). This fact can be generalized to all condensable E_m -algebras in any E_m -multi-fusion n -categories. We will develop the theory elsewhere. In this work, we simply take it for granted that all internal homs naturally constructed are condensable higher algebras, and vice versa. A condensable E_1 -algebra is a direct sum of indecomposable ones²⁷; and a condensable E_m -algebra for $m > 1$ is direct sum of simple ones. By our condensation theory and results in [KZ22b, KZ24], the category of E_m -modules over a condensable E_m -algebras in an E_m -multi-fusion n -category is automatically condensation complete and E_m -multi-fusion. The special case for $m = n = 2$ was proved in [DX24]. We also want to remind readers about Remark 1.2.8 for our assumptions on condensability or separability used throughout this work.

Remark 2.3.16. For physics oriented readers, it is helpful and harmless to ignore the subtleness in the definition of separability and take it for granted that all physically natural algebras or higher monoidal categories are automatically condensable or separable and unitary. Indeed, in all pictures, when we illustrate the physical intuitions, the unitarity is automatically assumed to avoid the framing issues. Under the unitary assumption, the left and right duals of a topological defect x are the same, i.e., $x^R = x^L = x^*$, which should be viewed as the anti-defect of x . $\diamond$

²⁷See the structure theorem of a condensable E_1 -algebra in a multi-fusion 1-category in Theorem 3.1.11 and a proposed structure theorem for such algebras in a multi-fusion n -category in Remark 3.1.12.

2.3.5 E_m -center and E_m -centralizer

The notion of the center of an algebraic structure is very common in both physics and mathematics. The familiar notions of center include the center of a group G , defined by $Z(G) := \{z \in G | zg = gz, \forall g \in G\}$; the center of a $\mathbb{C}$ -algebra A , defined by $Z(A) := \{z \in A | za = az, \forall a \in A\}$; the Drinfeld center of a fusion category; the Müger center of a braided fusion category; the full center in 1+1D rational CFTs. The reason that they are all called ‘centers’ because they satisfy the same universal properties (defined in different categories).

In this subsubsection, we review the definitions of and some useful results on E_m -centralizer and E_m -centers from [KZ24]. For physics oriented readers, the definition is quite abstract. We suggest readers to first focus on the results instead of the definitions.

Definition 2.3.17. For $\mathcal{A}, \mathcal{B} \in E_m \text{KarCat}_n^{\mathbb{C}}$ and a $\mathbb{C}$ -linear E_m -monoidal functor $F : \mathcal{A} \rightarrow \mathcal{B}$, the E_m -centralizer of F is the universal condensation-complete $\mathbb{C}$ -linear E_m -monoidal n -category $\mathfrak{Z}_m(F)$ equipped with a unital action $G : \mathfrak{Z}_m(F) \boxtimes \mathcal{A} \rightarrow \mathcal{B}$, i.e. a $\mathbb{C}$ -linear E_m -monoidal functor exhibiting the following diagram commutative (up to equivalences).

$$\begin{array}{ccc} & \mathfrak{Z}_m(F) \boxtimes \mathcal{A} & \\ \mathbb{1}_{\mathfrak{Z}_m(F)} \boxtimes \text{id}_{\mathcal{A}} \nearrow & & \searrow G \\ \mathcal{A} & \xrightarrow{F} & \mathcal{B} \end{array}$$

For convenience, when F is clear from the context, we also denote $\mathfrak{Z}_m(F)$ by $\mathfrak{Z}_m(\mathcal{A}, \mathcal{B})$. When $F = \text{id}_{\mathcal{A}}$, then $\mathfrak{Z}_m(\mathcal{A}) := \mathfrak{Z}_m(\text{id}_{\mathcal{A}}) = \mathfrak{Z}_m(\mathcal{A}, \mathcal{A})$ defines the E_m -center of $\mathcal{A}$. $\square$

Remark 2.3.18. We spell out the universal property of $\mathfrak{Z}_m(F)$ more explicitly. If $\mathcal{X} \in E_m \text{KarCat}_n^{\mathbb{C}}$ is equipped with a unital action $H : \mathcal{X} \boxtimes \mathcal{A} \rightarrow \mathcal{B}$ (i.e., exhibiting the bottom triangle in the following diagram commutative),

$$\begin{array}{ccc} & \mathfrak{Z}_m(F) \boxtimes \mathcal{A} & \\ \mathbb{1}_{\mathfrak{Z}_m(F)} \boxtimes \text{id}_{\mathcal{A}} \nearrow & \uparrow \exists! f \boxtimes \text{id}_{\mathcal{A}} & \searrow G \\ & \mathcal{X} \boxtimes \mathcal{A} & \\ \mathbb{1}_{\mathcal{A}} \boxtimes \text{id}_{\mathcal{X}} \nearrow & & \searrow H \\ \mathcal{A} & \xrightarrow{F} & \mathcal{B} \end{array} \quad (2.3.10)$$

then there is a unique $\mathbb{C}$ -linear E_m -monoidal functor $f : \mathcal{X} \rightarrow \mathfrak{Z}_m(F)$ rendering above diagram commutative. $\diamond$

Example 2.3.19. We want to emphasize that the diagram (2.3.10) defines the notion of a centralizer or center in any category. In other words, any thing satisfies the universal property defined by the diagram (2.3.10) in any (higher) categories is qualified to be called a ‘centralizer’ (or a ‘center’ for $F = \text{id}_{\mathcal{A}}$). We explain a few well known centers in terms of E_m -centers.

1. The center of a group is an E_1 -center in the category of sets. Note that a group can be viewed as an E_1 -algebra in the category of sets.
2. The center of a $\mathbb{C}$ -algebra A is an E_1 -center in the category of vector spaces over $\mathbb{C}$.
3. The Drinfeld center of a monoidal 1-category is an E_1 -center in the 2-category Cat_1 of 1-categories (as objects), functors (as 1-morphisms) and natural transformations (as 2-morphisms). An E_1 -algebra in Cat_1 is precisely a monoidal 1-category.
4. The Müger center of a braided monoidal 1-categories is an E_2 -center in the 2-category Cat_1 . An E_2 -algebra in Cat_1 is precisely a braided monoidal 1-category.

Example 2.3.20. In the category of potentially anomalous topological orders (or quantum liquids or QFT’s), a morphism $f : A^n \rightarrow B^n$ between A^n and B^n was introduced in [KWZ15, KWZ17] and is defined by a gapped domain wall f^n between $Z(A)^{n+1}$ and $Z(B)^{n+1}$ such that $f^n \boxtimes_{Z(A)^{n+1}} A^n = B^n$ as illustrated below.

$$\begin{array}{c} Z(B)^{n+1} \quad \xrightarrow{f^n} \quad Z(A)^{n+1} \quad \xrightarrow{\quad} \quad A^n \\ \hline f^n \boxtimes_{Z(A)^{n+1}} A^n = B^n \end{array}$$

By checking the universal property of the centralizer (2.3.10), it is an easy exercise to show that $f^n = Z(f)^n$, where $Z(f)^n$ denotes the centralizer²⁸ of f , and $Z(A)^n = Z(\text{id}_A)^n$ (see [KWZ15, KWZ17] for a proof of the later case and the same proof works for generic f). Similarly, in the category of fusion n -categories, for a monoidal functor $f : \mathcal{A} \rightarrow \mathcal{B}$, we have $\mathfrak{Z}_1(f) \simeq \text{Fun}_{\mathcal{A}|\mathcal{B}}(\mathcal{B}, \mathcal{B})$, where the left $\mathcal{A}$ -module structure on $\mathcal{B}$ is defined by the monoidal functor f . $\heartsuit$


Remark 2.3.21. There are more general notions of centers. For example, consider an algebra A in a monoidal 1-category $\mathcal{A}$. It is impossible to define the center of A within $\mathcal{A}$ because there is no structure of commutativity in $\mathcal{A}$. However, it is possible to define the notion of the center of A in the Drinfeld center of $\mathcal{A}$ [Dav10a, KYZ21]. $\diamond$

Example 2.3.22. For $F : \mathcal{A} \rightarrow \mathcal{B}$, $\mathfrak{Z}_0(F) = \mathfrak{Z}_0(\mathcal{A}, \mathcal{B}) = (\text{Fun}(\mathcal{A}, \mathcal{B}), F)$ because giving a unital action $\mathbb{C} \boxtimes \mathcal{A} \rightarrow \mathcal{B}$ is equivalent to giving a $\mathbb{C}$ -linear functor $\mathbb{C} \rightarrow \text{Fun}(\mathcal{A}, \mathcal{B})$ that maps $\mathbb{1}_{\mathbb{C}}$ to F . $\heartsuit$

Example 2.3.23. For $\mathcal{A} \in \text{Alg}_{\mathbb{E}_1}^{\mathbb{C}}((n+1)\text{Vec})$ and $\mathcal{M} \in \text{LMod}_{\mathcal{A}}((n+1)\text{Vec})$, by definition, $\mathcal{M}$ is equipped with a monoidal functor $F : \mathcal{A} \rightarrow \mathfrak{Z}_0(\mathcal{M})$. We have

$$\mathfrak{Z}_1(F) = \mathfrak{Z}_1(\mathcal{A}, \mathfrak{Z}_0(\mathcal{M})) \simeq \text{Fun}_{\mathcal{A}}(\mathcal{M}, \mathcal{M}). \quad (2.3.11)$$

In fact, giving a unital action $\mathcal{X} \boxtimes \mathcal{A} \rightarrow \mathfrak{Z}_0(\mathcal{M})$ is equivalent to giving a $\mathbb{C}$ -linear monoidal functor $\mathcal{X} \rightarrow \text{Fun}_{\mathcal{A}}(\mathcal{M}, \mathcal{M})$. The physical meaning of (2.3.11) is illustrated below.



$$\begin{aligned} \mathcal{M}^n &= (\mathcal{M}, m), \quad m \in \mathcal{M}, \\ \mathcal{B}^{\text{op}} &= \text{Fun}_{\mathcal{A}}(\mathcal{M}, \mathcal{M}) \simeq \mathfrak{Z}_1(\mathcal{A}, \mathfrak{Z}_0(\mathcal{M})), \\ \mathcal{A} &= \text{Fun}_{\mathcal{B}^{\text{op}}}(\mathcal{M}, \mathcal{M}) \simeq \mathfrak{Z}_1(\mathcal{B}^{\text{op}}, \mathfrak{Z}_0(\mathcal{M})), \end{aligned} \quad (2.3.12)$$

where the categories of all topological defects in two simple $n+1$ D topological order A^{n+1} and B^{n+1} are the fusion n -category $\mathcal{A}$ and $\mathcal{B}$, respectively, and the gapped domain wall M^n , as we show later, can be described by a pair $(\mathcal{M}, m)$, where $\mathcal{M} \in \text{BMod}_{\mathcal{A}|\mathcal{B}}((n+1)\text{Vec})$ is the category of wall conditions and m is an object in $\mathcal{M}$ specifying a wall condition. For example, when $\mathcal{A} = n\text{Vec}$, $\mathcal{M} \in (n+1)\text{Vec}$, if $F : n\text{Vec} \rightarrow \mathfrak{Z}_0(\mathcal{M})$ is the tensor unit of $\mathfrak{Z}_0(\mathcal{M})$, we obtain $\mathfrak{Z}_1(F) \simeq \mathfrak{Z}_0(\mathcal{M})$. $\heartsuit$

Example 2.3.24. The notion of a morphism between two (potentially anomalous) topological orders is defined by a gapped domain wall. For example, M^n in Figure 3 defines a morphism between A^{n+1} and B^{n+1} .²⁹ However, this notion only distinguish topological orders up to Morita equivalence. A more fundamental notion of a morphism between two topological orders should distinguish them up to isomorphisms. Such a notion was introduced [KWZ15] (see also [KWZ17] and Figure 3). For example, roughly speaking, B^{n+1} in the picture in (2.3.12) defines a morphism $F : A^{n+1} \rightarrow Z(\mathcal{M})^{n+1}$. This notion leads to a new symmetric monoidal higher category of $n+1$ D topological orders [KWZ15]. In this category, the bulk $Z(A)^{n+2}$ of A^{n+1} is indeed the center of A^{n+1} . This justifies the notation of the bulk. Moreover, one can also define the notion of centralizer in this category as in Definition 2.3.17. For example, we have $Z(F)^{n+1} = Z(A, Z(\mathcal{M}))^{n+1} = B^{n+1}$. $\heartsuit$

Proposition^o 2.3.25. Let $F : \mathcal{A} \rightarrow \Omega\mathcal{B}$ be a $\mathbb{C}$ -linear E_m -monoidal functor for $\mathcal{A} \in E_m\text{KarCat}_n^{\mathbb{C}}$ and $\mathcal{B} \in E_{m-1}\text{KarCat}_n^{\mathbb{C}}$. We have

$$\mathfrak{Z}_m(\mathcal{A}, \Omega\mathcal{B}) \simeq \Omega\mathfrak{Z}_{m-1}(\Sigma\mathcal{A}, \mathcal{B}). \quad (2.3.13)$$

Theorem^o 2.3.26 ([KZ24]). Let $F : \mathcal{A} \rightarrow \mathcal{B}$ be a $\mathbb{C}$ -linear E_m -monoidal functor for $\mathcal{A}, \mathcal{B} \in E_m\text{KarCat}_n^{\mathbb{C}}$. For $0 \leq k \leq m$, F induces canonically a functor $\Sigma^k F : \Sigma^k \mathcal{A} \rightarrow \Sigma^k \mathcal{B}$. We have

$$\mathfrak{Z}_m(F) = \mathfrak{Z}_m(\mathcal{A}, \mathcal{B}) = \Omega^k \mathfrak{Z}_{m-k}(\Sigma^k \mathcal{A}, \Sigma^k \mathcal{B}) = \Omega^m(\text{Fun}(\Sigma^m \mathcal{A}, \Sigma^m \mathcal{B}), \Sigma^m F). \quad (2.3.14)$$

As a special case, we have

$$\mathfrak{Z}_m(\mathcal{A}) \simeq \Omega\mathfrak{Z}_{m-1}(\Sigma\mathcal{A}) \simeq \Omega^m \mathfrak{Z}_0(\Sigma^m \mathcal{A}) \simeq \Omega^m \text{Fun}(\Sigma^m \mathcal{A}, \Sigma^m \mathcal{A}). \quad (2.3.15)$$

²⁸We prefer not to specify a subscript for the ‘Z’ in the centralizer $Z(f)^n$ of f . But we will provide a subscript for ‘Z’ as in Definition 2.3.17 when A^n and B^n are replaced by their categorical description $\mathcal{A}$ and $\mathcal{B}$ and $f : \mathcal{A} \rightarrow \mathcal{B}$ is a monoidal functor.

²⁹For intrigued readers, we leave it as an exercise to compute the centralizer of this morphism M^n . We will discuss it elsewhere.

Remark 2.3.27. By the results in [KZ22b, KZ24], the notion of a center or centralizer preserves the separability (or the condensation completeness). In particular, $\mathfrak{Z}_m(\mathcal{A})$ is automatically E_{m+1} -multi-fusion and $\mathfrak{Z}_m(\mathcal{A}, \mathcal{B})$ is E_m -multi-fusion if both $\mathcal{A}$ and $\mathcal{B}$ are E_m -multi-fusion. $\diamond$

Example 2.3.28. When $m = 1$, we obtain

$$\mathfrak{Z}_1(\mathcal{A}, \mathcal{B}) = \Omega(\text{Fun}(\Sigma\mathcal{A}, \Sigma\mathcal{B}), F) \simeq \text{Fun}_{\mathcal{A}|\mathcal{B}}(\mathcal{B}, \mathcal{B}) \simeq \text{Fun}_{\mathcal{A}|\mathcal{B}}(\mathcal{A}, \mathcal{B}).$$

Let $\mathcal{A}$ be an indecomposable multi-fusion n -category. By Proposition 2.2.34, we have the decomposition $\mathcal{A} \simeq \oplus_{i,j} \mathcal{A}_{ij}$ as separable n -categories. We obtain a useful result:

$$\mathfrak{Z}_1(\mathcal{A}) \simeq \Omega\mathfrak{Z}_0(\Sigma\mathcal{A}) \simeq \Omega\mathfrak{Z}_0(\Sigma\mathcal{A}_{ii}) \simeq \mathfrak{Z}_1(\mathcal{A}_{ii}).$$

Corollary 2.3.29 ([KZ24]). If $\mathcal{A}, \mathcal{B} \in \text{Alg}_{E_m}^c((n+1)\text{Vec})$ and $F : \mathcal{A} \rightarrow \mathcal{B}$ is $\mathbb{C}$ -linear E_m -monoidal, then $\mathfrak{Z}_m(F) = \mathfrak{Z}_m(\mathcal{A}, \mathcal{B}) \in \text{Alg}_{E_m}^c((n+1)\text{Vec})$.

Remark 2.3.30. The special cases of $n = 2$ and $m = 0, 1, 2, 3$ of Corollary 2.3.29 are rigorously proved recently in [Xu24a]. $\diamond$

Definition 2.3.31. We introduce the following notions (as working definitions for convenience³⁰).

1. An indecomposable E_1 -multi-fusion n -category $\mathcal{A}$ is called *non-degenerate* if $\mathfrak{Z}_1(\mathcal{A}) \simeq n\text{Vec}$. In this case, for $A \in \text{Alg}_{E_1}^c(\mathcal{A})$, A is called *Lagrangian* if $\text{Mod}_A^{E_1}(\mathcal{A}) \simeq n\text{Vec}$.
2. An E_2 -fusion n -category $\mathcal{B}$ is called *non-degenerate* if $\mathfrak{Z}_2(\mathcal{B}) \simeq n\text{Vec}$. In this case, for $B \in \text{Alg}_{E_2}^c(\mathcal{B})$, B is called *Lagrangian* if $\text{Mod}_B^{E_2}(\mathcal{B}) \simeq n\text{Vec}$.

Remark 2.3.32. An multi-fusion n -category $\mathcal{C}$ is non-degenerate if and only if $\mathcal{C}$ as a $\mathcal{C}$ - $\mathcal{C}$ -bimodule is closed, i.e., the canonical monoidal functor $\mathcal{C} \boxtimes \mathcal{C}^{\text{op}} \rightarrow \text{Fun}(\mathcal{C}, \mathcal{C})$ is an equivalence [KZ24]. The E_1 -center of a direct sum of two multi-fusion n -categories is the direct sum of the E_1 -centers of the summands. Moreover, the E_1 -center of an indecomposable multi-fusion n -category is an E_2 -fusion n -category. Therefore, a non-degenerate multi-fusion n -category is automatically an indecomposable multi-fusion n -category. $\diamond$

2.4 Center functor

The previous boundary-bulk relation can be generalized to include gapped domain walls. This more complete version of boundary-bulk relation can be formulated precisely in terms of the so-called center functor [KWZ15, KZ18a, KZ24]. In this subsection, we review the center functor and related results that are useful in later sections.

2.4.1 Complete boundary-bulk relation as a center functor

We have discussed the boundary-bulk relation in Section 2.1.3. It turns out that (2.1.19) is only a part of more complete boundary-bulk relation [KWZ15, KZ24], which is also very useful to our higher condensation theory. We review it now. Consider the physical configuration depicted in Figure 7, in which $A^{n+1}, B^{n+1}, C^{n+1}$ are potentially anomalous type-I topological orders (see Remark 2.4.4 for this ‘type-I’ assumption); M^n and N^n are gapped domain walls.

The orientation of the $Z(A)^{n+1}$ follows that in Figure 1 and that of A^{n+1} is the one induced from the bulk, and $Z(A)^{n+2}$ is on the left side of A^{n+1} if one follows the arrow on A^{n+1} , which is also the direction we fuse 1-codimensional defects in A^{n+1} . There are two different ways to look at M^n .

1. M^n is a gapped defect junction of A^{n+1} , B^{n+1} and $Z^{(1)}(M)^{n+1}$.

³⁰We will take a more systematic approach in [KZZ25] to define the notion of a Lagrangian E_m -algebra in a non-degenerate E_m -multi-fusion n -category.




Figure 7: the functoriality of the boundary-bulk relation

2. When M^n is viewed as gapped domain wall between A^{n+1} and B^{n+1} , it is potentially anomalous in the sense that the $n+1$ D topological order $Z^{(1)}(M)^{n+1}$, which can be viewed as the anomaly of M^{n+1} as a domain wall of A^{n+1} and B^{n+1} , can be non-trivial. According to [KW14, KWZ15], M^{n+1} uniquely determines $Z^{(1)}(M)^{n+1}$, called the *relative bulk* of M^n . This fact also explains the notation. Moreover, $Z^{(1)}(M)^{n+1}$ is also type-I. Note that the bulk $Z(M)^{n+1}$ of M^n is different from the relative bulk $Z^{(1)}(M)^{n+1}$ of M^n but can be expressed in terms of the later as follows:

$$Z(M)^{n+1} = \overline{B^{n+1}} \boxtimes_{Z(B)} Z^{(1)}(M)^{n+1} \boxtimes_{Z(A)} A^{n+1},$$

where $Z^{(1)}(M)^{n+1} \boxtimes_{Z(A)} A^{n+1}$ represents the fusion of $Z^{(1)}(M)^{n+1}$ and A^{n+1} along $Z(A)^{n+2}$.

Remark 2.4.1. In Figure 7, even when the defect junction N^n is gapless, its relative bulk $Z(N)^{n+1}$ is always gapped because any domain walls between two non-chiral topological orders $Z(A)^{n+2}$ and $Z(C)^{n+2}$ are necessarily gappable. $\diamond$

Theorem^{ph} 2.4.2 (Functoriality of Boundary-Bulk Relation [KWZ15]). The boundary-bulk relation, i.e., the assignment:

$$A^{n+1} \mapsto Z(A)^{n+1}, \quad M^n \mapsto Z^{(1)}(M)^{n+1},$$

gives a well-defined functor called *center functor* $Z : \text{TO}_{\text{gp-wall}}^{n+1} \rightarrow \text{afTO}_{\text{af-gp-wall}}^{n+2}$, where

- $\text{TO}_{\text{gp-wall}}^{n+1}$ is the 1-category³¹ of potentially anomalous $n+1$ D type-I topological orders as objects and potentially anomalous gapped domain walls as 1-morphisms;
- $\text{afTO}_{\text{af-gp-wall}}^{n+2}$ is the 1-category of anomaly-free simple $n+2$ D topological orders as objects and anomaly-free type-I gapped domain walls as 1-morphisms.

More precisely, in the physical situation depicted in Figure 7, we have an equivalence of type-I topological orders:

$$Z^{(1)}(N)^{n+1} \boxtimes_{Z(B)} Z^{(1)}(M)^{n+1} = Z^{(1)}(M \boxtimes_B N)^{n+1}, \quad (2.4.1)$$

where $\boxtimes_{Z(B)}$ means horizontally fuse $Z^{(1)}(M)^{n+1}$ with $Z^{(1)}(N)^{n+1}$ along $Z(B)^{n+2}$. $\square$

Remark 2.4.3. This functoriality or the equation (2.4.1) is a manifestation of a general principle of vanishing spatial fusion anomaly [KZ20, KYZ21] for 1-codimensional topological defects in an anomaly-free topological order. In the physical configuration depicted Fig. 7, with respect to the anomaly-free topological orders $Z(A)^{n+2}, Z(B)^{n+2}, Z(C)^{n+2}$, the domain walls $Z^{(1)}(L), Z^{(1)}(M), Z^{(1)}(N)$ are all 1-codimensional. This principle says that the fusion of the observables on two of such 1-codimensional defects (i.e., the left hand side of (2.4.1)) coincides with the observables on the fused defects (i.e., the right hand side of (2.4.1)). We believe that this principle applies to all quantum liquids and more general QFT's. We will provide the complete story of this principle elsewhere (see also Remark 4.2.11 and 6.2.2). $\diamond$

Remark 2.4.4. The functoriality of boundary-bulk relation for type-II topological orders is not natural (but possible by imposing some technical conditions). We do not need it. $\diamond$

³¹It is more natural to formulate the boundary-bulk relation as a functor between two higher categories. We only need this truncated version in this work. Some discussion about higher functor formulation can be found in [Kon13, DKR15, KWZ15].

Above result can be translated into a mathematical one. Note that the categories of topological defects $\mathcal{A}, \mathcal{B}, \mathcal{C}$ in the type-I topological orders $A^{n+1}, B^{n+1}, C^{n+1}$, respectively, are indecomposable multi-fusion n -categories. Let $\mathcal{M}$ be a $\mathcal{A}$ - $\mathcal{B}$ -bimodule and $\mathcal{N}$ a $\mathcal{B}$ - $\mathcal{C}$ -bimodule.

Theorem° 2.4.5 ([KWZ15, KZ18a, KZ24]). The following assignment (see Figure 7):

$$\mathcal{A} \mapsto \mathfrak{Z}_1(\mathcal{A}), \quad {}_{\mathcal{A}}\mathcal{M}_{\mathcal{B}} \mapsto \mathfrak{Z}_1^{(1)}(\mathcal{M}) := \text{Fun}_{\mathcal{A}|\mathcal{B}}(\mathcal{M}, \mathcal{M}) \quad (2.4.2)$$

defines the so-called *center functor* $\mathfrak{Z}_1 : \mathcal{MFus}_n^{\text{ind}} \rightarrow {}^{\text{ndg}}\mathcal{BFus}_n^{\text{cl}}$, where

- $\mathcal{MFus}_n^{\text{ind}}$ is the symmetric monoidal 1-category of indecomposable multi-fusion n -categories as objects and the equivalence classes of non-zero separable bimodules as 1-morphisms;
- ${}^{\text{ndg}}\mathcal{BFus}_n^{\text{cl}}$ is the symmetric monoidal 1-category of non-degenerate braided fusion n -categories as objects and the equivalence classes of closed multi-fusion $\mathcal{B}$ - $\mathcal{A}$ -modules as morphisms from $\mathcal{A}$ to $\mathcal{B}$.

In particular, this functoriality³² means that it respects the horizontal fusion:

$$\begin{aligned} \text{Fun}_{\mathcal{B}|\mathcal{C}}(\mathcal{N}, \mathcal{N}) \boxtimes_{\mathfrak{Z}_1(\mathcal{B})} \text{Fun}_{\mathcal{A}|\mathcal{B}}(\mathcal{M}, \mathcal{M}) &\xrightarrow{\simeq} \text{Fun}_{\mathcal{A}|\mathcal{C}}(\mathcal{M} \boxtimes_{\mathcal{B}} \mathcal{N}, \mathcal{M} \boxtimes_{\mathcal{B}} \mathcal{N}) \\ g \boxtimes_{\mathfrak{Z}_1(\mathcal{B})} f &\mapsto f \boxtimes_{\mathcal{B}} g. \end{aligned} \quad (2.4.3)$$

Remark 2.4.6. When $n = 1$, Theorem° 2.4.5 was rigorously proved in [KZ18a]. $\diamond$

It turns out that this center functor $\mathfrak{Z}_1(-)$ is a consequence of a simpler center functor $\mathfrak{Z}_0 : \text{Sep}_{n+1}^{\text{ind}} \rightarrow {}^{\text{ndg}}\mathcal{Fus}_{n+1}^{\text{cl}}$ [KZ24].

- The domain category $\text{Sep}_{n+1}^{\text{ind}}$ is the symmetric monoidal 1-category of indecomposable separable $(n+1)$ -categories, i.e., $\Sigma\mathcal{A}$ for an indecomposable multi-fusion n -category, as objects and the equivalence classes of non-zero $\mathbb{C}$ -linear functors as 1-morphisms.
- The codomain category ${}^{\text{ndg}}\mathcal{Fus}_{n+1}^{\text{cl}}$ is the symmetric monoidal 1-category of non-degenerate fusion $(n+1)$ -categories as objects and the equivalence classes of pairs $(\mathcal{M}, X)$ as morphisms from $\mathcal{A}$ to $\mathcal{B}$, where $\mathcal{M}$ is a closed $\mathcal{A}$ - $\mathcal{B}$ -bimodule and X is a non-zero object of $\mathcal{M}$. Two pairs $(\mathcal{M}, X)$ and $(\mathcal{N}, Y)$ are equivalent if there is a $\mathbb{C}$ -linear bimodule equivalence $F : \mathcal{M} \rightarrow \mathcal{N}$ such that $F(X) \simeq Y$.

The tensor product is given by $\boxtimes$. The center functor $\mathfrak{Z}_0 : \text{Sep}_{n+1}^{\text{ind}} \rightarrow {}^{\text{ndg}}\mathcal{Fus}_{n+1}^{\text{cl}}$ is defined by

$$\Sigma\mathcal{A} \mapsto \mathfrak{Z}_0(\Sigma\mathcal{A}) = \text{Fun}(\Sigma\mathcal{A}, \Sigma\mathcal{A}), \quad (\Sigma\mathcal{A} \xrightarrow{F} \Sigma\mathcal{B}) \mapsto \mathfrak{Z}_0^{(1)}(F) := (\text{Fun}(\Sigma\mathcal{A}, \Sigma\mathcal{B}), F), \quad (2.4.4)$$

and is symmetric monoidal. The center functor $\mathfrak{Z}_1$ is simply a consequence of the center functor $\mathfrak{Z}_0$ because the delooping functors

$$\Sigma : \mathcal{MFus}_n^{\text{ind}} \rightarrow \text{Sep}_n^{\text{ind}} \quad \text{and} \quad \Sigma : {}^{\text{ndg}}\mathcal{BFus}_n^{\text{cl}} \rightarrow {}^{\text{ndg}}\mathcal{Fus}_n^{\text{cl}} \quad (2.4.5)$$

are symmetric monoidal equivalences. More explicitly, we illustrate the relation between the $\mathfrak{Z}_0$ -functor and the $\mathfrak{Z}_1$ -functor in the following diagrams:

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{\mathfrak{Z}_1} & \mathfrak{Z}_1(\mathcal{A}) \\ \simeq \downarrow \Sigma & & \simeq \downarrow \Sigma \\ \Sigma\mathcal{A} & \xrightarrow{\mathfrak{Z}_0} & \mathfrak{Z}_0(\Sigma\mathcal{A}), \end{array} \quad \begin{array}{ccc} {}_{\mathcal{A}}\mathcal{M}_{\mathcal{B}} & \xrightarrow{\mathfrak{Z}_1^{(1)}} & \text{Fun}_{\mathcal{A}|\mathcal{B}}(\mathcal{M}, \mathcal{M}) \\ \downarrow & & \downarrow \\ F_{\mathcal{M}} & \xrightarrow{\mathfrak{Z}_0^{(1)}} & (\text{Fun}(\Sigma\mathcal{A}, \Sigma\mathcal{B}), F_{\mathcal{M}}), \end{array} \quad (2.4.6)$$

where $F_{\mathcal{M}} := - \boxtimes_{\mathcal{A}} \mathcal{M} \in \text{Fun}(\Sigma\mathcal{A}, \Sigma\mathcal{B})$.

³²This functoriality is stronger than the weak notion of a lax functor. This is a manifestation of an interesting physical principle, which will be discussed elsewhere.

2.4.2 Measure non-surjectivity and non-fully-faithfulness

It turns out that the deviations of the $\mathfrak{Z}_0$ -functor (or the $\mathfrak{Z}_1$ -functor) from being essentially surjective, full and faithful are very important in the study of topological orders. In the proof of Proposition* 3.13 in [KZ24], we actually proved that invertible separable higher categories precisely measure the non-surjectivity and non-fully-faithfulness of the center functor, a result which is a slightly refinement of the original statement of Proposition* 3.13 in [KZ24]. We state it more explicitly below.

Theorem° 2.4.7 ([KZ24]). We denote the full subcategory of $(n+1)\text{Vec}$ consisting of invertible objects (i.e., invertible separable n -categories) by $(n+1)\text{Vec}^\times$ and denote its underlying group by $\text{BrPic}(n\text{Vec})$.

- (1) $\mathfrak{Z}_0 : \text{Sep}_{n-1}^{\text{ind}} \rightarrow {}^{\text{ndg}}\mathcal{Fus}_{n-1}^{\text{cl}}$ is ‘essentially surjective’ up to invertible separable n -categories in the sense that
 - (a) $\mathcal{X} \in {}^{\text{ndg}}\mathcal{Fus}_{n-1}^{\text{cl}}$ if and only if $\Sigma\mathcal{X} \in (n+1)\text{Vec}^\times$;
 - (b) $\mathcal{X} \in \mathfrak{Z}_0(\text{Sep}_{n-1}^{\text{ind}})$ if and only if $\Sigma\mathcal{X} \simeq n\text{Vec} \in (n+1)\text{Vec}^\times$.
- (2) $\mathfrak{Z}_0 : \text{Sep}_n^{\text{ind}} \rightarrow {}^{\text{ndg}}\mathcal{Fus}_n^{\text{cl}}$ is ‘full’ up to invertible separable n -categories in the sense that, for any 1-morphism $\mathfrak{Z}_0(\mathcal{S}) \xrightarrow{(\mathcal{X}, \mathcal{X})} \mathfrak{Z}_0(\mathcal{S}')$ in ${}^{\text{ndg}}\mathcal{Fus}_n^{\text{cl}}$ and $\mathcal{S}, \mathcal{S}' \in \text{Sep}_n^{\text{ind}}$, there exists a $\mathcal{U} \in (n+1)\text{Vec}^\times$ and a $\mathbb{C}$ -linear functor $\mathcal{S} \xrightarrow{F} \mathcal{S}'$ such that $\mathcal{X} \simeq \mathfrak{Z}_0(F) \boxtimes \mathcal{U}$.
- (3) $\mathfrak{Z}_0 : \text{Sep}_{n+1}^{\text{ind}} \rightarrow {}^{\text{ndg}}\mathcal{Fus}_{n+1}^{\text{cl}}$ is ‘faithful’ up to invertible separable n -categories in the sense that any two 1-morphisms $F, G \in \text{Fun}(\mathcal{S}, \mathcal{S}')$ for $\mathcal{S}, \mathcal{S}' \in \text{Sep}_{n+1}^{\text{ind}}$ share the ‘same’ image, i.e., $\mathfrak{Z}_0^{(1)}(F) \simeq \mathfrak{Z}_0^{(1)}(G)$, if and only if there exists a $\mathcal{V} \in (n+1)\text{Vec}^\times$ such that $G(-) \simeq F(-) \boxtimes \mathcal{V}$.

The proof of Theorem 2.4.7 is identical to that of Proposition* 3.13 in [KZ24]. We do not repeat it here. Instead, we provide a physical proof of this result reformulated in terms of the $\mathfrak{Z}_1$ -functor below.

Theorem° 2.4.8 ([KZ24]). For readers convenience, we rewrite above theorem in terms of the $\mathfrak{Z}_1$ -functor based on the relation (2.4.6). See Figure 7 for the associated physical configuration.

- (1) $\mathfrak{Z}_1 : \mathcal{MFus}_{n-2}^{\text{ind}} \rightarrow {}^{\text{ndg}}\mathcal{BFus}_{n-2}^{\text{cl}}$ is ‘essentially surjective’ up to invertible separable n -categories, i.e.,
 - (a) $\mathcal{C} \in {}^{\text{ndg}}\mathcal{BFus}_{n-2}^{\text{cl}}$ if and only if $\Sigma^2\mathcal{C} \in (n+1)\text{Vec}^\times$;
 - (b) $\mathcal{C} \in \mathfrak{Z}_1(\mathcal{MFus}_{n-2}^{\text{ind}})$ if and only if $\Sigma^2\mathcal{C} \simeq n\text{Vec} \in (n+1)\text{Vec}^\times$.
- (2) $\mathfrak{Z}_1 : \mathcal{MFus}_{n-1}^{\text{ind}} \rightarrow {}^{\text{ndg}}\mathcal{BFus}_{n-1}^{\text{cl}}$ is ‘full’ up to invertible separable n -categories, i.e., for any 1-morphism $\mathcal{K} : \mathfrak{Z}_1(\mathcal{A}) \rightarrow \mathfrak{Z}_1(\mathcal{B})$ in ${}^{\text{ndg}}\mathcal{BFus}_{n-1}^{\text{cl}}$ and $\mathcal{A}, \mathcal{B} \in \mathcal{MFus}_{n-1}^{\text{ind}}$, there exists a $\mathcal{U} \in (n+1)\text{Vec}^\times$ and a non-zero $\mathcal{M} \in \text{BMod}_{\mathcal{A}|\mathcal{B}}((n+1)\text{Vec})$ such that $\Sigma\mathcal{K} \simeq \Sigma\mathfrak{Z}_1^{(1)}(\mathcal{M}) \boxtimes \mathcal{U}$.
- (3) $\mathfrak{Z}_1 : \mathcal{MFus}_n^{\text{ind}} \rightarrow {}^{\text{ndg}}\mathcal{BFus}_n^{\text{cl}}$ is ‘faithful’ up to invertible separable n -categories. That is, two 1-morphisms $\mathcal{A}\mathcal{M}_{\mathcal{B}}, \mathcal{A}\mathcal{N}_{\mathcal{B}}$ for $\mathcal{A}, \mathcal{B} \in \mathcal{MFus}_n^{\text{ind}}$ share the ‘same’ image, i.e., $\mathfrak{Z}_1^{(1)}(\mathcal{M}) \simeq \mathfrak{Z}_1^{(1)}(\mathcal{N})$, if and only if there exists an $\mathcal{V} \in (n+1)\text{Vec}^\times$ such that $\mathcal{N} \simeq \mathcal{M} \boxtimes \mathcal{V}$.

In the special case $\mathcal{A} = \mathcal{B} = n\text{Vec}$, two separable n -categories $\mathcal{M}$ and $\mathcal{N}$ share the same center, i.e., $\mathfrak{Z}_0(\mathcal{M}) \simeq \mathfrak{Z}_0(\mathcal{N})$, if and only if there exists a $\mathcal{V} \in (n+1)\text{Vec}^\times$ such that $\mathcal{N} \simeq \mathcal{M} \boxtimes \mathcal{V}$.

Proof. We provide a physical proof here. The proof of (1) is obvious. The proof of (2) is precisely given by that of Theorem^{ph} 2.2.26. Now we give a proof of (3). Look at the first picture in Figure 7. Note that stacking M^n with an anomaly-free nD topological order T^n followed by a condensation within $M^n \boxtimes T^n$ do not change anything around the defect junction. In particular, $Z^{(1)}(M)^{n+1}$ remains intact. Moreover, if a replacement of the junction M^n by a new junction N^n leave its neighborhood intact, then, by Theorem^{ph} 2.2.26, it is necessary that $\mathcal{N} \simeq \mathcal{M} \boxtimes \mathcal{V}$ for $\mathcal{V} \in (n+1)\text{Vec}^\times$. $\square$

Remark 2.4.9. The (2)-part of Theorem 2.4.8 first explicitly appeared (but formulated differently but equivalently) in Theorem^{ph} 3.2.13 in the first version of this paper (see also Theorem^{ph} 2.2.26 and 3.3.14 in this version). It also appeared later in the third version of [LYW24] and was formulated as in Theorem 2.2.42. $\diamond$

Corollary^o 2.4.10. This corollary is essentially the Example 3.14 in [KZ24]. There is no non-trivial invertible separable n -category only $n = 0, 1, 2$. For $n = 0, 1$, $\mathfrak{Z}_0 : \text{Sep}_n^{\text{ind}} \rightarrow \text{ndgFus}_n^{\text{cl}}$ is a symmetric monoidal equivalence; for $n = 2$, it is fully faithful; for $n = 3$, it is faithful but neither full nor essentially surjective; for $n > 3$, $\mathfrak{Z}_0$ is neither essentially surjective nor full nor faithful.

Remark 2.4.11. Theorem 2.4.5, 2.4.7 or 2.4.8 are important in that it can lead us to many new results. For example, for $\mathcal{A} \in \mathcal{MFus}_{n-1}^{\text{ind}}$, we have an exact sequence of group homomorphisms:


$$1 \rightarrow \text{BrPic}(n\text{Vec})_{\mathcal{A}} \hookrightarrow \text{BrPic}(n\text{Vec}) \xrightarrow{\mathcal{A} \boxtimes -} \text{BrPic}(\mathcal{A}) \xrightarrow{\mathfrak{Z}_1^{(1)}} \text{Aut}_0^{E_2}(\mathfrak{Z}_1(\mathcal{A})) \rightarrow 1.$$

- We denote the group of the invertible $\mathcal{A}$ - $\mathcal{A}$ -bimodules by $\text{BrPic}(\mathcal{A})$. We define $\text{BrPic}(n\text{Vec})_{\mathcal{A}} := \{\mathcal{V} \in \text{BrPic}(n\text{Vec}) \mid \mathcal{A} \boxtimes \mathcal{V} \simeq \mathcal{A}\}$ to be the stabilizer subgroup of $\text{BrPic}(n\text{Vec})$.
- Note that the group of the invertible objects in $\Sigma \mathfrak{Z}_1(\mathcal{A})$ is naturally embedded in that of the braided auto-equivalences of $\mathfrak{Z}_1(\mathcal{A})$. We denote its image by $\text{Aut}_0^{E_2}(\mathfrak{Z}_1(\mathcal{A}))$.

A detailed study of this topics and its relations to minimal modular extension (see [KZ24, Section 4.3]) will be given elsewhere. $\diamond$

The center functor $Z : \text{TO}_{\text{gp-wall}}^{n+1} \rightarrow \text{afTO}_{\text{af-gp-wall}}^{n+2}$ is neither full nor faithful in general. However, the center functor Z can be modified to give a full functor if we enlarge the domain category $\text{TO}_{\text{gp-wall}}^{n+1}$ to $\text{TO}_{\text{wall}}^{n+1}$ by allowing the 1-morphisms M^n and N^n to be gapless. Indeed, for a generic type-I gapped domain wall K^{n+1} between $Z(A)^{n+2}$ and $Z(B)^{n+2}$, the associated defect junction M^n is gapless in general (see the first picture in (2.4.7)).

- According to [KZ18b, KZ20, KZ21], M^n is holographically dual (via topological Wick rotation) to a pair (T^{n+1}, X^n) , where T^{n+1} is an anomaly-free $n+1$ D simple topological order and X^n is a gapped defect junction among $A^{n+1} \boxtimes T^{n+1}$, K^{n+1} and B^{n+1} , as illustrated below.



Note that the gapped wall K^{n+1} is necessarily Morita equivalent to $\overline{A^{n+1}} \boxtimes \overline{T^{n+1}} \boxtimes B^{n+1}$ for an anomaly-free T^{n+1} .

- Alternatively, using Theorem^{ph} 2.2.26, one can always find the anomaly-free topological order T^{n+1} , which is unique up to Morita equivalences, such that K^{n+1} is Morita equivalent to $\overline{A^{n+1}} \boxtimes \overline{T^{n+1}} \boxtimes B^{n+1}$.

This result, together with Theorem 2.4.7 and 2.4.8, leads us to the following result, which is useful in later sections.

Theorem^{ph} 2.4.12. The map $Z^{(1)} : X^n \mapsto Z^{(1)}(X)^{n+1}$ (or equivalently, ${}_{\mathcal{A} \boxtimes \mathcal{T}} \mathcal{X}_{\mathcal{B}} \mapsto \text{Fun}_{\mathcal{A} \boxtimes \mathcal{T} | \mathcal{B}}(\mathcal{X}, \mathcal{X})$) defines a one-to-one correspondence between:

- (1) $(\mathcal{A} \boxtimes \mathcal{T})$ - $\mathcal{B}$ -bimodules modulo the equivalence relation: $\mathcal{X} \sim \mathcal{Y}$ if $\mathcal{Y} \simeq \mathcal{X} \boxtimes \mathcal{V}$ for $\mathcal{V} \in (n+1)\text{Vec}^\times$;
- (2) type-I gapped walls between $Z(A)^{n+2}$ and $Z(B)^{n+2}$ in the Morita class $[\overline{A^{n+1}} \boxtimes \overline{T^{n+1}} \boxtimes B^{n+1}]$.

The wall is simple if and only if the associated $(\mathcal{A} \boxtimes \mathcal{T})$ - $\mathcal{B}$ -bimodule is indecomposable. When $B^{n+1} = \mathbf{1}^{n+1}$, the slightly modified map³³

$$X^n \mapsto \overline{Z^{(1)}(X)^{n+1}} \quad \text{or equivalently} \quad {}_{\mathcal{A} \boxtimes \mathcal{T}} \mathcal{X} \mapsto \text{Fun}_{\mathcal{A} \boxtimes \mathcal{T}}(\mathcal{X}, \mathcal{X})^{\text{op}} \quad (2.4.8)$$

defines a one-to-one correspondence between left $(\mathcal{A} \boxtimes \mathcal{T})$ -modules modulo the equivalence relation $\sim$ and gapped boundaries of $Z(A)^{n+2}$ in the Morita class $[A^{n+1} \boxtimes T^{n+1}]$. $\square$

³³Flipping the orientation of $Z^{(1)}(X)^{n+1}$ is necessary for the boundary to have the orientation induced from that of the bulk.

Remark 2.4.13. Note that the $n = 1$ case of Theorem^{ph} 2.4.12 recovers the well-known one-to-one correspondence between left indecomposable $\mathcal{A}$ -modules and simple gapped boundaries of the 3D non-chiral simple topological order $Z(\mathcal{A})^3$ [KK12, DMNO13, Kon14]. For $n \geq 1$, it provides a powerful tool to classify gapped boundaries of $Z(\mathcal{A})^{n+2}$ (or Lagrangian E_1 - or E_2 -algebras, see Corollary 3.3.39 and Corollary 4.2.23). $\diamond$

Remark 2.4.14. Note that it is possible that $\mathcal{X} \simeq \mathcal{X} \boxtimes \mathcal{V}$ for $\mathcal{V} \in (n+1)\text{Vec}^\times$. For example, when $\mathcal{A} = \mathcal{T} = \mathcal{B} = n\text{Vec}$, consider $\mathcal{V} \in (n+1)\text{Vec}^\times$ such that $\mathcal{V}^{\boxtimes m} \simeq n\text{Vec}$ (e.g., $(\Sigma\mathcal{I}s)^{\boxtimes 16} \simeq 3\text{Vec}$). We set $\mathcal{X} := \oplus_{i=0}^{m-1} \mathcal{V}^{\boxtimes i}$. Then we have $\mathcal{X} \boxtimes \mathcal{V} \simeq \mathcal{X}$. $\diamond$

Corollary^{ph} 2.4.15. When $\mathcal{A}$ is a fusion n -category and is connected as a separable n -category, i.e., $\mathcal{A} = \Sigma\Omega\mathcal{A}$, there is a one-to-one correspondence between

- (1) braided monoidal functors $\Omega\mathcal{A} \rightarrow \mathcal{O}$ for a non-degenerate braided fusion $(n-1)$ -category $\mathcal{O}$;
- (2) simple gapped boundaries of $Z(\mathcal{A})^{n+2}$.

Moreover, the Morita class of the boundary is precisely the Morita class $[\mathcal{A} \boxtimes \Sigma\mathcal{O}^{\text{op}}]$.

Proof. Let $\mathcal{T}$ be a non-degenerate fusion n -category $\mathcal{T}$. An indecomposable left $\mathcal{A} \boxtimes \mathcal{T}$ -module is again indecomposable as a separable n -category, i.e., $\Sigma\mathcal{Q}$ for a random choice of fusion $(n-1)$ -category $\mathcal{Q}$ within the same Morita class. Therefore, an indecomposable left $\mathcal{A} \boxtimes \mathcal{T}$ -module $\Sigma\mathcal{Q}$, which is defined by a monoidal functor $\mathcal{A} \boxtimes \mathcal{T} \rightarrow \mathfrak{Z}_0(\Sigma\mathcal{Q}) \simeq \Sigma\mathfrak{Z}_1(\mathcal{Q})$, determines a braided monoidal functor $\Omega\mathcal{A} \boxtimes \Omega\mathcal{T} \rightarrow \mathfrak{Z}_1(\mathcal{Q})$, which further induces a braided monoidal functor $\Omega\mathcal{A} \rightarrow \mathfrak{Z}_2(\Omega\mathcal{T}, \mathfrak{Z}_1(\mathcal{Q}))$ by the universal property of the E_2 -centralizer, where $\mathfrak{Z}_2(\Omega\mathcal{T}, \mathfrak{Z}_1(\mathcal{Q}))$ is non-degenerate as a braided fusion $(n-1)$ -category.

Conversely, given a braided monoidal functor $f : \Omega\mathcal{A} \rightarrow \mathcal{O}$ for a non-degenerate braided fusion $(n-1)$ -category $\mathcal{O}$ within the Witt class $[\Omega\mathcal{T}]^{-1}$, we obtain a monoidal functor

$$\mathcal{A} \boxtimes \mathcal{T} \xrightarrow{\Sigma f \boxtimes 1} \Sigma\mathcal{O} \boxtimes \mathcal{T} \simeq \mathfrak{Z}_0(\mathcal{M}), \quad \text{for some } \mathcal{M} \in (n+1)\text{Vec},$$

which further defines an indecomposable $\mathcal{A} \boxtimes \mathcal{T}$ -module $\mathcal{M}$ precisely up to the equivalence relation $\sim$ by Theorem 2.4.8 (3). Moreover, we have an equality of the Morita class $[\mathcal{A} \boxtimes \mathcal{T}] = [\mathcal{A} \boxtimes \Sigma\mathcal{O}^{\text{op}}]$. $\square$

Remark 2.4.16. Although we do have an explicit formula (recall (2.4.8)) for the gapped boundary associated to the braided monoidal functor $f : \Omega\mathcal{A} \rightarrow \mathcal{O}$, it does not reveal the meaning of f directly. When we identify each simple gapped boundary with a Lagrangian E_2 -algebra (see Corollary^{ph} 3.3.40), we see that the braided monoidal functor $f : \Omega\mathcal{A} \rightarrow \mathcal{O}$ precisely defines a Lagrangian E_2 -algebra in $\mathfrak{Z}_1(\mathcal{A})$ because we have $\mathfrak{Z}_1(\mathcal{A}) \simeq \mathfrak{Z}_1(\Sigma\Omega\mathcal{A}) \simeq \text{Mod}_{\Omega\mathcal{A}}^{E_2}(n\text{Vec})^{\text{op}}$ (see (4.2.13)). $\diamond$

2.4.3 Generalized center functors

In the center functor defined in Theorem^{ph} 2.4.2, the image of the functor are all non-chiral topological orders in the bulk. In order to include all chiral topological orders, we can either allow the boundary to be gapless, or somewhat ‘equivalently’, consider a relative version of the center functor.




Figure 8: a relative version of boundary-bulk relation

Consider the physical configuration depicted in Figure 8 (a). We use $Z^{(P)A}^{n+2}$ to denote the bulk of A^{n+1} relative to P^{n+2} . The notation $^P A$ does not have any extra meaning other than the pair (P, A) . However, this

notation has some advantages in addition to being short. It reminds us that P^{n+2} and A^{n+1} have different dimensions and P^{n+2} acts on A^{n+1} in some sense. Similarly, the notation ${}^{\Omega\mathcal{P}}\mathcal{A}$ also does not have any extra meaning other than the pair $(\Omega\mathcal{P}, \mathcal{A})$. Although it might cause confusion because we also use ${}^{\Omega\mathcal{P}}\mathcal{A}$ to represent an $\Omega\mathcal{P}$ -enriched category only in Section 6.3.2, this confusion is harmless and is guaranteed to give the same result by the idea of topological Wick rotation [KZ18b, KZ20, KZ21, KZ22b].

Theorem^{ph} 2.4.17. For type-I topological orders A^{n+1} , the assignment $A^{n+1} \mapsto Z(PA)^{n+2}$ and $M^n \mapsto Z^{(1)}(PM)$ is functorial in the sense that the following identity holds.

$$Z^{(1)}(PN)^{n+1} \boxtimes_{Z(PB)} Z^{(1)}(PM)^{n+1} = Z^{(1)}(P(M \boxtimes_B N))^{n+1}.$$

Mathematically, this amounts to the following assignment (see Figure 8 (b)):

$$\mathcal{A} \mapsto \mathfrak{Z}_1({}^{\Omega\mathcal{P}}\mathcal{A}) := \mathfrak{Z}_2(\Omega\mathcal{P}^{\text{op}}, \mathfrak{Z}_1(\mathcal{A})), \quad \mathcal{M} \mapsto \mathfrak{Z}_1^{(1)}({}^{\Omega\mathcal{P}}\mathcal{M}) := \text{Fun}_{\mathcal{A} \boxtimes_{\Omega\mathcal{P}} \mathcal{B}^{\text{op}}}(\mathcal{M}, \mathcal{M}),$$

which defines a relative version of the center functor, well-defined due to the monoidal equivalence:

$$\text{Fun}_{\mathcal{B} \boxtimes_{\Omega\mathcal{P}} \mathcal{C}^{\text{op}}}(\mathcal{N}, \mathcal{N}) \boxtimes_{\mathfrak{Z}_2(\Omega\mathcal{P}^{\text{op}}, \mathfrak{Z}_1(\mathcal{B}))} \text{Fun}_{\mathcal{A} \boxtimes_{\Omega\mathcal{P}} \mathcal{B}^{\text{op}}}(\mathcal{M}, \mathcal{M}) \simeq \text{Fun}_{\mathcal{A} \boxtimes_{\Omega\mathcal{P}} \mathcal{C}^{\text{op}}}(\mathcal{M} \boxtimes_{\mathcal{B}} \mathcal{N}, \mathcal{M} \boxtimes_{\mathcal{B}} \mathcal{N}).$$

We have not specified the (co-)domain category of this functor. We leave it to intrigued readers. $\square$

It is possible to generalize Theorem 2.4.7 or Theorem 2.4.8 to the generalized center functor defined in Theorem^{ph} 2.4.17. For the purpose of studying condensations in this work, however, we do not gain any truly new result because the folding trick reduces the problem of constructing or classifying gapped domain walls (or 1-codimensional defect condensations in later sections) between two Morita equivalent chiral topological orders to the problem of constructing or classifying gapped boundaries of a non-chiral topological order. More precisely, applying the folding trick to the physical configuration in the first picture in Figure 9, followed by a topological Wick rotation (TWR), we obtain a reformulation of Theorem^{ph} 2.4.12 for chiral topological orders as summarized below.




Figure 9: reduce a wall to a boundary by the folding trick followed by a TWR

Theorem^{ph} 2.4.18. For an anomaly-free simple $n+1$ D topological order T^{n+1} , there is a one-to-one correspondence between

- (1) left $((\mathcal{A} \boxtimes_{\Omega\mathcal{P}} \mathcal{B}^{\text{op}}) \boxtimes \mathcal{T})$ -modules modulo the equivalence relation: $\mathcal{X} \sim \mathcal{Y}$ if $\mathcal{Y} \simeq \mathcal{X} \boxtimes \mathcal{V}$ for $\mathcal{V} \in (n+1)\text{Vec}^\times$;
- (2) type-I gapped walls between two (potentially chiral) simple topological orders $Z(PA)^{n+2}$ and $Z(PB)^{n+2}$ in the Morita class $[(B \boxtimes_P A)^{n+1} \boxtimes \overline{T}^{n+1}]$

defined by $\mathcal{X} \mapsto \text{Fun}_{\mathcal{A} \boxtimes_{\Omega\mathcal{P}} \mathcal{B}^{\text{op}} \boxtimes \mathcal{T}}(\mathcal{X}, \mathcal{X})$. The gapped wall is simple, i.e., $\text{Fun}_{\mathcal{A} \boxtimes_{\Omega\mathcal{P}} \mathcal{B}^{\text{op}}}(\mathcal{X}, \mathcal{X})$ is a fusion n -category, if and only if the left $((\mathcal{A} \boxtimes_{\Omega\mathcal{P}} \mathcal{B}^{\text{op}}) \boxtimes \mathcal{T})$ -module $\mathcal{X}$ is indecomposable. $\square$

Remark 2.4.19. We show in Section 3.3.2 that a type-I gapped domain wall between two anomaly-free topological orders C^{n+2} and D^{n+2} is precisely an indecomposable condensable E_1 -algebra in $\mathcal{C}$, which defines a 1-codimensional defect condensation from C^{n+2} to D^{n+2} . Therefore, Theorem^{ph} 2.4.18 provides a powerful tool to classify 1-codimensional defect condensations from C^{n+2} to D^{n+2} . $\diamond$

Remark 2.4.20. In applications, one can choose a convenient P^{n+2} in order to simplify the calculation. For example, if we already know an example of gapped domain wall K^{n+1} between $C^{n+2} = Z(PA)^{n+2}$ and $D^{n+2} = Z(PB)^{n+2}$. We can choose $P^{n+2} = C^{n+2}$, $A^{n+1} = C^{n+1}$ and $B^{n+1} = K^{n+1}$. Then we obtain a one-to-one correspondence, defined by $\mathcal{X} \mapsto \text{Fun}_{\mathcal{K}^{\text{op}} \boxtimes \mathcal{T}}(\mathcal{X}, \mathcal{X})$, between

- (1) left $(\mathcal{K}^{\text{op}} \boxtimes \mathcal{T})$ -modules modulo the equivalence relation: $\mathcal{X} \sim \mathcal{Y}$ if $\mathcal{Y} \simeq \mathcal{X} \boxtimes \mathcal{V}$ for $\mathcal{V} \in (n+1)\text{Vec}^\times$;
- (2) $\frac{\text{type-I}}{\mathbb{T}^{n+1}}$ gapped walls between two (potentially chiral) simple C^{n+2} and D^{n+2} in the Morita class $[K^{n+1} \boxtimes \mathbb{T}^{n+1}]$. $\diamond$

Remark 2.4.21. Theorem^{ph} 2.4.18 and Remark 2.4.20 provide a powerful tool to construct and classify of 1-codimensional defect condensations between two chiral topological orders in higher dimensions. Unfortunately, we are not able to provide any concrete applications of this result in $n+1$ D for $n > 3$ in this work because we do not know the categorical description of any non-trivial chiral topological orders in $n+1$ D for $n > 3$. $\diamond$

Remark 2.4.22. Theorem^{ph} 2.4.2 is enough for the purpose of including chiral topological orders in the image of the center functor. However, it is even possible to further enlarge both the domain category and the codomain category to obtain an even more general center functor by allowing P^{n+2} in Figure 9 to vary for different A^{n+1} , B^{n+1} and M^n or even slightly beyond that. This further generalization is essentially equivalent to allowing A^{n+1} , B^{n+1} and M^n to be gapless. In [KZ21, Theorem 4.15], we introduced this generalized functor for $n = 2$, which can be generalized to higher dimensions tautologically by restricting to only the topological skeletons [KZ22b]. In particular, the illustrating picture of this functor in [KZ21, Figure 10] works for all dimensions tautologically. Moreover, it is even possible to further generalize the center functor such that the bulk phases $Z(A)^{n+2}$ and $Z(B)^{n+2}$ are also gapless [KYZ21, KZZZ25]. We do not need these generalized center functors in this work. We will come back to this issue elsewhere. $\diamond$

3 Condensations of 1-codimensional topological defects

In this section, we study the condensations of 1-codimensional topological defects. We start from reviewing the theory of the particle condensations in 1+1D (potentially anomalous) topological orders via an algebraic bootstrap approach [Kon14] in Section 3.1; then we redo it via a geometric approach [KS11a, FSV13] which is ready to be generalized to higher dimensional topological orders in Section 3.2; then we provide the general theory of the condensations of 1-codimensional topological defects for topological orders in all dimensions in Section 3.3.

3.1 Particle condensations in 1+1D: algebraic approach

In this subsection, we briefly review the bootstrap arguments in [Kon14] that lead to the particle condensation theory for potentially anomalous 1+1D topological orders.

3.1.1 Algebraic bootstrap

Consider a 1+1D (potentially anomalous) simple topological order C^2 as depicted in only the spatial dimension in Figure 10. Its particles form a fusion 1-category $\mathcal{C}$ (recall Theorem^{ph} 2.2.37). Now we consider a particle condensation happening in a large but connected region and producing a new 1+1D topological order D^2 , whose particles form a new fusion 1-category $\mathcal{D}$ (see Figure 10), and two gapped domain wall M^1 and $\overline{M}^1$, where $\overline{M}^1$ can be viewed as the time reversal of M . The bulk $Z(C)^2$ remains unchanged.




Figure 10: a condensation in the topological order C^2

Since D^2 is obtained from a condensation of C^2 , particles in $\mathcal{D}$ necessarily come from those in $\mathcal{C}$, i.e. $\text{ob}(\mathcal{D}) \subset \text{ob}(\mathcal{C})$. The hom space $\text{hom}_{\mathcal{C}}(a, b)$ for $a, b \in \mathcal{C}$ encodes the information of how many independent channels of fusing or splitting a into b . It is clear that the fusion-splitting channels between particles in $\mathcal{D}$ necessarily come from those between particles in $\mathcal{C}$, i.e. $\text{hom}_{\mathcal{D}}(a, b) \subset \text{hom}_{\mathcal{C}}(a, b)$. As a consequence, $\mathcal{D}$ is necessarily a sub-category of $\mathcal{C}$, i.e. $\mathcal{D} \subset \mathcal{C}$. In particular, the vacuum or the trivial particle in $\mathcal{D}$ is necessarily a (composite) particle $A \in \mathcal{C}$.

A particle condensation is triggered by introducing interactions among particles. For example, for $a, b \in \mathcal{D} \subset \mathcal{C}$, the consequence of introducing interactions between particles is that a sub-Hilbert space of the Hilbert space associated to $a \otimes_{\mathcal{C}} b$ (or simply $a \otimes b$ if no confusion is caused) becomes energy favorable. This sub-Hilbert space gives a physical definition of the fusion product $a \otimes_{\mathcal{D}} b$ in the condensed phase D^2 . In other words, the condensation process produces a family of projections (called condensation maps):

$$a \otimes_{\mathcal{C}} b \xrightarrow{p_{a,b}} a \otimes_{\mathcal{D}} b, \quad \forall a, b \in \mathcal{D}.$$

In particular, we have the following condensation maps:

$$\begin{aligned} \mu_A : A \otimes_{\mathcal{C}} A &\xrightarrow{p_{AA}} A \otimes_{\mathcal{D}} A \simeq A. \\ \forall x \in \mathcal{D}, \quad \mu_x^L : A \otimes_{\mathcal{C}} x &\xrightarrow{p_{Ax}} A \otimes_{\mathcal{D}} x \simeq x, \quad \mu_x^R : x \otimes_{\mathcal{C}} A \xrightarrow{p_{xA}} x \otimes_{\mathcal{D}} A \simeq x. \end{aligned}$$

Moreover, since the vacuum in $\mathcal{C}$ should condense into the vacuum in $\mathcal{D}$, we expect to have a morphism $\eta_A : \mathbb{1}_{\mathcal{C}} \rightarrow A$. The triple (A, μ_A, η_A) defines an algebra or an E_1 -algebra in $\mathcal{C}$ and the triple (x, μ_x^L, μ_x^R) defines an A - A -bimodule in $\mathcal{C}$. Moreover, μ_A splits as an A - A -bimodule map due to a natural condition on the stability of the vacuum of the condensed phase D^2 [Kon14]. Namely, there exists an A - A -bimodule map $e_A : A \rightarrow A \otimes_{\mathcal{C}} A$ such that $\mu_A \circ e_A = \text{id}_A$, which is the defining condition of a condensable (or separable) E_1 -algebra

in $\mathcal{C}$. In the unitary case, e_A can be chosen to be $\mu_A^\dagger$ and A is naturally a special $\dagger$ -Frobenius algebra³⁴. If $\dim \text{hom}_{\mathcal{C}}(\mathbb{1}_{\mathcal{C}}, A) = 1$, the condensable E_1 -algebra A is called simple. If a condensable E_1 -algebra A cannot be decomposed into a direct sum of algebras, then A is called indecomposable. In general, a condensable E_1 -algebra A in $\mathcal{C}$ is a direct sum of indecomposable ones. We postpone the structure theorem of a condensable E_1 -algebra to Section 3.1.2 after we review the notion of an internal hom.

Remark 3.1.1. As we explained in Section 2.3.1, $\mathcal{C} \in \text{Alg}_{E_1}(\text{2Vec})$ and there are infinite number of tensor products $\otimes_{(r,s)}$ on $\mathcal{C}$, which can be reduced to a single fusion product $\otimes_{\mathcal{C}} = \otimes$ (and $\otimes^{\text{op}}$) due to the Adiabatic Principle (2.3.1). Therefore, there is also infinitely many multiplication maps $A \otimes_{(r,s)} A \rightarrow A$, which endows A with the structure of an E_1 -algebra in $\mathcal{C}$. In this case, it turns out that an E_1 -algebra in $\mathcal{C}$ is equivalent to an associative algebra in $\mathcal{C}$. We use $\text{Alg}_{E_1}^{\mathcal{C}}(\mathcal{C})$ to denote the category of condensable E_1 -algebras in $\mathcal{C}$.

Similarly, the infinitely many A -action on $x \in \mathcal{D}$: $x \otimes_{(r,s)} A \rightarrow x$ endows A with a structure of an E_1 - A -module. In this case, it turns out that an E_1 - A -module in $\mathcal{C}$ is equivalent to an A - A -bimodule in $\mathcal{C}$, and an E_1 - A -module map is precisely an A - A -bimodule map. We denote the 1-category of E_1 - A -modules (as objects) and E_1 - A -module maps (as 1-morphisms) by $\text{Mod}_A^{E_1}(\mathcal{C})$. We have a canonical equivalence $\text{Mod}_A^{E_1}(\mathcal{C}) \simeq \text{BMod}_{A|A}(\mathcal{C})$. For simplicity, we simply set $\text{Mod}_A^{E_1}(\mathcal{C}) = \text{BMod}_{A|A}(\mathcal{C})$. $\diamond$

The following bootstrap result was obtained in [Kon14] under the name of ‘1-d condensation’.

Theorem^{ph} 3.1.2. Using some obvious physical intuitions and certain stability arguments, one can easily derive the following results.

- (1) The triple (A, μ_A, η_A) defines a condensable E_1 -algebra in $\mathcal{C}$.
- (2) For $x \in \mathcal{D}$, the triple (x, μ_x^L, μ_x^R) defines an A - A -bimodule in $\mathcal{C}$.
- (3) $\mathcal{D} = \text{Mod}_A^{E_1}(\mathcal{C})$, where $\text{Mod}_A^{E_1}(\mathcal{C}) = \text{BMod}_{A|A}(\mathcal{C})$ is the category of A - A -bimodules or E_1 - A -modules in $\mathcal{C}$ (as objects) and E_1 - A -module maps (as 1-morphisms). The fusion product $\otimes_{\mathcal{D}}$ can be identified with the relative tensor product $\otimes_A$, i.e. $\otimes_{\mathcal{D}} = \otimes_A$, and $\mathbb{1}_{\mathcal{D}} = A$. If $A = \mathbb{1}_{\mathcal{C}}$, then $\mathcal{D} = \mathcal{C}$ and the condensation is called a trivial condensation.

The gapped domain wall M^1 can also support particles. These particles form a separable 1-category $\mathcal{M}$. Since particles in $\mathcal{C}$ can fuse onto the wall from left and particles in $\mathcal{D}$ can fuse onto the wall from right, $\mathcal{M}$ is necessarily a left $\mathcal{C}$ -module and a right $\mathcal{D}$ -module. More precisely, a left $\mathcal{C}$ -module is a category $\mathcal{M}$ equipped with a left $\mathcal{C}$ -action functor $\odot : \mathcal{C} \times \mathcal{M} \rightarrow \mathcal{M}$ that is unital and associative, i.e. there exist the following natural isomorphisms:

$$\mathbb{1}_{\mathcal{C}} \odot x \simeq x, \quad a \odot (b \odot x) \simeq (a \otimes_{\mathcal{C}} b) \odot x, \quad \forall a, b \in \mathcal{C}, x \in \mathcal{M},$$

where we have used the following notation $\odot(a, x) := a \odot x$, satisfying some physically natural conditions (see [EGNO15] for a review). A right $\mathcal{D}$ -module is similar. Moreover, the two side actions on $\mathcal{M}$ are obviously commutative. This means that there are natural isomorphisms

$$a \odot (x \odot d) \simeq (a \odot x) \odot d, \quad \forall a \in \mathcal{C}, x \in \mathcal{M}, d \in \mathcal{D},$$

satisfying some natural conditions. These data and conditions are the defining data and axioms of a $\mathcal{C}$ - $\mathcal{D}$ -bimodule category. In summary, $\mathcal{M}$ is a $\mathcal{C}$ - $\mathcal{D}$ -bimodule in 2Vec . However, $\mathcal{M}$ does not encode the complete physical information associated to the wall M^1 . The physical domain wall M^1 must specify a particle $m \in \mathcal{M}$ (as a *wall condition*³⁵) such that fusing particles in $\mathcal{C}$ (or $\mathcal{D}$) onto the wall gives a well defined map $\mathcal{C} \rightarrow \mathcal{M}$ defined by $a \mapsto a \odot m$ for $a \in \mathcal{C}$, and a well define map $\mathcal{D} \rightarrow \mathcal{M}$ defined by $d \mapsto m \odot d$ for $d \in \mathcal{D}$. Therefore, the physical domain wall M^1 can be mathematically described by a pair $(\mathcal{M}, m)$. Such a pair defines a so-called E_0 -algebra in the symmetric monoidal 2-category 2Vec [Lur17]. A morphism between two E_0 -algebras $(\mathcal{M}, m)$ and $(\mathcal{M}', m')$ is a functor $f : \mathcal{M} \rightarrow \mathcal{M}'$, together with a morphism $g : f(m) \rightarrow m'$. We denote the category of E_0 -algebras in 2Vec by $\text{Alg}_{E_0}(\text{2Vec})$.

Remark 3.1.3. Similar to Remark 3.1.1, there are infinitely many $\mathcal{C}$ -action on $\mathcal{M}$ given by $\odot_{(r,s)} : \mathcal{C} \times \mathcal{M} \rightarrow \mathcal{M}$ for $r, s \in \mathbb{R}$ and $r \neq s$. When $\mathcal{C} = \mathcal{D}$, the category $\mathcal{M}$ is naturally equipped with the structure of an E_1 - $\mathcal{C}$ -module, i.e. $\mathcal{M} \in \text{Mod}_{\mathcal{C}}^{E_1}(\text{2Vec}) = \text{BMod}_{\mathcal{C}|\mathcal{C}}(\text{2Vec})$. $\diamond$

³⁴A special $\dagger$ -Frobenius algebra A in $\mathcal{C}$ is an algebra in $\mathcal{C}$ defined by the triple (A, μ, η) and a coalgebra defined by the triple (A, Δ, ϵ) such that $(\mu \otimes_{\mathcal{C}} \text{id}_A) \circ (\text{id}_A \otimes_{\mathcal{C}} \Delta) = \Delta \circ \mu = (\text{id}_A \otimes_{\mathcal{C}} \mu) \circ (\Delta \otimes_{\mathcal{C}} \text{id}_A)$, $\mu \circ \Delta = \text{id}_A$, $\Delta = \mu^\dagger$ and $\epsilon = \eta^\dagger$ (see for example [FRS02, Kon14]).

³⁵In other words, $\mathcal{M}$ is also the category of all wall conditions.

Since the domain wall M^1 is also created from a particle condensation of $\mathcal{C}^2$, the particles and the fusion-splitting channels in $\mathcal{M}$ should all come from $\mathcal{C}$. In other words, $\mathcal{M}$ must be a subcategory of $\mathcal{C}$. Moreover, the condensation again produces a condensation map $x \otimes_{\mathcal{C}} A \rightarrow x \odot A \simeq x$, which clearly defines a right A -module structure on x . Note that an A - A -module in $\mathcal{C}$ is clearly a deconfined particle living in $\mathcal{D}$, while a right A -module in $\mathcal{C}$ defines a particle that is necessarily confined to the domain wall. Therefore, we obtain $\mathcal{M} = \text{RMod}_A(\mathcal{C})$, where $\text{RMod}_A(\mathcal{C})$ denotes the category of right A -modules in $\mathcal{C}$ as objects and intertwiners (of the A -action) as morphisms. As a consequence, we have $a \odot x := a \otimes_{\mathcal{C}} x$ and $x \odot d := x \otimes_A d$ for $x \in \mathcal{M}$, $a \in \mathcal{C}$ and $d \in \mathcal{D}$. If we do not specify any particle in the condensed phase in $\mathcal{D}$, any site in the $\mathcal{D}^2$ -phase naturally has the trivial particle $\mathbb{1}_{\mathcal{D}} = A$ on it. Similarly, if we do not specify a particle on the wall M^1 , then the trivial particle A naturally lives on it. Therefore, we should have $M^1 = (\text{RMod}_A(\mathcal{C}), A)$.

For the other wall $\overline{M}^1$, by the similar reason, we should expect that the category of particles on this wall is given by $\mathcal{M}^{\text{op}} = \text{LMod}_A(\mathcal{C})$. As a $\mathcal{D}$ - $\mathcal{C}$ -bimodule $\mathcal{M}^{\text{op}}$, the action is defined by $d \odot m \odot c := c^* \odot m \odot d^*$, where we have used convention $d^* = d^L = d^R$ as in the unitary case to avoid framing issues.

We summarize the result below.

Theorem^{ph} 3.1.4 ([Kon14]). The gapped domain wall M can be mathematically characterized by the pair $(\mathcal{M}, A)$, where $\mathcal{M} = \text{RMod}_A(\mathcal{C})$ and $m \in \mathcal{M}$.

- (1) Particles in $\mathcal{C}$ move to the wall M according to the map $a \mapsto a \otimes_{\mathcal{C}} A, \forall a \in \mathcal{C}$;
- (2) particles in $\mathcal{D}$ move to the wall M according to the map $d \mapsto A \otimes_A d = d, \forall d \in \mathcal{D}$.

We also have $\overline{M} = (\mathcal{M}^{\text{op}}, A)$, where $\mathcal{M}^{\text{op}} \simeq \text{LMod}_A(\mathcal{C})$. □

3.1.2 Internal homs and structure theorem

These mathematical precise results rely heavily on an important categorical notion called an *internal hom*, which is very important to the condensation theory and to all QFTs as well (see [Dav10a, DKR15, KYZ21] and Remark 3.1.10). For this reason, we first review this notion (see more details in the book [EGNO15]).³⁶

Definition 3.1.5 ([Ost03a]). Let $\mathcal{C}$ be a multi-fusion 1-category and $\mathcal{M}$ be a separable left $\mathcal{C}$ -module³⁷, i.e. a category equipped with a (unital and associative) $\mathcal{C}$ -action $\odot : \mathcal{C} \times \mathcal{M} \rightarrow \mathcal{M}$. For $x, y \in \mathcal{M}$, the internal hom $[x, y]_{\mathcal{C}}$ is an object uniquely determined by the following isomorphisms:

$$\text{hom}_{\mathcal{M}}(a \odot x, y) \simeq \text{hom}_{\mathcal{C}}(a, [x, y]_{\mathcal{C}}), \quad \forall a \in \mathcal{C}, x, y \in \mathcal{M} \quad (3.1.1)$$

that are natural in the variable a , i.e. the following diagram:

$$\begin{array}{ccc} \text{hom}_{\mathcal{M}}(a' \odot x, y) & \xrightarrow{\simeq} & \text{hom}_{\mathcal{C}}(a', [x, y]_{\mathcal{C}}) \\ \downarrow - \circ (h \odot \text{id}_x) & & \downarrow - \circ h \\ \text{hom}_{\mathcal{M}}(a \odot x, y) & \xrightarrow{\simeq} & \text{hom}_{\mathcal{C}}(a, [x, y]_{\mathcal{C}}) \end{array}$$

is commutative for all $h : a \rightarrow a'$. We sometimes abbreviate $[x, y]_{\mathcal{C}}$ to $[x, y]$ for simplicity. □

There is a distinguished morphism $\text{ev} : [x, y]_{\mathcal{C}} \odot x \rightarrow y$ in $\mathcal{M}$ defined by the preimage of $\text{id}_{[x, y]_{\mathcal{C}}}$ under the isomorphism in (3.1.1). It allows us to give an equivalent definition of the internal hom via its universal property. Both definitions are useful and important in this work.

Definition 3.1.6. The internal hom can be equivalently defined by a pair $([x, y]_{\mathcal{C}}, \text{ev})$, where $\text{ev} : [x, y]_{\mathcal{C}} \odot x \rightarrow y$ is a morphism in $\mathcal{M}$, such that it is terminal among all such pairs. That is, given another pair (F, f) ,

³⁶Although it is very tempting to suggest physics oriented readers to skip this subsection for their first reading, we want to emphasize that the notion of internal hom is so important and natural that it is precisely the mathematical structure needed to catch many physical intuitions. Since it is so basic, natural and important to QFT's, we are sure that it will become one of the most basic concepts in physics and a powerful tool for the physicists in the next generation.

³⁷Our assumptions on $\mathcal{C}$ and $\mathcal{M}$ guarantee the existence of the internal homs.

where $F \in \mathcal{C}$ and $f : F \odot x \rightarrow y$ is a morphism in $\mathcal{M}$, there exists a unique morphism $g : F \rightarrow [x, y]_{\mathcal{C}}$ such that $\text{ev} \circ (g \odot \text{id}_x) = f$. This universal property is often expressed by the following commutative diagram:

$$\begin{array}{ccc} & [x, y]_{\mathcal{C}} \odot x & \\ \exists! g \odot \text{id}_x \nearrow & & \searrow \text{ev} \\ F \odot x & \xrightarrow{f} & y \end{array} \quad (3.1.2)$$

where the notation ‘ $\exists$ ’ represents ‘exists’ and the notation ‘!’ represents ‘unique’.

Example 3.1.7. We give two most useful examples, which also tell us how to compute internal homs.

1. when $\mathcal{M} = \mathcal{C}$ and $\odot = \otimes_{\mathcal{C}}$, we have $[x, y]_{\mathcal{C}} = y \otimes_{\mathcal{C}} x^L$;
2. when $\mathcal{M} = \text{RMod}_A(\mathcal{C})$ and $\odot = \otimes_{\mathcal{C}}$, we have $[x, y]_{\mathcal{C}} = (x \otimes_A y^R)^L$ [Ost03a]. It includes the first case as a special case when $A = \mathbb{1}_{\mathcal{C}}$.

Remark 3.1.8. The internal homs define an internal hom functor $[-, -]_{\mathcal{C}} : \mathcal{M}^{\text{op}} \times \mathcal{M} \rightarrow \mathcal{C}$, i.e. $(x, y) \mapsto [x, y]_{\mathcal{C}}$. As a consequence, the isomorphisms in (3.1.1) are natural in all three variables a, x, y . Moreover, the internal hom can be equivalently defined by stating that $[x, -] : \mathcal{M} \rightarrow \mathcal{C}$ is the right adjoint of the functor $- \odot x : \mathcal{C} \rightarrow \mathcal{M}$.

What is remarkable about the internal hom is that, by the universal property of the internal hom $[x, z]$, there is a distinguished morphism

$$[y, z] \otimes_{\mathcal{C}} [x, y] \xrightarrow{\text{ev}} [x, z] \quad (3.1.3)$$

determined by the composed morphism: $[y, z] \odot ([x, y] \odot x) \xrightarrow{\text{id}_{[y, z]} \otimes_{\mathcal{C}} \text{ev}} [y, z] \odot y \xrightarrow{\text{ev}} z$ and the universal property. Moreover, we have a canonical morphism $\eta : \mathbb{1}_{\mathcal{C}} \rightarrow [x, x]$ defined by $\mathbb{1}_{\mathcal{C}} \odot x \simeq x$.

Lemma 3.1.9. The triple $([x, x], \text{ev}, \eta)$ defines an E_1 -algebra in $\mathcal{C}$; the pair $([x, y], \text{ev})$, where $\text{ev} : [x, y] \otimes_{\mathcal{C}} [x, x] \rightarrow [x, y]$, defines a right $[x, x]$ -module structure on $[x, y]$; and $[x, y]$ is naturally a $[y, y]$ - $[x, x]$ -bimodule.

Remark 3.1.10. Internal homs play very important role in QFT’s. For example, in 1+1D rational CFT’s, all modular-invariant bulk CFT’s, boundary CFT’s and defect or wall CFT’s are internal homs. More precisely, let V be a rational vertex operator algebra (VOA) V . The category $\mathcal{C} := \text{Mod}_V$ is automatically a modular tensor category (MTC) [MS89a, Hua08]. A category of boundary conditions preserving the chiral symmetry V is defined by an indecomposable separable left $\mathcal{C}$ -module $\mathcal{M}$, i.e. each object in $\mathcal{M}$ represents a boundary condition preserving the chiral symmetry V .

1. The 0+1D boundary CFT associated to the boundary condition $x \in \mathcal{M}$ is precisely the internal hom $[x, x]_{\mathcal{C}}$. The algebraic structure defined by the triple $([x, x], \text{ev}, \eta)$ precisely encodes the mathematical structure of OPE among boundary fields [FRS02, HK04, KR09, DKR15].
2. The 0D domain wall between two boundary CFT’s associated to the boundary conditions x and y is given by the internal hom $[x, y]_{\mathcal{C}}$ [FFRS07, DKR15]. The morphism defined in (3.1.3) precisely encode the most general OPE between two fields living in two domain walls.
3. The modular-invariant bulk CFT is again the internal hom $[\text{id}_{\mathcal{M}}, \text{id}_{\mathcal{M}}]_{\mathfrak{Z}_1(\mathcal{C})}$ but living in $\mathfrak{Z}_1(\mathcal{C})$ (i.e. the Drinfeld center of $\mathcal{C}$) [FRS02, KR09, Dav10a], where $\text{id}_{\mathcal{M}} \in \text{Fun}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})$ is the identity functor, and $\text{Fun}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})$ is the category of $\mathcal{C}$ -module functors from $\mathcal{M}$ to $\mathcal{M}$ and, at the same time, a left $\mathfrak{Z}_1(\mathcal{C})$ -module defined by $(z \odot F)(-) := z \odot F(-)$ for $z \in \mathfrak{Z}_1(\mathcal{C})$ and $F \in \text{Fun}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})$. This internal hom $[\text{id}_{\mathcal{M}}, \text{id}_{\mathcal{M}}]_{\mathfrak{Z}_1(\mathcal{C})}$ is automatically a Lagrangian algebra in $\mathfrak{Z}_1(\mathcal{C})$ [KR09, DMNO13].

The following mathematical theorem tells us the basic structure of a condensable E_1 -algebra in a multi-fusion 1-category $\mathcal{A}$.

Theorem 3.1.11 ([KZ17]). A condensable E_1 -algebra A in a multi-fusion 1-category $\mathcal{A}$ is a direct sum of indecomposable condensable E_1 -algebras, each of which is a matrix algebra. More explicitly, assume A is indecomposable, let $A = \oplus_{i=1}^m x_i$ be the decomposition of the right A -module into simple ones. We have the following results.

1. $A \simeq \oplus_{i,j}^m [x_i, x_j]$;
2. the functor $[x_i, -] : \text{RMod}_A(\mathcal{A}) \rightarrow \text{RMod}_{[x_i, x_i]}(\mathcal{A})$ is an equivalence;
3. $[x_i, x_i]$ is a simple condensable E_1 -algebra;
4. $[x_i, x_j]$ is an invertible $[x_j, x_j]$ - $[x_i, x_i]$ -bimodule;
5. $[x_j, x_l] \otimes_{[x_j, x_j]} [x_i, x_j] \simeq [x_i, x_l]$ as $[x_l, x_l]$ - $[x_i, x_i]$ -bimodules.

Remark 3.1.12. Up to the standard assumptions about higher n -categories and their (co)-limit theory (see [GJF19]), the proof of this theorem in [KZ17] can be generalized to condensable E_1 -algebras in a multi-fusion n -category. In this work, we simply take it for granted that condensable E_1 -algebras in a multi-fusion n -category as the same structure theorem as the $n = 1$ case. $\diamond$

Now we slightly generalize the bootstrap results in Theorem^{ph} 3.1.2 and Theorem^{ph} 3.1.4 for a potentially composite topological order C^2 (i.e., $\mathcal{C}$ is a multi-fusion 1-category).

Theorem^{ph} 3.1.13. A condensable algebra A in a multi-fusion 1-category $\mathcal{C}$ is a direct sum of non-zero condensable algebras in each indecomposable summand of $\mathcal{C}$.

1. C^2 is type-I (i.e., $\mathcal{C}$ is an indecomposable multi-fusion 1-category):
 - (a) If A is indecomposable, then the condensed phase D^2 is type-I and simple (i.e., $\mathcal{D}$ is a fusion 1-category), and $\mathcal{M} = \text{RMod}_A(\mathcal{C})$ is an indecomposable left $\mathcal{C}$ -module.
 - (b) If A is a direct sum of indecomposable ones, i.e., $A = \oplus_{i=1}^l A_i$, then the condensed phase D^2 is type-I, $\text{gsd}(D^2) = l$, and $\mathcal{D}$ is an indecomposable multi-fusion 1-category, which can be decomposed as a direct sum of separable 1-categories: $\mathcal{D} = \oplus_{i,j=1}^l \text{BMod}_{A_i|A_j}(\mathcal{C})$. The domain wall $M^1 = (\mathcal{M}, A)$ is of type-II and we have the decomposition of left $\mathcal{C}$ -modules: $\mathcal{M} = \text{RMod}_A(\mathcal{C}) = \oplus_{i=1}^l \text{RMod}_{A_i}(\mathcal{C})$.
2. C^2 is type-II (i.e., $\mathcal{C} = \oplus_{i=1}^k \mathcal{C}_i$ and $\mathcal{C}_i$ are indecomposable multi-fusion 1-categories):
 - (a) In this case, we have $A = \oplus_{i=1}^k A_i$, where $A_i = A \cap \mathcal{C}_i \neq 0$.
 - (b) The condensed phase D^2 is also type-II, $\text{mult}(C^2) \simeq \text{mult}(D^2)$, and $\mathcal{D} \simeq \oplus_{i=1}^k \text{BMod}_{A_i|A_i}(\mathcal{C}_i)$.
 - (c) The domain wall M^1 is type-II, and we have the decomposition of left $\mathcal{C}$ -modules: $\mathcal{M} = \text{RMod}_A(\mathcal{C}) = \oplus_{i=1}^k \text{RMod}_{A_i}(\mathcal{C}_i)$. $\square$

Remark 3.1.14. Note that the number $\text{gsd}(C^2)$ is not preserved in a condensation but the number $\text{mult}(C^2)$ is preserved. More precisely, when C^2 is type-I, $\text{gsd}(D^2)$ only depends on the number of direct summands of algebras in A . For example, by Proposition 2.2.34, we have $\mathcal{C} = \oplus_{i,j}^m \mathcal{C}_{i,j}$ as separable 1-categories and $\mathbb{1}_{\mathcal{C}} = \oplus_{i=1}^m e_i$. If $A = e_1$, then $\mathcal{D} \simeq \mathcal{C}_{11}$ and $\text{gsd}(D^2) = 1$; if $A = e_1 \oplus e_2$, then $\mathcal{D} \simeq \oplus_{i,j=1}^2 \mathcal{C}_{i,j}$ and $\text{gsd}(D^2) = 2$. $\diamond$

3.1.3 Examples of particle condensations in 1+1D

Since the only simple 1+1D anomaly-free topological order is the trivial one, we provide examples of condensations in an anomalous 1+1D simple topological order C^2 . In this case, $\mathcal{C}$ is a fusion 1-category. Physically, such a condensation can be viewed either as a phase transition occurred on the gapped boundary of $Z(C)^3$ or, by topological Wick rotation or topological holography [KZ18b, KZ20, KZ21, KLWZZ20a, KLWZZ20b, KZ22b], as condensation among 1+1D gapped quantum liquids.

Example 3.1.15. The topological defects on the trivial 1-codimensional topological defect in the 2+1D $\mathbb{Z}_2$ -gauge theory form a fusion category $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$. By condensing simple condensable E_1 -algebras in $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$, we obtain all simple 1-codimensional topological defects, which are simple objects in the fusion 2-category $\Sigma\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$. This fusion 2-category has six simple objects, denoted by $\mathbb{1}, \vartheta, \text{rr}, \text{rs}, \text{sr}, \text{ss}$ [KZ22a] (recall Example 2.1.5).

Since $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2)) \simeq \text{Vec}_{\mathbb{Z}_2 \times \mathbb{Z}_2}$, the condensable E_1 -algebras in $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$ are classified by subgroups of $\mathbb{Z}_2 \times \mathbb{Z}_2$ and the corresponding cohomology groups. There are four simple objects in $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$ denoted by $1, e, m, f$.

1. The tensor unit 1 is a condensable E_1 -algebra. By condensing 1 we obtain the trivial 1-codimensional topological defect $\mathbb{1}$.
2. There are three subgroups of $\mathbb{Z}_2 \times \mathbb{Z}_2$ which are isomorphic to $\mathbb{Z}_2$. Since $H^2(\mathbb{Z}_2; U(1)) = 0$ is trivial, there are three corresponding condensable E_1 -algebra $1 \oplus e, 1 \oplus m, 1 \oplus f$. By condensing them we obtain the 1-codimensional topological defects rr, ss, ϑ , respectively.
3. Since $H^2(\mathbb{Z}_2 \times \mathbb{Z}_2; U(1)) \simeq \mathbb{Z}_2$, there are two condensable E_1 -algebra structures on the object $1 \oplus e \oplus m \oplus f$. By condensing them we obtain the 1-codimensional topological defects rs and sr , respectively. $\heartsuit$

Example 3.1.16. Recall the Ising fusion category $\mathcal{I}_s$ in Example 2.1.7. According to [FRS02], all condensable E_1 -algebras A in $\mathcal{I}_s$ are Morita equivalent to the trivial E_1 -algebras $\mathbb{1}$. More explicitly, one can show that they are the internal hom algebras: $[x, x] = x \otimes x^*, \forall x \in \mathcal{I}_s$ (see Section 3.1.2 for a brief introduction of this notion). As a consequence, we have $\text{BMod}_{A|\mathcal{I}_s}(\mathcal{I}_s) \simeq \mathcal{I}_s$ for all condensable E_1 -algebras A . In other words, all particle condensations in $\mathcal{I}_s$ along a line, which can be viewed as a boundary phase of the 2+1D double Ising topological order, only produces the same and unique boundary phase of the 2+1D double Ising topological order. This fact is compatible with the fact that there is only one Lagrangian algebra in $\mathfrak{Z}_1(\mathcal{I}_s)$ [KR08, DMNO13]. $\heartsuit$

Example 3.1.17. Let G be a finite group. Consider the 2+1D finite gauge theory GT_G^3 .

- (1) It has a gapped boundary whose topological defects form the fusion 1-category $\text{Rep}(G)$ (when $G = \mathbb{Z}_2$, this is the smooth boundary of the toric code model). The classification of condensable E_1 -algebras in $\text{Rep}(G)$ is given by Ostrik [Ost03a]. A condensable E_1 -algebra in $\text{Rep}(G)$ is a separable algebra A in Vec equipped with a G -action (i.e., a group homomorphism $G \rightarrow \text{Aut}(A)$), and it is simple if and only if G acts on the set of primitive (minimal) central idempotents of A transitively. We give some simple examples.
 - (a) When we condense $A = \text{Fun}(G)$ in the gapped boundary associated to $\text{Rep}(G)$, we obtain the condensed boundary associated to $\text{BMod}_{A|\mathcal{I}_s}(\text{Rep}(G)) \simeq \text{Vec}_G$, and the gapped domain wall $\mathcal{M} = \text{RMod}_A(\text{Rep}(G)) \simeq \text{Vec}$. When $G = \mathbb{Z}_2$, this condensation defines a boundary phase transition from the smooth boundary to the rough boundary.³⁸
 - (b) When we condense $A = \text{Fun}(G/H)$ for a subgroup $H \leq G$ in the gapped boundary associated to $\text{Rep}(G)$, we obtain a condensed boundary whose boundary particles form the fusion category:

$$\text{BMod}_{A|\mathcal{I}_s}(\text{Rep}(G)) \simeq \text{Fun}(\text{Vec}_{G/H}, \text{Vec}_{G/H})^G,$$

where $\text{Fun}(\text{Vec}_{G/H}, \text{Vec}_{G/H})^G$ denotes the G -equivariantization of $\text{Fun}(\text{Vec}_{G/H}, \text{Vec}_{G/H})$ [XZ24].³⁹

Now we give a general construction of simple condensable E_1 -algebras in $\text{Rep}(G)$ via internal homs in (3.1.4). By [BO04, Ost03a], an indecomposable finite semisimple left $\text{Rep}(G)$ -module is necessarily equivalent to $\text{Rep}'(\tilde{H})$, where $\tilde{H}$ is the central extension of a subgroup $H \leq G$ determined by a 2-cohomology class $[\alpha] \in H^2(H, \mathbb{C}^\times)$:

$$1 \rightarrow \mathbb{C}^\times \rightarrow \tilde{H} \rightarrow H \rightarrow 1$$

and $\text{Rep}'(\tilde{H})$ is the category of $\tilde{H}$ -representations such that $\mathbb{C}^\times$ acts via the identity character. Equivalently, $\text{Rep}'(\tilde{H})$ is equivalent to the category $\text{Rep}(H, \alpha)$ of α -twisted projective H -representations, where an object in $\text{Rep}(H, \alpha)$ is a vector space V equipped with a map $\rho: H \rightarrow \text{GL}(V)$ such that $\rho(g)\rho(h) = \alpha(g, h)\rho(gh)$ for all $g, h \in H$. The $\text{Rep}(G)$ -action on $\text{Rep}(H, \alpha)$ is defined by the usual vector space tensor product. For $V \in \text{Rep}(H, \alpha)$, the internal hom

$$A_V^{(H, \alpha)} := [V, V]_{\text{Rep}(G)} = \text{Ind}_H^G \text{End}(V) \quad (3.1.4)$$

³⁸According to topological Wick rotation [KZ22b, KWZ22], this condensation also defines the spontaneous symmetry-breaking from the symmetric phase to the symmetry-broken phase in the 1+1D Ising chain [KWZ22].

³⁹According to topological Wick rotation, this condensation defines the spontaneous symmetry-breaking from the symmetric phase to the symmetry-partially-broken (from G -symmetry to H -symmetry) phase in a 1+1D spin chain [XZ24].

defines a condensable E_1 -algebra in $\text{Rep}(G)$ such that $\text{Mod}_{A_V^{(H,\alpha)}}(\text{Rep}(G)) \simeq \text{Rep}(H, \alpha)$. In particular, it means that different choices of V define algebras in the same Morita class. Importantly, $\text{Rep}(H, \alpha)$ and $\text{Rep}(K, \beta)$ are equivalent as $\text{Rep}(G)$ -modules if and only if these two pairs (H, α) and (K, β) are conjugate in the sense that there exists an element $g \in G$ such that $gHg^{-1} = K$ and $[\alpha] = [g^*\beta] \in H^2(H; U(1))$, where $(g^*\beta)(x, y) := \beta(gxg^{-1}, gyg^{-1})$ for all $x, y \in H$. By condensing $A_V^{(H,\alpha)}$, we obtain a condensed phase, particles on which form the fusion 1-category:

$$\text{Mod}_{A_V^{(H,\alpha)}}^{E_1}(\text{Rep}(G)) \simeq \text{Fun}_{\text{Rep}(G)}(\text{Rep}(H, \alpha), \text{Rep}(H, \alpha))^{\text{op}} \simeq \mathfrak{Z}_1(\text{Rep}(G), \mathfrak{Z}_0(\text{Rep}(H, \alpha)))^{\text{op}}.$$

Note that $\text{Rep}(G, \alpha)$ is an invertible $\text{Rep}(G)$ - $\text{Rep}(G)$ -bimodule. Therefore, by condensing $A_V^{(G,\alpha)}$, we obtain the condensed phase $\text{Mod}_{A_V^{(G,\alpha)}}^{E_1}(\text{Rep}(G)) \simeq \text{Rep}(G)$.

- (2) It has another gapped boundary of GT_G^3 whose topological defects form a fusion category Vec_G (when $G = \mathbb{Z}_2$, this is the rough boundary of the toric code model). A condensable E_1 -algebra in Vec_G is a separable algebra A in Vec equipped with a G -grading, and it is simple if the trivial component $A_e = \mathbb{C}$ is trivial. The simple condensable E_1 -algebras in Vec_G are given by the twisted group algebras $\mathbb{C}[H, \alpha]$ equipped with the obvious G -grading, where $H \leq G$ is a subgroup and $\alpha \in Z^2(H; U(1))$ is a 2-cocycle. For $\chi \in C^1(H, \mathbb{C}^\times)$, $\mathbb{C}[H, \alpha]$ and $\mathbb{C}[H, \alpha + d\chi]$ are isomorphic as algebras. Two E_1 -condensable algebras $\mathbb{C}[H, \alpha]$ and $\mathbb{C}[K, \beta]$ are Morita equivalent if and only if the pairs (H, α) and (K, β) are conjugate.

By applying topological Wick rotation [KZ18b, KZ20, KZ21, KZ22b], condensations on the gapped boundaries of GT_G^3 can also be viewed as phase transitions, which are beyond Landau's spontaneous symmetry-breaking theory, among 1+1D gapped quantum liquids with the same onsite symmetry G [KZ22b]. $\heartsuit$

Example 3.1.18 ([FS03, Ost03a, Ost03b]). Consider the fusion 1-category Vec_G^ω for a finite group G and $\omega \in H^3(G, \mathbb{C}^\times)$. For each subgroup $H \leq G$ such that $\omega|_H = 1$ and $\psi \in H^2(H, \mathbb{C}^\times)$, the twist group algebra $\mathbb{C}[H, \psi]$ defines a simple condensable E_1 -algebra in Vec_G^ω with the G -grading supported on H . Conversely, any simple condensable E_1 -algebra in Vec_G^ω is isomorphic to $\mathbb{C}[H, \psi]$ for some H and ψ . Two algebras $\mathbb{C}[H, \psi]$ and $\mathbb{C}[H, \phi]$ are isomorphic if and only if $\phi = \psi + d\chi$ for $\chi \in C^1(H, \mathbb{C}^\times)$. Two algebras $\mathbb{C}[H, \psi]$ and $\mathbb{C}[K, \phi]$ are Morita equivalent if two pairs (H, ψ) and (K, ϕ) belong to the same conjugacy class. In other words, the indecomposable left Vec_G^ω -modules are classified by the conjugacy classes of the pairs (H, ψ) . Moreover, there is a one-to-one correspondence between indecomposable Vec_G^ω -modules (or equivalently, the Morita classes of simple condensable E_1 -algebras A in Vec_G^ω) and Lagrangian E_2 -algebras in $\mathfrak{Z}_1(\text{Vec}_G^\omega)$ [DMNO13] defined by $A \mapsto Z(A)$, where $Z(A)$ is the full center of A [FFRS07, KR08, Dav10a]. This recovers the classification of the Lagrangian E_2 -algebras in $\mathfrak{Z}_1(\text{Vec}_G^\omega)$ in [DS17]. $\heartsuit$

3.2 Particle condensations in 1+1D: geometric approach

In this subsection, we provide a geometric approach towards particle condensations. The key points in this approach had already appeared in [KS11a, FSV13]. Our presentation uses a generalization of the classical Eilenberg-Watts calculus, which is covered in Appendix A.2 and is also used in the geometric approach towards anyon condensation theory in 2+1D.

3.2.1 Physical intuitions

We provide a physical intuition of a particle condensation. Let $\mathcal{C}^2$ and $\mathcal{D}^2$ be two Morita equivalent 1+1D potentially anomalous type-I topological orders, i.e., $\mathcal{C}$ and $\mathcal{D}$ are indecomposable multi-fusion 1-categories. A condensation from $\mathcal{C}^2$ to $\mathcal{D}^2$ can be achieved by

- (1) first proliferating the $\mathcal{D}^2$ -phases inside the $\mathcal{C}^2$ -phase as illustrated below in spatial dimensions;

$$\begin{array}{ccccccccc} & \text{M}^1 & & \overline{\text{M}}^1 & & \text{M}^1 & & \overline{\text{M}}^1 & & \text{M}^1 & & \overline{\text{M}}^1 \\ \text{---} & \blacksquare & \text{---} & \blacksquare & \text{---} & \blacksquare & \text{---} & \blacksquare & \text{---} & \blacksquare & \text{---} & \blacksquare & \text{---} \\ \mathcal{C}^2 & & \mathcal{D}^2 & & \mathcal{C}^2 & & \mathcal{D}^2 & & \mathcal{C}^2 & & \mathcal{D}^2 & & \mathcal{C}^2 \end{array} \quad (3.2.1)$$

- (2) then shrinking each line segment $\overline{\text{M}}^1 \boxtimes_{\mathcal{C}} \text{M}^1$ produces a particle in $\mathcal{D}^2$;

(3) then annihilating this particle, i.e., mapping it to the trivial particle $\mathbb{1}_{\mathcal{D}}$ in $\mathcal{D}^2$.

This process defines the fusion of two green line segments into one. As a consequence, the entire 1d (spatial dimension) line turn into a green line.

Notice that shrinking a green line segment in (3.2.1), i.e., $M \boxtimes_{\mathcal{D}} \bar{M}$, defines a particle $A \in \mathcal{C}$. The step (2) and (3) simply produces a morphism $\mu_A : A \otimes_{\mathcal{C}} A \rightarrow A$. Moreover, we should have a canonical map $\eta_A : \mathbb{1}_{\mathcal{C}} \rightarrow A$, which creates the green line out of nothing. This map makes the proliferating possible in the first place. Since the physical meanings of 1-morphisms η_A and μ_A are instantons, we drew these instantons in the following spacetime pictures.

$$(3.2.2)$$

We provide later in (3.2.5) an algebraic definition of the instanton lying in the $\mathcal{D}^2$ -phase in the second picture in (3.2.2) and depicted as the “annihilation of the domain wall $M^1 = (\mathcal{M}, m)$ ”. Then the following identities of 1-morphisms in $\mathcal{C}$:

$$\mu_A \circ (\mu_A \otimes_{\mathcal{C}} \text{id}_A) = \mu_A \circ (\text{id}_A \otimes_{\mathcal{C}} \mu_A), \quad \mu_A \circ (\eta_A \otimes_{\mathcal{C}} \text{id}_A) = \text{id}_A = \mu_A \circ (\text{id}_A \otimes_{\mathcal{C}} \eta_A)$$

holds by the obvious physical intuitions. For example, the first identity simply says that first running the procedure (2) and (3) to a teal line the first, then running the same procedures to the teal line next to it makes no difference to running in the opposite order. As a consequence, the triple (A, μ_A, η_A) should define a condensable E_1 -algebra in $\mathcal{C}$, which defines the fusion of two green lines into one.

Remark 3.2.1. In the unitary case, by turning the pictures in (3.2.2) upside down, we obtain the counit ϵ_A and comultiplication $\Delta_A = \mu_A^\dagger$:

$$(3.2.3)$$

which is co-unital and co-associative. Moreover, $(A, \mu_A, \eta_A, \Delta_A, \epsilon_A)$ defines a Frobenius algebra [KS11a, FSV13]. Moreover, we have $\mu_A \circ \Delta_A = \text{id}_A$, i.e., A is a special Frobenius algebra in $\mathcal{C}$ [FRS02]. $\diamond$

A particle $d \in \mathcal{D}$ in the condensed phase $\mathcal{D}$ produces a particle in $\mathcal{C}$ by viewing the green line segment decorated by $d \in \mathcal{D}$ as a particle in $\mathcal{C}$ as illustrated below.

$$(3.2.3)$$

When $d = \mathbb{1}_{\mathcal{D}}$, this particle is $\mathbb{1}_{\mathcal{D}} = A \in \mathcal{C}$. Note that A is precisely $\mathbb{1}_{\mathcal{D}}$ viewed as an object in $\mathcal{C}$ because $\mathcal{D} \subset \mathcal{C}$. Therefore, for general $d \in \mathcal{D}$, we expect that this particle is precisely the same particle d viewed as an object in $\mathcal{C}$. Then it is clear that procedures (2) and (3) define an A - A -bimodule structure on the particle $d \in \mathcal{C}$. Therefore, we have recovered the same result $\mathcal{D} = \text{BMod}_{A|A}(\mathcal{C})$.

As we have shown in Section 3.1, the gapped domain wall M^1 can also support particles, which form a separable 1-category $\mathcal{M}$. First, $m \in \mathcal{M}$, m is necessarily an object in $\mathcal{C}$. Secondly, m is necessarily a right A -module. This suggests that $\mathcal{M} = \text{RMod}_A(\mathcal{C})$. The fact that the green line shrink to the particle in $\mathcal{C}$ and the fact that the teal line shrink to a particle in $\mathcal{D}$ simply suggests that

$$\mathcal{M}^{\text{op}} \boxtimes_{\mathcal{C}} \mathcal{M} \simeq \mathcal{D}, \quad \mathcal{M} \boxtimes_{\mathcal{D}} \mathcal{M}^{\text{op}} \simeq \mathcal{C}$$

Mathematically, it is given by the canonical equivalence $\mathcal{M} \boxtimes_{\mathcal{D}} \mathcal{M}^{\text{op}} \simeq \mathcal{C}$ defined by $x \boxtimes_{\mathcal{D}} y \mapsto [x, y]_{\mathcal{C}}^*$.

Moreover, according to (3.2.4) and (3.2.5), the procedure (2) and (3) defines a morphism $\mu_A : A \otimes_{\mathcal{C}} A \rightarrow A$ as the following composed map:

$$[m, m] \otimes_{\mathcal{C}} [m, m] \simeq [m, [m, m] \odot m] \xrightarrow{[m, \text{ev}_m]} [m, m], \quad (3.2.7)$$

where the first ‘ $\simeq$ ’ is due to the canonical isomorphism $a \otimes_{\mathcal{C}} [x, y] \simeq [x, a \otimes_{\mathcal{C}} y]$ [Ost03a]. It turns out that the composed map (3.2.7) coincides with the canonical morphism $[m, m] \otimes [m, m] \rightarrow [m, m]$ defined by the universal property of the internal hom. Moreover, it also coincides with the geometric construction of μ_A in (3.2.2)

Remark 3.2.2. Strictly speaking, that the algebraic construction (3.2.7) coincides with the geometric construction (3.2.2) is only intuitively clear instead of being rigorously proved. A rigorous proof needs a rigorous computation of the instanton μ_A in (3.2.2), which might be achieved by explicit computations in Turaev-Viro TQFT, or equivalently, by the mathematical theory of the so-called β -version of factorization homology (see Remark 6.2.2). $\diamond$

We have a canonical morphism from $\eta : \mathbb{1}_{\mathcal{C}} \rightarrow [m, m]$, which can be defined physically in two steps.

1. In the first step, we simply factorizes $\mathbb{1}_{\mathcal{C}}$ as follows:

$$\text{blue line } \mathbb{1}_{\mathcal{C}} = \int_{x \in \mathcal{M}}^{\mathcal{D}} \text{blue line } x \in \mathcal{M} \text{ and green line } x \in \mathcal{M}^{\text{op}}$$

That is, the canonical equivalence $\mathcal{C} \simeq \mathcal{M} \boxtimes_{\mathcal{D}} \mathcal{M}^{\text{op}}$ maps $\mathbb{1}_{\mathcal{C}}$ to an object $\int_{x \in \mathcal{M}}^{\mathcal{D}} x \boxtimes_{\mathcal{D}} x$ in $\mathcal{M} \boxtimes_{\mathcal{D}} \mathcal{M}^{\text{op}}$, where the integral defines a limit (called ‘end’) in the category $\mathcal{C}$. In Appendix A.2, we explain this notion as a generalization of the classical Eilenberg-Watts calculus. That $\mathbb{1}_{\mathcal{C}} \mapsto \int_{x \in \mathcal{M}}^{\mathcal{D}} x \boxtimes_{\mathcal{D}} x$ in $\mathcal{M} \boxtimes_{\mathcal{D}} \mathcal{M}^{\text{op}}$ is explained in Example A.2.12. Physically, it means that one can always split a teal line by inserting a green line segment, then all possible particles $x \in \mathcal{M}, x \in \mathcal{M}^{\text{op}}$ are accumulating at two ends in pairs. Such a pair of particles can be viewed as a pair of boundary conditions at the two ends of the green line segment. By integrating all these possible boundary conditions, we simply recover the trivial particle in $\mathcal{C}$. This step is completely invertible.

2. In the second step, we define η by “projecting out” all the x -components for $x \neq m$. Mathematically, this “projection” is precisely the defining morphism $\int_{x \in \mathcal{M}}^{\mathcal{D}} x \boxtimes_{\mathcal{D}} x \rightarrow m \boxtimes_{\mathcal{D}} m$ of the limit. It turns out that such physically defined morphism η is precisely the one defined by the universal property of the internal hom $[m, m]$.

Example 3.2.3. It is helpful to look at a special case $\mathcal{D} = \mathcal{C} = \mathcal{M}$. In this case, we have (see also Example A.2.10)

$$\int_{x \in \mathcal{M}}^{\mathcal{D}} x \boxtimes_{\mathcal{D}} x = \int_{x \in \mathcal{C}}^{\mathcal{C}} x \otimes_{\mathcal{C}} x^* \simeq \mathbb{1}_{\mathcal{C}}.$$

If we already know that $\mathcal{D}^2$ and $\mathcal{M}^1$ are obtained from $\mathcal{C}^2$ by condensing a condensable E_1 -algebra $A \in \mathcal{C}$. In this case, we have $\mathcal{D} = \text{BMod}_{A|A}(\mathcal{C})$, $\mathcal{M} = \text{RMod}_A(\mathcal{C})$, $\mathcal{M}^{\text{op}} = \text{LMod}_A(\mathcal{C})$ and $\mathcal{M} = (\mathcal{M}, A)$.

1. The bulk-to-wall map $\mathcal{C} \rightarrow \mathcal{M}$ is $a \mapsto a \otimes_{\mathcal{C}} A$, $\forall a \in \mathcal{C}$. Its right adjoint functor is $[A, -] : \mathcal{M} \rightarrow \mathcal{C}$ is precisely the forgetful functor $f : \text{RMod}_A(\mathcal{C}) \rightarrow \mathcal{C}$ defined by $x \mapsto x$ by forgetting the A -module structure on x . Therefore, $[A, A]_{\mathcal{C}} = A$.
2. The d -particle in $\mathcal{D}^2$, as depicted in (3.2.3), becomes the particle $[A, d \otimes_A A] = d$ in $\mathcal{C}$. This assignment $\mathcal{D} \ni d \mapsto d \in \mathcal{C}$ is precisely the forgetful functor $f : \text{RMod}_A(\mathcal{C}) \rightarrow \mathcal{C}$.

Very importantly, what we have shown is that the data on the wall $\mathcal{M}^1 = (\mathcal{M}, m)$, which should be viewed as a boundary condition, uniquely determines a condensation. This is compatible with the usual physical intuition that a condensation can be constructed by proliferating the green line segments with a chosen boundary condition on the wall $\mathcal{M}^1$. Moreover, in (3.2.6), the choice of the object $m \in \mathcal{M}$ is arbitrary. A different choice m' simply produces a different condensable E_1 -algebra $A' = [m', m']^{\text{op}}$, which defines a

- (1) first proliferating the D^{n+1} -phases inside the C^{n+1} -phase along the unique transversal direction x^1 as illustrated Figure 11, where the domain wall M^n is assumed to be gapped;




Figure 11: Condensation of 1-codimensional topological defects in C^{n+1}

- (2) then shrinking each teal region $\overline{M}^n \boxtimes_C M^n$ to a 1-codimensional topological defect in D^{n+1} ;
- (3) then annihilating this defect, i.e. mapping it to the trivial 1-codimensional topological defect $1_{\mathcal{D}}$ in D^{n+1} .

As a consequence, the entire phase C^{n+1} turn into a new phase D^{n+1} , which we assume to be indecomposable.

Remark 3.3.1. Such a condensation produces a new phase D^{n+1} that share the same gravitational anomaly with C^{n+1} as illustrated below.

$$\begin{array}{c} Z(C)^{n+2} = Z(D)^{n+2} \\ \hline \begin{array}{ccc} \xrightarrow{C^{n+1}} & \text{■} & \xrightarrow{D^{n+1}} \\ & M^n & \end{array} \end{array}$$

Therefore, one can also view this condensation as a purely boundary phase transition of $Z(C)^{n+2}$. We assume that the domain wall M^n is gapped just for convenience. One can see that if the domain wall M^n is gapless, the argument is parallel. This defines a condensation of liquid-like gapless defects in C^{n+1} , which is beyond the setting of this section. We postpone it to the last section. $\diamond$

One can see that condensing a 1-codimensional topological defect in an $n+1$ D topological order is completely parallel to that in a $1+1$ D topological order. We simply summarize the result below.

Theorem^{ph} 3.3.2. A condensation of a 1-codimensional topological defect in an $n+1$ D type-I topological order C^{n+1} can be mathematically described as follows (we set $\otimes_e = \otimes$):

1. Since the condensed phase D^{n+1} is condensed from C^{n+1} , the multi-fusion n -category $\mathcal{D}$ is necessarily a sub- n -category of $\mathcal{C}$. The trivial 1-codimensional topological defect $1_{\mathcal{D}}$ in $\mathcal{D}$ corresponds to a non-trivial 1-codimensional topological defect $A \in \mathcal{C}$, which is determined by shrinking a green strap in Figure 11 to a 1-codimensional topological defect in C^{n+1} .
2. The A -defect is naturally equipped with the structure of an algebra in $\mathcal{C}$ with the unit 1-morphism η_A and the multiplication 1-morphism μ_A defined by the following pictures in spatial dimensions⁴⁰.

$$\begin{array}{c} \begin{array}{c} \text{M} \quad \overline{\text{M}} \\ \text{η}_A = \text{[diagram of a blue arc in a green region]} \\ \text{x}^2 \uparrow \\ \text{x}^1 \rightarrow \end{array} \quad \begin{array}{c} \text{M} \quad \overline{\text{M}} \\ \text{μ}_A = \text{[diagram of two blue arcs meeting at a point in a green region]} \\ \overline{\text{M}} \quad \text{M} \end{array} \end{array} \quad (3.3.1)$$

⁴⁰Note that the vertical axis in (3.3.1) is a spatial direction, while that in (3.2.2) is the temporal direction.

Since the physical meaning of a 1-morphism in $\mathcal{C}$ is a 2-codimensional topological defect, both η_A and μ_A are defined by the 2-codimensional topological defects encircled by the dashed boxes in above picture. It is clear that we should have the associativity and the unital “properties” of μ_A and η_A . However, for $n > 1$, these “properties” are more than properties because higher morphisms (i.e., higher codimensional defects) are needed as defining data. For example, we should have the following 2-isomorphisms

$$\mu_A \circ (\mu_A \otimes \text{id}_A) \xrightarrow{\sim} \mu_A \circ (\text{id}_A \otimes \mu_A), \quad \mu_A \circ (\eta_A \otimes \text{id}_A) \xrightarrow{\sim} \text{id}_A = \mu_A \circ (\text{id}_A \otimes \eta_A).$$

If $n = 2$, these 2-isomorphisms should satisfy a pentagon condition and a triangle condition (see [D  c23b] for a precise mathematical definitions and see also [ZLZH⁺23]); if $n > 2$, there should be a 3-isomorphism associated to each pentagon and one associated to each triangle, so on and so forth. Our description here is mainly descriptive instead of being precise. A formal theory to take care all the higher coherence data can be found in [Lur17]. The higher coherence data can also be more efficiently encoded by internal homs as we show in Theorem 3.3.9. Moreover, we require such algebra to be condensable.

3. Similar to the particle condensations in 1+1D topological orders, 1-codimensional topological defects in $\mathcal{D}^{n+1}$ are precisely A - A -modules or E_1 - A -modules in $\mathcal{C}$ (see [D  c23b, ZLZH⁺23] for mathematical definitions for $n = 2$), i.e. $\mathcal{D} = \text{Mod}_A^{E_1}(\mathcal{C})$ (recall Remark 3.1.1), which is an indecomposable multi-fusion n -category (i.e., $\mathcal{D}^{n+1}$ is type-I). If A is indecomposable, then $\mathcal{D} = \text{Mod}_A^{E_1}(\mathcal{C})$ is a fusion n -category.
4. M^n can be described by a pair $(\mathcal{M}, m)$, where $\mathcal{M}$ is a separable n -category given by $\text{RMod}_A(\mathcal{C})$ and $m = A$ is an object in $\mathcal{M} = \text{RMod}_A(\mathcal{C})$ representing a 1-codimensional defect on the wall. The pair $(\mathcal{M}, m)$ can be viewed as an E_0 -algebra in $(n+1)\text{Vec}$.
5. Topological defects in $\mathcal{C}^{n+1}$ move onto the M^n -wall according to the left bulk-to-wall map (or functor)

$$L := - \otimes A : \mathcal{C} \rightarrow \mathcal{M} = \text{RMod}_A(\mathcal{C}) \quad \text{defined by } a \mapsto a \otimes A. \quad (3.3.2)$$

Topological defects in $\mathcal{D}^{n+1}$ move onto the M^n -wall according to the right bulk-to-wall map (or functor)

$$R := A \otimes_A - : \mathcal{D} \rightarrow \mathcal{M} = \text{RMod}_A(\mathcal{C}) \quad \text{defined by } d \mapsto A \otimes_A d. \quad (3.3.3)$$

□

6. Note that the right adjoint L^R of the functor $L = - \otimes A : \mathcal{C} \rightarrow \mathcal{M} = \text{RMod}_A(\mathcal{C})$ is the internal hom functor $[A, -]$ (see Definition 3.3.6), which, in this case, is precisely the forgetful functor $f : \text{RMod}_A(\mathcal{C}) \rightarrow \mathcal{C}$ by forgetting the right A -module structure, i.e.

$$L^R = [A, -] = f.$$

In particular, the condensable E_1 -algebra can be recovered as $L^R(A) = [A, A] = A$.

7. This condensation preserves the gravitational anomaly (recall Remark 3.3.1), i.e., $Z(\mathcal{C})^{n+2} = Z(\mathcal{D})^{n+2}$. In other words, $\mathcal{C}^{n+1}$ and $\mathcal{D}^{n+1}$ are Morita equivalent. Mathematically, it means that the indecomposable multi-fusion n -categories $\mathcal{C}$ and $\mathcal{D}$ are Morita equivalent, and we have the following braided equivalence:

$$\mathfrak{Z}_1(\mathcal{C}) \simeq \mathfrak{Z}_1(\text{Mod}_A^{E_1}(\mathcal{C})).$$

Remark 3.3.3. When $\mathcal{C}^{n+1} = \mathbf{1}^{n+1}$, Theorem^{ph} 3.3.2 (see also Theorem^{ph} 3.3.9) is precisely a reformulation of the layer constructions of the $n+1$ D topological order $\mathcal{D}^{n+1}$ from the trivial $n+1$ D topological order $\mathbf{1}^{n+1}$. In this sense, general condensations can be viewed as layer constructions on a non-trivial background phase $\mathcal{C}^{n+1}$. Notice that such (generalized) layer constructions can only create a new phase $\mathcal{D}^{n+1}$ that connects to $\mathcal{C}^{n+1}$ by a gapped domain wall. When $\mathcal{C}^{n+1} = \mathbf{1}^{n+1}$, all non-chiral $n+1$ D topological orders can be obtained from the condensations of 1-codimensional topological defects in $\mathbf{1}^{n+1}$. ◇

Then the boundary-bulk relation [KWZ15, KWZ17] immediately gives the following mathematical corollary.

Corollary^o 3.3.4. Let $\mathcal{C}$ be an indecomposable multi-fusion n -category, $A \in \text{Alg}_{E_1}^c(\mathcal{C})$, $\mathcal{D} = \text{Mod}_A^{E_1}(\mathcal{C})$ and $\mathcal{M} = \text{RMod}_A(\mathcal{C})$. We have a natural monoidal equivalence (with an illustration of its physical meaning):

$$\begin{array}{c} \boxed{\mathfrak{Z}_1(\mathcal{C}) \simeq \mathfrak{Z}_1(\mathcal{D})} \\ \xrightarrow{\mathcal{C}} \xrightarrow{\mathcal{M}} \xrightarrow{\mathcal{D}} \end{array} \quad \phi : \text{Mod}_A^{E_1}(\mathcal{C})^{\text{op}} \boxtimes_{\mathfrak{Z}_1(\mathcal{C})} \mathcal{C} \xrightarrow{\simeq} \mathfrak{Z}_0(\text{RMod}_A(\mathcal{C})). \quad (3.3.4)$$

or equivalently, we have

$$\text{Mod}_A^{E_1}(\mathcal{C})^{\text{op}} \simeq \mathfrak{Z}_1(\mathcal{C}, \mathfrak{Z}_0(\text{RMod}_A(\mathcal{C}))). \quad (3.3.5)$$

Remark 3.3.5. Corollary^{ph} 3.3.4 generalizes [DMNO13, Corollary 3.30]. When $n = 2$ and $\mathfrak{Z}_1(\mathcal{C}) \simeq 2\text{Vec}$, taking looping on both sides and applying the formula (2.3.13) and Theorem 5.1.4, we obtain [DMNO13, Corollary 3.30]. We provide another way to understand Corollary^{ph} 3.3.4 as the MME-gauging of a fusion n -category symmetry (see Remark 5.3.15). $\diamond$

Definition 3.3.6. Let $\mathcal{A}$ be a $\mathbb{C}$ -linear monoidal n -category and $\mathcal{M}$ be a $\mathbb{C}$ -linear left $\mathcal{A}$ -module. The internal hom $[x, y]$ for $x, y \in \mathcal{M}$, if exists, is defined to be the object of $\mathcal{A}$ representing the functor $\text{hom}_{\mathcal{M}}(- \odot x, y) : \mathcal{A}^{\text{op}} \rightarrow \text{Cat}_{n-1}^{\mathbb{C}}$. That is,

$$\text{hom}_{\mathcal{A}}(-, [x, y]) \simeq \text{hom}_{\mathcal{M}}(- \odot x, y). \quad (3.3.6)$$

We say that $\mathcal{M}$ is *enriched* in $\mathcal{A}$ if $[x, y]$ exists for all $x, y \in \mathcal{M}$. $\square$

Remark 3.3.7. We want to point out that the equivalence (or better adjoint equivalence for physical applications) in (3.3.6) is a defining data of the internal hom. The higher isomorphisms in the definition of the equivalence determine the higher isomorphisms in the definition of the algebraic structure on the internal hom. $\diamond$

Proposition^o 3.3.8 ([KZ24]). For a multi-fusion n -category $\mathcal{A}$, every separable left $\mathcal{A}$ -module (i.e. a left $\mathcal{A}$ -module in $(n+1)\text{Vec}$) is enriched in $\mathcal{A}$.

The next result provides explicit constructions of condensable E_1 -algebras in $\mathcal{C}$ via internal homs.

Theorem^{ph} 3.3.9. Given two $n+1$ D Morita equivalent type-I topological orders $\mathcal{C}^{n+1}$ and $\mathcal{D}^{n+1}$ connected by a gapped domain wall $\mathcal{M}^n$ with a specified boundary condition $m \in \mathcal{M}$, i.e. $\mathcal{M}^n = (\mathcal{M}, m)$.

$$\begin{array}{c} \boxed{Z(\mathcal{C})^{n+2} = Z(\mathcal{D})^{n+2}} \\ \xrightarrow{\mathcal{C}^{n+1}} \xrightarrow{\mathcal{M}^n} \xrightarrow{\mathcal{D}^{n+1}} \end{array}$$

We have the following results.


- (1) $\mathcal{M} \in \text{BMod}_{\mathcal{C}|\mathcal{D}}((n+1)\text{Vec})$ and the defining bimodule structure on $\mathcal{M}$ induces a monoidal equivalence:

$$\mathcal{D}^{\text{op}} \boxtimes_{\mathfrak{Z}_1(\mathcal{C})} \mathcal{C} \xrightarrow{\simeq} \mathfrak{Z}_0(\mathcal{M}) = \text{Fun}(\mathcal{M}, \mathcal{M}); \quad (3.3.7)$$

or equivalently,


$$\mathcal{D}^{\text{op}} \simeq \text{Fun}_{\mathcal{C}}(\mathcal{M}, \mathcal{M}) \simeq \mathfrak{Z}_1(\mathcal{C}, \mathfrak{Z}_0(\mathcal{M})). \quad (3.3.8)$$

- (2) By shrinking a $\mathcal{D}^{n+1}$ -phase strap in the $\mathcal{C}^{n+1}$ -phase, we obtain a 1-codimensional topological defect in $\mathcal{C}^{n+1}$, which means, in particular, $\mathcal{M} \boxtimes_{\mathcal{D}} \mathcal{M}^{\text{op}} \simeq \mathcal{C}$ as $\mathcal{C}$ - $\mathcal{C}$ -bimodules. This 1-codimensional topological defect in $\mathcal{C}^{n+1}$ is given by the internal hom algebra $[m, m]_{\mathcal{C}}$ in $\mathcal{C}$, which is automatically a condensable E_1 -algebra in $\mathcal{C}$.



We have $\mathcal{M} \simeq \text{RMod}_{[m, m]_{\mathcal{C}}}(\mathcal{C})$ and $\mathcal{D} \simeq \text{Mod}_{[m, m]_{\mathcal{C}}}^{E_1}(\mathcal{C})$. Moreover, all condensable E_1 -algebras in $\mathcal{C}$ arise as $[m, m]_{\mathcal{C}}$ for some $\mathcal{D}, \mathcal{M}, m$.

- (3) We can reverse the process. By shrinking a C^{n+1} -phase strap in the D^{n+1} -phase, we obtain a 1-codimensional topological defect in D^{n+1} , which means, in particular, $\mathcal{M}^{\text{op}} \boxtimes_{\mathcal{C}} \mathcal{M} \simeq \mathcal{D}$ as $\mathcal{D}$ - $\mathcal{D}$ -bimodules. This 1-codimensional topological defect in $\mathcal{D}$ is given by the internal hom algebra $[m, m]_{\mathcal{D}}$ in $\mathcal{D}$, which is automatically a condensable E_1 -algebra in $\mathcal{D}$.




We have $\mathcal{M} \simeq \text{LMod}_{[m, m]_{\mathcal{D}}}(\mathcal{D})$ and $\mathcal{C} \simeq \text{Mod}_{[m, m]_{\mathcal{D}}}^{E_1}(\mathcal{D})$. Moreover, all condensable E_1 -algebras in $\mathcal{D}$ arise as $[m, m]_{\mathcal{D}}$ for some $\mathcal{C}, \mathcal{M}, m$.

In particular, two $n+1$ D topological orders are Morita equivalent if and only if they can be obtained from each other by condensing a topological defect of codimension 1 (recall Theorem^{ph} 2.1.12). $\square$

Remark 3.3.10. When both C^{n+1} and D^{n+1} are anomaly-free, it is illuminating to rewrite the results in terms of $\Omega\mathcal{C}$ and $\Omega\mathcal{D}$ because they determine $\mathcal{C}$ and $\mathcal{D}$, respectively, by delooping. Note that $\Omega\mathcal{C}$ and $\Omega\mathcal{D}$ are non-degenerate braided fusion $(n-1)$ -categories. All topological defects of codimension 1 and higher on a gapped domain wall form a closed multi-fusion $\Omega\mathcal{D}$ - $\Omega\mathcal{C}$ -bimodule $\mathcal{X}$, i.e. a multi-fusion $(n-1)$ -category equipped with a braided equivalence $\phi_{\mathcal{X}} : \Omega\mathcal{D}^{\text{op}} \boxtimes \Omega\mathcal{C} \rightarrow \mathfrak{Z}_1(\mathcal{X})$. We obtain the unique closed $\mathcal{C}$ - $\mathcal{D}$ -bimodule $\mathcal{M}$ as $\Sigma\mathcal{X}$, i.e. $\mathcal{M} \simeq \Sigma\mathcal{X}$. Moreover, the condition that $\mathcal{X}$ is closed multi-fusion $\Omega\mathcal{C}$ - $\Omega\mathcal{D}$ -bimodule is equivalent to the condition that $\Sigma\mathcal{X}$ is the unique closed $\mathcal{C}$ - $\mathcal{D}$ -bimodule $\mathcal{M}$ mainly because $\Sigma\mathfrak{Z}_1(\mathcal{X}) \simeq \mathfrak{Z}_0(\Sigma\mathcal{X})$ [JF22, KIWZZ20b, KZ22b]. Note that $\Sigma\mathcal{X}$ is precisely the separable n -category of all possible gapped wall conditions between C^{n+1} and D^{n+1} within a single Morita class. An object $m \in \mathcal{M}$ represents a single gapped domain wall and $\text{hom}_{\mathcal{M}}(m, m)$ is the multi-fusion $(n-1)$ -category of topological defects of codimension 1 and higher on the m -wall. $\diamond$

Remark 3.3.11. Similar to the 1+1D cases, once we fixed the domain wall M^1 by a pair $(\mathcal{M}, m)$ (so is its time reversal $\overline{M}^1 = (\mathcal{M}^{\text{op}}, m)$), then the condensation process is uniquely fixed with the condensable algebra defined by the internal hom algebra⁴¹ $[m, m]_{\mathcal{C}} \in \mathcal{C}$. Different choices of m define different condensations microscopically. But the macroscopic result of the condensation, i.e. the condensed phase, is the same. Different m 's give different equivalences $\mathcal{D} \simeq \text{Mod}_{[m, m]_{\mathcal{C}}}^{E_1}(\mathcal{C})$ and $\mathcal{M} \simeq \text{RMod}_{[m, m]_{\mathcal{C}}}(\mathcal{C})$, which provide different relabelings of the objects in $\mathcal{D}$ and $\mathcal{M}$. $\diamond$

Remark 3.3.12. A 1-codimensional topological defect $d \in \mathcal{D}$ in the condensed phase D^{n+1} 1-codimensional topological defect in C^{n+1} as illustrated below.



When $d = \mathbb{1}_{\mathcal{D}}$, this 1-codimensional topological defect is $\mathbb{1}_{\mathcal{D}} = A \in \mathcal{C}$. Note that A is precisely $\mathbb{1}_{\mathcal{D}}$ viewed as an object in $\mathcal{C}$ because $\mathcal{D} \subset \mathcal{C}$. Therefore, for general $d \in \mathcal{D}$, we expect that this 1-codimensional topological defect is given by $[A, d]_{\mathcal{C}} \in \mathcal{C}$. When $\mathcal{D} = \text{Mod}_A^{E_1}(\mathcal{C})$ and $\mathcal{M} = \text{RMod}_A(\mathcal{C})$, $[A, -] : \text{RMod}_A(\mathcal{C}) \rightarrow \mathcal{C}$ is precisely the forgetful functor. $\diamond$

Remark 3.3.13. Recall that the separable $(n+1)$ -category $\Sigma\mathcal{C} \simeq \Sigma\mathcal{D}$ is a connected component of the category of gapped boundary conditions of $Z(\mathcal{C})^{n+2}$. In $\Sigma\mathcal{C}$, C^{n+1} is labeled by the distinguished object $\bullet \in \Sigma\mathcal{C}$; D^{n+1} is labeled by $\mathcal{M}^{\text{op}} \in \Sigma\mathcal{C} = \text{RMod}_{\mathcal{C}}((n+1)\text{Vec})$ and $\mathcal{D} = \Omega_{\mathcal{M}^{\text{op}}}(\Sigma\mathcal{C})$. In $\Sigma\mathcal{D}$, D^{n+1} is labeled by $\bullet \in \Sigma\mathcal{D}$; C^{n+1} is labeled by $\mathcal{M} \in \Sigma\mathcal{D} = \text{RMod}_{\mathcal{D}}((n+1)\text{Vec})$ and $\mathcal{C} = \Omega_{\mathcal{M}}(\Sigma\mathcal{D})$. Moreover, when d is a higher morphism in $\mathcal{D}$, the curly brace in above picture automatically represents a higher morphism $d \in \text{Mod}_A^{E_1}(\mathcal{C}) \subset \mathcal{C}$. When d is a 1-morphism in $\mathcal{D}$, the curly brace as a 1-morphism clearly intertwines the two-side A -actions. $\diamond$

⁴¹In 2-categories, the internal hom algebra is automatically equipped with not only the unit and associativity 1-morphisms but also necessary 2-morphisms, which are automatically included as the defining data of the universal property of the internal hom in 2-categories.

From a single C^{n+1} , one can obtain all D^{n+1} that are Morita equivalent to C^{n+1} . In particular, all anomaly-free non-chiral $n+1$ D topological orders can be obtained by condensing 1-codimensional topological defects in 1^{n+1} , or equivalently, by condensing condensable E_1 -algebras in $n\text{Vec}$. We generalize it to a more general situation. This generalized result is essentially a reformulation of a part of Proposition* 3.13 in [KZ24] (see also Remark 3.3.15), and a reformulation of Theorem^{ph} 2.2.26 and Theorem 2.2.42, and was obtained independently in the third arXiv version of [LYW24].

Theorem^{ph} 3.3.14. Any gapped boundary N^n of an $n+1$ D anomaly-free simple topological order C^{n+1} can be obtained from a single gapped boundary M^n by first stacking an anomaly-free n D topological order X^n to M^n then condensing a 1-codimensional topological defect in $M^n \boxtimes X^n$.

Mathematically, it means that two indecomposable multi-fusion $(n-1)$ -categories $\mathcal{M}$ and $\mathcal{N}$ for $n \geq 2$ share the same E_1 -center, i.e., $\mathfrak{Z}_1(\mathcal{M}) \simeq \mathfrak{Z}_1(\mathcal{N})$ if and only if $\mathcal{N}$ is Morita equivalent to $\mathcal{M} \boxtimes \mathcal{X}$ for a non-degenerate multi-fusion $(n-1)$ -category $\mathcal{X}$, or equivalently, $\Sigma\mathcal{N} \simeq \Sigma\mathcal{M} \boxtimes \mathcal{S}$ for an invertible separable n -category $\mathcal{S}$ (this reformulation first appeared in [LYW24]). $\square$

Remark 3.3.15. This result follows from the proof of (1) $\Rightarrow$ (3) part of Proposition* 3.13 in [KZ24], which says that the center functor $\text{Sep}_n^{\text{ind}} \rightarrow \mathcal{Fus}_n^{\text{cl}}$ is full up to a factor of invertible separable n -categories. We can further reformulate it in terms of MC^n and $\text{MC}^n(C^{n+1})$ as in Theorem^{ph} 2.2.26. $\diamond$

Remark 3.3.16. Theorem^{ph} 3.3.14 automatically covers the same result for the gapped domain walls between two topological orders because a wall is a boundary by the folding trick. Since gapped domain walls between two Morita equivalence type-I topological orders A^{n+1} and B^{n+1} determine condensations from A^{n+1} to B^{n+1} , the notion of Morita equivalence between two gapped domain walls naturally carries over to the notion of Morita equivalence between two condensable E_1 -algebras in $\mathcal{A}$. This leads to an interesting way to classify condensations from A^{n+1} to B^{n+1} via the center functor as we show in Theorem^{ph} 3.3.33 and Corollary^{ph} 3.3.36 and 3.3.37. $\diamond$

We have assumed ‘type-I’ condition on C^{n+1} in previous discussion. Relaxing this condition gives essentially the same result as Theorem^{ph} 3.1.13. For readers convenience, we write it down explicitly.

Theorem^{ph} 3.3.17. In general, C^{n+1} could be type-I or type-II and $\mathcal{C}$ is a multi-fusion n -category. A condensable algebra A in a multi-fusion n -category $\mathcal{C}$ is a direct sum of non-zero condensable algebras in each indecomposable summand of $\mathcal{C}$. Condensations preserve the multiplicity.

1. C^{n+1} is type-I (i.e., $\mathcal{C}$ is an indecomposable multi-fusion n -category):
 - (a) If A is indecomposable, then the condensed phase D^{n+1} is type-I and simple (i.e., $\mathcal{D}$ is a fusion n -category), and $\mathcal{M} = \text{RMod}_A(\mathcal{C})$ is an indecomposable left $\mathcal{C}$ -module.
 - (b) If A is a direct sum of indecomposable ones, i.e., $A = \oplus_{i=1}^l A_i$, then the condensed phase D^{n+1} is type-I composite, $\text{gsd}(D^{n+1}) = l$, and $\mathcal{D}$ is an indecomposable multi-fusion n -category, which can be decomposed as a direct sum of separable n -categories: $\mathcal{D} = \oplus_{i,j=1}^l \text{BMod}_{A_i|A_j}(\mathcal{C})$. The domain wall $M^n = (\mathcal{M}, A)$ is of type-II and we have the decomposition of left $\mathcal{C}$ -modules: $\mathcal{M} = \text{RMod}_A(\mathcal{C}) = \oplus_{i=1}^l \text{RMod}_{A_i}(\mathcal{C})$.
2. C^{n+1} is type-II (i.e., $\mathcal{C} = \oplus_{i=1}^k \mathcal{C}_i$ and $\mathcal{C}_i$ are indecomposable multi-fusion n -categories):
 - (a) In this case, we have $A = \oplus_{i=1}^k A_i$, where $A_i = A \cap \mathcal{C}_i \neq 0$.
 - (b) The condensed phase D^{n+1} is type-II, $\text{mult}(C^2) \simeq \text{mult}(D^2)$, and $\mathcal{D} \simeq \oplus_{i=1}^k \text{BMod}_{A_i|A_i}(\mathcal{C}_i)$.
 - (c) The domain wall M^n is type-II, and we have the decomposition of left $\mathcal{C}$ -modules: $\mathcal{M} = \text{RMod}_A(\mathcal{C}) = \oplus_{i=1}^k \text{RMod}_{A_i}(\mathcal{C}_i)$. $\square$

Since all the subtleness associated to the type-I or type-II composite property are summarized in above theorem, from now on, we mainly focus on the condensations in type-I or simple topological orders.

3.3.2 General constructions

In this subsection, we provide some general examples, in which we either provide a direct construction of condensable E_1 -algebras in $\mathcal{C}$ then determine $\mathcal{D}$ and $\mathcal{M}$ (as in Theorem^{ph} 3.3.2), or provide $\mathcal{C}$ and $\mathcal{M}$ with concrete coordinate systems such that the condensable E_1 -algebra $[m, m]_{\mathcal{C}}$ in $\mathcal{C}$ can be determined explicitly (as in Theorem^{ph} 3.3.9).

When $\mathcal{C}^{n+1} = \mathbf{1}^{n+1}$, a condensable E_1 -algebra in $n\text{Vec}$ is precisely a multi-fusion $(n-1)$ -category $\mathcal{A}$. By condensing it, we obtain the condensed phase $\mathcal{D}^{n+1}$ and a gapped domain wall M^n such that

$$\mathcal{D} = \text{Mod}_{\mathcal{A}}^{E_1}(n\text{Vec}), \quad (\mathcal{M}, m) = (\text{RMod}_{\mathcal{A}}(n\text{Vec}), \mathcal{A}) = (\Sigma\mathcal{A}, \mathcal{A}), \quad \mathcal{D}^{\text{op}} \simeq \mathfrak{Z}_0(\Sigma\mathcal{A}).$$

We want to understand this result from a different perspective in the cases (1) and (2) below.

(1). $\mathcal{C}^{n+1} = \mathcal{D}^{n+1} = \mathbf{1}^{n+1}$: It turns out that this seemingly “trivial” case can have non-trivial condensations. In this case, a gapped domain wall M^n between $\mathcal{C}^{n+1}$ and $\mathcal{D}^{n+1}$ is nothing but an anomaly-free nD topological order. Again, $M^n = (\mathcal{M}, m)$, where $\mathcal{M}$ is a separable n -category, m is a distinguished object in $\mathcal{M}$ and labels the nD anomaly-free topological order M^n as a boundary condition of $\mathbf{1}^{n+1}$. By boundary-bulk relation, we must have

$$\mathfrak{Z}_0(\mathcal{M}, m) = \text{Fun}(\mathcal{M}, \mathcal{M}) \simeq n\text{Vec}. \quad (3.3.9)$$

As a consequence, $\mathcal{M}$ is necessarily indecomposable (actually invertible in $(n+1)\text{Vec}$ by [KZ24, Corollary 2.10]). The category of topological defects on M^n is $\Omega_m\mathcal{M}$, which is necessarily an indecomposable multi-fusion $(n-1)$ -category. Since $\mathcal{M}$ is indecomposable, we have $\mathcal{M} = \Sigma\Omega_m\mathcal{M}$. By boundary-bulk relation, we have

$$\mathfrak{Z}_1(\Omega_m\mathcal{M}) \simeq (n-1)\text{Vec},$$

which is an immediate consequence of (3.3.9), $\mathcal{M} = \Sigma\Omega_m\mathcal{M}$ and (2.3.15).

Note that the equivalence $\mathcal{M} = \Sigma\Omega_m\mathcal{M} \simeq \text{RMod}_{\Omega_m\mathcal{M}}(n\text{Vec})$ provides a coordinate system that allows us to identify the condensable E_1 -algebra $[m, m]_{\mathcal{C}}$ in $\mathcal{C}$ explicitly. Indeed, in this coordinate system $m = \Omega_m\mathcal{M} \in \text{RMod}_{\Omega_m\mathcal{M}}(n\text{Vec})$, and the left bulk-to-wall functor can be rewritten as follows.

$$L_1 = - \odot m = - \otimes \Omega_m\mathcal{M} : n\text{Vec} \rightarrow \text{RMod}_{\Omega_m\mathcal{M}}(n\text{Vec}),$$

which is illustrated in the following picture.

Then its right adjoint functor is precisely the forgetful functor, i.e.,

$$L_1^R = f : \text{RMod}_{\Omega_m\mathcal{M}}(n\text{Vec}) \rightarrow n\text{Vec}.$$

Therefore, $L_1^R(\Omega_m\mathcal{M}) = [\Omega_m\mathcal{M}, \Omega_m\mathcal{M}]_{n\text{Vec}} = \Omega_m\mathcal{M}$ is precisely the indecomposable condensable E_1 -algebra in $n\text{Vec}$ that defines the condensation. Note that, in the case, the internal hom $[m, m]_{\mathcal{C}}$ is no longer an abstract nonsense. It is the indecomposable multi-fusion $(n-1)$ -category $\Omega_m\mathcal{M}$, which is precisely an indecomposable condensable E_1 -algebra in $\mathcal{C} = n\text{Vec}$. By Theorem^{ph} 3.3.9, we obtain

$$\mathcal{D} = n\text{Vec} \simeq \text{Mod}_{\Omega_m\mathcal{M}}^{E_1}(n\text{Vec}), \quad \mathcal{M} \simeq \text{RMod}_{\Omega_m\mathcal{M}}(n\text{Vec}),$$

where the second equivalence is tautological and can be viewed as a consistency check, but the first monoidal equivalence is non-trivial mathematically. We reformulate this result mathematically.

Proposition^o 3.3.18. A Lagrangian E_1 -algebra in $n\text{Vec}$ (recall Definition 2.3.31) is precisely a non-degenerate multi-fusion $(n-1)$ -category $\mathcal{A}$. In this case, we have

$$\text{Mod}_{\mathcal{A}}^{E_1}(n\text{Vec}) \simeq n\text{Vec}. \quad (3.3.10)$$

Remark 3.3.19. The subscript of L_1 represents that it is left bulk-to-wall map of 1-codimensional topological defects. We denote the left bulk-to-wall map of k -codimensional topological defects by $L_k : \Omega^{k-1}\mathcal{C} \rightarrow \Omega^{k-1}\mathcal{M}$ and the right bulk-to-wall map of k -codimensional topological defect by $R_k : \Omega^{k-1}\mathcal{D} \rightarrow \Omega^{k-1}\mathcal{M}$. $\diamond$

Remark 3.3.20. Proposition 3.3.18 is the mathematical formulation of the physical fact that condensing an anomaly-free n D topological order $\mathcal{M}^n$, viewed as 1-codimensional topological defects in the trivial phase $\mathbf{1}^{n+1}$, reproduces the trivial phase $\mathbf{1}^{n+1}$, and gapped domain wall of this condensation is precisely the anomaly-free n D topological order $\mathcal{M}^n$. $\diamond$

(2). $\mathcal{C}^{n+1} = \mathbf{1}^{n+1} \neq \mathcal{D}^{n+1}$: In this case, the discussion is completely parallel to the previous case. In particular, $\mathcal{M}$ is a separable n -category and $\Omega_m\mathcal{M}$ is a multi-fusion $(n-1)$ -category. If m is non-zero in each connected component of $\mathcal{M}$, then we obtain

$$\mathcal{D} \simeq \text{Mod}_{\Omega_m\mathcal{M}}^{\mathbf{E}_1}(n\text{Vec}) \simeq \text{Fun}(\mathcal{M}, \mathcal{M})^{\text{op}} \simeq \mathfrak{Z}_0(\mathcal{M}, m)^{\text{op}}, \quad \mathcal{M} = \Sigma\Omega_m\mathcal{M} = \text{RMod}_{\Omega_m\mathcal{M}}(n\text{Vec}).$$

We reformulate the associated mathematical result as a generalization of Proposition 3.3.18.

Proposition 3.3.21. A condensable $\mathbf{E}_1$ -algebra in $n\text{Vec}$ is precisely a multi-fusion $(n-1)$ -category $\mathcal{A}$. We have

$$\text{Mod}_{\mathcal{A}}^{\mathbf{E}_1}(n\text{Vec})^{\text{op}} \simeq \mathfrak{Z}_0(\Sigma\mathcal{A}). \quad (3.3.11)$$


If $\mathcal{A}$ is an indecomposable multi-fusion $(n-1)$ -category, then we have [KZ22b]

$$\text{Mod}_{\mathcal{A}}^{\mathbf{E}_1}(n\text{Vec})^{\text{op}} \simeq \mathfrak{Z}_0(\Sigma\mathcal{A}) \simeq \Sigma\mathfrak{Z}_1(\mathcal{A}). \quad (3.3.12)$$

Remark 3.3.22. The Levin-Wen model [LW05] can be viewed as a physical way to realize the condensation defined by condensing a simple condensable $\mathbf{E}_1$ -algebra $\mathcal{A}$ in 2Vec (i.e., $\mathcal{A}$ is a fusion 1-category) with a gapped domain wall $\mathcal{M}^2 = (\Sigma\mathcal{A}, \mathcal{A})$ constructed in [KK12]. Similarly, a higher dimensional generalization of Levin-Wen model based on indecomposable multi-fusion $(n-1)$ -category $\mathcal{A}$ (predicted and briefly described in [KZ22b, Remark 5.11]) should be viewed as a physical way to realize the condensation defined by the indecomposable condensable $\mathbf{E}_1$ -algebra $\mathcal{A}$ in $n\text{Vec}$. $\diamond$

Remark 3.3.23. The layer construction of 3+1D topological orders via layers of 2+1D topological orders [JQ14] provides a concrete realization of a condensation of 1-codimensional defects in $\mathbf{1}^4$. For example, the layer construction of the 3+1D $\mathbb{Z}_2$ topological order [ZLZH⁺23] should be viewed as a physical realization of the 1-codimensional condensation that produces the 3+1D $\mathbb{Z}_2$ topological order from $\mathbf{1}^4$. $\diamond$

(3). $\mathcal{Z}(\mathcal{C})^{n+2} = \mathbf{1}^{n+2}$: We further assume that $\mathcal{C}^{n+1}$ is simple. In this case, $\mathcal{C} = \Sigma\Omega\mathcal{C} = \text{RMod}_{\Omega\mathcal{C}}(n\text{Vec})$ provides a concrete coordinate system to the non-degenerate fusion n -category $\mathcal{C}$. By Remark 2.2.33, the fusion product in this coordinate system is defined by $\mathcal{K} \otimes^1 \mathcal{L} := \mathcal{K} \boxtimes_{\Omega\mathcal{C}} \mathcal{L}$ as depicted in the following picture.



The right $\Omega\mathcal{C}$ -module structures and tensor product:

- (1). $\mathcal{K} \times \Omega\mathcal{C} \xrightarrow{\circ} \mathcal{K}; \quad (k, a) \mapsto k \otimes^2 a;$
- (2). $\mathcal{L} \times \Omega\mathcal{C} \xrightarrow{\circ} \mathcal{L}; \quad (l, a) \mapsto l \otimes^2 a;$
- (3). $\boxtimes_{\Omega\mathcal{C}} : \mathcal{K} \times \mathcal{L} \rightarrow \mathcal{K} \boxtimes_{\Omega\mathcal{C}} \mathcal{L}, \quad (k, l) \mapsto k \boxtimes_{\Omega\mathcal{C}} l.$

(3.3.13)

In this case, it is possible to classify all the condensable $\mathbf{E}_1$ -algebras in $\mathcal{C}$ via the following equivalence⁴²:

$$\text{Alg}_{\mathbf{E}_1}^{\mathcal{C}}(\text{RMod}_{\Omega\mathcal{C}}(n\text{Vec})) \simeq \text{LMod}_{\Omega\mathcal{C}}(\text{Alg}_{\mathbf{E}_1}^{\mathcal{C}}(n\text{Vec})). \quad (3.3.14)$$

More explicitly, an indecomposable condensable $\mathbf{E}_1$ -algebra $\mathcal{A}$ in $\mathcal{C}$ is precisely an indecomposable multi-fusion $(n-1)$ -category $\mathcal{A}$ equipped with a monoidal action functor $\odot : \Omega\mathcal{C} \times \mathcal{A} \rightarrow \mathcal{A}$ that defines the structure of left $\Omega\mathcal{C}$ -module on $\mathcal{A}$, or equivalently, an indecomposable multi-fusion $(n-1)$ -category $\mathcal{A}$ equipped with a central functor $\Omega\mathcal{C} \rightarrow \mathcal{A}$ (or equivalently, a braided monoidal functor $\phi : \Omega\mathcal{C} \rightarrow \mathfrak{Z}_1(\mathcal{A})$)⁴³.

⁴²The proof of this equivalence was known for the $n = 2$ case in [BJS21]. The same proof works for general n . We provide a physical proof of this result in (3.3.15), (3.3.16) and (3.3.17).

⁴³When $n = 2$, such $\mathcal{A}$ is also called a left multi-fusion $\Omega\mathcal{C}$ -module [KZ18a] or a module tensor category [HPT16].

Example 3.3.24. For $A \in \text{Alg}_{\mathbb{E}_2}^c(\Omega\mathcal{C})$ (see Section 4.1), we have $\text{RMod}_A(\Omega\mathcal{C}) \in \text{LMod}_{\Omega\mathcal{C}}(\text{Alg}_{\mathbb{E}_1}^c(n\text{Vec}))$. This example is important later. $\heartsuit$


For an indecomposable condensable $\mathbb{E}_1$ -algebra $\mathcal{A}$ in $\mathcal{C}$, we can condense it and obtain a condensed phase $\mathcal{D}^{n+1}$, which is also simple. We have

$$\begin{aligned} \mathcal{D} &\simeq \text{Mod}_{\mathcal{A}}^{\mathbb{E}_1}(\mathcal{C}) \simeq \Sigma\mathfrak{Z}_2(\Omega\mathcal{C}, \mathfrak{Z}_1(\mathcal{A})), & \Omega\mathcal{D} &\simeq \mathfrak{Z}_2(\Omega\mathcal{C}, \mathfrak{Z}_1(\mathcal{A})); \\ \mathcal{M} &\simeq \text{RMod}_{\mathcal{A}}(\mathcal{C}) \simeq \text{RMod}_{\mathcal{A}}(n\text{Vec}), & m = \mathcal{A} &\in \text{RMod}_{\mathcal{A}}(\mathcal{C}), \end{aligned}$$

where the last “ $\simeq$ ” is due to the fact that the right $\Omega\mathcal{C}$ -action factors through the right $\mathcal{A}$ -action. Moreover, we have

$$\Omega_m\mathcal{M} = \text{hom}_{\text{RMod}_{\mathcal{A}}(\mathcal{C})}(\mathcal{A}, \mathcal{A}) \simeq \text{Fun}_{\mathcal{A}^{\text{op}}}(\mathcal{A}, \mathcal{A}) \simeq \mathcal{A}.$$


It means that $\mathcal{A}$ is precisely the category of topological defects on the gapped wall $(\mathcal{M}, m)$ between $\mathcal{C}^{n+1}$ and $\mathcal{D}^{n+1}$. This fact reveals the geometric relation between $\Omega\mathcal{C}, \mathcal{A}, \Omega\mathcal{D}$ as illustrated in the following picture.


(3.3.15)

Note that the 1-codimensional defect in $\mathcal{C}$ obtained by horizontally squeezing the $\Omega\mathcal{D}$ -region in the second picture in (3.3.15) is labeled by the object $\Omega_m\mathcal{M} \in \mathcal{C} = \text{RMod}_{\Omega\mathcal{C}}(n\text{Vec})$; and higher codimensional topological defects on it form the following multi-fusion $(n-1)$ -category:

$$\Omega_{\Omega_m\mathcal{M}}(\mathcal{C}) \simeq \Omega_m\mathcal{M} \boxtimes_{\Omega\mathcal{D}} \Omega_m\mathcal{M}^{\text{op}} \simeq \text{Fun}_{\Omega\mathcal{C}^{\text{op}}}(\Omega_m\mathcal{M}, \Omega_m\mathcal{M}).$$

In general, when an object $\Omega_m\mathcal{M}$ is viewed as an algebra in $\mathcal{C} = \text{RMod}_{\Omega\mathcal{C}}(n\text{Vec})$, its algebraic structure does not necessarily coincide with the monoidal structure on $\Omega_m\mathcal{M}^{\text{op}}$ defined by the particle fusions along the wall $(\mathcal{M}, m)$. However, in this case, two structures indeed coincide as illustrated in the following picture.


(3.3.16)

Indeed, the two top blue dots labeled by $\Omega_m\mathcal{M}$ (or better the top dashed box) represent the object $\Omega_m\mathcal{M} \boxtimes_{\Omega\mathcal{C}} \Omega_m\mathcal{M}$ in $\mathcal{C} = \text{RMod}_{\Omega\mathcal{C}}(n\text{Vec})$ because (recall Remark 2.2.33)

$$\text{hom}_{\mathcal{C}}(\mathbb{1}_{\mathcal{C}}, \Omega_m\mathcal{M} \boxtimes_{\Omega\mathcal{C}} \Omega_m\mathcal{M}) \simeq \text{Fun}_{\Omega\mathcal{C}^{\text{op}}}(\Omega\mathcal{C}, \Omega_m\mathcal{M} \boxtimes_{\Omega\mathcal{C}} \Omega_m\mathcal{M}) \simeq \Omega_m\mathcal{M} \boxtimes_{\Omega\mathcal{C}} \Omega_m\mathcal{M}.$$

The three lower small circles labeled by $1_m \in \Omega_m\mathcal{M}$ (or better the lower dashed box) define a 1-morphism $\Omega_m\mathcal{M} \boxtimes_{\Omega\mathcal{C}} \Omega_m\mathcal{M} \rightarrow \Omega_m\mathcal{M}$, which defines the algebraic structure on the 1-codimensional defect labeled by $\Omega_m\mathcal{M}$. Using the physical intuitions explained in Remark 2.2.33, we see that it induces on $\text{hom}_{\mathcal{C}}(\mathbb{1}_{\mathcal{C}}, \Omega_m\mathcal{M}) = \Omega_m\mathcal{M}$ an algebraic structure, i.e., $\Omega_m\mathcal{M} \boxtimes_{\Omega\mathcal{C}} \Omega_m\mathcal{M} \rightarrow \Omega_m\mathcal{M}$, defined by fusing two dashed boxes, which defines the following functor,

$$a \boxtimes_{\Omega\mathcal{C}} b \mapsto 1_m \otimes a \otimes 1_m \otimes b \otimes 1_m \simeq a \otimes b. \quad (3.3.17)$$

This means that such defined algebraic structure on $\Omega_m\mathcal{M}$ in $\mathcal{C} = \Sigma\Omega\mathcal{C}$ coincides with the monoidal structure (i.e., particle fusions) on $\Omega_m\mathcal{M}$. This fact provides a physical proof of the equivalence (3.3.14).

Remark 3.3.25. In this remark, we emphasize the significance of the physical proof of (3.3.14). This physical proof immediately provides us a geometric intuition which is ready to be generalized. Indeed, from the details of above arguments, it is clear that the geometric intuition of proof is so robust that it is insensitive to the assumption that all $\mathcal{C}^{n+1}, \mathcal{M}^n, \mathcal{D}^{n+1}$ are gapped. Therefore, we expect the equivalence (3.3.14) to be a general principle that also applies to gapless quantum phases (see Remark 6.3.7). $\diamond$

By choosing different $\mathcal{M}, m, \mathcal{D}$'s, we recover all possible condensations. It means that we can reverse the story. More explicitly, given $M^n = (\mathcal{M}, m)$, where $\mathcal{M}$ is a separable n -category and $\Omega_m \mathcal{M}$ is a multi-fusion $(n-1)$ -category. We assume that m is non-zero in each indecomposable left $\mathcal{C}$ -submodule of $\mathcal{M}$. By moving 2-codimensional topological defects in C^{n+1} onto the wall M^n , we obtain an action $\Omega \mathcal{C} \times \Omega_m \mathcal{M} \rightarrow \Omega_m \mathcal{M}$, which is automatically monoidal, i.e., $\Omega_m \mathcal{M}$ is monoidal and the monoidal structure is compatible with the $\Omega \mathcal{C}$ -action. As a consequence, we have

$$\Omega_m \mathcal{M} \in \text{LMod}_{\Omega \mathcal{C}}(\text{Alg}_{E_1}^{\mathcal{C}}(n\text{Vec})) \simeq \text{Alg}_{E_1}^{\mathcal{C}}(\text{RMod}_{\Omega \mathcal{C}}(n\text{Vec})).$$

In other words, $\Omega_m \mathcal{M}$ is a condensable E_1 -algebra in $\mathcal{C} = \Sigma \Omega \mathcal{C}$. Moreover, $[m, m]_{\mathcal{C}}$ can be identified with $\Omega_m \mathcal{M}$, and we have

$$\mathcal{D} \simeq \text{Mod}_{\Omega_m \mathcal{M}}^{E_1}(\mathcal{C}), \quad \mathcal{M} \simeq \text{RMod}_{\Omega_m \mathcal{M}}(\mathcal{C}) \simeq \text{RMod}_{\Omega_m \mathcal{M}}(n\text{Vec}). \quad (3.3.18)$$

In this case, we also have $\mathcal{D}^{\text{op}} \simeq \text{Mod}_{\Omega_m \mathcal{M}}^{E_1}(\mathcal{C})^{\text{op}} \simeq \mathfrak{Z}_1(\mathcal{C}, \mathfrak{Z}_0(\text{RMod}_{\Omega_m \mathcal{M}}(\mathcal{C})))$. We summarize the key point in this case in the following theorem.

Theorem^{ph} 3.3.26. When C^{n+1} is anomaly-free and simple, an indecomposable condensable E_1 -algebra in the fusion n -category $\mathcal{C} = \Sigma \Omega \mathcal{C}$ is precisely given by an indecomposable multi-fusion $(n-1)$ -category $\mathcal{A}$ equipped with a braided monoidal functor $\phi : \Omega \mathcal{C} \rightarrow \mathfrak{Z}_1(\mathcal{A})$ (or equivalently, by a type-I gapped domain wall between C^{n+1} and the condensed phase obtained by condensing $\mathcal{A}$). Moreover, we have the following equivalence of categories:

$$\text{Alg}_{E_1}^{\mathcal{C}}(\text{RMod}_{\Omega \mathcal{C}}(n\text{Vec})) \simeq \text{LMod}_{\Omega \mathcal{C}}(\text{Alg}_{E_1}^{\mathcal{C}}(n\text{Vec})). \quad (3.3.19)$$

By condensing $\mathcal{A}$ in $\mathcal{C}$, we obtain a condensed phase D^{n+1} and a gapped domain wall M^n such that

$$\begin{aligned} \mathcal{D} &\simeq \text{Mod}_{\mathcal{A}}^{E_1}(\mathcal{C}) \simeq \Sigma \mathfrak{Z}_2(\Omega \mathcal{C}, \mathfrak{Z}_1(\mathcal{A})), & \Omega \mathcal{D} &\simeq \mathfrak{Z}_2(\Omega \mathcal{C}, \mathfrak{Z}_1(\mathcal{A})); \\ \mathcal{M} &\simeq \text{RMod}_{\mathcal{A}}(\mathcal{C}) \simeq \text{RMod}_{\mathcal{A}}(n\text{Vec}), & m &= \mathcal{A} \in \text{RMod}_{\mathcal{A}}(\mathcal{C}), \end{aligned}$$


Moreover, $\mathcal{A} = \Omega_m \mathcal{M}$ provides the physical meaning of $\mathcal{A}$ as the $(n-1)$ -category of topological defects on the wall $M^n = (\mathcal{M}, m)$. If ϕ is an equivalence, then $\mathcal{A}$ is “Lagrangian” in the sense that $D^{n+1} = \mathbf{1}^{n+1}$. $\square$

Remark 3.3.27. When C^{n+1} is anomaly-free and simple, above theorem allows us to define the Morita equivalence between two indecomposable E_1 -algebras in $\mathcal{C}$ according to that between two type-I gapped domain walls. When C^{n+1} is anomalous and simple, in general, (3.3.19) only gives a subset of all condensable E_1 -algebras in $\mathcal{C}$ because $\Sigma \Omega \mathcal{C}$ is only a fusion full subcategory of $\mathcal{C}$. $\diamond$


Remark 3.3.28. In the orbifold of a 2+1D Reshetikhin-Turaev TQFT [CRS20], the initial data of this orbifold construction is called ‘orbifold data’, which consists of (1) $\mathcal{K}$ a modular tensor category that defines the initial 2+1D Reshetikhin-Turaev TQFT; (2) A is a Δ -separable symmetric Frobenius algebra in $\mathcal{K}$; (3) $T = {}_A T_{AA}$ is an A - $A \otimes A$ -bimodule in $\mathcal{K}$; (4) certain choices of morphisms, satisfying conditions in [CRS20, Eq. (3.16-3.20)]. We were informed by Ingo Runkel that it is possible to prove that this data, together with condition (3.16) & (3.17) in [CRS20], amounts to say that $\text{LMod}_{\mathcal{A}}(\mathcal{K}) \in \text{LMod}_{\mathcal{K}}(\text{Alg}_{E_1}^{\mathcal{C}}(2\text{Vec}))$. If this is true, it suggests that this ‘orbifold construction’ should be viewed as the counterpart (in the TQFT framework) of our 1-codimensional defect condensation theory. We hope to see more detailed clarification of their relation in the future. $\diamond$

By Theorem^{ph} 2.4.18 and Remark 2.4.20, we obtain a classification result for 1-codimensional defect condensations between two (potentially chiral) anomaly-free simple topological orders.

Theorem^{ph} 3.3.29. Given two Morita equivalent anomaly-free simple topological orders C^{n+1} and D^{n+1} connected by a gapped domain wall K^n as illustrated in the first picture below. Using the folding trick and topological Wick rotation (TWR) (recall Figure 9), we obtain a one-to-one correspondence between



$\xrightarrow{\text{folding + TWR}}$



$(3.3.20)$

- $(\mathcal{K} \boxtimes \mathcal{T})$ -modules modulo the equivalence relation: $\mathcal{X} \sim \mathcal{Y}$ if $\mathcal{Y} \simeq \mathcal{X} \boxtimes \mathcal{V}$ for $\mathcal{V} \in (n+1)\text{Vec}^\times$;
- indecomposable condensable E_1 -algebras in $\mathcal{C}$ that define 1-codimensional defect condensations from $\mathcal{C}^{n+1}$ to $\mathcal{D}^{n+1}$ in the Morita class $[K^n \boxtimes T^n]$.

defined by $\mathcal{X} \mapsto \text{Fun}_{\mathcal{K} \boxtimes \mathcal{T}}(\mathcal{X}, \mathcal{X})^{\text{op}}$. The condensable E_1 -algebra $A = \text{Fun}_{\mathcal{K} \boxtimes \mathcal{T}}(\mathcal{X}, \mathcal{X})^{\text{op}}$ is simple if and only if the $(\mathcal{K} \boxtimes \mathcal{T})$ -module $\mathcal{X}$ is indecomposable. $\square$

Remark 3.3.30. Theorem^{ph} 3.3.29 reduces the problem of classifying 1-codimensional defect condensations between two chiral topological orders to the problem of classifying gapped boundaries of a non-chiral topological order. Unfortunately, not much chiral topological orders in higher dimensions are known. We will return to the problem of constructing interesting chiral examples in the future. In this work, we focus on examples from non-chiral topological orders. $\diamond$

(4). $\mathcal{C}^{n+1} = \mathcal{Z}(\mathcal{B})^{n+1}$: In this case, $\mathcal{C}^{n+1}$ is anomaly-free and non-chiral, and admits a gapped boundary $\mathcal{B}^n$, which is assumed to be simple. Then $\mathcal{C}^{n+1}$ is also simple. Therefore, this case is just a special case of (3). However, in this case, $\mathcal{C}$ has another natural coordinate system given by

$$\mathcal{C} = \text{BMod}_{\mathcal{B}^{\text{op}}|\mathcal{B}^{\text{op}}}(n\text{Vec}) \simeq \text{BMod}_{\mathcal{B}|\mathcal{B}}(n\text{Vec})^{\text{op}},$$

where $\mathcal{B}$ is the fusion $(n-1)$ -category of all topological defects in $\mathcal{B}^n$. An indecomposable condensable E_1 -algebra in $\mathcal{C}$ is precisely an indecomposable multi-fusion $(n-1)$ -category $\mathcal{A}^{\text{op}}$ equipped with a monoidal functor $\psi : \mathcal{B} \rightarrow \mathcal{A}$. By condensing the condensable E_1 -algebra $\mathcal{A}^{\text{op}}$ in $\mathcal{C}$, we obtain the condensed phase $\mathcal{D}^{n+1}$ and $\mathcal{M}^n$ as follows:

$$\mathcal{D} = \text{Mod}_{\mathcal{A}^{\text{op}}}^{\mathcal{E}_1}(\mathcal{C}) \simeq \text{Mod}_{\mathcal{A}^{\text{op}}}^{\mathcal{E}_1}(n\text{Vec}) \simeq \Sigma \mathfrak{Z}_1(\mathcal{A}), \quad (3.3.21)$$

$$(\mathcal{M}, m) = (\text{LMod}_{\mathcal{A}}(\mathcal{C}), \mathcal{A}) = (\text{BMod}_{\mathcal{A}|\mathcal{B}}(n\text{Vec}), \mathcal{A}), \quad (3.3.22)$$

where the first “ $\simeq$ ” is due to the fact that the $\mathcal{B}$ -module structure factors through the monoidal functor ψ . As a consequence, we obtain

$$\Omega \mathcal{D} \simeq \mathfrak{Z}_1(\mathcal{A}), \quad \Omega_m \mathcal{M} \simeq \text{Fun}_{\mathcal{A}|\mathcal{B}}(\mathcal{A}, \mathcal{A}). \quad (3.3.23)$$

Note that, according to [KZ18a, Theorem 3.2.3][KZ22b], $\text{Fun}_{\mathcal{B}|\mathcal{A}^{\text{op}}}(\mathcal{A}, \mathcal{A})$ is precisely the closed $\Omega \mathcal{D}^{\text{op}}$ - $\Omega \mathcal{C}$ -bimodule that defines alternatively the monoidal functor $\psi : \mathcal{B}^{\text{op}} \rightarrow \mathcal{A}^{\text{op}}$ as the following composed functor (with an illustration of its physical meaning).

$$\mathcal{B}^{\text{op}} \rightarrow \mathcal{B}^{\text{op}} \boxtimes_{\mathfrak{Z}_1(\mathcal{B})} \text{Fun}_{\mathcal{A}|\mathcal{B}}(\mathcal{A}, \mathcal{A}) \simeq \text{Fun}_{\mathcal{A}}(\mathcal{A}, \mathcal{A}) \simeq \mathcal{A}^{\text{op}}. \quad (3.3.24)$$

Remark 3.3.31. It is very illuminating to see the connection between the two coordinate systems in this case. By [KLWZZ20b], these two coordinate systems $\mathcal{C} = \text{RMod}_{\Omega \mathcal{C}}(n\text{Vec})$ and $\mathcal{C} = \text{BMod}_{\mathcal{B}^{\text{op}}|\mathcal{B}^{\text{op}}}(n\text{Vec})$ are related by the following monoidal equivalence:

$$\begin{aligned} \text{RMod}_{\Omega \mathcal{C}}(n\text{Vec}) &\xrightarrow{\simeq} \text{BMod}_{\mathcal{B}^{\text{op}}|\mathcal{B}^{\text{op}}}(n\text{Vec}) \\ \mathcal{K} &\mapsto \mathcal{K} \boxtimes_{\mathfrak{Z}_1(\mathcal{B})} \mathcal{B}. \end{aligned}$$


Using this equivalence, we immediately see that the relation between $\Omega_m \mathcal{M} \in \text{Alg}_{E_1}^c(\text{RMod}_{\Omega \mathcal{C}}(n\text{Vec}))$ and its corresponding condensable E_1 -algebra $\mathcal{A}^{\text{op}} \in \text{Alg}_{E_1}^c(\text{BMod}_{\mathcal{B}^{\text{op}}|\mathcal{B}^{\text{op}}}(n\text{Vec}))$ is precisely the one illustrated in the picture in (3.3.24). $\diamond$

This result provides a powerful tool to construct condensable 1-codimensional topological defects in a non-chiral topological order, or equivalently, condensable E_1 -algebra in a fusion n -category obtained from the delooping of a non-degenerate and non-chiral braided fusion $(n-1)$ -category. We summarize it as a theorem.

Theorem^{ph} 3.3.32. For a non-chiral simple topological order $C^{n+1} = Z(B)^{n+1}$, all indecomposable condensable E_1 -algebras in the fusion n -category $\mathcal{C} = \text{BMod}_{\mathcal{B}|\mathcal{B}}(n\text{Vec})^{\text{op}}$ are given by an indecomposable multi-fusion n -category $\mathcal{A}^{\text{op}}$ equipped with a monoidal functor $\mathcal{B} \rightarrow \mathcal{A}$. By condensing $\mathcal{A}^{\text{op}}$ in $\mathcal{C}$, we obtain the condensed topological order $D^{n+1} = Z(A)^{n+1}$ and a gapped domain M^n such that

$$\begin{aligned} \mathcal{D} &\simeq \text{Mod}_{\mathcal{A}^{\text{op}}}^{E_1}(\mathcal{C}) \simeq \text{Mod}_{\mathcal{A}^{\text{op}}}^{E_1}(n\text{Vec}) \simeq \Sigma \mathfrak{Z}_1(\mathcal{A}), & \Omega \mathcal{D} &\simeq \mathfrak{Z}_1(\mathcal{A}); \\ M^n &= (\mathcal{M}, m) \simeq (\text{BMod}_{\mathcal{A}|\mathcal{B}}(n\text{Vec}), \mathcal{A}), & \Omega_m \mathcal{M} &= \text{Fun}_{\mathcal{A}|\mathcal{B}}(\mathcal{A}, \mathcal{A}). \end{aligned}$$

The relation between this classification and the one in Theorem^{ph} 3.3.26 is illustrated below.



Two definitions of the same algebra in $\mathcal{C}$:

- (1) the monoidal functor $\mathcal{B} \rightarrow \mathcal{A}$ defines $\mathcal{A}^{\text{op}} \in \text{Alg}_{E_1}^c(\text{BMod}_{\mathcal{B}|\mathcal{B}}(n\text{Vec})^{\text{op}})$;
- (2) the central functor $\mathfrak{Z}_1(\mathcal{B}) \rightarrow \text{Fun}_{\mathcal{A}|\mathcal{B}}(\mathcal{A}, \mathcal{A})$ defines $\text{Fun}_{\mathcal{A}|\mathcal{B}}(\mathcal{A}, \mathcal{A}) \in \text{Alg}_{E_1}^c(\text{RMod}_{\mathfrak{Z}_1(\mathcal{B})}(n\text{Vec}))$.


For $\mathcal{X} \in \text{LMod}_{\mathcal{B}}(n\text{Vec})$, the canonical monoidal functor $\mathcal{B} \rightarrow \text{Fun}(\mathcal{X}, \mathcal{X})$ defines a Lagrangian E_1 -algebra $\text{Fun}(\mathcal{X}, \mathcal{X})^{\text{op}}$ in $\mathcal{C} = \text{BMod}_{\mathcal{B}|\mathcal{B}}(n\text{Vec})^{\text{op}}$. $\square$

We provide more concrete constructions in the following theorem.

Theorem^{ph} 3.3.33. We fixed another non-chiral simple topological order $Z(P)^{n+1}$ with a simple gapped boundary P^n . We can construct indecomposable condensable E_1 -algebras in $\mathcal{C} = \text{BMod}_{\mathcal{B}|\mathcal{B}}(n\text{Vec})^{\text{op}}$ that define condensations from $Z(B)^{n+1}$ to $Z(P)^{n+1}$ more concretely in the following way. In this case, $\mathcal{P}$ is a fusion $(n-1)$ -category. Let $\mathcal{X} \in \text{BMod}_{\mathcal{B}|\mathcal{P}}(n\text{Vec})$. Then we have a canonical monoidal functor from $\mathcal{B}$ to an indecomposable multi-fusion n -category $\mathcal{A}_{\mathcal{X}} := \text{Fun}_{\mathcal{P}^{\text{op}}}(\mathcal{X}, \mathcal{X})$:

$$\mathcal{B} \rightarrow \text{Fun}_{\mathcal{B}|\mathcal{P}}(\mathcal{X}, \mathcal{X}) \boxtimes_{\mathfrak{Z}_1(\mathcal{B})} \mathcal{B} \simeq \text{Fun}_{\mathcal{P}^{\text{op}}}(\mathcal{X}, \mathcal{X}) = \mathcal{A}_{\mathcal{X}},$$

which defines an indecomposable condensable E_1 -algebra $\mathcal{A}_{\mathcal{X}}^{\text{op}}$ in $\text{BMod}_{\mathcal{B}|\mathcal{B}}(n\text{Vec})^{\text{op}}$. The physical meaning of each data is illustrated below.



Two definitions of the same algebra in $\mathcal{C}$:

- (1) the monoidal functor $\mathcal{B} \rightarrow \mathcal{A}_{\mathcal{X}}$ defines $\mathcal{A}_{\mathcal{X}}^{\text{op}} \in \text{Alg}_{E_1}^c(\text{BMod}_{\mathcal{B}|\mathcal{B}}(n\text{Vec})^{\text{op}})$;
- (2) the central functor $\mathfrak{Z}_1(\mathcal{B}) \rightarrow \text{Fun}_{\mathcal{B}|\mathcal{P}}(\mathcal{X}, \mathcal{X})^{\text{op}}$ defines $\mathcal{K}_{\mathcal{X}}^{\text{op}} := \text{Fun}_{\mathcal{B}|\mathcal{P}}(\mathcal{X}, \mathcal{X})^{\text{op}} \in \text{Alg}_{E_1}^c(\text{RMod}_{\mathfrak{Z}_1(\mathcal{B})}(n\text{Vec}))$.

By condensing $\mathcal{A}_{\mathcal{X}}^{\text{op}}$ in $\mathcal{C}$, we obtain the condensed topological order $D^{n+1} = Z(P)^{n+1}$ with $\mathcal{D} \simeq \text{Mod}_{\mathcal{A}_{\mathcal{X}}^{\text{op}}}^{E_1}(\mathcal{C})$ and a gapped domain wall $M^n = (\mathcal{M}, m) \simeq (\text{BMod}_{\mathcal{A}_{\mathcal{X}}|\mathcal{B}}(n\text{Vec}), \mathcal{A}_{\mathcal{X}})$. $\square$

Remark 3.3.34. However, this construction $\mathcal{X} \mapsto \mathcal{A}_{\mathcal{X}}$ (or $\mathcal{X} \mapsto \mathcal{K}_{\mathcal{X}}$) does not exhaust all possible condensations from $Z(B)^{n+1}$ to $Z(P)^{n+1}$ because the domain wall X^{n-1} can be gapless in general. If we allow X^{n-1} to be gapless⁴⁴, or equivalently, if we allow to stack anomaly-free Y^n onto the domain wall $Z(X)^n$ followed by condensations along the domain wall (recall Theorem 3.3.14), all indecomposable condensable E_1 -algebras in $\mathcal{C} = \text{BMod}_{\mathcal{B}|\mathcal{B}}(n\text{Vec})^{\text{op}}$ arise in this way (see Corollary^{ph} 3.3.37). $\diamond$

⁴⁴In this case, $\mathcal{X}$ is again a $\mathcal{B}$ - $\mathcal{P}$ -bimodule but living in the world of enriched separable higher categories [KZ22b].

Remark 3.3.35. When $P^n = 1^n$, the correspondence between $\mathcal{A}_{\mathcal{X}}$ and $\mathcal{K}_{\mathcal{X}}$ precisely explains the relation between two different but equivalent notions of gauging the fusion category symmetry $\mathcal{B}$. On the one hand, the gauging of the $\mathcal{B}$ -symmetry in the sense of [FFRS10, BT18, TW24, RSS23, LYW24] produces its dual symmetry $\mathcal{K}_{\mathcal{X}}^{\text{op}} = \text{Fun}_{\mathcal{B}}(\mathcal{X}, \mathcal{X})^{\text{op}}$ for a left $\mathcal{B}$ -module $\mathcal{X}$. It amounts to condensing $\mathcal{B}$ via the gapped domain wall $\mathcal{X}$, and $\text{Fun}_{\mathcal{B}}(\mathcal{X}, \mathcal{X})^{\text{op}}$ is the fusion category symmetry associated to the condensed phase. On the other hand, the gauging process in the sense of [KLWZZ20b, KLWZZ20a] maps the $\mathcal{B}$ -symmetry to an anomaly-free topological order. Mathematically, it amounts to a monoidal functor $\mathcal{B} \rightarrow \text{Fun}(\mathcal{X}, \mathcal{X}) = \mathcal{A}_{\mathcal{X}}$ for some separable $(n-1)$ -category $\mathcal{X}$. By [KZ18a, Theorem 3.2.3][KZ24, Remark 3.30], these two gauging processes are mathematically equivalent. Moreover, Theorem^{ph} 3.3.33 further relates these two gaugings to the condensation of the same 1-codimensional ‘Lagrangian’ topological defect in $Z(\mathcal{B})^{n+1}$ but in two different coordinate systems, respectively. $\diamond$

Using Theorem^{ph} 3.3.33 and [KZ21, Proposition 3.13], we obtain more precise classification results. For $n = 2$ (as in Theorem^{ph} 3.3.33), we recover an old result [KR08, DMNO13] but reformulated in a new context of 1-codimensional condensations; for $n \geq 3$, we obtain a new result.

Corollary^{ph} 3.3.36. When $n = 2$ (as in Theorem^{ph} 3.3.33), we obtain a one-to-one correspondence between (the equivalence classes of) $\mathcal{B}$ - $\mathcal{P}$ -bimodules and 1-codimensional defect condensations from $Z(\mathcal{B})^3$ to $Z(\mathcal{P})^3$ (i.e., indecomposable condensable E_1 -algebras in $\mathcal{C} = \Sigma\mathfrak{Z}_1(\mathcal{B})$ that produce $Z(\mathcal{P})^3$, or equivalently, type-I gapped domain walls between $Z(\mathcal{B})^3$ and $Z(\mathcal{P})^3$) defined by

$$\begin{aligned} \mathcal{B}\mathcal{X}_{\mathcal{P}} &\mapsto \mathcal{K}_{\mathcal{X}}^{\text{op}} := \text{Fun}_{\mathcal{B}|\mathcal{P}}(\mathcal{X}, \mathcal{X})^{\text{op}} \in \text{Alg}_{E_1}^c(\text{RMod}_{\mathfrak{Z}_1(\mathcal{B})}(2\text{Vec})), \\ \mathcal{B}\mathcal{X}_{\mathcal{P}} &\mapsto \mathcal{A}_{\mathcal{X}}^{\text{op}} := \text{Fun}_{\mathcal{P}|\mathcal{B}}(\mathcal{X}, \mathcal{X})^{\text{op}} \in \text{Alg}_{E_1}^c(\text{BMod}_{\mathcal{B}|\mathcal{B}}(2\text{Vec})^{\text{op}}) \end{aligned}$$

in two coordinate systems: $\Sigma\mathfrak{Z}_1(\mathcal{B}) = \text{RMod}_{\mathfrak{Z}_1(\mathcal{B})}(2\text{Vec})$ and $\Sigma\mathfrak{Z}_1(\mathcal{B}) = \text{BMod}_{\mathcal{B}|\mathcal{B}}(2\text{Vec})^{\text{op}}$, respectively.

Proof. It follows immediately from Theorem^{ph} 3.3.33 and Corollary 2.4.10. $\square$

Using the idea of topological Wick rotation [KZ18b, KZ20, KZ21, KZ22b], we obtain a generalization of Corollary^{ph} 3.3.36 to all dimensions. The following result is an immediate mathematical consequence of Theorem^{ph} 2.4.12.

Corollary^{ph} 3.3.37. For each non-degenerate fusion $(n-1)$ -category $\mathcal{T}$ for $n \geq 2$, there is a one-to-one correspondence (defined in two coordinate systems):

$$\mathcal{B} \boxtimes \mathcal{T} \mathcal{X}_{\mathcal{P}} \mapsto \mathcal{K}_{\mathcal{X}}^{\text{op}} := \text{Fun}_{\mathcal{B} \boxtimes \mathcal{T}|\mathcal{P}}(\mathcal{X}, \mathcal{X})^{\text{op}} \in \text{Alg}_{E_1}^c(\text{RMod}_{\mathfrak{Z}_1(\mathcal{B})}(n\text{Vec})), \quad (3.3.25)$$


$$\mathcal{B} \boxtimes \mathcal{T} \mathcal{X}_{\mathcal{P}} \mapsto \mathcal{A}_{\mathcal{X}}^{\text{op}} := \text{Fun}_{\mathcal{T}|\mathcal{P}}(\mathcal{X}, \mathcal{X})^{\text{op}} \in \text{Alg}_{E_1}^c(\text{BMod}_{\mathcal{B}|\mathcal{B}}(n\text{Vec})^{\text{op}}) \quad (3.3.26)$$

between the following two sets:

- $(\mathcal{B} \boxtimes \mathcal{T})$ - $\mathcal{P}$ -bimodules modulo the equivalence relation defined by $\mathcal{M} \sim \mathcal{N}$ if $\mathcal{N} \simeq \mathcal{M} \boxtimes \mathcal{V}$ for $\mathcal{V} \in n\text{Vec}^\times$;
- 1-codimensional defect condensations from $Z(\mathcal{B})^{n+1}$ to $Z(\mathcal{P})^{n+1}$, i.e., indecomposable condensable E_1 -algebras A in $\Sigma\mathfrak{Z}_1(\mathcal{B})$, that belong to the same Morita equivalence class as $(\mathcal{B} \boxtimes \mathcal{T}) \boxtimes \mathcal{P}^{\text{op}} \in \text{Alg}_{E_1}^c(\text{RMod}_{\mathfrak{Z}_1(\mathcal{B})}(n\text{Vec}))$ (recall Remark 3.3.16).

The condensable E_1 -algebra is simple if and only if the $(\mathcal{B} \boxtimes \mathcal{T})$ - $\mathcal{P}$ -bimodule is indecomposable. Note that the $n = 2$ case is precisely Corollary^{ph} 3.3.36 because the only non-degenerate fusion 1-category is the trivial one Vec .

Proof. This result follows immediately from Theorem^{ph} 2.4.12. Recall that the key idea of the proof is the holographic dual (via topological Wick rotation) between a potentially gapless defect junction Y^{n-1} and the pair (T^n, X^{n-1}) , where T^n is an anomaly-free nD topological order and X^{n-1} is a gapped defect junction among three anomalous topological orders $B^n \boxtimes T^n$, K^n and P^n . For readers convenience, we re-illustrate this idea in the pictures below with the corrected labels and dimensions.



It is worthwhile to note that those A 's in $\Sigma\mathfrak{Z}_1(\mathcal{B})$ are precisely indecomposable multi-fusion $(n-1)$ -categories that are Morita equivalent to $(\mathcal{B} \boxtimes \mathcal{T}) \boxtimes \mathcal{P}^{\text{op}}$, including those $(\mathcal{B} \boxtimes \mathcal{T}) \boxtimes \mathcal{P}^{\text{op}} \boxtimes \Sigma\mathfrak{Z}_1(\mathcal{Q})$ (for a fusion $(n-2)$ -category $\mathcal{Q}$), which is precisely the image of the $(\mathcal{B} \boxtimes \mathcal{T})$ - $\mathcal{P}$ bimodule $\mathcal{X} := \mathcal{B} \boxtimes \mathcal{T} \boxtimes \mathcal{P} \boxtimes \Sigma\mathcal{Q}$ under the one-to-one correspondence (3.3.25). $\square$

Remark 3.3.38. If we replace $\mathcal{T}$ by a Morita equivalent one $\mathcal{T}'$, it simply provides a new coordinate system for the domain of this one-to-one correspondence. $\diamond$

The following two results are reformulations of Theorem^{ph} 2.4.12 and 2.4.15 in terms of Lagrangian E_1 -algebras.

Corollary^{ph} 3.3.39. The map ${}_{\mathcal{B} \boxtimes \mathcal{T}}\mathcal{X} \mapsto \text{Fun}_{{}_{\mathcal{B} \boxtimes \mathcal{T}}}(\mathcal{X}, \mathcal{X})^{\text{op}}$ defines a one-to-one correspondence between:

- left indecomposable $\mathcal{B} \boxtimes \mathcal{T}$ -modules modulo the equivalence relation $\sim$,
- Lagrangian E_1 -algebras in $\Sigma\mathfrak{Z}_1(\mathcal{B})$ (or simple gapped boundaries of $Z(\mathcal{B})^{n+1}$) in the Morita class $[\mathcal{B} \boxtimes \mathcal{T}]$;

See Theorem^{ph} 2.4.12 and Corollary 4.2.23 for an equivalent result.

The following result generalizes an earlier result for fusion 2-categories [JFR23, DX24].

Corollary^o 3.3.40. When the fusion $(n-1)$ -category $\mathcal{B}$ is connected as a separable $(n-1)$ -category, i.e., $\mathcal{B} = \Sigma\Omega\mathcal{B}$, there is a one-to-one correspondence between

- braided monoidal functors $\Omega\mathcal{B} \rightarrow \mathcal{Q}$ for a non-degenerate braided fusion $(n-2)$ -category $\mathcal{Q}$;
- Lagrangian E_1 -algebra in $\Sigma\mathfrak{Z}_1(\mathcal{B})$.

Moreover, the Morita class of the Lagrangian E_1 -algebra is determined by the Morita class of $\Sigma\mathcal{Q}$.

Example 3.3.41. A typical example of $\mathcal{B} = \Sigma\Omega\mathcal{B}$ is $\mathcal{B} = (n-1)\text{Rep}(G)$. In this case, Corollary 3.3.40 provide a classification of Lagrangian E_1 -algebras in $\mathcal{G}\mathcal{T}_G^{n+1} \simeq \Sigma\mathfrak{Z}_1((n-1)\text{Rep}(G))$, or equivalently, that of the gapped boundaries of the finite gauge theory GT_G^{n+1} . $\heartsuit$

3.3.3 Examples in 2+1D: anomaly-free cases

Example 3.3.42. The topological defects in the 2+1D $\mathbb{Z}_2$ -gauge theory form the fusion 2-category $\mathcal{TC}$, which can be identified with the 2-category $\text{RMod}_{\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))}(2\text{Vec})$. By [BJS21, Déc23b] (or Theorem 3.3.26), a condensable E_1 -algebra in $\mathcal{TC}$ are precisely multi-fusion left $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$ -modules, i.e., a multi-fusion categories $\mathcal{A}$ equipped with a braided functors $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2)) \rightarrow \mathfrak{Z}_1(\mathcal{A})$. Here we list some examples.

1. The fusion category $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$ equipped with the identity functor $\text{id}_{\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))}$, which should be viewed as a central functor, defines a condensable E_1 -algebra $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$ in the coordinate system $\mathcal{TC} = \text{RMod}_{\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))}(2\text{Vec})$. It is the tensor unit of $\mathcal{TC}$.
2. The fusion category $\text{Rep}(\mathbb{Z}_2)$ equipped with the central functor $f : \mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2)) \rightarrow \text{Rep}(\mathbb{Z}_2)$ defines a condensable E_1 -algebra $\text{Rep}(\mathbb{Z}_2)^{\text{op}} = \text{Rep}(\mathbb{Z}_2)$ in $\text{RMod}_{\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))}(2\text{Vec})$. It corresponds to the 1-codimensional defect ss in $\mathcal{TC}$. We have $\text{Mod}_{\text{Rep}(\mathbb{Z}_2)}^{E_1}(\Sigma\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))) \simeq 2\text{Vec}$, which means that the condensed 2+1D topological order is trivial. This condensation defined by ss is the same as the anyon condensation defined by $1 \oplus m$. Therefore, ss is a Lagrangian condensable E_1 -algebra in $\mathcal{TC}$.
3. The fusion category $\text{Vec}_{\mathbb{Z}_2}$ is equipped with the canonical equivalence $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2)) \rightarrow \mathfrak{Z}_1(\text{Vec}_{\mathbb{Z}_2})$. It defines a condensable E_1 -algebra $\text{Vec}_{\mathbb{Z}_2}^{\text{op}} = \text{Vec}_{\mathbb{Z}_2}$, corresponding to the 1-codimensional defect rr in $\mathcal{TC}$. We have $\text{Mod}_{\text{Vec}_{\mathbb{Z}_2}}^{E_1}(\Sigma\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))) \simeq 2\text{Vec}$, which means that the condensed topological order is trivial. This condensation defined by rr is the same as the anyon condensation defined by $1 \oplus e$. Therefore, this condensable E_1 -algebra $\text{Vec}_{\mathbb{Z}_2}$ is also Lagrangian.

4. As we review in Example 4.1.13, the following object

$$A := \mathbb{1} \boxtimes \mathbb{1} \oplus \psi \boxtimes \psi \in \Omega\mathcal{I}s \boxtimes \Omega\mathcal{I}s^{\text{op}} \simeq \mathfrak{Z}_1(\Omega\mathcal{I}s) \quad (3.3.27)$$

has a canonical structure of a simple commutative separable algebra in $\mathfrak{Z}_1(\Omega\mathcal{I}s)$. Then fusion 1-category $\mathcal{K} := \text{RMod}_A(\mathfrak{Z}_1(\Omega\mathcal{I}s))^{\text{op}}$ is naturally a condensable E_1 -algebra in $\mathcal{TC}$. The category $\mathcal{K}$ has six simple objects $1, e, m, f, \chi_{\pm}$ (see Example 4.1.13). Using the coordinate systems $\mathcal{TC} \simeq \text{RMod}_{\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))}(2\text{Vec})$ and Table 1, we see immediately that

$$\mathcal{K} = 1 \oplus \vartheta \in \mathcal{TC}^3. \quad (3.3.28)$$

The algebraic structure on $1 \oplus \vartheta$ can be reconstructed as a special case of a more general construction (see Section 5.3). Condensing $\mathcal{K}$ in $\mathcal{TC}$ produces the condensed phase D^3 , which is precisely the 2+1D double Ising topological order, i.e.,

$$\text{Mod}_{\mathcal{K}}^{E_1}(\mathcal{TC}) \simeq \mathcal{I}s \boxtimes \mathcal{I}s^{\text{op}}$$

and a gapped domain wall $M^2 = (\mathcal{M}, m) = (\text{RMod}_{\mathcal{K}}(\mathcal{TC}), \mathcal{K}) = (\text{RMod}_{\mathcal{K}}(2\text{Vec}), \mathcal{K})$. We explain in Section 5.3 that condensing $\mathcal{K}$ in $\mathcal{TC}$ can also be understood as the process of gauging the e - m -duality in $\mathcal{TC}^3$ (or an equivariantization). $\heartsuit$

Example 3.3.43. We can choose a different coordinate system for $\mathcal{TC}$. Indeed, by [KK12], $\mathcal{TC}$ can be identified with the fusion 2-category $\text{BMod}_{\text{Rep}(\mathbb{Z}_2)|\text{Rep}(\mathbb{Z}_2)}(2\text{Vec})^{\text{op}}$. A condensable E_1 -algebra in $\mathcal{TC}$ is precisely a fusion 1-category $\mathcal{B}^{\text{op}}$ equipped with a monoidal functor $\text{Rep}(\mathbb{Z}_2) \rightarrow \mathcal{B}$. Here we list some examples:

1. The fusion category $\text{Rep}(\mathbb{Z}_2)$ equipped with the identity functor $\text{Rep}(\mathbb{Z}_2) \rightarrow \text{Rep}(\mathbb{Z}_2)$ is a condensable E_1 -algebra. It is the tensor unit of $\mathcal{TC}$.
2. The fusion category Vec equipped with the forgetful functor $\text{Rep}(\mathbb{Z}_2) \rightarrow \text{Vec}$ is a Lagrangian E_1 -algebra, which corresponds to the 1-codimensional topological defect rr in $\mathcal{TC}$.
3. The multi-fusion category $\text{Fun}(\text{Rep}(\mathbb{Z}_2), \text{Rep}(\mathbb{Z}_2))^{\text{op}}$ equipped with the canonical embedding

$$\text{Rep}(\mathbb{Z}_2) \rightarrow \text{Fun}(\text{Rep}(\mathbb{Z}_2), \text{Rep}(\mathbb{Z}_2))$$

is a Lagrangian E_1 -algebra, which corresponds to the 1-codimensional topological defect ss in $\mathcal{TC}$.

4. We set $\mathcal{K} := \text{RMod}_A(\mathfrak{Z}_1(\Omega\mathcal{I}s))$ for A defined in (3.3.27). There is a canonical monoidal functor

$$\text{Rep}(\mathbb{Z}_2) \rightarrow \mathcal{K} \boxtimes_{\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))} \text{Rep}(\mathbb{Z}_2) \simeq \Omega\mathcal{I}s := \mathcal{A},$$


where the monoidal equivalence ‘ $\simeq$ ’ follows from a general fact⁴⁵, defines a condensable E_1 -algebra $\mathcal{A}^{\text{op}}$ in $\mathcal{TC} = \text{BMod}_{\text{Rep}(\mathbb{Z}_2)|\text{Rep}(\mathbb{Z}_2)}(2\text{Vec})^{\text{op}}$. Condensing $\mathcal{A}^{\text{op}}$ produces the condensed topological order $Z(\text{Is})^3$:

$$\mathcal{I}s \boxtimes \mathcal{I}s^{\text{op}} \simeq \text{Mod}_{\mathcal{A}^{\text{op}}}^{E_1}(\mathcal{TC}),$$

and the gapped domain wall M^2 such that (using (3.3.22) and (3.3.23))

$$(\mathcal{M}, m) = (\text{BMod}_{\mathcal{A}|\mathcal{B}}(n\text{Vec}), \mathcal{A}) \simeq (\Sigma\mathcal{K}^{\text{op}}, \mathcal{K}).$$

The physical meaning of above data is depicted in the following picture.



⁴⁵For a non-degenerate braided fusion category $\mathcal{C}$ and a condensable E_2 -algebra A and a condensable E_2 -algebra B over A [DMNO13], we have a monoidal equivalence $\text{RMod}_A(\mathcal{C}) \boxtimes_{\text{Mod}_A^{E_2}(\mathcal{C})} \text{RMod}_B(\text{Mod}_A^{E_2}(\mathcal{C})) \simeq \text{RMod}_B(\mathcal{C})$.

We provide more examples of condensations in $\mathcal{TC}$ as the special cases of condensations in 2+1D finite gauge theories in Example 3.3.44. $\heartsuit$

Example 3.3.44. Consider 2+1D finite gauge theory GT_G^3 for a finite group G . We provide some examples of condensable E_1 -algebras in $\mathcal{GT}_G^3$ in two coordinate systems.

1. $\mathcal{GT}_G^3 = \text{RMod}_{\mathfrak{Z}_1(\text{Rep}(G))}(2\text{Vec})$:


- (a) Consider a simple condensable E_2 -algebra A (i.e., étale algebra) in $\mathfrak{Z}_1(\text{Rep}(G))$. Such algebras have been classified in [Dav10b] (see Example 4.1.11). The central functor

$$-\otimes A : \mathfrak{Z}_1(\text{Rep}(G)) \rightarrow \text{RMod}_A(\mathfrak{Z}_1(\text{Rep}(G))) \quad (3.3.29)$$

defines a condensable E_1 -algebra $B := \text{RMod}_A(\mathfrak{Z}_1(\text{Rep}(G)))$ in $\text{RMod}_{\mathfrak{Z}_1(\text{Rep}(G))}(2\text{Vec})$. By condensing it, we obtain a new 2+1D topological order $\mathcal{D}^3$ and a gapped domain wall M^2 given by

$$\begin{aligned} \mathcal{D} &\simeq \text{Mod}_B^{E_1}(\Sigma \mathfrak{Z}_1(\text{Rep}(G))) \simeq \Sigma(\text{Mod}_A^{E_2}(\mathfrak{Z}_1(\text{Rep}(G))), \\ (\mathcal{M}, m) &\simeq (\text{RMod}_B(\Sigma \mathfrak{Z}_1(\text{Rep}(G))), B). \end{aligned}$$

- i. When $A = \mathbb{1}_{\mathfrak{Z}_1(\text{Rep}(G))}$, $-\otimes A \simeq \text{id}_{\mathfrak{Z}_1(\text{Rep}(G))}$ defines the trivial condensable E_1 -algebra $\mathbb{1}_{\mathcal{GT}_G^3}$, which defines a trivial condensation.
 - ii. The group algebra $A = \mathbb{C}[G]$ can be viewed as a Lagrangian E_2 -algebra in $\mathfrak{Z}_1(\text{Vec}_G) \simeq \mathfrak{Z}_1(\text{Rep}(G))$ (see Example 4.1.11). In this case, the central functor $-\otimes A$ can be identified with the canonical central functor: the forgetful functor $\mathfrak{Z}_1(\text{Rep}(G)) \rightarrow \text{Rep}(G)$, which defines a Lagrangian E_1 -algebra $\text{Rep}(G)$ in $\mathcal{GT}_G^3$. By condensing it, we obtain $\mathbb{1}^3$ as the condensed phase and a gapped boundary M^2 such that $(\mathcal{M}, m) \simeq (2\text{Rep}(G), \mathbb{1})$.
 - iii. For a subgroup $H \leq G$, the algebra $\text{Fun}(G/H)$ is a condensable E_2 -algebra in $\text{Rep}(G) \hookrightarrow \mathfrak{Z}_1(\text{Rep}(G))$. In this case, the central functor $-\otimes A$ in (3.3.29) defines a condensable E_1 -algebra in $\mathcal{GT}_G^3$. By condensing it, we obtain GT_H^3 as the condensed phase and a gapped boundary M^2 such that $\mathcal{M} \simeq \Sigma \text{RMod}_A(\mathfrak{Z}_1(\text{Rep}(G)))$.
- (b) We can condense a condensable E_1 -algebra in $\mathcal{GT}_G^3$ to obtain any 2+1D simple non-chiral topological orders as shown in Theorem^{ph} 3.3.32. We give some concrete constructions.
- i. Let G_1 and G_2 be two finite groups and H a subgroup of both G_1 and G_2 . By restriction, we obtain two monoidal functors $\text{Rep}(G_1) \rightarrow \text{Rep}(H)$ and $\text{Rep}(G_2) \rightarrow \text{Rep}(H)$. Therefore, $\text{Rep}(H)$ is a $\text{Rep}(G_1)$ - $\text{Rep}(G_2)$ -bimodule. As a consequence, there is a natural central functor $\mathfrak{Z}_1(\text{Rep}(G_1)) \rightarrow \mathcal{K} := \text{Fun}_{\text{Rep}(G_1)|\text{Rep}(G_2)}(\text{Rep}(H), \text{Rep}(H))$, which defines a condensable E_1 -algebra $\mathcal{K}$ in $\mathcal{GT}_{G_1}^3$. By condensing the 1-codimensional topological defect $\mathcal{K}$ in $\text{RMod}_{\mathfrak{Z}_1(\text{Rep}(G_1))}(2\text{Vec})$, we obtain the $\text{GT}_{G_2}^3$ as the condensed 2+1D phase.
- For example, one can always choose H to be the trivial group. In this case, $\text{Rep}(H) = \text{Vec}$ and $\mathcal{K} \simeq \text{Vec}_{G_1 \times G_2}$ as monoidal 1-categories.
- ii. In this case, we can obtain a complete classification of all 1-codimensional defect condensations from $\text{GT}_{G_1}^3$ to $\text{GT}_{G_2}^3$ by applying Corollary^{ph} 3.3.36 as illustrated below.



By [KZ18a, Theorem 3.3.7], the functorial assignment $\mathcal{X} \rightarrow \mathcal{K}_{\mathcal{X}}$ is fully faithful. It means that there is a one-to-one correspondence between such simple condensable E_1 -algebras $\mathcal{K}_{\mathcal{X}}$ in $\text{RMod}_{\mathfrak{Z}_1(\text{Rep}(G_1))}(2\text{Vec})$ and indecomposable $\text{Rep}(G_1)$ - $\text{Rep}(G_2)$ -bimodules $\mathcal{X}$ in 2Vec , which is equivalent to indecomposable left $\text{Rep}(G_1 \times G_2)$ -modules in 2Vec and was classified in [BO04, Ost03a]. More precisely, an indecomposable left $\text{Rep}(G_1 \times G_2)$ -module $\mathcal{X}$ in 2Vec

is given by $\text{Rep}'(\tilde{H})$ or $\text{Rep}(H, \alpha)$, where $H \leq G_1 \times G_2$ is a subgroup, $[\alpha] \in H^2(H, \mathbb{C}^\times)$ is a 2-cohomology class and $\tilde{H}$ is the central extension of H determined by $[\alpha]$ (see Example 3.1.17). The associated simple condensable E_1 -algebra $\mathcal{K}_{(H, \omega)}$ in $\text{RMod}_{\mathfrak{Z}_1(\text{Rep}(G_1))}(2\text{Vec})$ is

$$\mathcal{K}_{(H, \omega)} = \text{Fun}_{\text{Rep}(G_1)|\text{Rep}(G_2)}(\text{Rep}'(\tilde{H}), \text{Rep}'(\tilde{H})).$$

When G_2 is the trivial group, all these condensable E_1 -algebras $\mathcal{K}_{(H, \omega)}^{\text{op}}$ are Lagrangian. This classification result coincides with the classification of Lagrangian E_2 -algebras in $\mathfrak{Z}_1(\text{Rep}(G_1))$ in [Dav10b] (see also the 5th case in Example 4.1.11).

2. $\mathcal{GT}_G^3 = \text{BMod}_{\text{Rep}(G)|\text{Rep}(G)}(2\text{Vec})^{\text{op}}$:

(a) The central functor (3.3.29) defines a monoidal functor

$$\text{Rep}(G) \rightarrow \text{Rep}(G) \boxtimes_{\mathfrak{Z}_1(\text{Rep}(G))} \text{RMod}_A(\mathfrak{Z}_1(\text{Rep}(G))) \simeq \text{RMod}_A(\text{Rep}(G)), \quad (3.3.30)$$

where the ‘ $\simeq$ ’ is proved in [KZ18a, XY25]. By condensing $\text{RMod}_A(\text{Rep}(G))^{\text{op}}$, we obtain a condensed phase $\mathcal{D}^3$ such that

$$\mathcal{D} \simeq \text{Mod}_{\text{RMod}_A(\text{Rep}(G))^{\text{op}}}^{E_1}(\mathcal{GT}_G^3) \simeq \Sigma \text{Mod}_A^{E_2}(\mathfrak{Z}_1(\text{Rep}(G))),$$

where the second ‘ $\simeq$ ’ is as a special case of Theorem 5.1.5 (see also Example 5.1.10).

When $A = \text{Fun}(G/H)$ for a subgroup $H \leq G$, the monoidal functor (3.3.30) is precisely the forgetful functor $\text{Rep}(G) \rightarrow \text{Rep}(H)$. By condensing $\text{Rep}(H)^{\text{op}}$, we obtain the condensed phase $\mathcal{D}^3$ such that $\mathcal{D} = \text{Mod}_{\text{Rep}(H)^{\text{op}}}^{E_1}(\mathcal{GT}_G^3) \simeq \text{Mod}_{\text{Rep}(H)^{\text{op}}}^{E_1}(2\text{Vec}) = \mathcal{GT}_H^3$.

(b) Using Corollary^{ph} 3.3.36, we obtain a complete classification of all 1-codimensional defect condensations from $\text{GT}_{G_1}^3$ to $\text{GT}_{G_2}^3$ in the new coordinate. More precisely, required simple condensable E_1 -algebras in $\text{BMod}_{\text{Rep}(G)|\text{Rep}(G)}(2\text{Vec})^{\text{op}}$ are one-to-one corresponding to indecomposable left $\text{Rep}(G_1 \times G_2)$ -modules in 2Vec , each of which is given by $\text{Rep}'(\tilde{H})$ or $\text{Rep}(H, \alpha)$, where $H \leq G_1 \times G_2$ is a subgroup, $[\alpha] \in H^2(H, \mathbb{C}^\times)$ is a 2-cohomology class and $\tilde{H}$ is the central extension of H determined by $[\alpha]$ (see Example 3.1.17) [BO04, Ost03a]. The associated simple condensable E_1 -algebra $\mathcal{A}_{(H, \omega)}^{\text{op}}$ in $\text{BMod}_{\text{Rep}(G)|\text{Rep}(G)}(2\text{Vec})^{\text{op}}$ is defined by the monoidal functor

$$\text{Rep}(G_1) \rightarrow \mathcal{K}_{(H, \omega)} \boxtimes_{\mathfrak{Z}_1(\text{Rep}(G_1))} \text{Rep}(G_1) \simeq \text{Fun}_{\text{Rep}(G_2)^{\text{op}}}(\text{Rep}'(\tilde{H}), \text{Rep}'(\tilde{H})) =: \mathcal{A}_{(H, \omega)}. \quad \heartsuit$$

When G_2 is the trivial group, all these condensable E_1 -algebras $\mathcal{A}_{(H, \omega)}^{\text{op}}$ are Lagrangian.

Remark 3.3.45. After the appearance of our paper, more concrete examples in 2+1D have been worked out in literature (see for example [BBDR25]). $\diamond$

3.3.4 Examples in 2+1D: anomalous cases

We provide some concrete examples of 1-codimensional defect condensations in 2+1D anomalous topological orders. By topological Wick rotation (or topological holography) [KZ18b, KZ20, KZ21, KIWZZ20a, KIWZZ20b, KZ22b], a 1-codimensional defect condensation in a 2+1D anomalous topological order can also be understood as a condensation between 2+1D gapped quantum liquids with the same finite onsite symmetry, and as the gauging of 0-form symmetries in a gapless 2+1D QFT.

Example 3.3.46. Let G be a finite group. The 3+1D G gauge theory GT_G^4 has a gapped boundary obtained by condensing all particle-like topological defects (symmetry charges). The topological defects on this gapped boundary form the fusion 2-category 2Vec_G .

By [Déc23b], the condensable E_1 -algebras in 2Vec_G are G -graded multi-fusion 1-categories, and the simple ones are G -graded fusion 1-categories. A G -graded (multi-)fusion 1-category is a (multi-)fusion 1-category $\mathcal{A}$ equipped with a G -grading $\mathcal{A} = \bigoplus_{g \in G} \mathcal{A}_g$ such that $x \otimes y \in \mathcal{A}_{gh}$ for any $x \in \mathcal{A}_g$ and $y \in \mathcal{A}_h$. In particular, the trivial component $\mathcal{A}_e$ is also a (multi-)fusion 1-category. We say that the G -grading is faithful if $\mathcal{A}_g \neq 0$ for all $g \in G$.

- For example, given a subgroup $H \leq G$ and a 3-cocycle $\omega \in Z^3(H, U(1))$, there is a G -graded fusion category Vec_H^ω with the obvious G -grading supported on H and the associator given by ω . Two G -graded fusion 1-categories Vec_H^ω and $\text{Vec}_K^{\omega'}$ are equivalent as condensable E_1 -algebras in 2Vec_G if and only if $H = K$ and $[\omega] = [\omega'] \in H^3(H, U(1))$.

For a G -graded fusion 1-category $\mathcal{A}$, we give a more explicit mathematical description of the left 2Vec_G -module $\text{RMod}_{\mathcal{A}}(2\text{Vec}_G)$. First, we define a 2-category $\text{B}_G\mathcal{A}$ as follows.

- The objects are elements in G .
- For $g, h \in G$, the hom 1-category $\text{Hom}(g, h)$ is $\mathcal{A}_{hg^{-1}}$. In particular, the endomorphism 1-category of every object is the fusion 1-category $\mathcal{A}_e$.
- The 2-category structure (including the composition, identity, associator, ...) is induced by the monoidal structure of $\mathcal{A}$.

Secondly, we define $\Sigma_G\mathcal{A}$ to be the condensation completion of $\text{B}_G\mathcal{A}$. Then $\Sigma_G\mathcal{A}$ is a finite semisimple 2-category (see [DR18, Construction 2.1.23] and references therein). Note that there is an obvious right translation G -action on $\text{B}_G\mathcal{A}$:

$$g \odot h := hg^{-1}, \quad \text{Hom}(g \odot h, g \odot k) = \text{Hom}(hg^{-1}, kg^{-1}) = \mathcal{A}_{kh^{-1}} = \text{Hom}(h, k).$$

It induces a G -action on $\Sigma_G\mathcal{A}$. Therefore, $\Sigma_G\mathcal{A}$ is a finite semisimple 2Vec_G -module.

Now we show that $\Sigma_G\mathcal{A} \simeq \text{RMod}_{\mathcal{A}}(2\text{Vec}_G)$ as 2Vec_G -modules. Denote the (representative of) simple objects in 2Vec_G by $\{\delta_g\}_{g \in G}$. There is a 2-functor $\text{B}_G\mathcal{A} \rightarrow \text{RMod}_{\mathcal{A}}(2\text{Vec}_G)$ defined by

- the map between objects: an object $g \in G$ is mapped to the free right $\mathcal{A}$ -module $\delta_{g^{-1}} \otimes \mathcal{A}$;
- the functors between hom 1-categories: the canonical equivalence

$$\text{Hom}_{\text{B}_G\mathcal{A}}(g, h) = \mathcal{A}_{hg^{-1}} \simeq \text{Hom}_{2\text{Vec}_G}(\delta_{g^{-1}}, \delta_{h^{-1}} \otimes \mathcal{A}) \simeq \text{Hom}_{\text{RMod}_{\mathcal{A}}(2\text{Vec}_G)}(\delta_{g^{-1}} \otimes \mathcal{A}, \delta_{h^{-1}} \otimes \mathcal{A}).$$

Moreover, this is a G -module 2-functor. Thus it induces a G -module 2-functor $\Sigma_G\mathcal{A} \rightarrow \text{RMod}_{\mathcal{A}}(2\text{Vec}_G)$. Note that every simple module $x \in \text{RMod}_{\mathcal{A}}(2\text{Vec}_G)$ can be condensed from the free module $x \otimes \mathcal{A}$. Thus the G -module 2-functor $\Sigma_G\mathcal{A} \rightarrow \text{RMod}_{\mathcal{A}}(2\text{Vec}_G)$ is an equivalence.

There are some immediate corollaries of this result. Let H be the support of the G -grading on $\mathcal{A}$. Then it is easy to see that $\pi_0(\Sigma_G\mathcal{A}) \simeq G/H$. In particular, $\text{RMod}_{\mathcal{A}}(2\text{Vec}_G) \simeq \Sigma_G\mathcal{A}$ is connected if and only if the G -grading on $\mathcal{A}$ is faithful. Moreover, when the G -grading on $\mathcal{A}$ is faithful, there is an equivalence of 2-categories $\Sigma_G\mathcal{A} \simeq \Sigma\mathcal{A}_e$. In this case, the 2Vec_G -action on $\Sigma\mathcal{A}_e$ is a monoidal 2-functor $2\text{Vec}_G \rightarrow \text{BMod}_{\mathcal{A}_e|\mathcal{A}_e}(2\text{Vec}) \simeq \text{End}(\Sigma\mathcal{A}_e)$, and it induces a 3-group homomorphism $G \rightarrow \text{BrPic}(\mathcal{A}_e)$, where the Brauer-Picard 3-group $\text{BrPic}(\mathcal{A}_e)$ is the maximal 3-group of $\text{BMod}_{\mathcal{A}_e|\mathcal{A}_e}(2\text{Vec})^\times$. By [ENO10], this 3-group homomorphism is equivalent to a faithful G -graded extension of $\mathcal{A}_e$, which is exactly the G -graded fusion 1-category $\mathcal{A}$.

As an example, for $\omega \in Z^3(G, U(1))$, we have $\text{RMod}_{\text{Vec}_G^\omega}(2\text{Vec}_G) \simeq 2\text{Vec}$, and the pentagonator of the 2Vec_G -action on 2Vec is given by ω . Equivalently, this 2Vec_G -action on 2Vec is induced from the monoidal 2-functor $2\text{Vec}_G \rightarrow 2\text{Vec}$ twisted by ω . Another example is the Ising fusion 1-category $\Omega\mathcal{I}s \simeq \text{Vec}_{\mathbb{Z}_2} \oplus \text{Vec}$ as a $\mathbb{Z}_2$ -graded fusion category. The $2\text{Vec}_{\mathbb{Z}_2}$ -action on $\text{RMod}_{\Omega\mathcal{I}s}(2\text{Vec}_{\mathbb{Z}_2}) \simeq \Sigma\text{Vec}_{\mathbb{Z}_2} \simeq 2\text{Rep}(\mathbb{Z}_2)$ is given by permuting two simple objects. $\heartsuit$

Example 3.3.47. The 3+1D G gauge theory GT_G^4 has another gapped boundary obtained by condensing all symmetry fluxes. The fusion 2-category of topological defects on this boundary is equivalent to $2\text{Rep}(G)$. There are two coordinate systems on $2\text{Rep}(G)$.

- (1) In the coordinate system, $2\text{Rep}(G) \simeq \text{Fun}(\text{BG}, 2\text{Vec})$, it is clear that objects in $2\text{Rep}(G)$ are separable 1-categories equipped with G -actions. A condensable E_1 -algebra in $2\text{Rep}(G)$ is precisely a multi-fusion 1-category $\mathcal{A}$ equipped with monoidal G -action, which is also called a G -equivariant multi-fusion 1-category. A condensable E_1 -algebra $\mathcal{A}$ in $2\text{Rep}(G)$ is simple if G acts transitively on the simple direct summands of the tensor unit of $\mathcal{A}$.

- (2) In the coordinate system $2\text{Rep}(G) \simeq \Sigma\text{Rep}(G) \simeq \text{RMod}_{\text{Rep}(G)}(2\text{Vec}) \simeq \text{LMod}_{\text{Rep}(G)}(2\text{Vec})$, by [BJS21, Déc23b], a condensable E_1 -algebra in $2\text{Rep}(G)$ is precisely a multi-fusion $\text{Rep}(G)$ -module, i.e., a multi-fusion 1-category $\mathcal{B}$ equipped with a braided monoidal functor $\text{Rep}(G) \rightarrow \mathfrak{Z}_1(\mathcal{B})$, and the simple ones are fusion $\text{Rep}(G)$ -modules. Such a multi-fusion $\text{Rep}(G)$ -module is also called a multi-fusion 1-category over $\text{Rep}(G)$. By [DMNO13, Lemma 3.5], a monoidal functor $\text{Rep}(G) \rightarrow \mathfrak{Z}_1(\mathcal{B}) \rightarrow \mathcal{B}$ factors through $\text{Rep}(G) \rightarrow \text{Rep}(H) \rightarrow \mathcal{B}$ for some subgroup $H \leq G$ (up to conjugation) such that $\text{Rep}(H) \rightarrow \mathcal{B}$ is an embedding (fully faithful).

The coordinate transformations between two coordinate systems of $2\text{Rep}(G)$ are given by the equivariantization and de-equivariantization (see [DGNO10, Section 4.1]). More precisely,

$$\begin{aligned} \{\mathbb{C}\text{-linear categories with } G\text{-action}\} &\longleftrightarrow \{\mathbb{C}\text{-linear categories with } \text{Rep}(G)\text{-action}\} \\ \mathcal{A} &\mapsto \mathcal{A}^G \\ \mathcal{B}_G := \text{RMod}_{\mathcal{A}}(\mathcal{B}) &\leftarrow \mathcal{B}, \end{aligned}$$

where $\mathcal{A}^G$ consists of G -invariant objects x (i.e., $g(x) \simeq x$ for all $g \in G$) and where $\mathcal{A} = \text{Fun}(G)$ is a condensable E_1 -algebra in $\text{Rep}(G)$. These coordinate transformations give one-to-one correspondences between condensable E_1 -algebras in different coordinates [DGNO10, Section 4.2]:

$$\{G\text{-equivariant multi-fusion 1-categories}\} \longleftrightarrow \{\text{multi-fusion 1-categories over } \text{Rep}(G)\}$$

Given a simple condensable E_1 -algebra $\mathcal{A} \in \text{LMod}_{\text{Vec}_G}(2\text{Vec}) \simeq \text{Fun}(BG, 2\text{Vec})$, the set of simple direct summands of the tensor unit is isomorphic to G/H for some subgroup H . In other words, its tensor unit spans a multi-fusion subcategory $\text{Fun}(G/H, \text{Vec}) \subseteq \mathcal{A}$. The equivariantization of the monoidal functors $\text{Vec} \rightarrow \text{Fun}(G/H, \text{Vec}) \hookrightarrow \mathcal{A}$ is

$$\text{Vec}^G \simeq \text{Rep}(G) \rightarrow \text{Rep}(H) \simeq \text{Fun}(G/H, \text{Vec})^G \hookrightarrow \mathcal{A}^G,$$

which gives a simple condensable E_1 -algebra in $\text{LMod}_{\text{Rep}(G)}(2\text{Vec}) \simeq \Sigma\text{Rep}(G)$. This is the one-to-one correspondence of simple condensable E_1 -algebras in the two coordinates of $2\text{Rep}(G)$. When $H = G$, $\mathcal{A}$ is a G -equivariant fusion 1-category. Under this one-to-one correspondence, G -equivariant fusion 1-categories are mapped to fusion 1-categories $\mathcal{B}$ equipped with braided functor $\text{Rep}(G) \rightarrow \mathfrak{Z}_1(\mathcal{B})$ such that the composite functor $\text{Rep}(G) \rightarrow \mathfrak{Z}_1(\mathcal{B}) \rightarrow \mathcal{B}$ is fully faithful [DGNO10, Theorem 4.18 (iii)].

We give some examples here.

1. For every subgroup $H \leq G$, the forgetful functor $\text{Rep}(G) \rightarrow \text{Rep}(H)$ is symmetric, hence factors through $\mathfrak{Z}_1(\text{Rep}(H))$. Thus $\text{Rep}(H)$ is a simple condensable E_1 -algebra in $\Sigma\text{Rep}(G)$. We also have $\text{RMod}_{\text{Rep}(H)}(\Sigma\text{Rep}(G)) \simeq \text{RMod}_{\text{Rep}(H)}(2\text{Vec}) \simeq \Sigma\text{Rep}(H)$, and the $\Sigma\text{Rep}(G)$ -action on $\Sigma\text{Rep}(H)$ is induced by the monoidal functor $\text{Rep}(G) \rightarrow \text{Rep}(H)$. After de-equivariantization, the corresponding simple condensable E_1 -algebra in $2\text{Rep}(G) \simeq \text{Fun}(BG, 2\text{Vec})$ is $\text{Fun}(G/H, \text{Vec})$ with the pointwise tensor product and left translation G -action.
2. For every subgroup $H \leq G$ and 3-cocycle $\omega \in Z^3(H, U(1))$, there is a braided embedding $\text{Rep}(H) \rightarrow \mathfrak{Z}_1(\text{Vec}_H^\omega)$. Indeed, these are all the minimal nondegenerate extensions of $\text{Rep}(H)$ [DGNO10, LKW16a]. Thus Vec_H^ω is a simple condensable E_1 -algebra in $\Sigma\text{Rep}(G)$.

For a fusion $\text{Rep}(G)$ -module $\mathcal{B}$, we have $\text{RMod}_{\mathcal{B}}(2\text{Rep}(G)) \simeq \text{RMod}_{\mathcal{B}}(2\text{Vec}) \simeq \Sigma\mathcal{B}$. The left $2\text{Rep}(G)$ -action on $\Sigma\mathcal{B}$ is defined by the monoidal 2-functor $\Sigma\text{Rep}(G) \rightarrow \Sigma\mathfrak{Z}_1(\mathcal{B}) \simeq \mathfrak{Z}_0(\Sigma\mathcal{B})$, which is induced from the braided monoidal functor $\text{Rep}(G) \rightarrow \mathfrak{Z}_1(\mathcal{B})$. As discussed above, this braided monoidal functor factors through $\text{Rep}(G) \rightarrow \text{Rep}(H) \rightarrow \mathfrak{Z}_1(\mathcal{B})$ and $\text{Rep}(H) \rightarrow \mathfrak{Z}_1(\mathcal{B})$ is a braided embedding. It follows that the indecomposable $2\text{Rep}(G)$ -modules one-to-one correspond to the pairs $(H, \mathcal{M})$, where $H \leq G$ is a subgroup of G defined up to conjugation and $\mathcal{M}$ is a nondegenerate extension of $\text{Rep}(H)$ such that $\mathcal{M}$ is Witt trivial (i.e., $\mathcal{M} \simeq \mathfrak{Z}_1(\mathcal{B})$ for some fusion category $\mathcal{B}$). This classification of indecomposable $2\text{Rep}(G)$ -module coincides with that of gapped boundaries of GT_G^4 that are Morita equivalent to $2\text{Rep}(G)$ (see also Example 4.2.41 and 4.2.42 and Theorem^{ph} 4.2.25). $\heartsuit$

Example 3.3.48. Let $\mathcal{T}$ be a nondegenerate braided fusion 1-category. We can stack the 2+1D gapped boundary of GT_G^4 associated the fusion 2-category $2\text{Rep}(G)$ with an anomaly-free 2+1D topological order corresponding to $\mathcal{T}$. Then the fusion 2-category of topological defects on this new 2+1D boundary topological order is $2\text{Rep}(G) \boxtimes \Sigma\mathcal{T} \simeq \Sigma(\text{Rep}(G) \boxtimes \mathcal{T})$.

Similar to the above example, the condensable E_1 -algebras in $\Sigma(\text{Rep}(G) \boxtimes \mathcal{T})$ are multi-fusion $\text{Rep}(G) \boxtimes \mathcal{T}$ -modules, and the simple ones are fusion $\text{Rep}(G) \boxtimes \mathcal{T}$ -modules, that is, a fusion category $\mathcal{B}$ equipped with a braided functor $\text{Rep}(G) \boxtimes \mathcal{T} \rightarrow \mathfrak{Z}_1(\mathcal{B})$. This is equivalent to two braided functors $\text{Rep}(G) \rightarrow \mathfrak{Z}_1(\mathcal{B})$ and $\mathcal{T} \rightarrow \mathfrak{Z}_1(\mathcal{B})$ such that their images in $\mathfrak{Z}_1(\mathcal{B})$ are symmetric to each other. Since $\mathcal{T}$ is nondegenerate, the braided functor $\mathcal{T} \rightarrow \mathfrak{Z}_1(\mathcal{B})$ is fully faithful [DMNO13, Corollary 3.26]. Denote the Müger centralizer of $\mathcal{T}$ in $\mathfrak{Z}_1(\mathcal{B})$ by $\mathcal{M} := \mathfrak{Z}_2(\mathcal{T}, \mathfrak{Z}_1(\mathcal{B}))$. Then $\mathfrak{Z}_1(\mathcal{B}) \simeq \mathcal{M} \boxtimes \mathcal{T}$ by [DGNO10, Theorem 3.13] (see also [Müg03c, Theorem 4.2] for modular case). So the fusion $\text{Rep}(G) \boxtimes \mathcal{T}$ -module structure on $\mathcal{B}$ is equivalent to a nondegenerate braided fusion category $\mathcal{M}$, a braided equivalence $\mathcal{M} \boxtimes \mathcal{T} \simeq \mathfrak{Z}_1(\mathcal{B})$ and a braided functor $\text{Rep}(G) \rightarrow \mathcal{M}$. Similar to the above example, the braided functor $\text{Rep}(G) \rightarrow \mathcal{M}$ factors through $\text{Rep}(G) \rightarrow \text{Rep}(H) \rightarrow \mathcal{M}$ such that $\text{Rep}(H) \rightarrow \mathcal{M}$ is fully faithful.

Given a fusion $(\text{Rep}(G) \boxtimes \mathcal{T})$ -module $\mathcal{B}$, we have

$$\text{RMod}_{\mathcal{B}}(2\text{Rep}(G) \boxtimes \Sigma\mathcal{T}) \simeq \text{RMod}_{\mathcal{B}}(2\text{Vec}) \simeq \Sigma\mathcal{B},$$

and the left $\Sigma(\text{Rep}(G) \boxtimes \mathcal{T})$ -action is induced from the braided monoidal functor

$$\text{Rep}(G) \boxtimes \mathcal{T} \rightarrow \text{Rep}(H) \boxtimes \mathcal{T} \hookrightarrow \mathcal{M} \boxtimes \mathcal{T} \simeq \mathfrak{Z}_1(\mathcal{B}).$$

It follows that the indecomposable $(2\text{Rep}(G) \boxtimes \Sigma\mathcal{T})$ -modules are one-to-one corresponding to the pairs $(H, \mathcal{M})$, where $H \leq G$ is a subgroup defined up to conjugation and $\mathcal{M}$ is a nondegenerate extension of $\text{Rep}(H)$ such that $\mathcal{M} \boxtimes \mathcal{T}$ is Witt trivial. This classification of indecomposable $2\text{Rep}(G)$ -module coincides with that of gapped boundaries of GT_G^4 that are Morita equivalent to $2\text{Rep}(G) \boxtimes \Sigma\mathcal{T}$ (see also Example 4.2.41 and 4.2.42 and Theorem^{ph} 4.2.25). $\heartsuit$

Example 3.3.49. Let $\pi \in Z^4(G, U(1))$. There is a gapped boundary of the 3+1D twisted G gauge theory $\text{GT}_{(G, \pi)}^4$ obtained by condensing all particle-like topological defects (symmetry charges). The fusion 2-category of topological defects on this 2+1D boundary topological order is 2Vec_G^π . It is the same as the 2Vec_G as 2-categories but the pentagonator is twisted by π .

Simple condensable E_1 -algebras in 2Vec_G^π are ‘ π -twisted G -graded fusion categories’: they have the same defining data as G -graded fusion categories, but the pentagon equation holds up to π . For example, given a subgroup $H \leq G$ and a 3-cochain $\pi \in C^3(H, U(1))$ such that $d\psi = \pi|_H$, there is a simple condensable E_1 -algebra $\text{Vec}_H^\psi \in 2\text{Vec}_G^\pi$ with the obvious G -grading supported on H and the associator twisted by ψ .

The underlying 2-category of $\text{RMod}_{\text{Vec}_H^\psi}(2\text{Vec}_G^\pi)$ is $2\text{Vec}_{G/H} = 2\text{Vec}^{\oplus G/H}$, and the left 2Vec_G^π -action is induced by the left translation of G on G/H . Its pentagonator is a 3-cocycle in $Z^3(G; \text{Fun}(G/H, U(1)))$, which is determined by ψ under the isomorphism $H^3(H; U(1)) \simeq H^3(G; \text{Fun}(G/H, U(1)))$ by Shapiro’s lemma for group cohomology. The underlying 2-category and the (partial) fusion rule of the bimodule 2-category $\text{BMod}_{\text{Vec}_H^\psi | \text{Vec}_H^\psi}(2\text{Vec}_G^\pi)$ can be found in [DY25].

By Corollary^{ph} 3.3.39, the Morita classes of simple condensable E_1 -algebras A in 2Vec_G^π , or equivalently, indecomposable left 2Vec_G^π -modules, classify Lagrangian E_1 -algebras in $\Sigma\mathfrak{Z}_1(2\text{Vec}_G^\pi)$ in the Morita class of $[2\text{Vec}_G^\pi]$. We believe that this should lead us to more concrete classification result of Lagrangian E_2 -algebras in $\mathfrak{Z}_1(2\text{Vec}_G^\pi)$ (as the full center of A) obtained recently in [DHJFN⁺24, Xu24b]. $\heartsuit$

3.3.5 Examples in 3+1D

We have already seen condensations of 1-codimensional topological defects in $\mathbf{1}^{n+1}$ in Section 3.3.2. In this subsection, we provide some examples in finite gauge theories based on Theorem^{ph} 3.3.32.

Example 3.3.50. For any fusion n -categories $\mathcal{C}$, we have $\mathcal{C} \simeq \mathcal{C} \boxtimes n\text{Vec}$. Therefore, for $n \geq 2$, there are infinitely many condensable E_1 -algebras in any fusion n -categories $\mathcal{C}$ because there are infinitely many multi-fusion $(n-1)$ -categories as condensable E_1 -algebras in $n\text{Vec}$. $\heartsuit$

Example 3.3.51. We denote the 3+1D G -gauge theory for a finite group G by GT_G^4 . In this case, we have $\Omega\mathcal{GT}_G^4 = \mathfrak{Z}_1(2\text{Vec}_G)$ [KTZ20b], and there are two coordinate systems of $\mathcal{GT}_G^4$. We provide examples only in the following coordinate systems:

$$\mathcal{GT}_G^4 = \text{BMod}_{2\text{Rep}(G) | 2\text{Rep}(G)}(3\text{Vec})^{\text{op}}.$$

We leave the same constructions in the other coordinate system as exercises.

1. Let $f : 2\text{Rep}(G) \rightarrow 2\text{Vec}$ be the forgetful functor, which is symmetric monoidal. Then $2\text{Vec}^{\text{op}} = 2\text{Vec}$ is a condensable E_1 -algebra in $\text{BMod}_{2\text{Rep}(G)|2\text{Rep}(G)}(3\text{Vec})^{\text{op}}$. By condensing it, we obtain the trivial phase $D^4 = \mathbf{1}^4$ and a gapped boundary M^3 with

$$(\mathcal{M}, m) = (\text{RMod}_{2\text{Rep}(G)}(3\text{Vec}), 2\text{Vec}), \quad \Omega_m \mathcal{M} = \text{Fun}_{2\text{Rep}(G)}(2\text{Vec}, 2\text{Vec}) \simeq 2\text{Vec}_G.$$

This condensation of 1-codimensional topological defect $2\text{Vec} \in \mathcal{GT}_G^4$ is equivalent to a condensation of 3-codimensional topological defect (i.e., a particle) $\text{Fun}(G) \in \text{Rep}(G) \simeq \Omega^2 \mathcal{GT}_G^4$. When $G = \mathbb{Z}_2$, this particle is precisely the e -particle in 3+1D toric code model [HZW05, KTZ20a, ZLZH⁺23]. When $G = \mathbb{Z}_2$ the boundary M^3 is called the rough boundary [ZLZH⁺23], which can also be obtained by condensing the $\mathbb{1}_c$ -string [ZLZH⁺23].

2. Let $\psi : 2\text{Rep}(G) \rightarrow \mathcal{A} = \text{Fun}(2\text{Rep}(G), 2\text{Rep}(G))$ be the canonical monoidal functor defined by the left multiplication $x \otimes -$. Then $\mathcal{A}^{\text{op}}$ is a condensable E_1 -algebra in $\text{BMod}_{2\text{Rep}(G)|2\text{Rep}(G)}(3\text{Vec})^{\text{op}}$. By condensing it, we obtain the trivial phase $D^4 = \mathbf{1}^4$ and a gapped boundary M^3 with

$$(\mathcal{M}, m) = (\text{RMod}_{2\text{Rep}(G)|\mathcal{A}}(3\text{Vec}), \mathcal{A}), \\ \Omega_m \mathcal{M} = \text{Fun}_{2\text{Rep}(G)|\mathcal{A}}(\mathcal{A}, \mathcal{A}) \simeq 2\text{Rep}(G).$$

When $G = \mathbb{Z}_2$, this gapped boundary of GT_G^4 is called the smooth boundary [KTZ20a], which can also be obtained from condensing m -string or the Lagrangian E_2 -algebra $1 \oplus m$ in $\mathfrak{Z}_1(2\text{Vec}_G)$ [ZLZH⁺23]. Note that $\mathcal{A}^{\text{op}}$ is a Lagrangian E_1 -algebra in $\mathcal{GT}_G^4$.

Since $\mathfrak{Z}_1(2\text{Vec}_G)$ has a non-trivial braided auto-equivalence α [ZLZH⁺23], we can twist ψ by α as follows:

$$\psi^\alpha : 2\text{Rep}(G) \rightarrow 2\text{Rep}(G) \boxtimes_{\mathfrak{Z}_1(2\text{Vec}_G)} \mathfrak{Z}_1(2\text{Vec}_G) \boxtimes_{\mathfrak{Z}_1(2\text{Vec}_G)} 2\text{Rep}(G) \\ \xrightarrow{\text{id} \boxtimes_{\mathfrak{Z}_1(2\text{Vec}_G)} \varphi \boxtimes_{\mathfrak{Z}_1(2\text{Vec}_G)} \text{id}} 2\text{Rep}(G) \boxtimes_{\mathfrak{Z}_1(2\text{Vec}_G)} \mathfrak{Z}_1(2\text{Vec}_G) \boxtimes_{\mathfrak{Z}_1(2\text{Vec}_G)} 2\text{Rep}(G) \simeq \mathcal{A}.$$

Then ψ^α endows a different algebraic structure on $\mathcal{A}$ denoted by $\mathcal{A}^\alpha$. By condensing $(\mathcal{A}^\alpha)^{\text{op}}$, we obtained the condensed phase $D^4 = \mathbf{1}^4$ and the so-called twist smooth boundary of GG_G^4 [ZLZH⁺23].

3. Let $H \leq G$ be a subgroup. The forgetful functor $f : 2\text{Rep}(G) \rightarrow 2\text{Rep}(H)$ is symmetric monoidal [KZ24], which defines the condensable E_1 -algebra in $\mathcal{GT}_G^4$. By condensing $2\text{Rep}(H)^{\text{op}}$, we obtain the finite gauge theory GT_H^4 as the condensed phase, i.e.,

$$\mathcal{GT}_H^4 \simeq \text{BMod}_{2\text{Rep}(H)|2\text{Rep}(H)}(\mathcal{GT}_G^4)^{\text{op}} \simeq \Sigma \mathfrak{Z}_1(2\text{Rep}(H)),$$

and a gapped domain wall M^3 such that

$$(\mathcal{M}, m) = (\text{BMod}_{2\text{Rep}(H)|2\text{Rep}(G)}(3\text{Vec}), 2\text{Rep}(H)).$$

4. By [KTZ20b], there is a canonical braided monoidal embedding $\psi_G : 2\text{Rep}(G) \rightarrow \mathfrak{Z}_1(2\text{Vec}_G^\omega)$ for $\omega \in Z^4(G, U(1))$. Therefore, $\mathfrak{Z}_1(2\text{Vec}_G^\omega) \in \text{Alg}_{E_1}^c(\text{BMod}_{2\text{Rep}(G)|2\text{Rep}(G)}(3\text{Vec}))$. More generally, for a subgroup $H \leq G$ and $\sigma \in H^4(H, U(1))$, the following monoidal functor:

$$2\text{Rep}(G) \xrightarrow{f} 2\text{Rep}(H) \xrightarrow{\psi_H} \mathfrak{Z}_1(2\text{Vec}_H^\sigma)$$

endows the category $\mathfrak{Z}_1(2\text{Vec}_H^\sigma)$ with a condensable E_1 -algebra structure in $\mathcal{GT}_G^4$. By condensing it, we obtain the double of a 3+1D twisted finite gauge theory as the condensed topological order D^4 , i.e.,

$$\Omega \mathcal{D} \simeq \mathfrak{Z}_1(2\text{Vec}_H^\sigma) \boxtimes \mathfrak{Z}_1(2\text{Vec}_H^\sigma)^{\text{op}}$$

as braided fusion 2-categories, and a gapped domain wall M^3 such that

$$(\mathcal{M}, m) = (\text{BMod}_{\mathfrak{Z}_1(2\text{Vec}_H^\sigma)|2\text{Rep}(G)}(3\text{Vec}), \mathfrak{Z}_1(2\text{Vec}_H^\sigma)).$$

This gapped domain wall was not known before.

5. We can also construct condensations from $\mathcal{GT}_{G_1}^4$ to any 3+1D twisted finite gauge theory $\mathcal{GT}_{(G_2, \rho)}^4$ for $\rho \in H^4(G_2, U(1))$. We give some examples.

(a) For a subgroup $H \leq G$ and $\sigma \in H^4(H, U(1))$, the following composed monoidal functor:

$$2\text{Rep}(G) \xrightarrow{f} 2\text{Rep}(H) \xrightarrow{\psi_H} \mathfrak{Z}_1(2\text{Vec}_H^\sigma) \xrightarrow{f} 2\text{Vec}_H^\sigma$$

endows the category 2Vec_H^σ with a condensable E_1 -algebra structure in $\mathcal{GT}_G^4$. By condensing it, we obtain the 3+1D twist finite gauge theory $\mathcal{GT}_{(H, \omega)}^4$.


(b) The following monoidal functor

$$2\text{Rep}(G_1) \xrightarrow{f} 2\text{Vec} \simeq 2\text{Vec} \boxtimes 2\text{Vec} \rightarrow 2\text{Vec} \boxtimes 2\text{Vec}_{G_2}^\rho \simeq 2\text{Vec}_{G_2}^\rho$$

defines condensable E_1 -algebra $2\text{Vec}_{G_2}^\rho$ in $\mathcal{GT}_{G_1}^4$. By condensing it, we obtain the 3+1D twist finite gauge theory $\mathcal{GT}_{(G_2, \rho)}^4$. This condensation can be viewed as a two-step condensation: first condense $\mathcal{GT}_{G_1}^4$ to $\mathbf{1}^4$ then condense $\mathbf{1}^4$ to $\mathcal{GT}_{(G_2, \rho)}^4$. Of course, this two-step condensation applies to condensations between any two non-chiral topological orders.

(c) One can always use Theorem^{ph} 3.3.33 and Corollary^{ph} 3.3.37 to construct (or classify) examples systematically. In this way, one can partially reduce the problem to the construction (or classification) of 2Vec_{G_1} - $2\text{Vec}_{G_2}^\rho$ -bimodules. $\heartsuit$

Using Theorem^{ph} 3.3.33 and Corollary^{ph} 3.3.37, we can construct (or even classify to some extent) condensations from $\mathcal{GT}_{G_1}^4$ to $\mathcal{GT}_{G_2}^4$ by specifying a $2\text{Rep}(G_1)$ - $2\text{Rep}(G_2)$ -bimodule $\mathcal{X}$ (as illustrated in the following picture, in which we set $\mathcal{K}_{\mathcal{X}} := \text{Fun}_{2\text{Rep}(G_1)|2\text{Rep}(G_2)}(\mathcal{X}, \mathcal{X})$).



More precisely, given such a bimodule $\mathcal{X}$, we obtain a condensable E_1 -algebra $\mathcal{A}_{\mathcal{X}}^{\text{op}}$ in $\mathcal{GT}_{G_1}^4$ by the following monoidal functor

$$2\text{Rep}(G_1) \rightarrow 2\text{Rep}(G_1) \boxtimes_{\mathfrak{Z}_1(2\text{Rep}(G_1))} \mathcal{K}_{\mathcal{X}} \simeq \text{Fun}_{2\text{Rep}(G_2)^{\text{op}}}(\mathcal{X}, \mathcal{X}) =: \mathcal{A}_{\mathcal{X}}$$

such that condensing $\mathcal{A}_{\mathcal{X}}^{\text{op}}$ produces $\mathcal{GT}_{G_2}^4$ as the condensed phase. For example, such $\mathcal{X}$ can be constructed from a monoidal $\text{Rep}(G_1)$ - $\text{Rep}(G_2)$ -bimodules, such as $\text{Rep}(G)$ for a subgroup $G \leq G_1 \times G_2$, via delooping, i.e., $\mathcal{X} = 2\text{Rep}(G)$ is a $2\text{Rep}(G_1)$ - $2\text{Rep}(G_2)$ -bimodule.

More generally, a $2\text{Rep}(G_1)$ - $2\text{Rep}(G_2)$ -bimodule $\mathcal{X}$ is just a right $2\text{Rep}(G_1 \times G_2)$ -module, i.e., objects in $3\text{Rep}(G_2 \times G_2)$. A simple object in $3\text{Rep}(G_2 \times G_2)$ is precisely $\Sigma\mathcal{L}$ for a fusion 1-category $\mathcal{L}$ over $\text{Rep}(G_2 \times G_2)$ (i.e., a fusion 1-category $\mathcal{L}$ equipped with a braided functor $\phi : \text{Rep}(G_2 \times G_2) \rightarrow \mathfrak{Z}_1(\mathcal{L})$) [KZZZ25]. Note that any indecomposable multi-fusion 1-category $\mathcal{Q}$ Morita equivalent to $\mathcal{L}$ produces the same simple $2\text{Rep}(G_1)$ - $2\text{Rep}(G_2)$ -bimodule $\mathcal{X}$, i.e., $\Sigma\mathcal{L} \simeq \Sigma\mathcal{Q}$.

For example, let $G \leq G_1 \times G_2$ be a subgroup. Then $\text{Rep}(G)$ is fusion 1-category over $\text{Rep}(G_2 \times G_2)$ and $\mathcal{X} = \Sigma\text{Rep}(G)$ is a right $2\text{Rep}(G_1 \times G_2)$ -module⁴⁶. If we require ϕ to factor through the embedding $\text{Rep}(G_1 \times G_2) \rightarrow \mathfrak{Z}_1(\text{Rep}(G_1 \times G_2))$, such ϕ 's can be classified, and it is easy to see that such obtained $\mathcal{X}$ are all of type $\Sigma\text{Rep}(G) \boxtimes \mathcal{S}$ for a separable 2-category $\mathcal{S}$.

⁴⁶An indecomposable left $\text{Rep}(G)$ -module is $\text{Rep}(H, \alpha)$ for a subgroup $H \leq G$ and $\alpha \in Z^2(H, U(1))$ (Recall Example 3.1.17). We have the following composed braided monoidal functor $\text{Rep}(G_2 \times G_2) \xrightarrow{f} \text{Rep}(G) \hookrightarrow \mathfrak{Z}_1(\text{Fun}_{\text{Rep}(G)}(\text{Rep}(H, \alpha), \text{Rep}(H, \alpha))^{\text{op}})$ which defines the same indecomposable $2\text{Rep}(G_1)$ - $2\text{Rep}(G_2)$ -bimodule $\Sigma\text{Rep}(G)$.

By Corollary^{ph} 3.3.36 (2), the assignment (3.3.25) of equivalence classes, i.e.,

$$\mathcal{X} \mapsto \text{Fun}_{2\text{Rep}(G_1)|2\text{Rep}(G_2)}(\mathcal{X}, \mathcal{X})^{\text{op}} \in \text{Alg}_{E_1}^c(\text{RMod}_{\mathfrak{Z}_1(2\text{Rep}(G_1))}(3\text{Vec}))$$

is bijective if we restrict the codomain to the equivalence classes within the Morita class of

$$2\text{Rep}(G_1) \boxtimes 2\text{Rep}(G_2)^{\text{op}} \in \text{Alg}_{E_1}^c(\text{RMod}_{\mathfrak{Z}_1(2\text{Rep}(G_1))}(3\text{Vec}))$$

(recall Remark 3.3.16).

More generally, an indecomposable condensable E_1 -algebra in $\mathcal{GT}_{G_1}^4 = \text{RMod}_{\mathfrak{Z}_1(2\text{Rep}(G_1))}(3\text{Vec})$ that condenses to $\mathcal{GT}_{G_2}^4$ is precisely an indecomposable multi-fusion 2-category $\mathcal{K}$ that is Morita equivalent to $\mathcal{K}_{\mathcal{X}} \boxtimes \mathcal{T} \in \text{Alg}_{E_1}^c(\text{RMod}_{\mathfrak{Z}_1(2\text{Rep}(G_1))}(3\text{Vec}))$ for a bimodule $\mathcal{X}$ and a non-degenerate fusion 2-category $\mathcal{T}$ (or equivalently, using the characterization of $\mathcal{K}$ given in Corollary^{ph} 3.3.37). In the coordinate system $\mathcal{GT}_{G_1}^4 = \text{BMod}_{2\text{Rep}(G)|2\text{Rep}(G)}(3\text{Vec})^{\text{op}}$, this algebra $\mathcal{K}$ is precisely the algebra $\mathcal{A}$ defined by the monoidal functor

$$2\text{Rep}(G_1) \rightarrow \mathcal{K} \boxtimes_{\mathfrak{Z}_1(2\text{Rep}(G_1))} 2\text{Rep}(G_1) =: \mathcal{A}.$$

In this way, we have covered all condensations from $\mathcal{GT}_{G_1}^4$ to $\mathcal{GT}_{G_2}^4$.

3.3.6 Examples in higher dimensions

Example 3.3.52. Consider $n+2D$ finite gauge theories GT_G^{n+2} . We use the following coordinate.

$$\mathcal{GT}_G^{n+2} = \text{BMod}_{n\text{Rep}(G)|n\text{Rep}(G)}((n+1)\text{Vec})^{\text{op}}.$$

An indecomposable condensable E_1 -algebra in $\text{BMod}_{n\text{Rep}(G)|n\text{Rep}(G)}((n+1)\text{Vec})$ is precisely given by an indecomposable multi-fusion n -category $\mathcal{A}^{\text{op}}$ equipped with a monoidal functor $n\text{Rep}(G) \rightarrow \mathcal{A}$.

1. The canonical monoidal functors $f : n\text{Rep}(G) \rightarrow n\text{Vec}$ and $n\text{Rep}(G) \rightarrow \text{Fun}(n\text{Rep}(G), n\text{Rep}(G))$ define two Lagrangian E_1 -algebras in $\mathcal{GT}_G^{n+2}$, which determine the rough boundary (i.e., $\Omega_m \mathcal{M} = n\text{Vec}_G$) and the smooth boundary (i.e., $\Omega_m \mathcal{M} = n\text{Rep}(G)$), respectively. The canonical forgetful functor $f : n\text{Rep}(G) \rightarrow n\text{Rep}(H)$ for a subgroup $H \leq G$ [KZZ24] defines a condensable E_1 -algebra in $\mathcal{GT}_G^{n+2}$, which produces the finite gauge theory GT_H^{n+2} as the condensed phase. This example can be recovered as a condensation of higher codimensional defects in Example 5.2.25. All above examples can be viewed as special cases of more general constructions explained in Part 4 in this Example.
2. By [KTZ20b], there is a canonical braided monoidal embedding $\psi_G : n\text{Rep}(G) \rightarrow \mathfrak{Z}_1(n\text{Vec}_G^\omega)$ for $\omega \in Z^{n+2}(G, U(1))$. Therefore, $\mathfrak{Z}_1(n\text{Vec}_G^\omega)^{\text{op}} \in \text{Alg}_{E_1}^c(\text{BMod}_{n\text{Rep}(G)|n\text{Rep}(G)}((n+1)\text{Vec})^{\text{op}})$. By condensing it, we obtain the double of twisted finite gauge theory $\text{GT}_{(G,\omega)}^{n+1} \boxtimes \overline{\text{GT}_{(G,\omega)}^{n+1}}$ as the condensed phase D^{n+2} , i.e.,

$$\mathcal{D} \simeq \text{Mod}_{\mathfrak{Z}_1(n\text{Vec}_G^\omega)}^{E_1}((n+1)\text{Vec})^{\text{op}}, \quad \Omega \mathcal{D} \simeq \mathfrak{Z}_1(n\text{Vec}_G^\omega) \boxtimes \mathfrak{Z}_1(n\text{Vec}_G^\omega)^{\text{op}}$$

where the first equivalence is monoidal and the second one is braided monoidal, and a gapped domain wall M^n such that

$$(\mathcal{M}, m) = (\text{BMod}_{\mathfrak{Z}_1(n\text{Vec}_G^\omega)|n\text{Rep}(G)}((n+1)\text{Vec}), \mathfrak{Z}_1(n\text{Vec}_G^\omega)).$$

3. For $H \leq G$ and $\sigma \in H^{n+2}(H, U(1))$, the composed braided monoidal functor

$$n\text{Rep}(G) \xrightarrow{f} n\text{Rep}(H) \xrightarrow{\psi_H} \mathfrak{Z}_1(n\text{Vec}_H^\sigma)$$

defines a condensable E_1 -algebra $\mathfrak{Z}_1(n\text{Vec}_H^\sigma)^{\text{op}}$ in $\text{BMod}_{n\text{Rep}(G)|n\text{Rep}(G)}((n+1)\text{Vec})^{\text{op}}$. By condensing it, we obtain the double of twisted finite gauge theory $\text{GT}_{(H,\sigma)}^{n+1} \boxtimes \overline{\text{GT}_{(H,\sigma)}^{n+1}}$ as the condensed phase D^{n+2} , and a gapped domain wall M^n such that

$$(\mathcal{M}, m) = (\text{BMod}_{\mathfrak{Z}_1(n\text{Vec}_H^\sigma)|n\text{Rep}(G)}((n+1)\text{Vec}), \mathfrak{Z}_1(n\text{Vec}_H^\sigma)).$$

The following composed monoidal functor

$$n\text{Rep}(G) \xrightarrow{f} n\text{Rep}(H) \xrightarrow{\psi_H} \mathfrak{Z}_1(n\text{Vec}_H^\sigma) \xrightarrow{f} n\text{Vec}_H^\sigma$$

defines a condensable E_1 -algebra $(n\text{Vec}_H^\sigma)^{\text{op}}$, which determines $\text{GT}_{(H,\sigma)}^{n+1}$ as the condensed phase.

4. By Theorem^{ph} 3.3.33, we can construct a 1-codimensional defect condensation from $\mathrm{GT}_{G_1}^{n+2}$ to $\mathrm{GT}_{G_2}^{n+2}$ by specifying an $n\mathrm{Rep}(G_1)$ - $n\mathrm{Rep}(G_2)$ -bimodule $\mathcal{X}$. More precisely, given such a $\mathcal{X}$, we obtain a condensable E_1 -algebra $\mathcal{A}_{\mathcal{X}}^{\mathrm{op}}$ in $\mathcal{GT}_{G_1}^{n+1}$ by the following monoidal functor

$$n\mathrm{Rep}(G_1) \rightarrow \mathrm{Fun}_{n\mathrm{Rep}(G_2)^{\mathrm{op}}}(\mathcal{X}, \mathcal{X}) =: \mathcal{A}_{\mathcal{X}}$$

such that condensing $\mathcal{A}_{\mathcal{X}}^{\mathrm{op}}$ produces $\mathrm{GT}_{G_2}^{n+2}$ as the condensed phase. Examples of such $\mathcal{X}$ can be constructed as follows.

- (a) The forgetful functor $f : n\mathrm{Rep}(G_1 \times G_2) \rightarrow n\mathrm{Rep}(H)$ for a subgroup $H \leq G_1 \times G_2$ [KZ24] defines an $n\mathrm{Rep}(G_1)$ - $n\mathrm{Rep}(G_2)$ -bimodule $\mathcal{X} = n\mathrm{Rep}(H)$.
- (b) A simple $n\mathrm{Rep}(G_1)$ - $n\mathrm{Rep}(G_2)$ -bimodule $\mathcal{X}$ is precisely a simple object in $(n+1)\mathrm{Rep}(G_1 \times G_2)$, which is of the form $\Sigma\mathcal{L}$ for a fusion $(n-1)$ -category $\mathcal{L}$ over $(n-1)\mathrm{Rep}(G_1 \times G_2)$ [KZZZ25]. If we choose $\mathcal{L} = (n-1)\mathrm{Rep}(H)$, we recover the construction in (4.a). Note that any indecomposable multi-fusion $(n-1)$ -category $\mathcal{Q}$ that is Morita equivalent to $\mathcal{L}$ produces the same simple $n\mathrm{Rep}(G_1)$ - $n\mathrm{Rep}(G_2)$ -bimodule $\mathcal{X}$, i.e., $\Sigma\mathcal{L} \simeq \Sigma\mathcal{Q}$. Note that $\mathcal{X} = n\mathrm{Rep}(H) \boxtimes \mathcal{S}$ for any separable n -category $\mathcal{S}$ gives more but obvious examples. $\heartsuit$

When G_2 is the trivial group, such constructed condensable E_1 -algebras $\mathcal{A}_{\mathcal{X}}^{\mathrm{op}}$ are all Lagrangian.

Remark 3.3.53. From all these examples and from the results of Theorem^{ph} 3.3.26, 3.3.32 and 3.3.33 and Corollary^{ph} 3.3.36 and 3.3.37, we can see that the mathematical theory of the condensations of 1-codimensional defects are much simpler than that of the condensations of anyons. It is also much simpler in constructing examples or sometimes even obtaining a classification. As we show later, it also provides some powerful tools in the study of condensable/Lagrangian E_2 -algebras in (non-degenerate) braided fusion higher categories. $\diamond$

Remark 3.3.54. Since our knowledge of (braided) fusion higher categories are very limited, only explicit examples of condensations we can provide are those in the twisted finite gauge theories in higher dimensions. Actually, there are a few other workable examples from fermionic modular 2-categories [LW19, JF20]. Examples of condensations can certainly be found there. We encourage intrigued readers to work on these cases. $\diamond$

Before we end this section, we give a few examples of 1-codimensional defect condensations in anomalous $n+2$ D topological orders. More precisely, we consider condensable E_1 -algebras in $(n+1)\mathrm{Vec}_G$, $(n+1)\mathrm{Rep}(G)$ and $(n+1)\mathrm{Vec}_G^\pi$.

Example 3.3.55. For a finite group G , $(n+1)\mathrm{Vec}_G$ is the fusion $(n+1)$ -category of topological defects on a gapped boundary of the finite gauge theory GT_G^{n+3} . The condensable E_1 -algebras in $(n+1)\mathrm{Vec}_G$ are G -graded multi-fusion n -categories, and the simple ones are G -graded fusion n -categories. A G -graded (multi-)fusion n -category is a (multi-)fusion n -category $\mathcal{A}$ equipped with a G -grading, i.e., $\mathcal{A} = \bigoplus_{g \in G} \mathcal{A}_g$, such that $x \otimes y \in \mathcal{A}_{gh}$ for any $x \in \mathcal{A}_g$ and $y \in \mathcal{A}_h$. For example, for a subgroup $H \subseteq G$ and $\omega \in Z^{n+2}(H, U(1))$, the fusion n -category $n\mathrm{Vec}_H^\omega$ is a G -graded fusion n -category with the G -grading supported on H .

For a G -graded fusion n -category $\mathcal{A}$, we give a more explicit mathematical description of the left $2\mathrm{Vec}_G$ -module $\mathrm{RMod}_{\mathcal{A}}(2\mathrm{Vec}_G)$. First, similar to Example 3.3.46, we define an $(n+1)$ -category $B_G\mathcal{A}$ as follows: (1) the objects are elements in G ; (2) the hom n -categories are $\mathrm{Hom}_{B_G\mathcal{A}}(g, h) := \mathcal{A}_{hg^{-1}}$; (3) the composition n -functors

$$\mathrm{Hom}_{B_G\mathcal{A}}(h, k) \times \mathrm{Hom}_{B_G\mathcal{A}}(g, h) = \mathcal{A}_{kh^{-1}} \times \mathcal{A}_{hg^{-1}} \rightarrow \mathcal{A}_{kg^{-1}} = \mathrm{Hom}_{B_G\mathcal{A}}(g, k)$$

are given by the tensor product functor of $\mathcal{A}$. We denote the condensation completion of $B_G\mathcal{A}$ by $\Sigma_G\mathcal{A}$, which is a separable $(n+1)$ -category. Similar to Example 3.3.46, there is an equivalence of $(n+1)\mathrm{Vec}_G$ -modules $\mathrm{RMod}_{\mathcal{A}}((n+1)\mathrm{Vec}_G) \simeq \Sigma_G\mathcal{A}$. $\heartsuit$

Example 3.3.56. For a finite group G , $(n+1)\mathrm{Rep}(G)$ is the fusion $(n+1)$ -category of topological defects on another gapped boundary of the finite gauge theory GT_G^{n+3} . There are two coordinates on $(n+1)\mathrm{Rep}(G)$.

- $(n+1)\mathrm{Rep}(G) \simeq \mathrm{Fun}(B_G, (n+1)\mathrm{Vec}) \simeq \mathrm{LMod}_{n\mathrm{Vec}_G}((n+1)\mathrm{Vec})$: The objects are separable n -categories equipped with G -action. A condensable E_1 -algebra in $(n+1)\mathrm{Rep}(G)$ is precisely a multi-fusion n -category $\mathcal{A}$ equipped with a monoidal G -action (i.e., a monoidal n -functor $G \rightarrow \mathrm{Aut}^{\otimes}(\mathcal{A})$). Such $\mathcal{A}$ is also called a G -equivariant multi-fusion n -category. A condensable E_1 -algebra $\mathcal{A}$ is simple if and only if G acts transitively on the simple direct summands of the tensor unit of $\mathcal{A}$.

- $(n+1)\text{Rep}(G) \simeq \Sigma n\text{Rep}(G) \simeq \text{RMod}_{n\text{Rep}(G)}((n+1)\text{Vec}) \simeq \text{LMod}_{n\text{Rep}(G)}((n+1)\text{Vec})$: The objects are separable n -categories equipped with an $n\text{Rep}(G)$ -action. The condensable E_1 -algebras are multi-fusion $n\text{Rep}(G)$ -module $\mathcal{B}$, i.e., $\mathcal{B}$ is a multi-fusion n -category equipped with a braided monoidal n -functor $n\text{Rep}(G) \rightarrow \mathfrak{Z}_1(\mathcal{B})$. A condensable E_1 -algebra $\mathcal{B}$ is simple if and only if it is fusion.

These two coordinates should again be related by equivariantization and de-equivariantization. Both notions should be properly generalized to higher categories. For example, the equivariantization of $\mathcal{A}$, denoted by $\mathcal{A}^G$, should consist of G -equivariant objects of $\mathcal{A}$. A G -equivariant object in $\mathcal{A}$ is an object $X \in \mathcal{A}$ equipped with an isomorphism $u_g : F_g(X) \simeq X$ for $g \in G$ and F_g is the image of g in $\text{Aut}^\otimes(\mathcal{A})$ such that the following diagram

$$\begin{array}{ccc} F_g(F_h(X)) & \xrightarrow{F_g(u_h)} & F_g(X) \\ \simeq \downarrow & & \downarrow u_g \\ F_{gh}(X) & \xrightarrow{u_{gh}} & X \end{array}$$

commutes up to higher isomorphisms. A 1-morphism should be defined by a 1-morphism in $\mathcal{A}$ that commutes with $u_g, \forall g \in G$. However, one can see that it is impossible to avoid the higher coherence data and the higher morphisms in $\mathcal{A}^G$ explicitly. It is easier to define the de-equivariantization: $\mathcal{B} \mapsto \mathcal{B}_G$. We define $\mathcal{B}_G := \text{RMod}_{[\mathbb{1}, \mathbb{1}]}(\mathcal{B})$, where $\mathbb{1}$ is the tensor unit of $n\text{Vec}$ and $[\mathbb{1}, \mathbb{1}]$ is the internal hom algebra in $n\text{Rep}(G)$ for the $n\text{Rep}(G)$ -module $n\text{Vec}$. Therefore, the notion of a G -equivariant multi-fusion n -category can also be understood as the de-equivariantization of a multi-fusion $n\text{Rep}(G)$ -module $\mathcal{B}$.

Given a condensable E_1 -algebra $\mathcal{B} \in \text{LMod}_{n\text{Rep}(G)}((n+1)\text{Vec}) \simeq (n+1)\text{Rep}(G)$, there is an equivalence of $(n+1)$ -categories $\text{RMod}_{\mathcal{B}}((n+1)\text{Rep}(G)) \simeq \text{RMod}_{\mathcal{B}}((n+1)\text{Vec}) \simeq \Sigma \mathcal{B}$. The left $(n+1)\text{Rep}(G)$ -action on $\Sigma \mathcal{B}$ is defined by a monoidal $(n+1)$ -functor

$$(n+1)\text{Rep}(G) \simeq \Sigma n\text{Rep}(G) \rightarrow \Sigma \mathfrak{Z}_1(\mathcal{B}) \simeq \mathfrak{Z}_0(\Sigma \mathcal{B}),$$

which is induced from the braided monoidal functor $n\text{Rep}(G) \rightarrow \mathfrak{Z}_1(\mathcal{B})$. Recall that such braided monoidal functors $n\text{Rep}(G) \rightarrow \mathcal{M}$ for a non-degenerate braided fusion n -category $\mathcal{M}$ classify Lagrangian E_1 -algebras in $\mathcal{GT}_G^{n+2}$ by Corollary 3.3.40. $\heartsuit$

Example 3.3.57. For a finite group G and $\pi \in Z^{n+3}(G, U(1))$, $(n+1)\text{Vec}_G^\pi$ is the fusion $(n+1)$ -category of topological defects on a gapped boundary of the twisted finite gauge theory $\text{GT}_{(G, \pi)}^{n+3}$. It is the same $(n+1)$ -category as the $(n+1)\text{Vec}_G$ but with the associativity twisted by π in its highest coherence data.

A simple condensable E_1 -algebra in $(n+1)\text{Vec}_G^\pi$ is precisely a ‘ π -twisted G -graded fusion n -category’, which has the same defining data as a G -graded fusion n -category but with the associativity twisted by π in its highest coherence data.

- For example, given a subgroup $H \leq G$ and an $(n+3)$ -cochain $\pi \in C^{n+3}(H, U(1))$ such that $d\psi = \pi|_H$, there is a simple condensable E_1 -algebra $n\text{Vec}_H^\psi \in (n+1)\text{Vec}_G^\pi$ with the obvious G -grading supported on H and the associativity twisted by ψ in its highest coherence data. $\heartsuit$

The underlying $(n+1)$ -category of $\text{RMod}_{n\text{Vec}_H^\psi}((n+1)\text{Vec}_G^\pi)$ is $(n+1)\text{Vec}_{G/H} \simeq (n+1)\text{Vec}^{\oplus G/H}$. The left $(n+1)\text{Vec}_G^\pi$ -action on it is defined by the left translation of G on G/H . The associativity of the action is twisted by an $(n+3)$ -cocycle in $Z^{n+3}(G; \text{Fun}(G/H, U(1)))$, which is determined by ψ under the isomorphism $H^{n+3}(H; U(1)) \simeq H^{n+3}(G; \text{Fun}(G/H, U(1)))$.

4 Condensations of 2-codimensional topological defects

In previous section, we have seen that a condensation of 1-codimensional topological defects defines a phase transition from C^{n+1} to a new phase D^{n+1} , where both phase C^{n+1} and D^{n+1} are potentially anomalous $n+1$ D topological orders. When we restrict to the case both C^{n+1} and D^{n+1} are anomaly-free and simple, both categories of topological defects $\mathcal{C}$ and $\mathcal{D}$ are determined by $\Omega\mathcal{C}$ and $\Omega\mathcal{D}$, respectively. In this case, it is possible to define a condensation in C^{n+1} by condensing a 2-codimensional topological defect in $\Omega\mathcal{C}$ directly. When $n = 2$, such a condensation is precisely what was known as an anyon condensation (or boson condensation) defined in the non-degenerate braided fusion 1-category $\Omega\mathcal{C}$ [MS89c, BSS02, BSS03, BS09, BSH09, BSS11, KS11a, Lev13, BJQ13, Kon14]. When C^{n+1} is anomalous, it is still possible to define a condensation of 2-codimensional topological defect A by first condensing it to a condensed defect ΣA of codimension 1 followed by a condensation of ΣA . We postpone the study of this two-step condensation to Section 5.

In this section, we first review the theory of anyon condensations in 2+1D anomaly-free simple topological orders with some new details and from some new perspectives; then we generalize it to higher dimensions.

4.1 Anyon condensations in 2+1D

In Section 4.1.1, we review the main results of an algebraic bootstrap approach towards anyon condensation [Kon14] but from a new perspective of higher algebras and higher representations. We also introduce necessary new notations along the way. In Section 4.1.2, we provide a lot of concrete examples of anyon condensations. In Section 4.1.3, we introduce a geometric approach, which can be generalized automatically to higher dimensions.

4.1.1 Algebraic approach

We re-derive the main results in [Kon14] but from a new perspective explained in Section 2.3.2. Assume that an anyon condensation occurs in a disk-like region within a 2+1D anomaly-free simple topological order C^3 as depicted in Figure 12. We denote the condensed topological order by D^3 . We assume that D^3 is also simple⁴⁷. We denote the gapped domain wall by M^2 . The categories of all topological defects in C^3 and D^3 are $\mathcal{C}$ and $\mathcal{D}$, respectively; and that of topological defects of codimension 2 (or anyons) are $\Omega\mathcal{C}$ and $\Omega\mathcal{D}$, respectively. Mathematically, $M^2 = (\mathcal{M}, m)$, where $\mathcal{M}$ is the category of walls conditions and m labels a single wall condition and topological particles on M^2 form a multi-fusion 1-category $\Omega_m\mathcal{M}$.




Figure 12: an anyon condensation in C^3 depicted in the spatial dimension

It is possible to derive a precise relation among the categories $\Omega\mathcal{C}, \Omega\mathcal{D}, \Omega_m\mathcal{M}$ via natural physical requirements as shown in [Kon14].

1. Anyons in D^3 come from those in C^3 . In other words, an anyon in D^3 is automatically an anyon in C^3 , i.e. $\text{ob}(\Omega\mathcal{D}) \subset \text{ob}(\Omega\mathcal{C})$. All fusion-splitting channels among anyons in $\Omega\mathcal{D}$ come from those in $\Omega\mathcal{C}$, i.e. $\text{hom}_{\Omega\mathcal{D}}(x, y) \subset \text{hom}_{\Omega\mathcal{C}}(x, y)$ for $x, y \in \text{ob}(\Omega\mathcal{D})$. As a consequence, $\Omega\mathcal{D}$ is a sub-category of $\Omega\mathcal{C}$.
2. The trivial particle $1_{\mathbb{1}_{\mathcal{D}}}$ in D^3 , i.e., the tensor unit of $\Omega\mathcal{D}$, is a non-trivial particle A in $\Omega\mathcal{C}$ unless the condensation is trivial.

⁴⁷If D^3 is not simple, it simply means that the associated condensable E_2 -algebra A in $\Omega\mathcal{C}$ is not simple. The result can be easily derived from the A -being-simple case. Note that, in this case, the fact $\mathcal{D} \neq \Sigma\Omega\mathcal{D}$ adds some slight inconvenience to state the results. So we simply leave the non-simple cases to intrigued readers.

3. Similar to 1+1D cases, particle condensation is triggered by introducing interactions among particles. For $a, b \in \Omega\mathcal{D} \subset \Omega\mathcal{C}$, a sub-Hilbert space $a \otimes_{\Omega\mathcal{D}} b$ of the Hilbert space associated to $a \otimes_{\Omega\mathcal{C}} b$ becomes energy favorable due to the interaction. In other words, the condensation process produces a family of projections (called condensation maps):

$$a \otimes_{\Omega\mathcal{C}} b \xrightarrow{P_{a,b}} a \otimes_{\Omega\mathcal{D}} b, \quad \forall a, b \in \Omega\mathcal{D}.$$

In particular, we have the following special condensation maps:

$$\begin{aligned} \mu_A : A \otimes_{\Omega\mathcal{C}} A &\xrightarrow{P_{AA}} A \otimes_{\Omega\mathcal{D}} A \simeq A. \\ \forall x \in \Omega\mathcal{D}, \quad \mu_x^L : A \otimes_{\Omega\mathcal{C}} x &\xrightarrow{P_{Ax}} A \otimes_{\Omega\mathcal{D}} x \simeq x, \quad \mu_x^R : x \otimes_{\Omega\mathcal{C}} A \xrightarrow{P_{xA}} x \otimes_{\Omega\mathcal{D}} A \simeq x. \end{aligned}$$

Moreover, since the trivial anyon $1_{\mathbb{1}_\mathcal{C}}$ in $\Omega\mathcal{C}$ should condense into the trivial anyon in $\Omega\mathcal{D}$, we expect to have a morphism $\eta_A : 1_{\mathbb{1}_\mathcal{C}} \rightarrow A$.

4. Similar to the 1+1D cases, the triple (A, μ_A, η_A) defines a separable algebra in $\Omega\mathcal{C}$ satisfying a new property: A is commutative. We do not repeat the reason provided for this commutativity in [Kon14]. Instead, we provide a new proof from the perspective of Section 2.3.2. Consider the physical configuration depicted left picture in Figure 13. By the same arguments in 1+1D case, there are two algebraic structures on A . One algebraic structure is defined for the horizontal fusion product $\otimes^1$, i.e. $\mu_A^1 : A \otimes^1 A \rightarrow A$. The other is defined for the vertical fusion product $\otimes^2$, i.e. $\mu_A^2 : A \otimes^2 A \rightarrow A$. The following diagram should be automatically commutative.

$$\begin{array}{ccc} (A \otimes^1 A) \otimes^2 (A \otimes^1 A) & \xrightarrow[\simeq]{\delta_{AAAA}} & (A \otimes^2 A) \otimes^1 (A \otimes^2 A) \\ \mu_A^1 \otimes^2 \mu_A^1 \downarrow & & \downarrow \mu_A^2 \otimes^1 \mu_A^2 \\ A \otimes^2 A & & A \otimes^1 A \\ \mu_A^2 \searrow & & \swarrow \mu_A^1 \\ & A & \end{array} \quad (4.1.1)$$

Notice that $(\mu_A^2 \otimes^1 \mu_A^2) \circ \delta_{AAAA}$ defines an algebraic structure on $A \otimes^1 A$ and $(\mu_A^1 \otimes^2 \mu_A^1) \circ \delta_{AAAA}^{-1}$ defines an algebraic structure on $A \otimes^2 A$. Then above commutativity diagram simply says that both morphisms $\mu_A^2 : A \otimes^2 A \rightarrow A$ and $\mu_A^1 : A \otimes^1 A \rightarrow A$ are algebra homomorphisms.

The unit morphism $\eta_A : 1_{\mathbb{1}_\mathcal{C}} \rightarrow A$ is the same for both algebraic structures. We obtain a commutative diagram:

$$\begin{array}{ccccc} 1_{\mathbb{1}_\mathcal{C}} \otimes^1 1_{\mathbb{1}_\mathcal{C}} & \xrightarrow{\eta_A \otimes^1 1} & A \otimes^1 1_{\mathbb{1}_\mathcal{C}} & \xrightarrow{1 \otimes \eta_A} & A \otimes^1 A \\ \simeq \downarrow & & \downarrow \simeq & \swarrow \mu_A^1 & \\ 1_{\mathbb{1}_\mathcal{C}} & \xrightarrow{\eta_A} & A & & \end{array}$$

where the commutativity of the left square is due to the naturalness of the vertical $\simeq$ and that of the right triangle is due to the unit property of (A, μ_A^1, η_A) . Then the commutativity of the outer-diagram means that $\eta_A : 1_{\mathbb{1}_\mathcal{C}} \rightarrow (A, \mu_A^1, \eta_A)$ is an algebraic homomorphism. Similarly, one can show that $\eta_A : 1_{\mathbb{1}_\mathcal{C}} \rightarrow (A, \mu_A^2, \eta_A)$ is also an algebra homomorphism.

In summary, both $\mu_A^2 : A \otimes^2 A \rightarrow A$ and η_A are morphisms in $\text{Alg}_{E_1}(\Omega\mathcal{C})$. Therefore, (A, μ_A^2, η_A) defines an algebra in $\text{Alg}_{E_1}(\Omega\mathcal{C})$. In other words, $A \in \text{Alg}_{E_1}(\text{Alg}_{E_1}(\Omega\mathcal{C})) =: \text{Alg}_{E_2}(\Omega\mathcal{C})$. This fact automatically implies the compatibility between μ_A^1 and μ_A^2 and the commutativity of A . Indeed, by composing the diagram (4.1.1) with the unit morphism $\eta_A : 1_{\mathbb{1}_\mathcal{C}} \rightarrow A$ that are compatible with (2.3.5) and (2.3.6), we obtain two commutative diagrams:

$$\begin{array}{ccc} A \otimes^2 A & \xrightarrow{\simeq} & A \otimes^1 A \\ \mu_A^2 \searrow & & \swarrow \mu_A^1 \\ & A & \end{array} \quad \begin{array}{ccc} A \otimes^1 A & \xrightarrow{\simeq} & A \otimes^1 A \\ \mu_A^1 \searrow & & \swarrow \mu_A^1 \\ & A & \end{array} \quad (4.1.2)$$



Figure 13: a deconfined particle x as an E_2 - A -module in $\Omega\mathcal{C}$

where the ‘ $\simeq$ ’ in the first diagram is defined in (2.3.5), and the ‘ $\simeq$ ’ in the second diagram is precisely the braiding defined in (2.3.6).

Moreover, we require the commutative algebra A to be separable (i.e., there exists a bimodule map $\Delta_A : A \rightarrow A \otimes A$ such that $\mu_A^1 \circ \Delta_A = \text{id}_A$). Mathematically, this condition is needed for $\Omega\mathcal{D}$ and $\Omega_m\mathcal{M}$ to be separable 1-categories. Physically, this condition is a stability condition of the vacuum A as explained in [Kon14]. A commutative separable algebra in $\Omega\mathcal{C}$ is called a *condensable E_2 -algebra* in $\Omega\mathcal{C}$. Such an algebra is also equipped naturally with a Frobenius algebra structure (see [FRS02, Kon14]). We use the notation $A \in \text{Alg}_{E_2}^c(\Omega\mathcal{C})$ to represent the statement that A is a condensable E_2 -algebra in $\Omega\mathcal{C}$. Since D^3 is simple, A must be simple (i.e., $\dim \text{hom}_{\mathcal{C}}(1_{\mathbb{1}_{\mathcal{C}}}, A) = 1$). A general condensable E_2 -algebra in a braided fusion 1-category is a direct sum of simple ones.

5. Anyons in $\Omega\mathcal{D}$ are deconfined particles, which are necessarily endowed with a 2-dimensional action of the trivial particle $1_{\mathbb{1}_{\mathcal{D}}} = A$ in $\Omega\mathcal{D}$. Mathematically, it amounts to say that anyons in $\Omega\mathcal{D}$ are E_2 - A -modules (or local A -modules) in $\Omega\mathcal{C}$. We explain this fact in a few steps from the perspective of Section 2.3.2.

- For $A, B \in \text{Alg}_{E_2}^c(\Omega\mathcal{C})$, both categories $\text{LMod}_A(\Omega\mathcal{C})$ and $\text{RMod}_A(\Omega\mathcal{C})$ are monoidal with the monoidal structures defined as follows (see Remark 4.1.2 for another monoidal structure).

$$(4.1.3)$$

- (a) The tensor unit in $\text{LMod}_A(\Omega\mathcal{C})$ is A and the tensor product is given by $\otimes_A^1$, which needs an explanation. Note that the left A -action $\mu_A^1 : A \otimes^2 x \rightarrow x$ induces two horizontal A -actions on x (denoted by $\mu_A^{\leftarrow}$ and $\mu_A^{\rightarrow}$) by adiabatic moves as illustrated in the picture in (4.1.3). These two horizontal A -actions induces a tensor product $x \otimes_A^1 x'$ defined by the coequalizer of $x \otimes^1 A \otimes^1 x' \rightrightarrows x \otimes^1 x'$ for $x, x' \in \text{LMod}_A(\Omega\mathcal{C})$. Note that $x \otimes_A^1 x'$ has a structure of left A -module induced from that of x .
- (b) The tensor unit in $\text{RMod}_B(\Omega\mathcal{C})$ is B and the tensor product is given by $\otimes_B^1$, which needs an explanation. Note that the right B -action $\nu_B^1 : B \otimes^2 y \rightarrow y$ induces two horizontal B -actions on y (denoted by $\nu_B^{\leftarrow}$ and $\nu_B^{\rightarrow}$) by adiabatic moves as illustrated in the picture in (4.1.3). These two horizontal B -actions induces a tensor product $y \otimes_B^1 y'$ defined by the coequalizer of $y \otimes^1 B \otimes^1 y' \rightrightarrows y \otimes^1 y'$ for $y, y' \in \text{RMod}_B(\Omega\mathcal{C})$. Note that $y \otimes_B^1 y'$ has a structure of right A -module induced from that of y' .

As a consequence, the category $\text{BMod}_{A/B}(\Omega\mathcal{C})$ inherits a natural monoidal structure due to the following equivalence of 2-categories:

$$\text{BMod}_{A/B}(\Omega\mathcal{C}) \simeq \text{LMod}_A(\Omega\mathcal{C}) \boxtimes_{\Omega\mathcal{C}} \text{RMod}_B(\Omega\mathcal{C}). \quad (4.1.4)$$

More explicitly, the tensor product in $\text{BMod}_{A|B}(\Omega\mathcal{C})$, denoted by $\otimes_{A|B}^1$, is the one induced from the tensor product $\otimes_A^1 \boxtimes_{\Omega\mathcal{C}} \otimes_B^1$ in $\text{LMod}_A(\Omega\mathcal{C}) \boxtimes_{\Omega\mathcal{C}} \text{RMod}_B(\Omega\mathcal{C})$. The tensor unit in $\text{BMod}_{A|B}(\Omega\mathcal{C})$ is $A \otimes^2 B$, which is induced from the tensor unit $A \boxtimes_{\Omega\mathcal{C}} B$ in $\text{LMod}_A(\Omega\mathcal{C}) \boxtimes_{\Omega\mathcal{C}} \text{RMod}_B(\Omega\mathcal{C})$. The physical meaning of this tensor product is illustrated in a special case in (4.1.5).

- We are only interested in the case $B = A$. In this case, $\text{Mod}_A^{\text{E}_1}(\Omega\mathcal{C}) = \text{BMod}_{A|A}(\Omega\mathcal{C})$ is equipped with the monoidal structure with the tensor product $\otimes_{A|A}^1$ and the tensor unit $A \otimes^2 A$. Note that the E_2 -algebra structure on $A \in \text{Alg}_{\text{E}_2}^c(\Omega\mathcal{C})$ endows A with a structure of an E_1 -algebra in $\text{Mod}_A^{\text{E}_1}(\Omega\mathcal{C})$. In particular, the multiplication is given by $\mu_A^1 : A \otimes^1 A \rightarrow A$, and the unit morphism is given by $\mu_A^2 : A \otimes^2 A \rightarrow A$, which is an algebra homomorphism. Namely, $A \in \text{Alg}_{\text{E}_1}^c(\text{Mod}_A^{\text{E}_1}(\Omega\mathcal{C}))$. Note that, in this case, the tensor unit $A \otimes^2 A$ is not simple, and A is a direct summand of the tensor unit. By the structure theorem of a multi-fusion 1-category, $\text{Mod}_A^{\text{E}_1}(\text{Mod}_A^{\text{E}_1}(\Omega\mathcal{C}))$ is a full subcategory of $\text{Mod}_A^{\text{E}_1}(\Omega\mathcal{C})$.
- For $x \in \text{Mod}_A^{\text{E}_1}(\Omega\mathcal{C})$, the left $(A \otimes^2 A)$ -action on x is illustrated below.

$$(A \otimes^2 A) \otimes_{A|B}^1 (x) = \begin{array}{c} \Omega\mathcal{C} \quad A \otimes_B^1 B \\ \quad \quad \quad \otimes_B^2 \\ \quad \quad \quad x \\ \quad \quad \quad \otimes_A^2 \\ A \otimes_A^1 A \end{array} \xrightarrow{\sim} x \quad (4.1.5)$$

$x^2 \uparrow \quad x^1 \rightarrow$

The right $(A \otimes^2 A)$ -actions on x is similar. Note that $x \in \text{Mod}_A^{\text{E}_1}(\text{Mod}_A^{\text{E}_1}(\Omega\mathcal{C}))$ if and only if the left (resp. right) $(A \otimes^2 A)$ -actions on x factor through $A = A \otimes_A^2 A$. This condition is equivalent to the conditions that $\mu_x^\rightarrow = \nu_x^\rightarrow$ (resp. $\mu_x^\leftarrow = \nu_x^\leftarrow$). These conditions are further equivalent to the commutativity of the following diagram (where we set $\mu_x^\downarrow = \nu_x^\downarrow$):

$$\begin{array}{ccccc} & & x \otimes^2 A & & \\ & \nearrow \simeq & \downarrow \mu_x^\downarrow & \searrow \simeq & \\ A \otimes^1 x & \xrightarrow{\mu_x^\rightarrow} & x & \xleftarrow{\mu_x^\leftarrow} & x \otimes^1 A \\ & \nwarrow \simeq & \uparrow \mu_x^\uparrow & \swarrow \simeq & \\ & & A \otimes^2 x & & \end{array} \quad (4.1.6)$$

where all ' $\simeq$ ' are defined by (2.3.5) or its inverse. The composition of adjacent two such $\simeq$'s defines a braiding as in (2.3.6) and (2.3.7). Such a module x , i.e. equipped with a 2-dimensional A -actions, is called an E_2 - A -module. As a consequence, we have provided a physical proof of the following equivalence of 1-categories:

$$\text{Mod}_A^{\text{E}_2}(\Omega\mathcal{C}) \simeq \text{Mod}_A^{\text{E}_1}(\text{Mod}_A^{\text{E}_1}(\Omega\mathcal{C})), \quad (4.1.7)$$

which is a special case of a more general result (see Corollary 5.2.5). An E_2 - A -module in $\Omega\mathcal{C}$ is precisely a local A -module in the usual sense [KO02, FFRS06, Kon14]. It is clear that A itself is an E_2 - A -module.


6. Note that $\text{Mod}_A^{\text{E}_1}(\Omega\mathcal{C})$ has another natural monoidal structure with the tensor product $\otimes_A^2$ and the tensor unit A . The category $\text{Mod}_A^{\text{E}_1}(\text{Mod}_A^{\text{E}_1}(\Omega\mathcal{C}))$ is naturally equipped with two tensor products $\otimes_A^2$ and $\otimes_A^1$ that defines an E_2 -monoidal structure on $\text{Mod}_A^{\text{E}_1}(\text{Mod}_A^{\text{E}_1}(\Omega\mathcal{C}))$.
7. If an x -particle does allow A -actions from all directions but the commutativity of the diagram in (4.1.6) does not hold, then this non-commutativity causes a strong interference effect around the x -particle that confines the particle. Such confined particles are necessarily confined to the gapped domain wall M^2 between C^3 and D^3 . As a consequence, particles on the wall $M^2 = (\mathcal{M}, m)$ form a fusion 1-category $\Omega_m \mathcal{M} = \text{RMod}_A(\Omega\mathcal{C})$.

Remark 4.1.1. Note that the notion of an E_2 - A -module for an E_2 -algebra A in $\Omega\mathcal{C}$ is defined by four actions $\mu_x^\rightarrow, \mu_x^\downarrow, \mu_x^\leftarrow, \mu_x^\uparrow$ in four directions such that they are mutually commuting in the sense that the following diagram:

$$\begin{array}{ccc}
 A \otimes^2 (A \otimes^1 x) & \xrightarrow{\simeq} & A \otimes^1 (A \otimes^2 x) \\
 \downarrow 1_A \otimes^2 \mu_A^\rightarrow & & \downarrow \mu_A^\uparrow \\
 A \otimes^2 x & & A \otimes^1 x \\
 \searrow \mu_x^\uparrow & & \swarrow \mu_x^\rightarrow \\
 & x &
 \end{array} \tag{4.1.8}$$

and other similar diagrams commute, and are isomorphic in the sense of (4.1.6). In other words, there is essentially only one 2-dimensional A -action on x . This notion immediately generalizes to the notion of an E_k - B -module for an E_k -algebra B in an E_k -monoidal category (see Definition 5.2.4 and Corollary 5.2.5). $\diamond$


Remark 4.1.2. Actually, $\text{BMod}_{A|A}(\Omega\mathcal{C})$ is equipped with the second monoidal structure with the usual tensor product $\otimes_A^2$ and the tensor unit A . These two monoidal structures are not equivalent. They can be obtained from two different fusion processes, respectively, as illustrated below.



Note that the first monoidal structure is Morita equivalent to the fusion 1-category $\text{Mod}_A^{E_2}(\Omega\mathcal{C})$ and the second one is Morita equivalent to the fusion 1-category $\Omega\mathcal{C}$. Therefore, these two monoidal structures are not even Morita equivalent when $\dim A > 1$. We want to emphasize that the existence of multiple higher monoidal structures on a single higher category is a ubiquitous and important feature of higher categories and higher condensation theory. $\diamond$

In summary, we obtain the following categorical description of an anyon condensation.

Theorem^{ph} 4.1.3 ([Kon14]). If an anyon condensation in a 2+1D anomaly-free simple topological order $\mathcal{C}^3$ produces an anomaly-free simple condensed phase $\mathcal{D}^3$ and a gapped domain wall $\mathcal{M}^2$ (see Figure 12 and the picture below).



1. The vacuum particle in $\Omega\mathcal{D}$ can be identified with a simple condensable E_2 -algebra A in $\Omega\mathcal{C}$, which is a composite boson⁴⁸ (see Remark 4.1.5). Moreover, $\Omega\mathcal{D}$ consists of all deconfined particles and can be identified with the category $\text{Mod}_A^{E_2}(\Omega\mathcal{C})$ of E_2 - A -modules (or equivalently, local A -modules) in $\mathcal{C}$, i.e., $\Omega\mathcal{D} = \text{Mod}_A^{E_2}(\Omega\mathcal{C})$.
2. Particles on the wall $\mathcal{M}^2 = (\mathcal{M}, m)$ include all confined and deconfined particles, and can be identified with the fusion 1-category $\Omega_m \mathcal{M} = \text{RMod}_A(\Omega\mathcal{C})$.
3. Anyons in $\Omega\mathcal{C}$ move to the wall according to the central functor⁴⁹ $-\otimes A: \Omega\mathcal{C} \rightarrow \text{RMod}_A(\Omega\mathcal{C})$ defined by $x \mapsto x \otimes A$ for all $x \in \Omega\mathcal{C}$.

⁴⁸It is really a boson condensation [BSS03, BS09] instead of an ‘anyon condensation’, a term which is wildly used in literature.

⁴⁹Being a central functor means that $-\otimes A$ factors through a braided monoidal functor $g: \Omega\mathcal{C} \rightarrow \text{RMod}_A(\Omega\mathcal{C})$, i.e. it coincides with the composed functor $\Omega\mathcal{C} \xrightarrow{g} \mathfrak{Z}_1(\text{RMod}_A(\Omega\mathcal{C})) \xrightarrow{f} \text{RMod}_A(\Omega\mathcal{C})$, where f is the forgetful functor.

4. Anyons in $\Omega\mathcal{D}$ move to the wall according to the embedding $\Omega\mathcal{D} = \text{Mod}_A^{E_2}(\Omega\mathcal{C}) \hookrightarrow \text{RMod}_A(\Omega\mathcal{C})$, then can move out to the D^3 -side freely.

If A is also Lagrangian, i.e. $(\dim A)^2 = \dim(\Omega\mathcal{C})$ [KR09, DMNO13], we have $\Omega\mathcal{D} = \text{Mod}_A^{E_2}(\Omega\mathcal{C}) \simeq \text{Vect}$ [DMNO13]⁵⁰. In this case, M^2 is a gapped boundary of C^3 and consists of all confined particles and deconfined particles. $\square$

Remark 4.1.4. Different from the condensation of 1-codimensional defects (recall Theorem^{ph} 3.2.4), such defined anyon condensation process can not be reversed mathematically. More precisely, there is no condensable E_2 -algebra in $\Omega\mathcal{D}$ such that its condensation in D^3 reproduces C^3 . The simplest way to see this fact is through the following formula of quantum dimensions [DMNO13]:

$$\dim \text{Mod}_A^{E_2}(\Omega\mathcal{C}) = \frac{\dim \Omega\mathcal{C}}{(\dim A)^2}. \quad (4.1.9)$$

Since $\dim A > 1$ if $A \neq 1_{\mathcal{C}}$, a non-trivial particle condensation in $\Omega\mathcal{C}$ always reduces the quantum dimension. Therefore, there is no way to find a condensable E_2 -algebra in $\Omega\mathcal{D}$ to reproduce $\Omega\mathcal{C}$. On the other hand, the phase transition can always run in the reversed direction. This shortcoming can be fixed by taking all 1-codimensional topological defects in C^3 into account as in Section 3. $\diamond$

Remark 4.1.5. When we derive the properties of the condensable E_2 -algebra A in $\mathcal{C}$, we have ignored some properties that have played important roles in [Kon14].

1. We do not emphasize the unitary structure and the Frobenius algebra structures (as in Remark 3.2.1), which allows us to obtain a lot of graphic constructions of ingredients of the condensed phases, such as the screening operator [Kon14, Eq. (20)]. The reason is that the mathematical theory of unitary higher categories (see [FHJFK⁺24] for a recent progress) and that of Frobenius algebras in higher monoidal categories are still underdeveloped.
2. Another important property of A is that it should preserves the twist (or the topological spin) [Kon14, Eq. (12)]. As a consequence, A must have a trivial twist, i.e., $\theta_A = \text{id}_A$. It further means that A is a composite boson. This, of course, is a fundamental physical result. In 2+1D, however, a simple condensable E_2 -algebra A is automatically equipped with a structure of a simple special symmetric Frobenius algebra [FRS02, Kon14], which automatically implies that $\theta_A = \text{id}_A$. In higher dimensions, it is not clear to us how to generalize the notion of a twist to higher categories. We believe that this is a very important question because it allows us to define or characterize the statistical properties of higher dimensional topological defects. We hope to come back to this point in the future. $\diamond$

Remark 4.1.6. We give a brief remark on later developments. After the categorical description of an anyon condensation was fully established in [Kon14] (see Theorem^{ph} 4.1.3), some further discussions and explanations of this categorical description appeared in literature [ERB14, HW15, NHvKSB16a, NHvKSB16b] (see [Bur18] for a physical review), and concrete examples were computed recently in [LY23, Kik23b, Kik23a, KKH24, Kik24]. The anyon condensation theory for 2+1D topological orders was used and generalized to 2+1D SPT/SET orders [HW14, LKW16a, LKW16b, JR17, BJLP19]. Since the anyon condensation theory summarized in Theorem^{ph} 4.1.3 is model independent, it is very important to construct lattice models to realize a given anyon condensation. The microscopic or physical constructions of anyon condensations appeared first in the tensor network approach [DIHVS17, GRIPG17, XZZ20]. The lattice model constructions of anyon condensations appeared only recently in [HHHW22, ZHWHW23, CGHP23, LB24, ZW25b, ZW25a]. We do expect more progress in this direction in the coming future. Recently, anyon condensation theory has been used systematically in the study of topological phase transitions in 1+1D [CJKYZ20, CW23, LY23, CJW25, CAW24]. It was also used recently to study many different physical systems [CGS23, ZVW25, CGS24, Tak25, CGSO25, HJSWZ25]. $\diamond$

Remark 4.1.7. A 2+1D topological order is well-defined on an open 2-disk (of large size) in space dimension. Roughly speaking, a 3D topological order can be viewed as a 3D fully dualizable TQFT on an open disk (recall Remark 1.2.10). Therefore, it is natural to ask if an anyon condensation can be defined for a 3D TQFT

⁵⁰This fact follows immediately from the formula (4.1.9) of quantum dimensions.

defined on non-trivial spacetime manifolds. Parallel to anyon condensations in 2+1D topological orders, there is a program of ‘orbifold construction’ (or ‘gauging defects’) in 2+1D Reshetikhin-Turaev TQFT [RT91] developed by Carqueville, Runkel, Schaumann and Mulevicius in a series of works [CR16, CRS18, CRS20, CMRSS21, Mul24, MR23, CM23, CMRSS24] (see [Car23] for a survey). More precisely, they gave a state-sum construction of a new 2+1D Reshetikhin-Turaev TQFT from an old one based on the so-called ‘an orbifold datum’. This program is not limited to 2+1D [CRS19]. $\diamond$

4.1.2 Examples

Example 4.1.8. Take C^3 to be the 2+1D $\mathbb{Z}_2$ topological order, which is realizable by the 2+1D toric code model [Kit03]. In this case, the category of particles in C^3 is $\Omega C = \mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$, which is the Drinfeld center of $\text{Rep}(\mathbb{Z}_2)$. It contains four simple anyons $1, e, m, f$ with the fusion rule

$$e \otimes e \simeq m \otimes m \simeq f \otimes f \simeq 1, \quad f \simeq e \otimes m \simeq m \otimes e.$$

The double braiding of e with m is -1 . The self-braiding of e and m are trivial, and the self-braiding of f is -1 . There are two nontrivial condensable E_2 -algebra $A_e = 1 \oplus e$ and $A_m = 1 \oplus m$, both of which are Lagrangian.

1. The multiplication $\mu : A_e \otimes A_e \rightarrow A_e$ of A_e is given by

$$A_e \otimes A_e = 1 \otimes 1 \oplus e \otimes e \oplus 1 \otimes e \oplus e \otimes 1 = (1 \oplus 1) \oplus (e \oplus e) \xrightarrow{(1 \oplus 1) \oplus (1 \oplus 1)} 1 \oplus e = A_e,$$

and the unit of A_e is $1 \xrightarrow{1 \oplus 0} 1 \oplus e = A_e$. It is separable because the following morphism $\Delta : A_e \rightarrow A_e \otimes A_e$ is an A_e - A_e -bimodule map:

$$A_e = 1 \oplus e \xrightarrow{(\frac{1}{2} \oplus \frac{1}{2}) \oplus (\frac{1}{2} \oplus \frac{1}{2})} (1 \oplus 1) \oplus (e \oplus e) = 1 \otimes 1 \oplus e \otimes e \oplus 1 \otimes e \oplus e \otimes 1 = A_e \otimes A_e,$$

and splits the multiplication, i.e., $\mu \circ \Delta = \text{id}_{A_e}$. Since $\dim \mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2)) = 4$ and $\dim A_e = 2$, A_e is Lagrangian, i.e. $\text{Mod}_{A_e}^{E_2}(\Omega C) \simeq \text{Vec}$. Condensing A_e produces the trivial 2+1D topological order and a gapped 1+1D boundary of C^3 , which is called the rough boundary [BK98]. There are two simple anyons on the rough boundary, which correspond to two simple right A_e -modules: one is A_e itself, and the other one is the free module $M_e := m \otimes A_e = f \otimes A_e$. The fusion rule is given by $M_e \otimes_{A_e} M_e \simeq A_e$ and the associator is trivial. Hence $\text{RMod}_{A_e}(\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))) \simeq \text{Rep}(\mathbb{Z}_2)$. The right module M_e is not local because the double braiding of e and m is -1 . Thus A_e itself is the only simple local A_e -module.

2. Similarly, the multiplication of A_m is given by

$$A_m \otimes A_m = 1 \otimes 1 \oplus m \otimes m \oplus 1 \otimes m \oplus m \otimes 1 = (1 \oplus 1) \oplus (m \oplus m) \xrightarrow{(1 \oplus 1) \oplus (1 \oplus 1)} 1 \oplus m = A_m,$$

and the unit of A_m is $1 \xrightarrow{1 \oplus 0} 1 \oplus m = A_m$. Again A_m is Lagrangian. Condensing A_m gives the trivial phase and also produces a gapped boundary of the 2+1D $\mathbb{Z}_2$ topological order, which is called the smooth boundary. There are two simple anyons on the smooth boundary, which correspond to two simple right A_m -modules: one is A_m itself, and the other one is the free module $M_m := e \otimes A_m = f \otimes A_m$. The fusion rule is given by $M_m \otimes_{A_m} M_m \simeq A_m$ and the associator is trivial. Hence $\text{RMod}_{A_m}(\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))) \simeq \text{Vec}_{\mathbb{Z}_2}$. The right module A_m is local but M_m is not. $\heartsuit$

Example 4.1.9. Consider the 2+1D twisted $\mathbb{Z}_2$ topological order $\text{GT}_{(\mathbb{Z}_2, \omega)}^3$, where $\omega \in Z^3(\mathbb{Z}_2; U(1))$ represents the nontrivial cohomology class in $H^3(\mathbb{Z}_2; U(1)) \simeq \mathbb{Z}_2$. More explicitly we take $\omega(1, 1, 1) = -1$ and $\omega(a, b, c) = 1$ if one of a, b, c is 0. Note that $\text{GT}_{(\mathbb{Z}_2, \omega)}^3$ can be realized by the double semion model. In this case, the category of anyons is given by the Drinfeld center of $\text{Vec}_{\mathbb{Z}_2}^\omega$, i.e., $\Omega \mathcal{GT}_{(\mathbb{Z}_2, \omega)} = \mathfrak{Z}_1(\text{Vec}_{\mathbb{Z}_2}^\omega)$. There are four simple anyons $1, s, \bar{s}, s\bar{s}$ in $\mathfrak{Z}_1(\text{Vec}_{\mathbb{Z}_2}^\omega)$ with the following fusion rules:

$$s \otimes s \simeq \bar{s} \otimes \bar{s} \simeq 1, \quad s\bar{s} \simeq s \otimes \bar{s} \simeq \bar{s} \otimes s.$$

The double braiding of s and $\bar{s}$ is trivial. The self-braiding of s is i , the self-braiding of $\bar{s}$ is $-i$, and the self-braiding of $s\bar{s}$ is 1.

There is only one nontrivial condensable E_2 -algebra $A = \mathbb{1} \oplus s\bar{s}$. The multiplication is given by

$$A \otimes A = \mathbb{1} \otimes \mathbb{1} \oplus s\bar{s} \otimes s\bar{s} \oplus \mathbb{1} \otimes s\bar{s} \oplus s\bar{s} \otimes \mathbb{1} = (\mathbb{1} \oplus \mathbb{1}) \oplus (s\bar{s} \otimes s\bar{s}) \xrightarrow{(\mathbb{1} \oplus \mathbb{1}) \oplus (\mathbb{1} \oplus \mathbb{1})} \mathbb{1} \oplus s\bar{s} = A.$$

Condensing A produces a gapped boundary of $\text{GT}_{(\mathbb{Z}_2, \omega)}^3$. There are two simple anyons on this boundary corresponding to two simple right A -modules: one is just A itself, and the other is the free module $M_s := s \otimes A \simeq \bar{s} \otimes A$. The fusion rule is given by $M_s \otimes_A M_s \simeq A$ but the associator is non-trivial. We have $\text{RMod}_A(\mathfrak{Z}_1(\text{Vec}_{\mathbb{Z}_2}^\omega)) \simeq \text{Vec}_{\mathbb{Z}_2}^\omega$. The right module M_s is not a local module because the double braiding of s and $s\bar{s}$ is i . $\heartsuit$

Example 4.1.10. A 2+1D topological order C^3 is called *abelian* if for every simple anyon x there is an anyon x^* satisfying $x \otimes x^* \simeq \mathbb{1} \simeq x^* \otimes x$. Thus the category $\Omega\mathcal{C}$ of anyons is *pointed* in the sense that every simple object is invertible under the tensor product. In other words, the simple anyons form a finite abelian group under the fusion, denoted by G . Then for every $x, y \in G$ we define $q(x) \in U(1)$ to be the self-braiding of x and $b(x, y) \in U(1)$ to be the double braiding of x with y . Then we have

$$b(x, y) = \frac{q(xy)}{q(x)q(y)}$$

and $b: G \times G \rightarrow U(1)$ is a bi-character, i.e., $b(xy, z) = b(x, z)b(y, z)$ for all $x, y, z \in G$. We say that $q: G \rightarrow U(1)$ is a *quadratic form* and (G, q) is a *metric group*. The pointed braided fusion category $\Omega\mathcal{C}$ is completely determined by the metric group (G, q) consisting of simple anyons and self-braidings.

The condensable E_2 -algebras in $\Omega\mathcal{C}$ are classified by subgroups $H \leq G$ such that $q(x) = 1$ for all $x \in H$. Such a subgroup is called an *isotropic subgroup*. We denote the corresponding algebra by A_H . The multiplication of A_H is given by

$$A_H \otimes A_H = \bigoplus_{g, h \in H} g \otimes h \xrightarrow{\delta_{gh, k} \cdot \phi(g, h)} \bigoplus_{k \in H} k = A_H$$

for some 2-cochain $\phi \in C^2(G; U(1))$ such that $d\phi$ coincides with the associator of simple anyons in H . Indeed, the associator of simple anyons in H must be a coboundary by the isotropic condition. All simple right A_H -modules are free modules, i.e., of the form $x \otimes A_H$ for some $x \in G$. Two simple modules $x \otimes A_H$ and $y \otimes A_H$ are isomorphic if and only if x and y lie in the same coset of H in G , i.e., $x^{-1}y \in H$. The fusion rule of simple right A_H -modules is given by the multiplication of the quotient group G/H . Therefore, the fusion category $\text{RMod}_{A_H}(\Omega\mathcal{C})$ is a pointed fusion category with the group of simple objects being G/H . The associator can be found in [DS18, Theorem A.1.4]. A simple right A_H -module $x \otimes A_H$ is local if and only if x has trivial double braiding with all $y \in H$. We define $H^\perp := \{x \in G \mid b(x, y) = 1, \forall y \in H\}$. Then the local A_H -modules form a pointed modular tensor category with the group of simple objects being $H^\perp/H$, and the self-braiding is the restriction of q on $H^\perp/H$. Hence A_H is Lagrangian if and only if $H^\perp = H$. A subgroup $H \leq G$ is called *Lagrangian* if $H = H^\perp$. Lagrangian algebras in $\Omega\mathcal{C}$ one-to-one correspond to Lagrangian subgroups of G .

Both the 2+1D $\mathbb{Z}_2$ topological order and twisted $\mathbb{Z}_2$ topological order are abelian topological orders (see Example 4.1.8 and Example 4.1.9).

1. The $\mathbb{Z}_2$ topological order corresponds to $G = \mathbb{Z}_2 \times \mathbb{Z}_2$ and $q(a, b) = (-1)^{ab}$. There are two nontrivial isotropic subgroup generated by $(1, 0) = e$ and $(0, 1) = m$, and both of them are Lagrangian.
2. The twisted $\mathbb{Z}_2$ topological order corresponds to $G = \mathbb{Z}_2 \times \mathbb{Z}_2$ and $q(a, b) = i^{a^2 - b^2}$. The only nontrivial isotropic subgroup is generated by $(1, 1) = s\bar{s}$ and it is Lagrangian. $\heartsuit$

Example 4.1.11. Let G be a finite group. Anyons in the 2+1D G -gauge theory GT_G^3 (realizable by the Kitaev's quantum double model [Kit03]) form a non-degenerate braided fusion 1-category $\mathfrak{Z}_1(\text{Vec}_G) \simeq \mathfrak{Z}_1(\text{Rep}(G))$, which is the Drinfeld center of Vec_G or $\text{Rep}(G)$. An object in $\mathfrak{Z}_1(\text{Vec}_G)$ is a finite-dimensional G -graded vector space $V = \bigoplus_{g \in G} V_g$ equipped with a G -action $\rho: G \rightarrow \text{GL}(V)$ such that $\rho(g)(V_h) \subseteq V_{ghg^{-1}}$ for all $g, h \in G$. The tensor product in $\mathfrak{Z}_1(\text{Vec}_G)$ is the tensor product of G -graded vector spaces with the diagonal G -action, and the tensor unit is $1_e \in \text{Vec}_G$ with the trivial G -action. The braiding $c_{X, Y}: X \otimes Y \rightarrow Y \otimes X$ is defined by $c_{X, Y}(x \otimes y) = g(y) \otimes x$ for $x \in X_g, y \in Y$ [Dav10b]. The condensable E_2 -algebras in $\mathfrak{Z}_1(\text{Vec}_G)$ are completely classified in [Dav10b]. We list some examples here.

1. Let $\text{Fun}(G)$ be the space of $\mathbb{C}$ -valued functions on G . We equipped $\text{Fun}(G)$ with the left translation G -action:

$$(g \triangleright f)(h) := f(g^{-1}h), \quad g, h \in G, f \in \text{Fun}(G).$$

We can also equip $\text{Fun}(G)$ with the right translation G -action $(g \triangleright f)(h) := f(hg)$, but these two G -representations are isomorphic. Then we equip $\text{Fun}(G)$ with the G -grading concentrating on the trivial degree, i.e.,

$$\text{Fun}(G)_g = \begin{cases} \text{Fun}(G), & g = e, \\ 0, & g \neq e. \end{cases}$$

Then $\text{Fun}(G)$ can be viewed as an object in $\mathfrak{Z}_1(\text{Vec}_G)$. It is also a condensable E_2 -algebra in $\mathfrak{Z}_1(\text{Vec}_G)$ with the point-wise multiplication:

$$(f \cdot f')(g) := f(g)f'(g), \quad f, f' \in \text{Fun}(G), g \in G.$$

Let $\delta_g \in \text{Fun}(G)$ be the delta function on $g \in G$. Then $\{\delta_g\}_{g \in G}$ is a basis of $\text{Fun}(G)$ and the multiplication is given by $\delta_g \cdot \delta_h = \delta_{g,h} \delta_g$. The simple right $\text{Fun}(G)$ -modules are $\{M_g\}_{g \in G}$ where M_g is the same as $\text{Fun}(G)$ as G -representations but equipped with the different G -grading:

$$\delta_h \in (M_g)_{ghg^{-1}}, \quad g, h \in G.$$

In particular, M_e is the regular $\text{Fun}(G)$ -modules. The fusion rule is $M_g \otimes_{\text{Fun}(G)} M_h \simeq M_{gh}$ and the fusion category $\text{RMod}_{\text{Fun}(G)}(\mathfrak{Z}_1(\text{Vec}_G))$ is equivalent to Vec_G . The only local $\text{Fun}(G)$ -module is $\text{Fun}(G)$ itself. Thus $\text{Fun}(G) \in \mathfrak{Z}_1(\text{Vec}_G)$ is a Lagrangian algebra.

2. For any subgroup $H \leq G$, the space $\text{Fun}(G/H)$ of $\mathbb{C}$ -valued functions on the coset space G/H equipped with the left translation G -action, the trivial G -grading and the point-wise multiplication is a condensable E_2 -algebra. The modular tensor category of local $\text{Fun}(G/H)$ -modules is equivalent to $\mathfrak{Z}_1(\text{Vec}_H)$, and the fusion category of right $\text{Fun}(G/H)$ -modules is equivalent to the E_1 -centralizer of Vec_H in Vec_G . Two algebras $\text{Fun}(G/H)$ and $\text{Fun}(G/K)$ are isomorphic if and only if two subgroups H and K are conjugate to each other in G .
3. Let $\mathbb{C}[G]$ be group algebra of G . It has a basis $\{\tau_g\}_{g \in G}$ with the multiplication defined by $\tau_g \cdot \tau_h = \tau_{gh}$ for all $g, h \in G$. We equip $\mathbb{C}[G]$ with the canonical G -grading that $\tau_g \in \mathbb{C}[G]_g$ and the conjugation G -action that $g \triangleright \tau_h := \tau_{ghg^{-1}}$. Then $\mathbb{C}[G]$ is a condensable E_2 -algebra in $\mathfrak{Z}_1(\text{Vec}_G)$. Note that the multiplication of $\mathbb{C}[G]$ is not commutative in Vec when G is non-abelian, but is commutative in $\mathfrak{Z}_1(\text{Vec}_G)$ because the self-braiding of $\mathbb{C}[G]$ in $\mathfrak{Z}_1(\text{Vec}_G)$ is nontrivial due to the conjugation G -action. Every simple right $\mathbb{C}[G]$ -module is isomorphic to $V \otimes \mathbb{C}[G]$ for some irreducible G -representation V , viewed as an object in $\mathfrak{Z}_1(\text{Vec}_G)$ with the G -grading concentrating on the trivial degree. The fusion category $\text{RMod}_{\mathbb{C}[G]}(\mathfrak{Z}_1(\text{Vec}_G))$ is equivalent to $\text{Rep}(G)$. The only local $\mathbb{C}[G]$ -module is $\mathbb{C}[G]$ itself. Thus $\mathbb{C}[G] \in \mathfrak{Z}_1(\text{Vec}_G)$ is a Lagrangian algebra.
4. For any $\alpha \in Z^2(G; U(1))$, the twisted group algebra $\mathbb{C}[G, \alpha]$ is generated by $\{\tau_g\}_{g \in G}$ subject to the relation that $\tau_g \cdot \tau_h = \alpha(g, h) \tau_{gh}$ for all $g, h \in G$. We equip $\mathbb{C}[G, \alpha]$ with the canonical G -grading and the conjugation G -action:

$$g \triangleright \tau_h = \frac{\alpha(g, h)}{\alpha(ghg^{-1}, g)} \tau_{ghg^{-1}}.$$

Then $\mathbb{C}[G, \alpha] \in \mathfrak{Z}_1(\text{Vec}_G)$ is a Lagrangian algebra and we still have $\text{RMod}_{\mathbb{C}[G, \alpha]}(\mathfrak{Z}_1(\text{Vec}_G)) \simeq \text{Rep}(G)$.

5. The Lagrangian algebras in $\mathfrak{Z}_1(\text{Vec}_G)$ are classified by pairs (H, α) , where $H \leq G$ is a subgroup and $\alpha \in Z^2(H; U(1))$ is a 2-cocycle. We denote the corresponding algebra by $A_{H, \alpha}$. It is the induced representation of the twisted group algebra:

$$A_{H, \alpha} = \text{Ind}_H^G \mathbb{C}[H, \alpha].$$

Two Lagrangian algebras $A_{H, \alpha}$ and $A_{K, \beta}$ are isomorphic if and only if there exists an element $g \in G$ such that $gHg^{-1} = K$ and $[\alpha] = [g^* \beta] \in H^2(H; U(1))$, where $g^*: Z^2(K; U(1)) \rightarrow Z^2(H; U(1))$ is the pullback morphism induced by the conjugation of g .

For example, when $G = S_3$, there are 4 subgroups up to conjugation: $\{e\}$, $\mathbb{Z}_2$, $\mathbb{Z}_3$ and S_3 . For each subgroup K the cohomology group $H^2(K, U(1))$ is trivial. Therefore, there are 4 Lagrangian algebras in $\mathfrak{Z}_1(\text{Rep}(S_3)) \simeq \mathfrak{Z}_1(\text{Vec}_{S_3})$ [Dav10b, Section 4.2]. See also [CCW16, Section 3.8] for the condensation of these Lagrangian algebras.

6. When G is abelian, we can give a more explicit construction of the Lagrangian algebras $A_{H,\alpha}$. In this case $\mathfrak{Z}_1(\text{Vec}_G)$ is pointed, and the corresponding metric group is $(G \times \hat{G}, \chi)$, where $\hat{G} := \text{Hom}(G; U(1))$ is the dual group of G and $\chi(g, \phi) := \phi(g)$. Given a 2-cocycle $\alpha \in Z^2(G; U(1))$, its anti-symmetrization α_a is defined by

$$\alpha_a(g, h) := \frac{\alpha(g, h)}{\alpha(h, g)}, \quad g, h \in G.$$

The cocycle condition implies that $\alpha_a: G \times G \rightarrow U(1)$ is a bi-character. We denote the space of anti-symmetric bi-characters on G by $\text{Alt}^2(G; U(1))$. Then $\alpha \mapsto \alpha_a$ defines a group homomorphism $Z^2(G; U(1)) \rightarrow \text{Alt}^2(G; U(1))$. Moreover, if α is a coboundary, then α_a is trivial. Hence we get a group homomorphism $H^2(G; U(1)) \rightarrow \text{Alt}^2(G; U(1))$. Indeed, this is an isomorphism because every symmetric 2-cocycle must be a coboundary. The Lagrangian subgroups of $(G \times \hat{G}, q)$ are parametrized by $(H, [\alpha])$, where $H \leq G$ is a subgroup and $[\alpha] \in H^2(H; U(1))$. The corresponding Lagrangian subgroup of $(G \times \hat{G}, \chi)$ is

$$L_{H, [\alpha]} := \{(h, \phi) \in G \times \hat{G} \mid h \in H, \phi|_H = \alpha_a(h, -)\}.$$

This gives all Lagrangian algebras in $\mathfrak{Z}_1(\text{Vec}_G)$. ♡

Example 4.1.12 ([DS17]). We consider 2+1D twist finite gauge theory $\text{GT}_{(G, \omega)}^3$. Its category of anyons is given by $\Omega\mathcal{GT}_{(G, \omega)} = \mathfrak{Z}_1(\text{Vec}_G^\omega)$. The classification of simple condensable E_2 -algebras and Lagrangian algebras in $\mathfrak{Z}_1(\text{Vec}_G^\omega)$ are studied in [DS17]. More precisely, the Lagrangian algebras in $\mathfrak{Z}_1(\text{Vec}_G^\omega)$ are classified by the pairs $(H, [\alpha])$ up to conjugation, where $H \leq G$ is a subgroup such that $[\omega]|_H \in H^3(H, U(1))$ is trivial and $[\alpha] \in H^2(H, U(1))$. ♡

Example 4.1.13 ([BS09, CJKYZ20]). A very classical example of anyon condensation is the condensation from double Ising to toric code. It was first studied by Bais and Slingerland in [BS09]. The complete details of this example had been worked out in [CJKYZ20]. We briefly summarize the result below. The object

$$A = 1 \boxtimes 1 \oplus \psi \boxtimes \psi \in \Omega\mathcal{I}s \boxtimes \Omega\mathcal{I}s^{\text{op}} \simeq \mathfrak{Z}_1(\Omega\mathcal{I}s)$$

has a canonical structure of a condensable E_2 -algebra in $\mathfrak{Z}_1(\Omega\mathcal{I}s)$. It can be explicitly defined (see for example [CJKYZ20, Example 7]). There are six simple right A -modules in $\Omega\mathcal{I}s$:

$$\begin{aligned} 1 &:= A, & e &:= (\sigma \boxtimes \sigma), & m &:= (\sigma \boxtimes \sigma)^{\text{tw}}, & f &:= (\psi \boxtimes 1) \otimes A, \\ \chi_+ &:= (1 \boxtimes \sigma) \otimes A, & \chi_- &:= (\sigma \boxtimes 1) \otimes A, \end{aligned} \tag{4.1.10}$$

among which $1, e, m, f$ are E_2 - A -modules (or local A -modules) and $\chi_{\pm}$ are non-local A -modules. Moreover, the fusion and braiding structures on $1, e, m, f$ are the same as those on the 4 simple anyons in the 2+1D toric model, i.e. $\text{Mod}_A^{E_2}(\mathfrak{Z}_1(\Omega\mathcal{I}s)) \simeq \mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$. ♡

Example 4.1.14. Let V be a rational vertex operator algebra (VOA). Then the category Mod_V of V -modules is a modular tensor category [Hua08]. The simple condensable E_1 -algebras in Mod_V are open-string VOAs over V [HK04], and the simple condensable E_2 -algebras in Mod_V are VOA extensions of V [HKL15]. If $V \hookrightarrow W$ is an extension of VOAs, then $\text{Mod}_W^{E_2}(\text{Mod}_V) \simeq \text{Mod}_W$ as modular tensor categories. More generally, if V_L, V_R are rational VOAs, the simple condensable E_2 -algebras in $\text{Mod}_{V_L \otimes V_R} \simeq \text{Mod}_{V_L} \boxtimes \text{Mod}_{V_R}$ are full field algebras over $V_L \otimes V_R$ [Kon07], and the Lagrangian algebras are precisely modular invariant CFTs over $V_L \otimes V_R$ [KR09]. When $V_L = V_R = V$, there is a well-known Lagrangian algebra

$$\bigoplus_{m \in \text{Irr}(\text{Mod}_V)} m \boxtimes m^* \in \text{Mod}_V \boxtimes \overline{\text{Mod}_V} \simeq \text{Mod}_{V \otimes \bar{V}},$$

where $\text{Irr}(\text{Mod}_V)$ denotes the set of simple objects in Mod_V . This algebra is also a modular invariant closed CFT called the “charge conjugation construction” or the “Cardy case”. Under the canonical equivalence $\text{Mod}_V \boxtimes \overline{\text{Mod}_V} \simeq \mathfrak{Z}_1(\text{Mod}_V)$, this Lagrangian algebra is also the full center [FFRS06, KR09, Dav10a] of the tensor unit $1 = V \in \text{Mod}_V$. In this case, Lagrangian algebras in $\mathfrak{Z}_1(\text{Mod}_V)$ one-to-one correspond to indecomposable Mod_V -modules in 2Vec [KR08]. ♡

Example 4.1.15. Let k be a non-negative integer and $\mathcal{C}_k := \mathcal{C}(\widehat{\mathfrak{sl}}_2, k)$ be the category of integrable modules over $\widehat{\mathfrak{sl}}_2$ of level k . It is also equivalent to the semisimple part of the category of finite-dimensional representations over the quantum group $U_q(\mathfrak{sl}_2)$ with $q = \exp(\frac{i\pi}{k+2})$. Then $\mathcal{C}_k$ is a modular tensor category and it has $k+1$ simple objects, denoted by $V_0, \dots, V_k$.

The indecomposable finite semisimple modules over $\mathcal{C}_k$ are one-to-one corresponding to simply laced Dynkin diagrams of type A, D, E with Coxeter number $k+2$ [Ost03a], and the simple objects of the module correspond to the vertices of the Dynkin diagram. Moreover, each module of type A_n, D_{2n}, E_6, E_8 is equivalent to the category of right modules over a unique simple condensable E_2 -algebra in $\mathcal{C}_k$, and this is also a complete classification of simple condensable E_2 -algebras in $\mathcal{C}_k$ [KO02].

- Type A_n for $n = k+1$. The corresponding module is $\mathcal{C}_k$ itself as a regular $\mathcal{C}_k$ -module. It is equivalent to the category of right modules over the tensor unit $\mathbb{1} = V_0$.
- Type D_n for $2n = k+4$. In this case k must be even. Then $V_0 \oplus V_k \in \mathcal{C}_k$ is a simple condensable E_1 -algebra. The corresponding module is $\text{RMod}_{V_0 \oplus V_k}(\mathcal{C}_k)$. The algebra $V_0 \oplus V_k$ is commutative if and only if k is divided by 4 (i.e., n is even).
- Type E_6 . In this case $k = 10$. There is a VOA extension $(\widehat{\mathfrak{sl}}_2)_{10} \hookrightarrow (\widehat{\mathfrak{sp}}_4)_1$, which gives a simple condensable E_1 -algebra in $\mathcal{C}_{10}$ with the underlying object $V_0 \oplus V_6$. The corresponding module over $\mathcal{C}_{10}$ is $\text{RMod}_{V_0 \oplus V_6}(\mathcal{C}_{10})$.
- Type E_7 . In this case $k = 16$. There is no VOA extension of $(\widehat{\mathfrak{sl}}_2)_{16}$, but an VOA extension $(\widehat{\mathfrak{sl}}_2)_{16} \otimes (\widehat{\mathfrak{sl}}_3)_6 \hookrightarrow (E_8)_1$. The corresponding module over $\mathcal{C}_{16}$ can be constructed from this VOA extension. See [Ost03a, Proof of Theorem 6] for more details.
- Type E_8 . In this case $k = 28$. There is a VOA extension $(\widehat{\mathfrak{sl}}_2)_{28} \hookrightarrow (G_2)_1$, which gives a simple condensable E_1 -algebra in $\mathcal{C}_{28}$ with the underlying object $V_0 \oplus V_{10} \oplus V_{18} \oplus V_{28}$. The corresponding module over $\mathcal{C}_{28}$ is $\text{RMod}_{V_0 \oplus V_{10} \oplus V_{18} \oplus V_{28}}(\mathcal{C}_{28})$.

Equivalently, this also gives the complete classification of Lagrangian algebras in $\mathfrak{Z}_1(\mathcal{C}_k)$. As explained in Example 4.1.14, these Lagrangian algebras are modular invariant CFTs over $(\widehat{\mathfrak{sl}}_2)_k \otimes (\widehat{\mathfrak{sl}}_2)_k$. The partition functions of these CFTs, or the underlying objects of these Lagrangian algebras, can be found in [DFMS97, Chapter 17] for example. $\heartsuit$

Remark 4.1.16. Some examples of anyon condensations from para-fermions to finite gauge theories can be found in [LY23]. More examples in lower rank braided fusion 1-categories can be found in [Kik23b, Kik23a, KKH24, Kik24]. $\diamond$

4.1.3 Geometric approach

Now we take a geometric approach towards the anyon condensation theory in (2+1)D. The physical or geometric intuitions behind it naturally generalize to higher dimensions.


The setup is the same one given in Figure 12. Recall that the anyon condensation is determined by a condensable E_2 -algebra A in $\Omega\mathcal{C}$, and we have $\Omega\mathcal{D} \simeq \text{Mod}_A^{\mathcal{E}_2}(\Omega\mathcal{C})$, $\Omega_m\mathcal{M} \simeq \text{RMod}_A(\Omega\mathcal{C})$. We show that the algebra A can be constructed geometrically. We start from recalling a known result.

Lemma 4.1.17 ([Zhe17]). Let $\Omega\mathcal{C}, \Omega\mathcal{D}$ be non-degenerate braided fusion 1-categories and $\Omega_m\mathcal{M}$ a closed multi-fusion $\Omega\mathcal{C}$ - $\Omega\mathcal{D}$ -bimodule. We have canonical monoidal equivalences

$$\Omega_m\mathcal{M} \boxtimes_{\Omega\mathcal{D}} \Omega_m\mathcal{M}^{\text{op}} \xrightarrow{\simeq} \text{Fun}_{\Omega\mathcal{C}}(\Omega_m\mathcal{M}, \Omega_m\mathcal{M}), \quad x \boxtimes_{\Omega\mathcal{D}} y \mapsto x \otimes_{\Omega_m\mathcal{M}} y - \otimes_{\Omega_m\mathcal{M}} y \quad (4.1.11)$$

$$\Omega_m\mathcal{M} \boxtimes_{\Omega\mathcal{C}} \Omega_m\mathcal{M}^{\text{op}} \xrightarrow{\simeq} \text{Fun}_{\Omega\mathcal{D}}(\Omega_m\mathcal{M}, \Omega_m\mathcal{M}), \quad x \boxtimes_{\Omega\mathcal{C}} y \mapsto x \otimes_{\Omega_m\mathcal{M}} y - \otimes_{\Omega_m\mathcal{M}} y. \quad (4.1.12)$$

This Lemma says that if we squeeze $\Omega_m\mathcal{M}$ and $\Omega_m\mathcal{M}^{\text{op}}$ along $\Omega\mathcal{C}$ in the following physical configuration,



$$\begin{array}{c} \text{Diagrammatic equation (4.1.13):} \\ \text{Left side: } \Omega\mathcal{D} \text{ and } \Omega\mathcal{C} \text{ boxes with } \Omega_m\mathcal{M} \text{ and } \Omega_m\mathcal{M}^{\text{op}} \text{ lines.} \\ \xrightarrow{\text{Squeeze}} \text{Middle side: } \Omega\mathcal{D} \text{ and } \Omega\mathcal{C} \text{ boxes with } \Omega_m\mathcal{M} \text{ and } \Omega_m\mathcal{M}^{\text{op}} \text{ lines, including } \text{Fun}_{\Omega\mathcal{D}}(\Omega_m\mathcal{M}, \Omega_m\mathcal{M}). \\ = \text{Right side: } \Omega\mathcal{D} \text{ box with } \Omega_m\mathcal{M} \text{ and } \Omega_m\mathcal{M}^{\text{op}} \text{ lines, including } (\Omega_m\mathcal{M})^{\text{op}}. \end{array} \quad (4.1.13)$$

we obtain a domain wall whose particles form the fusion 1-category $\text{Fun}_{\Omega\mathcal{D}}(\Omega_m\mathcal{M}, \Omega_m\mathcal{M})$. The “=” is due to the equivalence $\text{Fun}_{\Omega\mathcal{D}}(\Omega_m\mathcal{M}, \Omega_m\mathcal{M}) \simeq (\Omega_m\mathcal{M})^{\text{op}} \boxtimes_{\Omega\mathcal{D}} \mathcal{M}$, which maps $\text{id}_{\Omega_m\mathcal{M}}$ to $\int_{x \in \mathcal{M}}^{\Omega\mathcal{D}} x^R \boxtimes_{\Omega\mathcal{D}} x$. The notation $\int_{x \in \mathcal{M}}^{\Omega\mathcal{D}}$ is called an ‘ $\Omega\mathcal{D}$ -module end’ and is explained in Example A.2.12 in Appendix A.2. The domain wall $\text{Fun}_{\Omega\mathcal{D}}(\Omega_m\mathcal{M}, \Omega_m\mathcal{M})$ without specifying a particle on it can be understood as the particle is trivial. Then we obtain the following physically equivalent configurations.

$$(4.1.14)$$

Now we show that the condensable E_2 -algebra A can be constructed geometrically as a D^3 -bubble in C^3 as illustrated below.

$$(4.1.15)$$

Viewed from far away, the bubble shrink to a particle in $\Omega\mathcal{C}$, it was shown in [AKZ17] that this particle is precisely A as an object in $\Omega\mathcal{C}$. We recall this result and give a new proof.

Lemma 4.1.18 ([AKZ17]). For $d \in \Omega\mathcal{D}$ and $y \in \Omega_m\mathcal{M}$, the following bubble defines a particle in $\Omega\mathcal{C}$ given by $[1_m, d \odot y]_{\Omega\mathcal{C}} \in \Omega\mathcal{C}$.

$$[1_m, d \odot y]_{\Omega\mathcal{C}} =$$

Proof. First squeeze the bubble vertically to a line then horizontally to a point. We obtain

$$[1_m, d \odot y]_{\Omega\mathcal{C}} =$$

The second “=” is due to the monoidal equivalence $(\Omega_m\mathcal{M})^{\text{op}} \boxtimes_{\mathcal{E}} \Omega_m\mathcal{M} \simeq \text{Fun}_{\mathcal{E}}(\Omega_m\mathcal{M}, \Omega_m\mathcal{M}) \simeq \Omega\mathcal{C}$, where $\mathcal{E} := \text{Fun}_{\Omega\mathcal{C}}(\Omega_m\mathcal{M}, \Omega_m\mathcal{M})$. This monoidal equivalence maps $1_m \boxtimes_{\mathcal{E}} y$ to

$$[- , 1_m]_{\mathcal{E}}^R \odot y \mapsto \int_{x \in \Omega_m\mathcal{M}}^{\mathcal{E}} [x, [x, 1_m]_{\mathcal{E}}^R \odot y]_{\Omega\mathcal{D}} \simeq \int_{\mathcal{E}}^{x \in \Omega_m\mathcal{M}} [x, 1_m] \odot x, y]_{\Omega\mathcal{C}} \simeq [1_m, y]_{\Omega\mathcal{C}}. \quad \square$$

The identity (4.1.15) is only an identity of objects. The algebraic structure on the bubble has not been constructed. We do that now. The multiplication morphisms $\mu_A : A \otimes A \rightarrow A$ can be defined geometrically by fusing two such bubbles into one bubble as follows.

$$(4.1.16)$$

where the arrow is defined by the canonical projection $\int_{x \in \Omega_m\mathcal{M}}^{\Omega\mathcal{C}} x^R \boxtimes_{\Omega\mathcal{D}} x \rightarrow 1_m \boxtimes_{\Omega\mathcal{D}} 1_m$. In the rest of this subsection, we show that this algebraic structure on A coincides with that of $[1_m, 1_m]_{\Omega\mathcal{C}}$.


Theorem 4.1.19. The algebraic structure on A defined in (4.1.16) coincides with that on $[1_m, 1_m]_{\Omega\mathcal{C}}$.

Proof. We only sketch the idea of the proof and leave the remaining details to Appendix A.1. The algebraic structure on the internal hom $[1_m, 1_m]_{\Omega\mathcal{C}}$ follows from the universal property of the internal hom. This universal construction, however, is not physical or geometrical. In order to find a proof, it is enough to find a geometric construction of the algebraic structure on $[1_m, 1_m]_{\Omega\mathcal{C}}$ and compare it with (4.1.16).

The idea is that we rewrite the internal hom as follows:

$$[1_m, 1_m]_{\Omega\mathcal{C}} = [1_m, -]_{\Omega\mathcal{C}}(1_m) = L_2^R(1_m),$$

where $L_2^R = [1_m, -]_{\Omega\mathcal{C}}$ is precisely the right adjoint of the functor $L_2 := - \odot 1_m : \Omega\mathcal{C} \rightarrow \Omega_m\mathcal{M}$ by the definition of internal hom (see Definition 3.1.5). Note that L has a clear physical meaning, i.e., moving bulk anyons to the wall, and is called bulk-to-wall map. Its right adjoint functor L^R also has a clear physical meaning as a physical process of creating a bubble from the wall M^2 to the bulk of $\mathcal{C}^3$ as illustrated below.




The algebraic structure on $[1_m, 1_m]_{\Omega\mathcal{C}} = L^R(1_m)$ is defined in terms of the defining data of the adjoint pair L_2 and L_2^R as follows.

$$L_2^R(1_m) \otimes L_2^R(1_m) \rightarrow L_2^R L_2(L_2^R(1_m) \otimes L_2^R(1_m)) \simeq L_2^R(L_2 L_2^R(1_m) \otimes L_2 L_2^R(1_m)) \rightarrow L_2^R(1_m). \quad (4.1.17)$$

It remains to show that the morphism (4.1.17) coincides with the one defined in (4.1.16). We postpone it to Appendix A.1. $\square$

Remark 4.1.20. Similar to the 1-codimensional situation (3.2.2), by considering the spacetime picture, there is a natural way to define an algebraic structure on A geometrically as follows:



Similar to Remark 3.2.2, it is intuitively clear that the geometric definition in the second picture in (4.1.18) coincides with the geometric definition (4.1.16). The unit morphism $\mathbb{1}_{\Omega\mathcal{C}} \rightarrow A = [1_m, 1_m]_{\Omega\mathcal{C}}$ defined by the universal property of the internal hom should also coincide with the first picture in (4.1.18). When the algebra A is indecomposable, this coincidence should follow automatically from that of μ_A and the unit property because $\text{hom}_{\Omega\mathcal{C}}(\mathbb{1}_{\Omega\mathcal{C}}, A) \simeq \mathbb{C}$.

In principle, a rigorous proof of these two coincidences can be achieved by computing this instanton carefully within Reshetikhin-Turaev TQFT enriched by gapped domain walls, or equivalently, by the so-called β -version of factorization homology (see Remark 6.2.2). $\diamond$


Remark 4.1.21. It is worth mentioning that the algebra we obtained by creating D^3 -bubbles (taking internal homs) is a condensable algebra in $\Omega\mathcal{C}$. This fact does not rely on the fact that D^3 is obtained from $\mathcal{C}^3$ via a particle condensation, so it applies to any D^3 that are Morita equivalent to $\mathcal{C}^3$. However, condensing this algebra does not reproduce the phase D^3 in general. For example, in the same setting as Figure 12, we create a $\mathcal{C}^3$ -bubble in D^3 , which is the trivial particle in $\Omega\mathcal{D}$, i.e.,

$$\begin{array}{c} \text{M}^2 \\ \downarrow \\ \boxed{\Omega\mathcal{C}} \\ \downarrow \\ \Omega\mathcal{D} \end{array} = [A, A]_{\Omega\mathcal{D}} = A = \mathbb{1}_{\Omega\mathcal{D}}. \quad (4.1.19)$$

Clearly, condensing this algebra (vacuum in $\Omega\mathcal{D}$) does not produce C^3 as the condensed phase. $\diamond$

Remark 4.1.22. The picture in (4.1.19) can be obtained by rolling up the domain wall M^2 to the C^3 -side. It creates a bubble in D^3 . Similarly, as illustrated in (4.1.15), rolling up the domain wall M^2 to the D^3 -side creates a bubble in C^3 , which turns out to be a condensable E_2 -algebra A in $\Omega\mathcal{C}$. This physical process naturally generalizes to higher dimension cases, in which one can roll up potentially higher codimensional defects over higher dimensional spheres or other manifolds. We use this intuition without a proof in later sections. To write down a proof explicitly is a non-trivial task. However, we expect that the proof should be a higher dimensional analogue of the one in this subsection. $\diamond$

Similar to the physical process of condensing 1-codimensional topological defects described in Section 3.2 and 3.3.1, we can provide a description of the physical process of a 2-codimensional defect condensation, which naturally generalizes to higher dimensional topological orders.



- (1) First proliferating the D^3 -phase in a disk-like region inside the C^3 -phase;
- (2) then shrinking the C^3 -phase within the region and merging two D^3 -disks into a larger D^3 -disk but with extra particles on the gapped boundary according to (4.1.13) and (4.1.14);
- (3) then annihilating the extra particles on the boundary according to (4.1.16).

Remark 4.1.23. Notice that the process of merging two D^3 -disks into a larger D^3 -disk can split into merging vertically then merging horizontally. This leads us to a general theory of the condensation of k -codimensional defects (see Section 5). $\diamond$

4.2 Condensations of 2-codimensional defects in $n+1D$

In this subsection, we discuss the theory of the condensations of 2-codimensional topological defects in $n+1D$ anomaly-free simple topological orders. The setup of the question remains the same as the 2+1D cases. More precisely, consider a condensation of 2-codimensional topological defect occurring in an $n+1D$ anomaly-free simple topological order C^{n+1} . It creates a new simple topological order D^{n+1} , which is also anomaly-free, and a gapped domain wall M^n as depicted in Figure 14.




Figure 14: a condensation of 2-codimensional topological defect in C^{n+1}

4.2.1 General theory

The bootstrap analysis in $n+1D$ is possible and is similar to the 2+1D cases. However, it becomes much harder in higher dimensions due to the complexity of the higher coherence relations. Instead, we use the geometric intuition of anyon condensation in an anomaly-free 2+1D simple topological order, which generalizes to that of condensations of 2-codimensional topological defects in an $n+1D$ anomaly-free simple topological order tautologically.


Definition 4.2.1. For a braided multi-fusion n -category $\mathcal{A}$ and $A \in \text{Alg}_{E_2}^c(\mathcal{A})$, roughly speaking, an E_2 - A -module in $\mathcal{A}$ is an object $x \in \mathcal{A}$ equipped with a 2-dimensional actions, i.e., four A -actions $\mu_x^\rightarrow, \mu_x^\downarrow, \mu_x^\leftarrow, \mu_x^\uparrow$ in four directions such that the diagram (4.1.6) is commutative. Recall (4.1.7), one can take

$$\text{Mod}_A^{E_2}(\mathcal{A}) \simeq \text{Mod}_A^{E_1}(\text{Mod}_A^{E_1}(\mathcal{A}))$$

as a working definition of an E_2 - A -module and the category $\text{Mod}_A^{E_2}(\mathcal{A})$ of E_2 - A -modules. $\square$

Now we are ready to state the result about the condensed phase and the gapped domain wall for a condensation of a 2-codimensional topological defect in C^{n+1} .

Theorem^{ph} 4.2.2. If an $n+1$ D anomaly-free simple topological order D^{n+1} and a gapped domain wall M^n are obtained from an anomaly-free simple topological order C^{n+1} via a condensation of 2-codimensional topological defect (see Figure 14 and the picture below).



1. The trivial 2-codimensional topological defect $1_{\mathbb{1}_{\mathcal{D}}}$ in $\Omega\mathcal{D}$ can be identified with a simple condensable E_2 -algebra A in $\Omega\mathcal{C}$. Moreover, $\Omega\mathcal{D}$ consists of all deconfined topological defects of codimension 2 and higher, and it can be identified with the category $\text{Mod}_A^{E_2}(\Omega\mathcal{C})$ of E_2 - A -modules (or equivalently, local A -modules) in $\mathcal{C}$, i.e., $\Omega\mathcal{D} = \text{Mod}_A^{E_2}(\Omega\mathcal{C})$.
2. Topological defects on the wall M^n include all confined and deconfined 1-codimensional defects, and can be identified with the fusion 1-category $\text{RMod}_A(\Omega\mathcal{C})$.
3. Defects in $\Omega\mathcal{C}$ move to the wall according to the central functor⁵¹ $L_2 := - \otimes A: \Omega\mathcal{C} \rightarrow \text{RMod}_A(\Omega\mathcal{C})$ defined by $x \mapsto x \otimes A$ for all $x \in \Omega\mathcal{C}$.
4. Defects in $\Omega\mathcal{D}$ move to the wall according to the canonical forgetful functor⁵² $R_2: \text{Mod}_A^{E_2}(\Omega\mathcal{C}) \rightarrow \text{RMod}_A(\Omega\mathcal{C})$.

The condensable E_2 -algebra A in $\Omega\mathcal{C}$ is called Lagrangian if $\text{Mod}_A^{E_2}(\Omega\mathcal{C}) \simeq \text{Vect}$. In this case, M^n becomes a gapped boundary of C^{n+1} and consists of all confined and deconfined defects. $\square$

Theorem^{ph} 4.2.3. By the folding trick and the boundary-bulk relation, we obtain a braided equivalence:

$$\Omega\mathcal{C} \boxtimes \text{Mod}_A^{E_2}(\Omega\mathcal{C})^{\text{op}} \simeq \mathfrak{Z}_1(\text{RMod}_A(\Omega\mathcal{C})).$$

This result is equivalent to the following braided equivalence:

$$\text{Mod}_A^{E_2}(\Omega\mathcal{C})^{\text{op}} \simeq \mathfrak{Z}_2(\Omega\mathcal{C}, \mathfrak{Z}_1(\text{RMod}_A(\Omega\mathcal{C}))), \quad (4.2.1)$$


which is ready to be generalized to higher algebras and higher representations (see Theorem 5.2.11). $\square$

Remark 4.2.4. If a topological defect is confined (resp. deconfined), then its condensation descendants is also confined (resp. deconfined). This natural physical intuition follows immediately from (4.2.1) and results in [KZ22b, KZ24] (recall Remark 2.3.27). The $n = 2$ case was proved in [DX24]. $\diamond$

⁵¹Being a central functor means that $- \otimes A$ factors through a braided monoidal functor $g: \Omega\mathcal{C} \rightarrow \mathfrak{Z}_1(\text{RMod}_A(\Omega\mathcal{C}))$, i.e., it coincides with the composed functor $\Omega\mathcal{C} \xrightarrow{g} \mathfrak{Z}_1(\text{RMod}_A(\Omega\mathcal{C})) \xrightarrow{f} \text{RMod}_A(\Omega\mathcal{C})$, where f is the forgetful functor.


⁵²The “ E_2 -module condition” for a right A -module is a property for $n = 2$ but a structure plus properties for $n > 2$. This functor R_2 is fully faithful for $n = 2$ but not fully faithful for $n > 2$.

Let C^{n+1} and D^{n+1} be any two $n+1$ D (potentially anomalous and composite) topological orders connected by a gapped domain wall $M^n = (\mathcal{M}, m)$, where $\mathcal{M}$ is a separable n -category and a $\mathcal{C}\text{-}\mathcal{D}$ -bimodule and m is an object in $\mathcal{M}$ represents a distinguished gapped wall condition.



There is a canonical functor $L_2 : \Omega\mathcal{C} \rightarrow \Omega_m \mathcal{M}$ from the category $\Omega\mathcal{C}$ of topological defects of codimension 2 and higher in C^{n+1} to that of topological defects on M^n , which is defined simply by moving topological defects in C^{n+1} onto M^n . Similarly, there is also a canonical functor $R_2 : \Omega\mathcal{D} \rightarrow \Omega_m \mathcal{M}$. It is clear that these two functors are both monoidal (i.e. preserving the fusion product). We denote their right adjoint functor by L_2^R and R_2^R , respectively.

By rolling up the right half (i.e. M^n and D^{n+1}) in the spatial dimension, we obtain a solid cylinder $D^2 \times \mathbb{R}^{n-1}$ with the interior filled with the D^{n+1} -phase and the boundary $S^1 \times \mathbb{R}^{n-1}$ cylinder decorated by M^n .⁵³ Similarly, we can roll up the left half to obtain a solid cylinder in D^{n+1} -phase as illustrated below.



Viewed from far away, these two solid cylinders become two 2-codimensional topological defects in C^{n+1} and D^{n+1} , respectively. The next result tells us how to compute these defects.

Theorem^{ph} 4.2.5. Such obtained 2-codimensional topological defects in C^{n+1} and D^{n+1} are given by

$$L_2^R(1_m) = [1_m, 1_m]_{\Omega\mathcal{C}} \in \Omega\mathcal{C}, \quad R_2^R(1_m) = [1_m, 1_m]_{\Omega\mathcal{D}} \in \Omega\mathcal{D}, \quad (4.2.3)$$

both of which are condensable E_2 -algebra in $\Omega\mathcal{C}$ and $\Omega\mathcal{D}$, respectively. Moreover, all condensable E_2 -algebra in $\Omega\mathcal{C}$ can be obtained in this way by properly choosing D^{n+1} and M^n . The Morita equivalence among gapped domain walls naturally defines the so-called 2-Morita equivalence relation among condensable E_2 -algebras (see Definition 6.1.6). $\square$

Proof. We sketch a proof. Note that $\Omega\mathcal{C}$ acts on $\Omega_m \mathcal{M}$ in the obvious way. We denote the action functor by $\odot : \Omega\mathcal{C} \times \Omega_m \mathcal{M} \rightarrow \Omega_m \mathcal{M}$. Therefore, the internal hom is well defined [KZ22b, Proposition 3.27]. Moreover, the action $\odot$ preserves the fusion product, i.e. $\odot$ is monoidal. By the universal property of the center, this action factors through the canonical action $\mathfrak{Z}_1(\Omega_m \mathcal{M}) \times \Omega_m \mathcal{M} \rightarrow \Omega_m \mathcal{M}$. Equivalently, 2-codimensional topological defects in C^{n+1} can be half-braided with 1-codimensional topological defects in M^n . This half-braiding structure upgrades the action $\odot$ to an action $\odot : \text{Alg}_{E_1}^c(\Omega\mathcal{C}) \times \text{Alg}_{E_1}^c(\Omega_m \mathcal{M}) \rightarrow \text{Alg}_{E_1}^c(\Omega_m \mathcal{M})$. As a consequence, the internal hom $[1_m, 1_m]_{\Omega\mathcal{C}} \in \text{Alg}_{E_1}^c(\text{Alg}_{E_1}^c(\Omega\mathcal{C})) = \text{Alg}_{E_2}^c(\Omega\mathcal{C})$.

For $A \in \text{Alg}_{E_2}^c(\Omega\mathcal{C})$, let D^{n+1} and M^n be the condensed phase and gapped domain wall obtained by condensing A (as in Theorem^{ph} 4.2.2), i.e., $\Omega_m \mathcal{M} = \text{RMod}_A(\Omega\mathcal{C})$ and $\Omega\mathcal{D} = \text{Mod}_A^{E_2}(\Omega\mathcal{C})$. In this case, $L_2 = - \otimes A$ and $L_2^R = [A, -]$, where $[A, -]$ is the forgetful functor $f : \text{RMod}_A(\Omega\mathcal{C}) \rightarrow \Omega\mathcal{C}$. Therefore, in this case, we obtain

$$L_2^R(1_m) = [1_m, 1_m]_{\Omega\mathcal{C}} = A \in \Omega\mathcal{C}; \quad (4.2.4)$$

Therefore, all condensable E_2 -algebras in $\Omega\mathcal{C}$ arise in this way. $\square$

In general, condensing $L_2^R(1_m)$ in $\Omega\mathcal{C}$ does not give $\Omega\mathcal{D}$, and condensing $R_2^R(1_m)$ in $\Omega\mathcal{D}$ does not give $\Omega\mathcal{C}$.

1. Let C^3 be the $\mathbb{Z}_2$ topological order and D^3 be the double semion topological order and M^2 be any gapped domain wall. Then $L_2^R(1_m) = 1_{\mathbb{1}_C}$ and $R_2^R(1_m) = 1_{\mathbb{1}_D}$ are both trivial.

⁵³That is equivalent to compactify the domain wall M^n from the spacetime manifold $\mathbb{R}^1 \times \mathbb{R}^{n-1}$ to $S^1 \times \mathbb{R}^{n-1}$.

2. When C^{n+1} and D^{n+1} are both anomaly-free and simple, if D^{n+1} and M^n are obtained from C^{n+1} by condensing a non-trivial simple condensable E_2 -algebra $A \in \Omega\mathcal{C}$, we expect that this condensation is not reversible in general. For example, for $n \geq 3$ and a non-trivial finite group G , let $C^{n+1} = \text{GT}_G^{n+1}$, $M^n = (\mathcal{M}, m) = (n\text{Vec}_G, \mathbb{1})$ and $D^{n+1} = \mathbf{1}^{n+1}$. In this case, the internal hom $[1_m, 1_m]_{\Omega\mathcal{C}}$ is a Lagrangian algebra in $\Omega\mathcal{C} = \mathfrak{Z}_1(n\text{Vec}_G)$, condensing which produces $\mathbf{1}^{n+1}$ as the condensed phase. However, the internal hom $[1_m, 1_m]_{\Omega\mathcal{D}} = \mathbb{1}_{\Omega\mathcal{D}}$ is the trivial algebra in $\Omega\mathcal{D} = (n-1)\text{Vec}$, condensing which does not reproduce C^{n+1} . However, for $n \geq 3$, there are some interesting exceptions (see for example Theorem 4.2.15 and Remark 4.2.17).

This phenomenon is very different from the condensation of 1-codimensional topological defects (recall Theorem^{ph} 3.3.9). The reason is that all 1-codimensional topological defects are missing from $\Omega\mathcal{C}$. If we use $A \in \Omega\mathcal{C}$ to define a condensed 1-codimensional topological defects, denoted by ΣA in $\mathcal{C}$, then the condensation of ΣA in $\mathcal{C}$ is reversible. We discuss this phenomenon in Section 5.1.

A special case is very important to us. If $D^{n+1} = \mathbf{1}^{n+1}$ is obtained from C^{n+1} by condensing $A \in \text{Alg}_{E_2}^c(\Omega\mathcal{C})$, then A is a Lagrangian algebra in $\Omega\mathcal{C}$, M^n is a gapped boundary of C^{n+1} (i.e. C^{n+1} is non-chiral). Therefore, we obtain a map:

$$\{\text{Lagrangian algebras in } \Omega\mathcal{C}\} \xrightarrow{\phi} \{\text{gapped boundaries of } C^{n+1}\}.$$

In this case, $C^{n+1} = Z(M)^{n+1}$ catches all the information of the gravitational anomaly of the gapped boundary M^n [KW14, KWZ15]. The non-degenerate braided fusion $(n-1)$ -category $\Omega\mathcal{C}$, as the center of $\Omega_m\mathcal{M}$, is maximal (or universal) in a precise sense. The internal hom $[1_m, 1_m]_{\Omega\mathcal{C}}$ is again maximal by the universal property of the internal hom. Therefore, it is natural to expect that it can recover M^n . Moreover, we expect that all gapped boundaries are the consequences of the condensation of 2-codimensional topological defects. We summarize this discussion as the following result.

Theorem^{ph} 4.2.6. When C^{n+1} is anomaly-free, non-chiral and simple, ϕ is bijective and its inverse ϕ^{-1} is defined by

$$M^n = (\mathcal{M}, m) \mapsto L_2^R(1_m) = [1_m, 1_m]_{\Omega\mathcal{C}} = \begin{array}{c} \text{C}^{n+1} \\ \text{M}^n \text{ } \mathbf{1}^{n+1} \\ L_2^R(1_m) \end{array} \quad (4.2.5)$$

Two Lagrangian E_2 -algebras A and B are called 2-Morita equivalent if the associated gapped boundaries (or the fusion n -categories $\Sigma A := \text{RMod}_A(\Omega\mathcal{C})$ and $\Sigma B := \text{RMod}_B(\Omega\mathcal{C})$) are Morita equivalent (see Definition 6.1.6). We identify the Morita classes $[A] = [\text{RMod}_A(\Omega\mathcal{C})]$. $\square$

Remark 4.2.7. In a general situation, when two arbitrary anomaly-free simple $n+1$ D topological orders C^{n+1} and D^{n+1} are connected by a gapped domain wall, the information of M^n can be encoded in the internal hom $[1_m, 1_m]_{\Omega\mathcal{C} \boxtimes \Omega\mathcal{D}^{\text{op}}}$, which is a Lagrangian algebra in $\mathfrak{Z}_1(\Omega_m\mathcal{M})$. While the following two condensable E_2 -algebras:

$$[1_m, 1_m]_{\Omega\mathcal{C}} = [1_m, 1_m]_{\Omega\mathcal{C} \boxtimes \Omega\mathcal{D}^{\text{op}}} \cap \Omega\mathcal{C} \quad \text{and} \quad [1_m, 1_m]_{\Omega\mathcal{D}} = [1_m, 1_m]_{\Omega\mathcal{C} \boxtimes \Omega\mathcal{D}^{\text{op}}} \cap \Omega\mathcal{D}^{\text{op}}$$

where both “ $\cap$ ” provide a heuristic understanding of the left hand sides, encode only some incomplete information of M^n . $\diamond$

Remark 4.2.8. There are infinitely many Lagrangian algebras or gapped boundaries in a non-chiral $n+1$ D topological order C^{n+1} for $n > 2$. For example, stacking a gapped boundary of C^{n+1} with an anomaly-free n D topological order gives another gapped boundary. The Lagrangian algebras in $\mathbf{1}^{n+1}$ associated to an anomaly-free n D topological order is given more explicitly in Section 4.2.2 (1) (see for example Remark 4.2.14). $\diamond$

Since the rolling-up process plays a very important role in our theory, we would like to introduce some useful notations that are inspired by the theory of factorization homology (see for example [Lur17, AF15, AFT17, AFT16, AFR18, AKZ17, AF20]). Note that, in the first picture in (4.2.2), one can view $\Omega\mathcal{C}$ as observables living on the space manifold $(\mathbb{R}^2 \setminus D^2) \times \mathbb{R}^{n-2}$, and view $\Omega\mathcal{D}$ as observables living in the interior of the solid cylinder $D^2 \times \mathbb{R}^{n-2}$ (in spatial dimension), denoted by $\mathring{D}^2 \times \mathbb{R}^{n-2}$, i.e.,

$$\text{Obs}((\mathbb{R}^2 \setminus D^2) \times \mathbb{R}^{n-2}) = \Omega\mathcal{C}, \quad \text{Obs}(\mathring{D}^2 \times \mathbb{R}^{n-2}) = \Omega\mathcal{D}.$$

Since $\Omega\mathcal{C}$ is an E_2 -algebra in $n\text{Vec}$. It makes perfect sense to integrate it over the 2-manifold $\mathring{D}^2$. Similarly, $\Omega_m\mathcal{M}$ describes the observable living on the boundary of the solid cylinder, i.e.,

$$\text{Obs}(\partial D^2 \times \mathbb{R}^{n-2}) = \Omega_m\mathcal{M}.$$


Since $\Omega_m\mathcal{M}$ is monoidal, i.e. an E_1 -algebra in $n\text{Vec}$, it makes perfect sense to integrate it along the 1-manifold ∂D^2 . By integrating all the observables in an open neighborhood of the solid cylinder in the space⁵⁴ manifold $\mathbb{R}^2 \times \mathbb{R}^{n-2}$, we obtain

$$\begin{aligned} \text{Obs}(\mathbb{R}^2 \times \mathbb{R}^{n-2}) &= \int_{\mathbb{R}^2 \times \mathbb{R}^{n-2}} (\text{Obs}((\mathbb{R}^2 \setminus D^2) \times \mathbb{R}^{n-2}), \text{Obs}(\mathring{D}^2 \times \mathbb{R}^{n-2}), \text{Obs}(\partial D^2 \times \mathbb{R}^{n-2})) \\ &= \int_{\mathbb{R}^2 \times \mathbb{R}^{n-2}} (\Omega\mathcal{C}|_{(\mathbb{R}^2 \setminus D^2) \times \mathbb{R}^{n-2}}, \Omega\mathcal{D}|_{\mathring{D}^2 \times \mathbb{R}^{n-2}}, \Omega_m\mathcal{M}|_{\partial D^2 \times \mathbb{R}^{n-2}}) = (\Omega\mathcal{C}, L_2^R(1_m)). \end{aligned}$$

However, from the physical point of view, choosing $\Omega\mathcal{D}$ and $\Omega_m\mathcal{M}$ instead of the complete data $\mathcal{D}$ and $(\mathcal{M}, m)$ is based on certain conventions of throwing away certain topological defects. Although it is possible to make the convention or rule explicitly, it is much more convenient to simply set:

$$\text{Obs}((\mathbb{R}^2 \setminus D^2) \times \mathbb{R}^{n-2}) = \mathcal{C}, \quad \text{Obs}(\mathring{D}^2 \times \mathbb{R}^{n-2}) := \mathcal{D}, \quad \text{Obs}(\partial D^2 \times \mathbb{R}^{n-2}) := (\mathcal{M}, m).$$

Then we have

$$\begin{aligned} \text{Obs}(\mathbb{R}^2 \times \mathbb{R}^{n-2}) &= \int_{D^2 \times \mathbb{R}^{n-2}} (\mathcal{C}|_{(\mathbb{R}^2 \setminus D^2) \times \mathbb{R}^{n-2}}, \mathcal{D}|_{\mathring{D}^2 \times \mathbb{R}^{n-2}}, (\mathcal{M}, m)|_{\partial D^2 \times \mathbb{R}^{n-2}}) \\ &= (\mathcal{C}, \mathbb{1}_{\mathcal{C}}, L_2^R(1_m)) = \end{aligned} \quad (4.2.6)$$


where the triple $(\mathcal{C}, \mathbb{1}_{\mathcal{C}}, L_2^R(1_m))$, i.e., a separable n -category $\mathcal{C}$, together with a distinguished 0-morphism $\mathbb{1}_{\mathcal{C}} \in \mathcal{C}$ and a distinguished 1-morphism $L_2^R(1_m) \in \Omega_{\mathbb{1}_{\mathcal{C}}} \mathcal{C}$, can be viewed as an E_{-1} -algebra object in $(n+1)\text{Vec}$ [KWZ15, Remark 2.24].


Remark 4.2.9. Note that there are extra consistency in our convention in (4.2.6). More precisely, $\mathcal{D} \in \text{Alg}_{E_1}^{\mathcal{C}}((n+1)\text{Vec})$ and $(\mathcal{M}, m) \in \text{Alg}_{E_0}^{\mathcal{C}}((n+1)\text{Vec})$. Integrating them over a 2-manifold and a 1-manifold, respectively, amount to reduce the level of commutativity by 2 and 1, respectively. As a consequence, both cases give the triple in (4.2.6) that can be viewed as an E_{-1} -algebra [KWZ15, Remark 2.24] after the integration. More generally, one can view

$$(\mathcal{A}, a_1, a_2, a_3, \dots, a_k) \in \text{Alg}_{E_{-k+1}}^{\mathcal{C}}((n+1)\text{Vec}), \quad (4.2.7)$$

where $\mathcal{A}$ is a separable n -category, $a_1 \in \mathcal{A}$ is a 0-morphism, $a_2 \in \Omega_{a_1} \mathcal{A}$ is a 1-morphism, $a_3 \in \Omega_{a_2}(\Omega_{a_1} \mathcal{A})$ is a 2-morphism, $a_k \in \Omega_{a_{k-1}}(\dots \Omega_{a_2}(\Omega_{a_1} \mathcal{A}) \dots)$ is a $(k-1)$ -morphism. This notation is quite useful when we study the integration of $\mathcal{D}$ and $(\mathcal{M}, m)$ over $D^k \times \mathbb{R}^{n-k}$ in next section. We suspect that our theory might be related to the so-called β -version of factorization homology [AFT17, AFT16, AFR18], our understanding of which is too limited to say anything precise. $\diamond$

Remark 4.2.10. The notations introduced in (4.2.6) are used later. Although we introduced the integral notation heuristically, it turns out that this integral has a mathematical foundation, which is called factorization homology [Lur17, AF15] (see [AF20] for a recent review). We hope to clarify this point in future publications. $\diamond$

Consider two $n+1$ D anomaly-free topological orders $C^{n+1}, D^{n+1}, E^{n+1}$ and two gapped domain walls M^n, N^n (as illustrated in the left picture in Eq. (4.2.8)) such that

$$\begin{aligned} & \begin{array}{|c|c|c|} \hline C^{n+1} & D^{n+1} & E^{n+1} \\ \hline M^n & & N^n \\ \hline \end{array} \xrightarrow{\text{rolling up}} \begin{array}{|c|} \hline C^{n+1} \\ \hline L_2^R(L_2^R(1_n)) \\ \hline \end{array} \end{aligned} \quad (4.2.8)$$


⁵⁴We generalize it to an integral on spacetime manifold in Section 6.2.

- D^{n+1} and M^n are obtained from C^{n+1} by condensing the 2-codimensional topological defect $A \in \text{Alg}_{E_2}^c(\Omega\mathcal{C})$, i.e., $\Omega\mathcal{D} = \text{Mod}_A^{E_2}(\Omega\mathcal{C})$ and $\Omega_m\mathcal{M} = \text{RMod}_A(\Omega\mathcal{C})$;
- E^{n+1} and N^n are obtained from D^{n+1} by condensing the 2-codimensional topological defect $B \in \text{Alg}_{E_2}^c(\Omega\mathcal{D})$, i.e., $\Omega\mathcal{E} = \text{Mod}_B^{E_2}(\Omega\mathcal{D})$ and $\Omega_n\mathcal{N} = \text{RMod}_B(\Omega\mathcal{D})$.

The left bulk-to-wall functors associated to M^n and N^n are $L_2 = - \otimes A$ or $L_2 = - \otimes_A B$, respectively. By rolling up two domain walls as in (4.2.8), we obtain a condensable E_2 -algebra in $\Omega\mathcal{C}$ given by $L_2^R(L_2^R(B)) = B$, where $B \in \text{Alg}_{E_2}^c(\text{Mod}_A^{E_2}(\Omega\mathcal{C}))$ and can be automatically viewed as an E_2 -algebra in $\Omega\mathcal{C}$. By condensing $B \in \Omega\mathcal{C}$, we obtain E^{n+1} as the condensed phase and a gapped domain wall $M^n \boxtimes_{D^{n+1}} N^n$, i.e.

$$\Omega\mathcal{E} = \text{Mod}_B^{E_2}(\Omega\mathcal{C}), \quad \Omega_{m \boxtimes_{D^n} n}(\mathcal{M} \boxtimes_{\mathcal{D}} \mathcal{N}) \simeq \Omega_m\mathcal{M} \boxtimes_{\Omega\mathcal{D}} \Omega_n\mathcal{N} = \text{RMod}_B(\Omega\mathcal{C}). \quad (4.2.9)$$

Remark 4.2.11. Above discussion has an immediate consequence Corollary^{ph} 4.2.30 as we will show later. The ‘ $\simeq$ ’ in (4.2.9) is a special case of a general principle of vanishing spatial fusion anomaly [KZ20, KYZ21] for 1-codimensional topological defects. We will discuss it elsewhere. $\diamond$

4.2.2 General construction

In this subsection, we provide some general constructions of the condensations of topological defects of codimension 2.

(1) $C^{n+1} = D^{n+1} = \mathbf{1}^{n+1}$: In this case, the gapped domain wall M^n is precisely an anomaly-free nD topological order $M^n = (\mathcal{M}, m)$, where $\mathcal{M}$ is an indecomposable separable n -category and m is a distinguished object in $\mathcal{M}$ that labels the anomaly-free nD topological order. The indecomposable multi-fusion $(n-1)$ -category $\Omega_m\mathcal{M}$ is the category of topological defects in the nD topological order labeled by m . All objects in $\mathcal{M}$ label anomaly-free nD topological orders that are all Morita equivalent. Mathematically, $\Omega_m\mathcal{M}$ and $\Omega_{m'}\mathcal{M}$ are Morita equivalent for $m, m' \in \mathcal{M}$, and we have $\mathcal{M} = \Sigma\Omega_m\mathcal{M}$. If we further assume that m is simple, then $\Omega_m\mathcal{M}$ is a non-degenerate fusion $(n-1)$ -category and $\Omega_m^2\mathcal{M} := \Omega(\Omega_m\mathcal{M})$ is a non-degenerate braided fusion $(n-2)$ -category, i.e., $\mathfrak{Z}_2(\Omega_m^2\mathcal{M}) \simeq (n-2)\text{Vec}$. Moreover, we have

$$\Omega_m\mathcal{M} = \Sigma\Omega_m^2\mathcal{M} = \text{RMod}_{\Omega_m^2\mathcal{M}}((n-1)\text{Vec}),$$

which provides a coordinate system for $\Omega_m\mathcal{M}$ that are very useful for computation.

Recall that all 2-codimensional topological defects in $\mathbf{1}^{n+1}$ form the trivial braided fusion $(n-1)$ -category $(n-1)\text{Vec}$. In this case, note that the first level of the left bulk-to-wall functor $L_1 : \mathcal{C} \rightarrow \mathcal{M}$ is defined by $- \odot m$, where $\odot$ denotes the natural $\mathcal{C}$ -action (of 1-codimensional topological defects) on $\mathcal{M}$. We also have the second level of left bulk-to-wall map that maps 2-codimensional topological defects in $\mathbf{1}^{n+1}$ to M^n defined as follows:

$$\begin{array}{ccc} (n-1)\text{Vec} & \xrightarrow{L_2 = - \boxtimes \Omega_m^2\mathcal{M}} & \Omega_m\mathcal{M} = \text{RMod}_{\Omega_m^2\mathcal{M}}((n-1)\text{Vec}) \\ a & \mapsto & a \odot 1_m = a \odot \Omega_m^2\mathcal{M}. \end{array}$$

Its physical meaning is illustrated below.

$$\begin{array}{ccc} \mathbf{1}^{n+1} & & \mathbf{1}^{n+1} \\ & \searrow & \\ (n-1)\text{Vec} & \xrightarrow{L_2 = - \otimes 1_m} \Omega_m\mathcal{M} & \begin{array}{l} \Omega_m\mathcal{M} = \text{RMod}_{\Omega_m^2\mathcal{M}}((n-1)\text{Vec}) \\ 1_m = \Omega_m^2\mathcal{M} \in \text{RMod}_{\Omega_m^2\mathcal{M}}((n-1)\text{Vec}) \end{array} \end{array}$$

The right adjoint L_2^R of L_2 is precisely the forgetful functor $f : \text{RMod}_{\Omega_m^2\mathcal{M}}((n-1)\text{Vec}) \rightarrow (n-1)\text{Vec}$. Therefore, we obtain $L_2^R(1_m) = \Omega_m^2\mathcal{M} \in \text{Alg}_{E_2}^c((n-1)\text{Vec})$. Moreover, by Theorem^{ph} 4.2.6, $L_2^R(1_m)$ is a Lagrangian algebra in $(n-1)\text{Vec}$, i.e. $\text{Mod}_{\Omega_m^2\mathcal{M}}^{E_2}((n-1)\text{Vec}) \simeq (n-1)\text{Vec}$. This result can be reformulated as a mathematical result.


Proposition^o 4.2.12. A non-degenerate braided fusion n -category $\mathcal{A}$ is automatically a Lagrangian algebra in $(n+1)\text{Vec}$ (recall Definition 2.3.31), i.e.

$$\text{Mod}_{\mathcal{A}}^{\text{E}_2}((n+1)\text{Vec}) \simeq (n+1)\text{Vec}.$$

Remark 4.2.13. When $n=2$, the above result was proved in mathematical literature [DN21, JF22, DX24]. Our results provide a physical meaning to this $n=2$ mathematical result and naturally generalize it to all dimensions. $\diamond$

Remark 4.2.14. Recall Proposition 3.3.18 and Remark 3.3.20, Proposition 4.2.12 is the second mathematical reformulation of the fact that condensing an anomaly-free $n\text{D}$ topological order in the trivial $n+1\text{D}$ topological order $\mathbf{1}^{n+1}$ reproduces $\mathbf{1}^{n+1}$. More precisely,

1. When an anomaly-free $n\text{D}$ simple topological order M^n is viewed as a 1-codimensional topological defect in $\mathbf{1}^{n+1}$, we can condense it to obtain $\mathbf{1}^{n+1}$ as the condensed phase and a gapped domain wall, which is nothing but M^n characterized by the pair $(\mathcal{M}, m)$. Mathematically, it amounts to condense the simple condensable E_1 -algebra $\Omega_m \mathcal{M}$ in $n\text{Vec}$.
2. When we rollup an anomaly-free $n\text{D}$ topological order M^n in one direction, it becomes a cylinder $S^1 \times \mathbb{R}^{n-2}$ in spatial n -dimensions $\mathbb{R}^n$. By shrinking the size of S^1 , it becomes a 2-codimensional topological defects in $\mathbf{1}^{n+1}$.



As an object in $(n-1)\text{Vec}$, it is precisely the non-degenerate braided fusion $(n-2)$ -category $\Omega_m^2 \mathcal{M}$, which is automatically a Lagrangian E_2 -algebra in $(n-1)\text{Vec}$. By condensing this 2-codimensional topological defects, we obtain $\mathbf{1}^{n+1}$ as the condensed phase.

Note that the second reformulation is possible because $\Omega_m^2 \mathcal{M}$ gives a complete characterization of all topological defects M^n up to condensation descendants. $\diamond$

(2) $C^{n+1} = \mathbf{1}^{n+1} \neq D^{n+1}$: We have seen that condensing a Lagrangian algebra $\mathcal{A}$ in $(n-1)\text{Vec}$, i.e. a non-degenerate braided fusion $(n-2)$ -category, produces the trivial phase $\mathbf{1}^{n+1}$ as the condensed phase. If $\mathcal{A}$ is not Lagrangian, i.e., a braided fusion $(n-2)$ -category, it is precisely a simple condensable E_2 -algebra in $(n-1)\text{Vec}$. By condensing $\mathcal{A}$ in $(n-1)\text{Vec}$, we obtain a condensed phase D^{n+1} and a gapped domain wall $M^n = (\mathcal{M}, m)$ such that

$$\Omega \mathcal{D} \simeq \text{Mod}_{\mathcal{A}}^{\text{E}_2}((n-1)\text{Vec}) \quad \text{and} \quad \Omega_m \mathcal{M} \simeq \text{RMod}_{\mathcal{A}}((n-1)\text{Vec}) = \Sigma \mathcal{A}. \quad (4.2.10)$$

Note that we have $\mathcal{D} = \Sigma \Omega \mathcal{D}$ and $\mathcal{M} = \Sigma \Omega_m \mathcal{M} = \Sigma^2 \mathcal{A}$. By the boundary-bulk relation, we obtain

$$\begin{aligned} \text{Mod}_{\mathcal{A}}^{\text{E}_2}((n-1)\text{Vec}) &\simeq \mathfrak{Z}_1(\text{RMod}_{\mathcal{A}}((n-1)\text{Vec}))^{\text{op}}, \\ \mathcal{D} = \Sigma \text{Mod}_{\mathcal{A}}^{\text{E}_2}((n-1)\text{Vec}) &\simeq \mathfrak{Z}_0((\mathcal{M}, m)) = \text{Fun}(\Sigma^2 \mathcal{A}, \Sigma^2 \mathcal{A}) \simeq \text{BMod}_{\Sigma \mathcal{A} | \Sigma \mathcal{A}}(n\text{Vec}). \end{aligned} \quad (4.2.11)$$

Moreover, we have $L^R(1_m) = L^R(\mathcal{A}) = [\mathcal{A}, \mathcal{A}]_{(n-1)\text{Vec}} = \mathcal{A}$, or equivalently,

$$\int_{\mathbb{R}^2 \times \mathbb{R}^{n-2}} (n\text{Vec}|_{(\mathbb{R}^2 \setminus D^2) \times \mathbb{R}^{n-2}}, \mathcal{D}|_{D^2 \times \mathbb{R}^{n-2}}, (\mathcal{M}, m)|_{\partial D^2 \times \mathbb{R}^{n-2}}) = (n\text{Vec}, \mathbb{1}, \mathcal{A}). \quad (4.2.12)$$

One can see immediately that D^{n+1} can be obtained from C^{n+1} by condensing 1-codimensional topological defect $\Sigma \mathcal{A}$, which is a condensable E_1 -algebra in $(n+2)\text{Vec}$. We restudy this phenomenon in details in Section 5. We summarize above results as a mathematical result.

Theorem° 4.2.15. For $m \geq 1$, a simple condensable E_2 -algebra in $(m+1)\text{Vec}$ is precisely a braided fusion m -category $\mathcal{A}$. We have

$$\text{Mod}_{\mathcal{A}}^{E_2}((m+1)\text{Vec})^{\text{op}} \simeq \mathfrak{Z}_1(\text{RMod}_{\mathcal{A}}((m+1)\text{Vec})) \simeq \mathfrak{Z}_1(\Sigma\mathcal{A}), \quad (4.2.13)$$

$$\Sigma \text{Mod}_{\mathcal{A}}^{E_2}((m+1)\text{Vec})^{\text{op}} \simeq \text{Fun}(\Sigma^2\mathcal{A}, \Sigma^2\mathcal{A}) \simeq \text{BMod}_{\Sigma\mathcal{A}|\Sigma\mathcal{A}}((m+2)\text{Vec}). \quad (4.2.14)$$

Example 4.2.16. Let $\mathcal{A} = n\text{Rep}(G)$ viewed as a condensable E_2 -algebra in $(n+1)\text{Vec}$, which is the category of topological defects of codimension 2 and higher in $\mathbf{1}^{n+3}$. By condensing $n\text{Rep}(G)$ in $(n+1)\text{Vec}$, we obtain

$$\text{Mod}_{n\text{Rep}(G)}^{E_2}((n+1)\text{Vec}) \simeq \mathfrak{Z}_1((n+1)\text{Rep}(G)).$$

Therefore, we obtain GT_G^{n+3} as the condensed phase. $\heartsuit$

Remark 4.2.17. The $n = 3$ case of (4.2.13) was first proved in [DN21, Theorem 4.11]. In this case, Décoppet proved that every fusion 2-category $\mathcal{B}$ is Morita equivalent to a connected one, i.e., $\Sigma\mathcal{A}$ for a braided fusion 1-category $\mathcal{A}$ [Déc23a]. As a consequence, every non-chiral 3+1D topological order can be obtained by condensing a simple condensable E_2 -algebra in 2Vec , or equivalently, by condensing a 2-codimensional topological defect in $\mathbf{1}^4$. However, it is not true that all gapped boundaries of a non-chiral 4D topological order can be obtained in this way⁵⁵. In general, a simple gapped boundary M^n of a simple non-chiral $n+1$ D topological order D^{n+1} can be obtained from $C^{n+1} = \mathbf{1}^{n+1}$ via the condensation of a topological defect of codimension 2 if and only if the fusion $(n-1)$ -category $\Omega_m\mathcal{M}$ is connected as a separable 2-category, i.e., $\Omega_m\mathcal{M} \simeq \Sigma\mathcal{A}$ for a braided fusion $(n-2)$ -category. Moreover, we do not expect that Décoppet's result generalizes to fusion k -categories for $k \geq 3$. The $k = 3$ case is special because 3+1D topological orders are very rare [LKW18, LW19, JF20]⁵⁶. $\diamond$

Remark 4.2.18. Similar to Section 3.3.2, one can also discuss general examples of the condensation of 2-codimensional topological defects. We postpone it to Section 5.2.2 as a special case of the condensations of k -codimensional topological defects. $\diamond$

(3) C^{n+1} is non-trivial and non-chiral: The following result provides a physical way to construct Lagrangian E_2 -algebras in $\Omega\mathcal{C}$ from a single one. It is an immediate consequence of Theorem^{ph} 3.3.14.

Theorem^{ph} 4.2.19. Given an anomaly-free non-chiral simple $n+1$ D topological order C^{n+1} and an anomaly-free n D topological order N^n . If A is a Lagrangian E_2 -algebra in $\Omega\mathcal{C}$, then $A \boxtimes \Omega N \in \Omega\mathcal{C} \boxtimes (n-1)\text{Vec} \simeq \Omega\mathcal{C}$ is also a Lagrangian E_2 -algebra in $\Omega\mathcal{C}$. Moreover, we can obtain all Lagrangian E_2 -algebras in $\Omega\mathcal{C}$ from a single one by the following three-step process:

- (1) First, stacking an anomaly-free n D topological order N^n to a given gapped boundary X^n of C^{n+1} , we obtain a new gapped boundary $X^n \boxtimes N^n$ of C^{n+1} ;
- (2) Secondly, condense $X^n \boxtimes N^n$ to a new but Morita equivalent gapped boundary Y^n ;
- (3) Thirdly, rolling up the boundary Y^n to a 2-codimensional topological defect A_Y in C^{n+1} , it gives a Lagrangian E_2 -algebra in $\Omega\mathcal{C}$.

All Lagrangian E_2 -algebras in $\Omega\mathcal{C}$ arise in this way from a single gapped boundary X^n . $\square$

Remark 4.2.20. One has to treat the roll-up process with caution. It is possible to switch the order the second step and the third step, i.e., obtaining A_Y from $A_{X \boxtimes N}$ by a condensation restricted on the 2-codimensional topological defect. It is because the roll-up process indeed maps a gapped domain wall between $X^n \boxtimes N^n$ and Y^n to a gapped domain wall between $A_{X \boxtimes N}$ and A_Y . However, such obtained condensations on $A_{X \boxtimes N}$ belongs to a very limited sub-family of all possible condensations. A generic condensations on $A_{X \boxtimes N}$ produces a condensation descendant, which is not necessarily an algebra. Also notice that the roll-up process sometimes maps a gapless domain wall between two gapped boundaries to a gappable domain wall between two 2-codimensional topological defects. For example, by rolling up a 1+1D anomaly-free rational CFT, we obtain a gapless 0+1D defect in $\mathbf{1}^3$ that is gappable. $\diamond$

⁵⁵All gapped boundaries of a non-chiral 4D topological order can be obtained from $\mathbf{1}^4$ by condensing a 2-codimensional topological defect followed by a condensation on the boundary. This combined process was called a two-step condensation in [Kon14].

⁵⁶By topological Wick rotation [KZ18b, KZ20, KZ21], this fact also suggests that 2+1D “fully dualizable” CFT's are rare comparing to 1+1D CFT's.

We recall a well-known result.

Theorem 4.2.21 ([KR08, DMNO13]). For a fusion 1-category $\mathcal{A}$, there is a one-to-one correspondence between Lagrangian E_2 -algebras in $\mathfrak{Z}_1(\mathcal{A})$ and indecomposable left $\mathcal{A}$ -modules, or equivalently, the Morita classes of indecomposable condensable E_1 -algebras in $\mathcal{A}$. Moreover, given an indecomposable condensable E_1 -algebra A in $\mathcal{A}$, the associated Lagrangian E_2 -algebra in $\mathfrak{Z}_1(\mathcal{A})$ is the full center of A .

Example 4.2.22. On the one hand, Lagrangian E_2 -algebras in $\mathfrak{Z}_1(\text{Rep}(G))$ are classified by a pair (H, ω) , where H is a subgroup of G and $\omega \in Z^2(H, \mathbb{C}^\times)$. On the other hand, left indecomposable $\text{Rep}(G)$ -modules are also classified by the same pairs (H, ω) . $\heartsuit$

We have a mathematical reformulation of Corollary^{ph} 3.3.39 in terms of Lagrangian E_2 -algebras as shown below. It is a generalization of above result for the $n = 1$ case and the $n = 2$ case [JFR23, DX24] to all $n > 2$.

Theorem^o 4.2.23. Let $\mathcal{T}$ be a non-degenerate fusion n -category $\mathcal{T}$, i.e., $\mathcal{T} = \Sigma\Omega\mathcal{T}$ and $\Omega\mathcal{T}$ is a non-degenerate braided fusion $(n-1)$ -category. For an indecomposable multi-fusion n -category $\mathcal{B}$, there are one-to-one correspondences among the following three sets:


- (1) Lagrangian E_2 -algebras in $\mathfrak{Z}_1(\mathcal{B})$ (up to isomorphisms) in the 2-Morita class $[\mathcal{B} \boxtimes \mathcal{T}]$ (recall Theorem^{ph} 4.2.6);
- (2) left indecomposable $\mathcal{B} \boxtimes \mathcal{T}$ -modules modulo the equivalence relation: $\mathcal{M} \sim \mathcal{N}$ if and only if $\mathcal{N} \simeq \mathcal{M} \boxtimes \mathcal{V}$ for $\mathcal{V} \in (n+1)\text{Vec}^\times$.
- (3) indecomposable condensable E_1 -algebras in $\mathcal{B} \boxtimes \mathcal{T}$ modulo the equivalence relation: $A \sim B$ if B is Morita equivalent to $A \boxtimes L \in (\mathcal{B} \boxtimes \mathcal{T}) \boxtimes n\text{Vec} \simeq \mathcal{B} \boxtimes \mathcal{T}$ for a Lagrangian E_1 -algebra L in $n\text{Vec}$ (i.e., L is a non-degenerate multi-fusion $(n-1)$ -category).

Remark 4.2.24. Similar to $n = 1$ case, we expect that, for an indecomposable condensable E_1 -algebra A in $\mathcal{B} \boxtimes \mathcal{T}$, the associated Lagrangian E_2 -algebra in $\mathfrak{Z}_1(\mathcal{B})$ is given by the full center of A . The notion of the full center of an algebra in a fusion n -category can be defined by the universal property generalizing that of the full center of an algebra in fusion 1-categories [Dav10a, KYZ21]. We will introduce this notion as a special case of the E_k -center of an E_k -algebra in an E_k -monoidal higher category in [KZZZ25]. $\diamond$

Now the following result provides a physical meaning to the classification result in Corollary^{ph} 2.4.15.

Theorem^{ph} 4.2.25. For $n \geq 2$ and a braided fusion $(n-1)$ -category $\mathcal{A}$, a condensable E_2 -algebra in $\mathfrak{Z}_1(\Sigma\mathcal{A})^{\text{op}} \simeq \text{Mod}_{\mathcal{A}}^{E_2}(n\text{Vec})$ (recall (4.2.13)) is precisely given by a braided multi-fusion $(n-1)$ -category $\mathcal{B}$ equipped with a braided monoidal functor $\mathcal{A} \rightarrow \mathcal{B}$. The condensable E_2 -algebra $\mathcal{B}$ is simple if and only if $\mathcal{B}$ is fusion, and it is Lagrangian if and only if $\mathcal{B}$ is non-degenerate. The 2-Morita class of the Lagrangian E_2 -algebra is determined by the Morita class of $\Sigma\mathcal{B}$.

Moreover, by condensing $\mathcal{B}$ in $\mathfrak{Z}_1(\Sigma\mathcal{A})^{\text{op}}$, where $\mathcal{B}$ is viewed as a condensable defect of codimension 2 in the anomaly-free topological order $\mathcal{C}^{n+1}$ such that $\mathcal{C} = \mathfrak{Z}_0(\Sigma^2\mathcal{A})^{\text{op}} \simeq \text{Mod}_{\Sigma\mathcal{A}}^{E_1}((n+1)\text{Vec})$ and $\Omega\mathcal{C} = \mathfrak{Z}_1(\Sigma\mathcal{A})^{\text{op}}$ as illustrated below,



we obtain a condensed phase $\mathcal{D}^{n+1}$ and a gapped domain wall $\mathcal{M}^n = (\mathcal{M}, m)$ such that

$$\begin{aligned} \mathcal{D} &\simeq \text{Mod}_{\Sigma\mathcal{B}}^{E_1}(\text{Mod}_{\Sigma\mathcal{A}}^{E_2}((n+1)\text{Vec})), \quad (\mathcal{M}, m) = (\text{BMod}_{\Sigma\mathcal{A}|\Sigma\mathcal{B}}((n+1)\text{Vec}), \Sigma\mathcal{B}), \\ \Omega\mathcal{D} &\simeq \text{Mod}_{\mathcal{B}}^{E_2}(\text{Mod}_{\mathcal{A}}^{E_2}(n\text{Vec})) \simeq \text{Mod}_{\mathcal{B}}^{E_2}(n\text{Vec}) \simeq \mathfrak{Z}_1(\Sigma\mathcal{B})^{\text{op}}, \\ \Omega_m\mathcal{M} &= \text{Fun}_{\Sigma\mathcal{A}|\Sigma\mathcal{B}}(\Sigma\mathcal{B}, \Sigma\mathcal{B}) \simeq \text{RMod}_{\mathcal{B}}(\text{Mod}_{\mathcal{A}}^{E_2}(n\text{Vec})). \end{aligned} \quad (4.2.15)$$

Moreover, set $\mathcal{L}^n = (\mathcal{L}, l) = (\Sigma^2\mathcal{A}, \Sigma\mathcal{A})$. We have

$$\begin{aligned} \Omega_l\mathcal{L} \boxtimes_{\Omega\mathcal{C}} \Omega_m\mathcal{M} &= \Sigma\mathcal{A} \boxtimes_{\text{Mod}_{\mathcal{A}}^{E_2}(n\text{Vec})} \text{RMod}_{\mathcal{B}}(\text{Mod}_{\mathcal{A}}^{E_2}(n\text{Vec})) \simeq \Sigma\mathcal{B}. \\ \mathcal{L} \boxtimes_{\mathcal{C}} \mathcal{M} &= \Sigma^2\mathcal{A} \boxtimes_{\Sigma\text{Mod}_{\mathcal{A}}^{E_2}(n\text{Vec})} \Sigma(\text{RMod}_{\mathcal{B}}(\text{Mod}_{\mathcal{A}}^{E_2}(n\text{Vec}))) \simeq \Sigma^2\mathcal{B}. \end{aligned} \quad (4.2.16) \quad \square$$

Remark 4.2.26. The $n = 2$ case of Theorem^{ph} 4.2.25 was obtained in [JFR23, DX24, Xu24a]. When $\mathcal{A} = \text{Rep}(G)$, this braided functor $\text{Rep}(G) \rightarrow \mathcal{B}$ necessarily factors through $\text{Rep}(G) \xrightarrow{f} \text{Rep}(H) \hookrightarrow \mathcal{B}$ [DMNO13]. This result also says that non-minimal non-degenerate extensions of $\text{Rep}(H)$ also have physical meanings. $\diamond$

Remark 4.2.27. Note that Theorem^{ph} 4.2.25 of a 2-codimensional defect condensation is an analogue of Theorem^{ph} 3.3.32 of a 1-codimensional defect condensation, and can be generalized further to higher codimensional defect condensations (see Theorem^{ph} 5.2.20). $\diamond$

4.2.3 Witt equivalence

Proposition^{ph} 4.2.28. If two anomaly-free $n+1$ D topological order C^{n+1} and D^{n+1} are Morita equivalent (i.e., admitting a gapped anomaly-free domain wall), then there exists the third anomaly-free $n+1$ D topological order A^{n+1} such that C^{n+1} and D^{n+1} can be obtained from A^{n+1} via two different condensations of 2-codimensional topological defects.

Proof. It follows from above Lemma by choosing $A^{n+1} = C^{n+1} \boxtimes Z(P)^{n+1} = D^{n+1} \boxtimes Z(Q)^{n+1}$. Note that $A_{\mathcal{P}} = [\mathbb{1}_{\mathcal{P}}, \mathbb{1}_{\mathcal{P}}]_{\mathfrak{Z}_1(\mathcal{P})}$ is a Lagrangian E_2 -algebra in $\mathfrak{Z}_1(\mathcal{P})$, and $A_Q = [\mathbb{1}_Q, \mathbb{1}_Q]_{\mathfrak{Z}_1(Q)}$ is a Lagrangian E_2 -algebra in $\mathfrak{Z}_1(Q)$. By condensing the condensable E_2 -algebra $A_1 = \mathbb{1}_{\mathcal{P}} \boxtimes A_{\mathcal{P}} \in \Omega\mathcal{C} \boxtimes \mathfrak{Z}_1(\mathcal{P})$, we obtain C^{n+1} as the condensed phase. By condensing the condensable E_2 -algebra $A_2 = \mathbb{1}_{\mathcal{D}} \boxtimes A_Q \in \Omega\mathcal{D} \boxtimes \mathfrak{Z}_1(Q)$, we obtain D^{n+1} as the condensed phase. $\square$

Remark 4.2.29. When $n = 2$, it was proved in [DMNO13] there exists an anomaly-free $n+1$ D topological order B^{n+1} such that B^{n+1} can be obtained from both C^{n+1} and D^{n+1} via condensations of 2-codimensional topological defects. It is natural to ask if it is true for $n > 2$ (see a related question in Remark 4.2.38). $\diamond$

The next result was first obtained in [Xu24a].

Corollary^{ph} 4.2.30. Two Morita equivalent anomaly-free topological order C^{3+1} and D^{3+1} can be obtained from each other by condensing 2-codimensional topological defects.

Proof. This is an immediate consequence of Proposition^{ph} 2.2.23, Remark 4.2.17 and discussion around (4.2.8). $\square$

Remark 4.2.31. Similar to Remark 4.2.17, not every gapped domain wall between C^4 and D^4 can be obtained from condensing a 2-codimensional topological defect in either C^4 or D^4 . $\diamond$

The following notion is a direct generalization of that for 1-categories introduced in [DMNO13].

Definition 4.2.32. Two non-degenerate braided fusion n -categories $\mathcal{A}$ and $\mathcal{B}$ are called *Witt equivalent* if there exist two fusion n -categories $\mathcal{P}$ and $\mathcal{Q}$ and a braided equivalence:

$$\mathcal{A} \boxtimes \mathfrak{Z}_1(\mathcal{P}) \simeq \mathcal{B} \boxtimes \mathfrak{Z}_1(\mathcal{Q}).$$

The physical result Proposition^{ph} 2.2.23 immediately implies the following mathematical result.

Corollary^o 4.2.33. Two non-degenerate braided fusion n -categories $\mathcal{A}$ and $\mathcal{B}$ are Witt equivalent if and only if there exists a fusion n -category $\mathcal{M}$ and braided equivalence:

$$\mathcal{A} \boxtimes \mathcal{B}^{\text{op}} \simeq \mathfrak{Z}_1(\mathcal{M}).$$

Corollary^o 4.2.34. Two non-degenerate braided fusion n -categories $\mathcal{A}$ and $\mathcal{B}$ are Witt equivalent if and only if there exists a third non-degenerate braided fusion n -category $\mathcal{C}$ and $A, B \in \text{Alg}_{E_2}^c(\mathcal{C})$ such that

$$\mathcal{A} \simeq \text{Mod}_A^{E_2}(\mathcal{C}) \quad \text{and} \quad \mathcal{B} \simeq \text{Mod}_B^{E_2}(\mathcal{C}).$$

Proposition^o 4.2.35. The notion of Witt equivalence is clearly an equivalence relation. We denote the Witt equivalence class of $\mathcal{C}$ by $[\mathcal{C}]$. The Witt equivalence classes of non-degenerate braided fusion n -categories clearly form a group, called the Witt group and denoted by Witt^n , with the multiplication defined by $[\mathcal{C}] \star [\mathcal{D}] := [\mathcal{C} \boxtimes \mathcal{D}]$, the unit defined by $n\text{Vec}$ and the inverse given by $[\mathcal{C}]^{-1} = [\mathcal{C}^{\text{op}}]$.

Remark 4.2.36. It was known that the map $[\mathcal{C}] \mapsto \Sigma^2 \mathcal{C}$ defines a group isomorphism from the Witt group Witt^n to the group $(n+3)\text{Vec}^\times$ of invertible separable $(n+2)$ -categories [JF22, KZ24]. $\diamond$

We generalize the notion of a completely anisotropic non-degenerate braided fusion 1-category [DMNO13] to higher categories.

Definition 4.2.37. A non-degenerate braided fusion n -category $\mathcal{B}$ is called *completely anisotropic* if every condensable E_2 -algebra in $\mathcal{B}$ is isomorphic to $\mathbb{1}_{\mathcal{B}} \boxtimes A \in \mathcal{B} \boxtimes n\text{Vec} \simeq \mathcal{B}$, where A is a condensable E_2 -algebra in $n\text{Vec}$. $\square$

Remark 4.2.38. An obvious question is if there exists a unique completely anisotropic non-degenerate braided fusion n -category in each Witt equivalence class as in the $n=1$ case [DMNO13]. We hope to explore this question in the future. $\diamond$

4.2.4 Examples in 3+1D

We have already seen the condensations of 2-codimensional topological defects in 1^4 . In this subsection, we provide some examples in 3+1D $\mathbb{Z}_2$ -gauge theory.


Example 4.2.39. Consider 3+1D $\mathbb{Z}_2$ gauge theory $\mathcal{GT}_{\mathbb{Z}_2}^4$. Then $\Omega\mathcal{GT}_{\mathbb{Z}_2}^4$ is a nondegenerate braided fusion 2-category and we have $\Omega\mathcal{GT}_{\mathbb{Z}_2}^4 \simeq \mathfrak{Z}_1(2\text{Rep}(\mathbb{Z}_2)) \simeq \mathfrak{Z}_1(2\text{Vec}_{\mathbb{Z}_2})$. This braided fusion 2-category was studied in [KTZ20b] and via the 3+1D toric code model in [KTZ20a]. We use the notation in [KTZ20a, ZLZH⁺23]. There are 4 simple objects $\mathbb{1}, \mathbb{1}_c, m, m_c$ in $\Omega\mathcal{GT}_{\mathbb{Z}_2}^4$ with the fusion rule

$$m \otimes m \simeq \mathbb{1}, \quad \mathbb{1}_c \otimes \mathbb{1}_c \simeq \mathbb{1}_c \oplus \mathbb{1}_c, \quad m \otimes \mathbb{1}_c \simeq \mathbb{1}_c \otimes m \simeq m_c.$$

The hom categories are depicted in the following diagram:



We also denote the simple objects in the hom categories in the following diagram:



The composition rules are given by $e \circ e = \mathbb{1}, z \circ z = \mathbb{1}_c, x \circ y = \mathbb{1}_c \oplus z$ and $y \circ x = \mathbb{1} \oplus e$.

We review three examples of Lagrangian E_2 -algebras in $\Omega\mathcal{GT}_{\mathbb{Z}_2}^4$ constructed in [ZLZH⁺23].

1. $A_e = \mathbb{1}_c$: Its multiplication morphism is given by

$$\mathbb{1}_c \otimes \mathbb{1}_c = \mathbb{1}_c \oplus \mathbb{1}_c \xrightarrow{\mathbb{1}_c \oplus 0} \mathbb{1}_c.$$

and the unit $x: \mathbb{1} \rightarrow \mathbb{1}_c$, together with the identity 2-associators, 2-unitors and 2-commutators is a Lagrangian algebra. There are two simple modules $\mathbb{1}_c$ and m_c , and only $\mathbb{1}_c$ itself is a local module. Condensing $\mathbb{1}_c$ leads to the rough boundary of the 3+1D toric code model. The topological defects on the boundary form a fusion 2-category $2\text{Vec}_{\mathbb{Z}_2}$. In the light of Theorem^{ph} 4.2.25, $\Omega\mathcal{GT}_{\mathbb{Z}_2}^4$ is equipped with another coordinate system:

$$\Omega\mathcal{GT}_{\mathbb{Z}_2}^4 \simeq \mathfrak{Z}_1(\Sigma\text{Rep}(G)) \simeq \text{Mod}_{\text{Rep}(G)}^{E_2}(2\text{Vec})^{\text{op}}.$$

In this coordinate system, the Lagrangian E_2 -algebra A_e is the one associated to the forgetful functor $\text{Rep}(\mathbb{Z}_2) \rightarrow \text{Vec}$.

2. $A_m = \mathbb{1} \oplus m$: The algebra A_m has the multiplication 1-morphism defined component-wise by

$$\begin{array}{cccc}
\mathbb{1} \otimes \mathbb{1} & \mathbb{1} \otimes m & m \otimes \mathbb{1} & m \otimes m \\
\downarrow 1_{\mathbb{1}} & \downarrow 1_m & \downarrow 1_m & \downarrow 1_{\mathbb{1}} \\
\mathbb{1} & m & m & \mathbb{1}
\end{array}$$

and the unit 1-morphism $\mathbb{1} \xrightarrow{1 \oplus 0} \mathbb{1} \oplus m$. The 2-associator and 2-units are identity 2-morphisms. The 2-commutator is trivial on all components except $\beta_{m,m} = \pm 1$. Two different choices of the 2-commutator define two commutative algebra structures, but they are equivalent to each other. There are two simple A_m -modules A_m and $\mathbb{1}_c \otimes A_1$, and only A_1 itself is a local module. Condensing A_m leads to the smooth boundary of the 3+1D toric code model. The topological defects on the boundary form a fusion 2-category $2\text{Rep}(\mathbb{Z}_2)$. In the light of Theorem^{ph} 4.2.25,

this Lagrangian E_2 -algebra is the one associated to the trivial minimal modular extension $\text{Rep}(\mathbb{Z}_2) \rightarrow \mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$.

3. $A_m^{\text{tw}} = (1 \oplus m)^{\text{tw}}$: There is another Lagrangian algebra structure on $1 \oplus m$, denoted by A_m^{tw} . It has the same multiplication 1-morphism and unit 1-morphism as those of A_m . The 2-associator has only one nontrivial component:

$$\begin{array}{ccc} \begin{array}{c} m \quad m \quad m \\ \diagdown \quad \diagup \quad | \\ \diagup \quad \diagdown \quad | \\ | \end{array} & \xRightarrow{\alpha_{m,m,m} = -1} & \begin{array}{c} m \quad m \quad m \\ | \quad \diagdown \quad \diagup \\ \diagup \quad \diagdown \quad | \\ | \end{array} \end{array}$$

The 2-commutator β is trivial on all components except $\beta_{m,m} = \pm i$. Two different choices of the 2-commutator define two commutative algebra structures, but they are equivalent to each other. Condensing A_m^{tw} leads to a ‘twisted’ smooth boundary. The topological defects on this boundary still form $2\text{Rep}(\mathbb{Z}_2)$, but the bulk-to-boundary map is different. In the light of Theorem^{ph} 4.2.25, this Lagrangian E_2 -algebra is the one associated to the non-trivial minimal modular extension of $\text{Rep}(\mathbb{Z}_2)$ given by the double semion MTC.

Recall that there are infinitely many condensable E_2 -algebras in $\Omega\mathcal{T}_{\mathbb{Z}_2}^4 = \mathfrak{Z}_1(2\mathrm{Rep}(\mathbb{Z}_2)) \simeq \mathrm{Mod}_{\mathrm{Rep}(\mathbb{Z}_2)}^{E_2}(2\mathrm{Vec})$, each of which is given by a braided fusion 1-category $\mathcal{Q}$ equipped with a braided monoidal functor $\mathrm{Rep}(\mathbb{Z}_2) \rightarrow \mathcal{Q}$. Moreover, such a condensable E_2 -algebra is Lagrangian if and only if $\mathcal{Q}$ is non-degenerate. By [DMNO13], this braided functor is either (1) a braided embedding $\mathrm{Rep}(\mathbb{Z}_2) \hookrightarrow \mathcal{Q}$ or (2) factors as $\mathrm{Rep}(\mathbb{Z}_2) \rightarrow \mathrm{Vec} \hookrightarrow \mathcal{Q}$. The case (1) covers A_m and A_m^{tw} ; and the case (2) are all of the form $A_e \boxtimes \mathcal{Q} \in \mathfrak{Z}_1(2\mathrm{Rep}(\mathbb{Z}_2)) \boxtimes 2\mathrm{Vec} \simeq \mathfrak{Z}_1(2\mathrm{Rep}(\mathbb{Z}_2))$. $\heartsuit$

Example 4.2.40. Consider 3+1D finite gauge theory GT_G^4 . The 2-category of topological defects of codimension 2 and higher is $\mathfrak{Z}_1(2\text{Vec}_G)$ [KTZ20b]. For a subgroup $H \leq G$, the forgetful functor $f : \text{Rep}(G) \rightarrow \text{Rep}(H)$ defines a condensable E_2 -algebra in $2\text{Rep}(G)$ thus in $\mathfrak{Z}_1(2\text{Vec}_G)$. By condensing the algebra $\text{Rep}(H)$ in $\mathfrak{Z}_1(2\text{Vec}_G)$, we obtain GT_H^4 as the condensed phase, i.e.,

$$\mathrm{Mod}_{\mathrm{Rep}(H)}^{\mathrm{E}_2}(\mathfrak{Z}_1(2\mathrm{Vec}_G)) \simeq \mathfrak{Z}_1(2\mathrm{Vec}_H).$$

The proof of this fact is given in Example 5.2.23.

Example 4.2.41. By [DX24], the condensable E_2 -algebras in $\mathfrak{Z}_1(2\mathrm{Vec}_G)$ are G -crossed braided multi-fusion 1-categories. A G -crossed braided multi-fusion 1-category [Tur00, Kir01, Müg04, DGNO10] consists of

- a G -graded multi-fusion 1-category $\mathcal{B} = \bigoplus_{g \in G} \mathcal{B}_g$;
- a monoidal G -action on $\mathcal{B}$ such that $g \odot \mathcal{B}_h \subseteq \mathcal{B}_{ghg^{-1}}$;
- a G -crossed braiding: $c_{x,y} : x \otimes y \rightarrow (g \odot y) \otimes x$ for $x \in \mathcal{C}_g, y \in \mathcal{B}$.

These data also satisfy some coherence condition. The half-braiding of $\mathcal{B} \in \mathfrak{Z}_1(2\mathrm{Vec}_G)$ is induced from the G -action. In particular, the trivial component $\mathcal{B}_e$ is a braided multi-fusion 1-category equipped with a braided

monoidal G -action, that is, a G -equivariant braided multi-fusion category. Similar to Example 3.3.47, a G -crossed braided multi-fusion 1-category is a simple condensable E_2 -algebra in $\mathfrak{Z}_1(2\text{Vec}_G)$ if and only if G acts on the simple direct summands of the tensor unit transitively. It follows that the simple condensable E_2 -algebras in $\mathfrak{Z}_1(2\text{Vec}_G)$ are classified by pairs $(H, \mathcal{B}_0)$, where $H \leq G$ is a subgroup defined up to conjugations and $\mathcal{B}_0$ is an H -crossed braided fusion category. Given such a pair $(H, \mathcal{B}_0)$, the corresponding E_2 -algebra is the induced category $\mathcal{B} := \text{Ind}_H^G \mathbb{C}_0 := \text{Fun}_H(G, \mathcal{B}_0)$, and it is Lagrangian if and only if the H -grading on $\mathcal{B}_0$ is faithful and the trivial component of $\mathcal{B}_0$ is a nondegenerate braided fusion 1-category [Xu24b, Wen24]. We list some examples here.

1. The category $\text{Fun}(G, \text{Vec})$ with the pointwise tensor product and left translation G -action is a braided multi-fusion category equipped with a G -action, which can be viewed as a G -crossed braided fusion category with only the trivial component. It is a Lagrangian algebra in $\mathfrak{Z}_1(\text{Vec}_G)$ and $\text{RMod}_{\text{Fun}(G, \text{Vec})}(\mathfrak{Z}_1(2\text{Vec}_G)) \simeq \text{Vec}_G$. When $G = \mathbb{Z}_2$, this is the Lagrangian algebra $\mathbb{1}_c$ in the 3+1D toric code model (see Example 4.2.39).
2. More generally, for every subgroup $H \leq G$, the category $\text{Fun}(G/H, \text{Vec})$ is a condensable E_2 -algebra in $\mathfrak{Z}_1(2\text{Vec}_G)$ and $\text{Mod}_{\text{Fun}(G/H, \text{Vec})}^{E_2}(\mathfrak{Z}_1(2\text{Vec}_G)) \simeq \mathfrak{Z}_1(2\text{Vec}_H)$.
3. For every 3-cocycle $\omega \in Z^3(G, U(1))$, there is a (unique up to unique isomorphism) G -crossed braided fusion 1-category structure on Vec_G^ω with the obvious G -grading and the conjugation G -action [DGNO10]. Then Vec_G^ω is a Lagrangian algebra in $\mathfrak{Z}_1(\text{Vec}_G)$ and $\text{RMod}_{\text{Vec}_G^\omega}(\mathfrak{Z}_1(2\text{Vec}_G)) \simeq 2\text{Rep}(G)$. When $G = \mathbb{Z}_2$, we have $H^3(\mathbb{Z}_2, U(1)) \simeq \mathbb{Z}_2$ and the two Lagrangian algebras $\text{Vec}_{\mathbb{Z}_2}^\omega$ are the two Lagrangian algebra structure on $\mathbb{1} \oplus m$ in the 3+1D toric code model (see Example 4.2.39).
4. In general, for a faithfully graded nondegenerate G -crossed braided fusion category $\mathcal{B}$, we have $\text{RMod}_{\mathcal{B}}(\mathfrak{Z}_1(2\text{Vec}_G)) \simeq \Sigma(\mathcal{B}_e)^G$ [Wen24, Wen25].
5. For every subgroup $H \leq G$ and 3-cocycle $\omega \in Z^3(H, U(1))$, the induced category $\text{Vec}_H^\omega := \text{Fun}_H(G, \text{Vec}_H^\omega)$ is a Lagrangian algebra in $\mathfrak{Z}_1(2\text{Vec}_G)$. ♥

Example 4.2.42. There are also two coordinate systems on $\mathfrak{Z}_1(2\text{Rep}(G))$.

- The condensable E_2 -algebras in $\mathfrak{Z}_1(2\text{Rep}(G)) \simeq \mathfrak{Z}_1(\text{LMod}_{\text{Vec}_G}(2\text{Vec}))$ are also G -crossed braided multi-fusion categories, on which the half-braidings are equivalent to the G -gradings.
- By [JFR23, DX24], the condensable E_2 -algebras in $\mathfrak{Z}_1(\Sigma\text{Rep}(G)) \simeq \text{Mod}_{\text{Rep}(G)}^{E_2}(2\text{Vec})$ are braided multi-fusion categories $\mathcal{M}$ equipped with braided monoidal functor $\text{Rep}(G) \rightarrow \mathcal{M}$, which are simple if and only if $\mathcal{M}$ is a braided fusion category. By [DMNO13, Lemma 3.5 and Remark 3.23], such a braided functor factors through $\text{Rep}(G) \rightarrow \text{Rep}(H) \rightarrow \mathcal{M}$ for some subgroup $H \leq G$ such that $\text{Rep}(H) \rightarrow \mathcal{M}$ is a braided embedding.

These two coordinate systems are related by the equivariantization and de-equivariantization. In particular, there is a one-to-one correspondence between G -crossed braided fusion categories and braided fusion categories containing $\text{Rep}(G)$ [DGNO10, Theorem 4.44].

Let $\text{Rep}(G) \rightarrow \mathcal{M}$ be a simple condensable E_2 -algebra in $\mathfrak{Z}_1(\Sigma\text{Rep}(G))$. We also denote it by $\mathcal{M}$ for simplicity. According to [DX24] (see also Theorem^{ph} 4.2.25), by condensing $\mathcal{M}$ as 2-codimensional topological defect in GT_G^4 , we obtain a condensed phase D^4 and a gapped domain wall M^3 such that

$$\Omega\mathcal{D} \simeq \text{Mod}_{\mathcal{M}}^{E_2}(\text{Mod}_{\text{Rep}(G)}^{E_2}(2\text{Vec})) \simeq \text{Mod}_{\mathcal{M}}^{E_2}(2\text{Vec}) \simeq \mathfrak{Z}_1(\Sigma\mathcal{M}), \quad \Omega_m\mathcal{M} \simeq \text{RMod}_{\mathcal{M}}(\text{Mod}_{\text{Rep}(G)}^{E_2}(2\text{Vec})).$$

It follows that the Lagrangian E_2 -algebras in $\mathfrak{Z}_1(\Sigma\text{Rep}(G))$ are nondegenerate braided fusion categories $\mathcal{M}$ equipped with braided functors $\text{Rep}(G) \rightarrow \mathcal{M}$. We list some examples here.

1. For every $\omega \in Z^3(G, U(1))$, there is a braided embedding $\text{Rep}(G) \rightarrow \mathfrak{Z}_1(\text{Vec}_G^\omega)$, which defines a Lagrangian E_2 -algebra in $\mathfrak{Z}_1(\Sigma\text{Rep}(G))$. As a consequence, the condensed phase $D^4 = \mathbb{1}^4$ is trivial and the gapped domain wall is such that

$$\Omega_m\mathcal{M} \simeq \text{RMod}_{\mathfrak{Z}_1(\text{Vec}_G^\omega)}(\mathfrak{Z}_1(\Sigma\text{Rep}(G))) \simeq 2\text{Rep}(G).$$

When $G = \mathbb{Z}_2$, we have $H^3(\mathbb{Z}_2, U(1)) \simeq \mathbb{Z}_2$ and the two minimal nondegenerate extensions correspond to the two different Lagrangian E_2 -algebra structures on $\mathbb{1} \oplus m$ in the 3+1D toric code model (see Example 4.2.39).

2. When $\mathcal{M}$ is a nondegenerate extension of $\text{Rep}(G)$, the 2-category $\text{RMod}_{\mathcal{M}}(\mathfrak{Z}_1(\Sigma\text{Rep}(G)))$ is connected and equivalent to $\Sigma\mathfrak{Z}_2(\text{Rep}(G) \rightarrow \mathcal{M})$.
3. More generally, for every subgroup $H \leq G$ and $\omega \in Z^3(H, U(1))$, the composite functor $\text{Rep}(G) \rightarrow \text{Rep}(H) \rightarrow \mathfrak{Z}_1(\text{Vec}_H^\omega)$ is a Lagrangian algebra in $\mathfrak{Z}_1(\Sigma\text{Rep}(G))$.
4. In particular, when H is the trivial subgroup, the forgetful functor $\text{Rep}(G) \rightarrow \text{Vec}$ is a Lagrangian algebra in $\mathfrak{Z}_1(\Sigma\text{Rep}(G))$, and $\text{RMod}_{\mathcal{M}}(\mathfrak{Z}_1(\Sigma\text{Rep}(G)))$ is equivalent to 2Vec_G . When $G = \mathbb{Z}_2$, this is the Lagrangian algebra 1_c in the 3+1D toric code model (see Example 4.2.39). $\heartsuit$

Example 4.2.43 ([Xu24b, Wen24]). Let $\pi \in Z^4(G, U(1))$. For the 3+1D π -twisted G gauge theory, the condensable E_2 -algebras in $\mathfrak{Z}_1(2\text{Vec}_G^\pi)$ are precisely π -twisted G -crossed braided multi-fusion 1-categories $\mathcal{A}$ [Xu24b, Lemma 5.8]. The simple condensable E_2 -algebras are given in Theorem 5.11, and Lagrangian E_2 -algebras are given in Corollary 5.12 in [Xu24b]. When $\mathcal{A}$ is a braided fusion 1-category with a G -action, then the 2-category of right $\mathcal{A}$ -modules in $\mathfrak{Z}_1(2\text{Vec}_G^\pi)$ was computed in Theorem 5.17 in [Xu24b], and the 2-category of E_2 - $\mathcal{A}$ -modules (or local $\mathcal{A}$ -modules) was computed in Theorem 5.20 in [Xu24b]. $\heartsuit$

4.2.5 Examples in higher dimensions

Example 4.2.44. Consider finite gauge theory GT_G^{n+2} for a finite group G . We have $\Omega\mathcal{GT}_G^{n+2}$ has two coordinate systems:

$$\Omega\mathcal{GT}_G^{n+2} \simeq \mathfrak{Z}_1((n+1)\text{Vec}_G), \quad \Omega\mathcal{GT}_G^{n+2} \simeq \mathfrak{Z}_1((n+1)\text{Rep}(G)) \simeq \text{Mod}_{n\text{Rep}(G)}^{E_2}((n+1)\text{Vec}),$$

where the last $\simeq$ is due to (4.2.13). We discuss condensable E_1 -algebras in $\Omega\mathcal{GT}_G^{n+2}$ in both coordinates.

- (1) A condensable E_2 -algebras in $\mathfrak{Z}_1((n+1)\text{Vec}_G)$ is precisely a G -crossed braided multi-fusion n -category. Inspired by the definition of a G -crossed braided multi-fusion 1-category [Tur00, Kir01, Müg04, DGN010], we expect a G -crossed braided multi-fusion n -category to consist of

- a G -graded multi-fusion 1-category $\mathcal{A} = \bigoplus_{g \in G} \mathcal{A}_g$;
- a monoidal G -action on $\mathcal{A}$ such that $g \odot \mathcal{A}_h \subseteq \mathcal{A}_{ghg^{-1}}$;
- a G -crossed braiding: $c_{x,y} : x \otimes y \rightarrow (g \odot y) \otimes x$ for $x \in \mathcal{C}_g, y \in \mathcal{A}$.

satisfying some (higher) coherence conditions. Higher coherence properties are always difficult in high category theory. We try to avoid higher coherence in this work. Instead, we give a convenient or almost physical way to define this notion. A G -crossed braided fusion n -category $\mathcal{A}$ can be understood as a monoidal $(n+1)$ -category $B_G\mathcal{A}$ consists of $|G|$ objects labeled by the elements in G with the monoidal structure defined by the multiplication of G and $\text{hom}_{B_G\mathcal{A}}(g, h) = \mathcal{A}_{hg^{-1}}$ (recall Example 3.3.46). When $n = 1$, this construction can be found in [Cui19, DR18, JPR22]. We can understand this construction physically. If we view the condensation completion of $B_G\mathcal{A}$, denoted by $\Sigma_G\mathcal{A}$, as the category of topological defects of an anomalous $n+1$ D topological order X^{n+1} , then the object $g \in G = \text{Ob}(B_G\mathcal{A})$ can be viewed as an invertible 1-codimensional defect in X^{n+1} , and $\text{hom}_{B_G\mathcal{A}}(g, h)$ can be viewed as the n -category of 2-codimensional defects living on the domain wall between two 1-codimensional defect $g, h \in G$.

Now we show that $\Sigma_G\mathcal{A} \simeq \text{RMod}_{\mathcal{A}}((n+1)\text{Vec}_G)$ as $(n+1)\text{Vec}_G$ -modules. Denote the simple objects in $(n+1)\text{Vec}_G$ by $\{\delta_g\}_{g \in G}$. There is a $(n+1)$ -functor $F : B_G\mathcal{A} \rightarrow \text{RMod}_{\mathcal{A}}((n+1)\text{Vec}_G)$ defined by

- an object $g \in G$ is mapped to the free right $\mathcal{A}$ -module $\delta_{g^{-1}} \otimes \mathcal{A}$;
- the functors between hom n -categories are defined by the canonical equivalences:

$$\text{Hom}_{B_G\mathcal{A}}(g, h) = \mathcal{A}_{hg^{-1}} \simeq \text{Hom}_{2\text{Vec}_G}(\delta_{g^{-1}}, \delta_{h^{-1}} \otimes \mathcal{A}) \simeq \text{Hom}_{\text{RMod}_{\mathcal{A}}(2\text{Vec}_G)}(\delta_{g^{-1}} \otimes \mathcal{A}, \delta_{h^{-1}} \otimes \mathcal{A}).$$

Moreover, this is a G -module $(n+1)$ -functor. Thus it induces a G -module $(n+1)$ -functor $\Sigma_G\mathcal{A} \rightarrow \text{RMod}_{\mathcal{A}}((n+1)\text{Vec}_G)$. Note that every simple module $x \in \text{RMod}_{\mathcal{A}}((n+1)\text{Vec}_G)$ can be condensed from the free module $x \otimes \mathcal{A}$. Thus the G -module $(n+1)$ -functor $\Sigma_G\mathcal{A} \rightarrow \text{RMod}_{\mathcal{A}}((n+1)\text{Vec}_G)$ is an equivalence.

- (2) A condensable E_2 -algebra in $\mathfrak{Z}_1((n+1)\text{Rep}(G))$ is precisely given by a braided monoidal functor $n\text{Rep}(G) \rightarrow \mathcal{B}$ for a braided multi-fusion n -category $\mathcal{B}$. It is simple if $\mathcal{B}$ is fusion.

These two ways of defining condensable E_2 -algebras in $\Omega\mathcal{GT}_G^{n+2}$ are related again by equivariantization $\mathcal{A} \mapsto \mathcal{A}^G$ and de-equivariantization $\mathcal{B} \mapsto \mathcal{B}_G$ (recall Example 3.3.56). In particular, $\mathcal{B}_G := \text{RMod}_{[\mathbb{1}, \mathbb{1}]}(\mathcal{B})$, where $\mathbb{1}$ is the tensor unit of $n\text{Vec}$ and $[\mathbb{1}, \mathbb{1}]$ is the internal hom algebra in $n\text{Rep}(G)$ for the $n\text{Rep}(G)$ -module $n\text{Vec}$. In other words, the notion of a G -crossed braided fusion n -category can also be understood as the de-equivariantization of a braided monoidal functor $n\text{Rep}(G) \rightarrow \mathcal{B}$.

Recall Theorem^{ph} 4.2.25, by condensing the condensable E_2 -algebra $\mathcal{B}$ in $\mathfrak{Z}_1((n+1)\text{Rep}(G))$, we obtain a condensed phase $\mathcal{D}^{n+2}$ such that $\Omega\mathcal{D} \simeq \mathfrak{Z}_1(\Sigma\mathcal{B})$. ♡

5 Condensations of higher codimensional topological defects

In this section, we give the general theory of the condensations of k -codimensional topological defects in an $n+1$ D topological order. We first study the relation between particle condensations and string condensations in 2+1D, then we generalize it to higher dimensions.

5.1 Particle condensations in 2+1D via a two-step process


In 2+1D, 1-codimensional defects are strings and 2-codimensional defects are particles. In Section 3.3 and 3.3.3, we have studied the string condensation in 2+1D; in Section 4.1, we have studied the particle (or anyon) condensation in 2+1D. In this subsection, we study their relation, which can be generalized to higher dimensions.

5.1.1 General theory

Consider a 2+1D anomaly-free simple topological orders $\mathcal{C}^3$. In this case, there is a natural coordinate system of $\mathcal{C}$, i.e., $\mathcal{C} = \Sigma\Omega\mathcal{C} = \text{RMod}_{\Omega\mathcal{C}}(2\text{Vec})$.

Consider a particle condensation in $\mathcal{C}^3$ defined by a simple condensable E_2 -algebra A in $\Omega\mathcal{C}$. It produces a condensed 2+1D topological order $\mathcal{D}^3$ and a gapped domain wall $M^2 = (\mathcal{M}, m)$ such that $\Omega\mathcal{D} \simeq \text{Mod}_A^{E_2}(\Omega\mathcal{C})$ and $\Omega_m\mathcal{M} = \text{RMod}_A(\Omega\mathcal{C})$. Now we show that the same phase transition can be realized in a different way, which splits into two steps.

1. In the first step, we condense the particle A along a line in space as illustrated below. It produces a condensed string denoted by ΣA .



In this step, one can forget about the E_2 -algebra structure on A but only remember its E_1 -algebra structure in the x^2 -direction. Recall Section 3.1 [Kon14], condensing A on a line produces a condensed string, particles on which form the fusion 1-category $\text{Mod}_A^{E_1}(\mathcal{C})$, i.e.

$$\text{hom}_{\mathcal{C}}(\Sigma A, \Sigma A) \simeq \text{Mod}_A^{E_1}(\mathcal{C}).$$

However, this result does not tell us what ΣA is as an object in $\mathcal{C}$. Now we show that ΣA can be identified with $\text{RMod}_A(\Omega\mathcal{C})$ as an object in $\mathcal{C} = \text{RMod}_{\Omega\mathcal{C}}(2\text{Vec})$.

- (a) First, we have already shown in Theorem^{ph} 3.3.26 that $\text{RMod}_A(\Omega\mathcal{C}) \in \text{Alg}_{E_1}^{\mathcal{C}}(\text{RMod}_{\Omega\mathcal{C}}(2\text{Vec})) \simeq \text{LMod}_{\Omega\mathcal{C}}(\text{Alg}_{E_1}^{\mathcal{C}}(2\text{Vec}))$ because the functor $- \otimes A : \Omega\mathcal{C} \rightarrow \text{RMod}_A(\Omega\mathcal{C})$ is a central functor.
- (b) It remains to show that ΣA can be identified with $\text{RMod}_A(\Omega\mathcal{C})$ as an object in $\mathcal{C} = \text{RMod}_{\Omega\mathcal{C}}(2\text{Vec})$. One can see this directly from the following picture⁵⁷:

$$\begin{array}{ccc}
 \text{RMod}_A(\Omega\mathcal{C}) \boxtimes_{\Omega\mathcal{D}} \text{RMod}_A(\Omega\mathcal{C})^{\text{op}} \simeq \text{Mod}_A^{E_1}(\Omega\mathcal{C}) & & \Sigma A \\
 \begin{array}{c} \text{RMod}_A(\Omega\mathcal{C}) \quad \Omega\mathcal{D} \quad \Omega\mathcal{C} \\ \text{RMod}_A(\Omega\mathcal{C}) \end{array} & \rightsquigarrow & \begin{array}{c} \Omega\mathcal{C} \\ \Sigma A \end{array}
 \end{array}
 \quad (5.1.1)$$

$\text{hom}_{\mathcal{C}}(\Sigma A, \Sigma A) \simeq \text{Mod}_A^{E_1}(\Omega\mathcal{C})$
 $\Sigma A = \text{RMod}_A(\Omega\mathcal{C}) = \text{hom}_{\mathcal{C}}(\mathbb{1}_{\mathcal{C}}, \Sigma A)$

where the green region shrink horizontally to the condensed string ΣA , and the wall between two colored region has particles which form a fusion 1-category given by $\text{RMod}_A(\Omega\mathcal{C})$. We recover the multi-fusion 1-category of particles on the condensed string by the following equivalences:

$$\text{RMod}_A(\Omega\mathcal{C}) \boxtimes_{\Omega\mathcal{D}} \text{RMod}_A(\Omega\mathcal{C})^{\text{op}} \simeq \text{RMod}_A(\Omega\mathcal{C}) \boxtimes_{\text{Mod}_A^{E_2}(\Omega\mathcal{C})} \text{LMod}_A(\Omega\mathcal{C}) \simeq \text{Mod}_A^{E_1}(\Omega\mathcal{C}),$$

⁵⁷In the second picture, the reason why both the blue line and the blue dot are labeled by ΣA was explained in Remark 2.2.33.

where we have used the natural equivalence $\text{RMod}_A(\Omega\mathcal{C}) \simeq \text{LMod}_A(\Omega\mathcal{C})^{\text{op}}$ defined by $x \mapsto x^R = x^*$ in the first ‘ $\simeq$ ’ and the second ‘ $\simeq$ ’ is a mathematical fact (see for example [KZ18a]). Remember $\text{hom}_{\mathcal{C}}(x, y), \forall x, y \in \mathcal{C}$ are physical observables in spacetime, but the label of a string can not be observed directly in spacetime. However, one can identify an object $x \in \mathcal{C}$ in the coordinate system $\mathcal{C} = \text{RMod}_{\Omega\mathcal{C}}(2\text{Vec})$ with $\text{hom}_{\mathcal{C}}(\Omega\mathcal{C}, x)$ because $\text{hom}_{\text{RMod}_{\Omega\mathcal{C}}(2\text{Vec})}(\Omega\mathcal{C}, x) \simeq x$ in $\text{RMod}_{\Omega\mathcal{C}}(2\text{Vec})$. From the physical configuration depicted in (5.1.1), we immediately obtain

$$\Sigma A \simeq \text{hom}_{\mathcal{C}}(\Omega\mathcal{C}, \Sigma A) \simeq \text{RMod}_A(\Omega\mathcal{C}) \in \text{RMod}_{\Omega\mathcal{C}}(2\text{Vec}) = \mathcal{C},$$

because $\text{hom}_{\mathcal{C}}(\Omega\mathcal{C}, \Sigma A)$ is precisely the observables living on the domain wall between the trivial 1-codimensional topological defect $\mathbb{1}_{\mathcal{C}} = \Omega\mathcal{C}$ and ΣA .

From now on, for simplicity, we set $\Sigma A := \text{RMod}_A(\Omega\mathcal{C}) \in \text{RMod}_{\Omega\mathcal{C}}(2\text{Vec}) = \mathcal{C}$ and, at the same time, ΣA is also the abstract label of the condensed defect if we do not choose any coordinate system.

2. In the second step, we condense ΣA . We have already shown in Theorem^{ph} 3.3.26 that ΣA is a condensable E_1 -algebra in $\mathcal{C}$. This fact can also be viewed in another way. Since the ΣA -string can be obtained by shrinking the green region horizontally as depicted in (5.1.1), by our theory of the condensations of 1-codimensional topological defects (Theorem^{ph} 3.3.2 and 3.3.9), the ΣA -string has a canonical structure of a condensable E_1 -algebra in $\mathcal{C}$. The algebraic structure on ΣA is defined in (3.3.1). In this argument, the E_2 -algebra structure on A is implicitly used. Moreover, this algebraic structure on ΣA coincides with the monoidal structure on $\text{RMod}_A(\Omega\mathcal{C})$ (recall (3.3.16)). Therefore, we can further condensing the ΣA -string thus produces a new phase. It is physically obvious that this new phase is precisely D^3 . In other words, we obtain

$$\mathcal{D} \simeq \text{Mod}_{\Sigma A}^{E_1}(\mathcal{C}). \quad (5.1.2)$$

It immediately implies the following equivalences:

$$\text{Mod}_{\Sigma A}^{E_1}(\mathcal{C}) \simeq \Sigma \text{Mod}_A^{E_2}(\Omega\mathcal{C}), \quad (5.1.3)$$

$$\Omega \text{Mod}_{\Sigma A}^{E_1}(\mathcal{C}) \simeq \text{Mod}_A^{E_2}(\Omega\mathcal{C}). \quad (5.1.4)$$

where the first equivalence is a monoidal equivalence and the second one is a braided monoidal equivalence. Note that (5.1.4) is an immediate consequence of (5.1.3) and is proved directly in Theorem 5.1.4. The gapped domain wall M^2 between C^3 and D^3 can be described by the pair

$$(\text{RMod}_{\Sigma A}(\mathcal{C}), \Sigma A).$$

The multi-fusion 1-category of particles on this domain wall is given by

$$\text{hom}_{\text{RMod}_{\Sigma A}(\mathcal{C})}(\Sigma A, \Sigma A) \simeq \text{Fun}_{\text{RMod}_A(\Omega\mathcal{C})^{\text{op}}}(\text{RMod}_A(\Omega\mathcal{C}), \text{RMod}_A(\Omega\mathcal{C})) \simeq \text{RMod}_A(\Omega\mathcal{C}),$$

where the first ‘ $\simeq$ ’ is due to the fact that a functor intertwining the ΣA -action automatically intertwines the $\Omega\mathcal{C}$ -action, which is defined via the monoidal functor $- \otimes A : \Omega\mathcal{C} \rightarrow \text{RMod}_A(\Omega\mathcal{C})$.

Note that the monoidal equivalences in (5.1.3) summarize precisely the relation between the direct condensation of a 2-codimensional topological defect A and that of the 1-codimensional topological defect ΣA .

All of above arguments automatically generalize to higher codimensional defects except a key point, which deserves to be reinvestigated. We see the key point here is to argue that ΣA is automatically a condensable E_1 -algebra in $\mathcal{C}$ if A is a condensable E_2 -algebra in $\Omega\mathcal{C}$. We have used the physical or geometric intuition established in Theorem^{ph} 3.3.2 and 3.3.9 (in particular, (3.3.1)). Unfortunately, this physical or geometric intuition stops to make sense for higher codimensional topological defects because k -codimensional topological defects for $k > 2$ do not give a complete mathematical characterization of a topological order. Whenever we want to apply the geometric intuition of a topological order C^{n+1} we should automatically include all topological defects of codimension 1 or higher if C^{n+1} is anomalous; or include all topological defect of codimension (at least) 2 or higher if C^{n+1} is anomaly-free. Therefore, in order to generalize this key point to higher codimensional defects, we need more fundamental arguments that are ready to be generalized.


Now we explain mathematically why ΣA is a condensable E_1 -algebra in $\mathcal{C}$. First, recall that $\otimes^2$ represents the fusion product in the vertical direction (see Figure 6 and 13), and $(\Omega\mathcal{C}, \otimes^2)$ is monoidal. Therefore, it

makes sense to talk about the 1-category $\text{Alg}_{E_1}(\Omega\mathcal{C})$ based on the monoidal structure $(\Omega\mathcal{C}, \otimes^2)$. It turns out that the second tensor product $\otimes^1$ on $\Omega\mathcal{C}$ (i.e., the horizontal fusion) endow both triples

$$(\text{Alg}_{E_1}(\Omega\mathcal{C}), \otimes^1, 1_{\mathbb{1}_{\mathcal{C}}}) \quad \text{and} \quad (\text{RMod}_{\Omega\mathcal{C}}(2\text{Vec}), \boxtimes_{\Omega\mathcal{C}}, \Omega\mathcal{C})$$

with the structure of a monoidal 1-category and a monoidal 2-category, respectively. We explain this fact below.

1. The tensor product in $\text{Alg}_{E_1}(\Omega\mathcal{C})$ is given by the tensor product $\otimes^1$ in $\Omega\mathcal{C}$. For $A, B \in \text{Alg}_{E_1}(\Omega\mathcal{C})$, $A \otimes^1 B$ is a well-defined E_1 -algebra in $\Omega\mathcal{C}$ with the algebraic structure defined as follows:



$$(A \otimes^1 B) \otimes^2 (A \otimes^1 B) \xrightarrow{\delta_{AA,B,B}} (A \otimes^2 A) \otimes^1 (B \otimes^2 B) \xrightarrow{\mu_A \mu_B} A \otimes^1 B.$$

2. A right $\Omega\mathcal{C}$ -module X with a right $\Omega\mathcal{C}$ -action $\odot : X \times \Omega\mathcal{C} \rightarrow X$ is automatically a left $\Omega\mathcal{C}$ -module with the left $\Omega\mathcal{C}$ -action defined by $a \odot x := x \odot a$ for $a \in \Omega\mathcal{C}, x \in X$. This is a well-defined left $\Omega\mathcal{C}$ -module due to the existence of the braiding in $\Omega\mathcal{C}$ as shown below:

$$a \odot (b \odot x) := (x \odot b) \odot a \simeq x \odot (b \otimes^1 a) \xrightarrow{1_x c_{b,a}^1} x \odot (a \otimes^1 b) = (a \otimes^1 b) \odot x.$$

where the braiding $c_{b,a}^1$ is defined in (2.3.6).

Note that there is a functor

$$\Sigma(-) = \text{RMod}_-(\Omega\mathcal{C}) : \text{Alg}_{E_1}(\Omega\mathcal{C}) \rightarrow \text{RMod}_{\Omega\mathcal{C}}(2\text{Vec}) = \mathcal{C} \quad (5.1.5)$$


defined by

$$\begin{array}{ccc} A & \longmapsto & \text{RMod}_A(\Omega\mathcal{C}) \\ f \downarrow & & \downarrow -\otimes_A f B \\ B & \longmapsto & \text{RMod}_B(\Omega\mathcal{C}) \end{array}$$

where $f B$ is the A - B -bimodule with the right A -module structure induced from the algebra map $f : A \rightarrow B$. One can show that $\text{RMod}_-(\Omega\mathcal{C}) : \text{Alg}_{E_1}(\Omega\mathcal{C}) \rightarrow \text{RMod}_{\Omega\mathcal{C}}(2\text{Vec})$ is a lax-monoidal functor with two defining 1-morphisms:

$$1_{\mathbb{1}_{\mathcal{C}}} \mapsto \text{RMod}_{1_{\mathbb{1}_{\mathcal{C}}}}(\Omega\mathcal{C}) \simeq \Omega\mathcal{C}; \quad \text{RMod}_A(\Omega\mathcal{C}) \boxtimes_{\Omega\mathcal{C}} \text{RMod}_B(\Omega\mathcal{C}) \xrightarrow{\tilde{F}} \text{RMod}_{A \otimes^1 B}(\Omega\mathcal{C}).$$

We give a construction of $\tilde{F}$. The functor $\text{RMod}_A(\Omega\mathcal{C}) \boxtimes \text{RMod}_B(\Omega\mathcal{C}) \xrightarrow{F} \text{RMod}_{A \otimes B}(\Omega\mathcal{C})$ defined by $m \boxtimes n \rightarrow m \otimes^1 n$, where the $A \otimes^1 B$ -action on $m \otimes^1 n$ is defined as follows:



$$(m \otimes^1 n) \otimes^2 (A \otimes^1 B) \xrightarrow{\delta_{m,n,A,B}} (m \otimes^2 A) \otimes^1 (n \otimes^2 B) \xrightarrow{\mu_m \mu_n} m \otimes^1 n.$$

From above picture, it is clear that this functor F is $\Omega\mathcal{C}$ -balanced, i.e., $(m \otimes^1 a) \otimes^1 n \simeq m \otimes^1 (a \otimes^1 n)$. Therefore, it induces a functor $\tilde{F} : \text{RMod}_A(\Omega\mathcal{C}) \boxtimes_{\Omega\mathcal{C}} \text{RMod}_B(\Omega\mathcal{C}) \rightarrow \text{RMod}_{A \otimes B}(\Omega\mathcal{C})$. This functor $\tilde{F}$ endows the functor $\Sigma(-) : \text{Alg}_{E_1}(\Omega\mathcal{C}) \rightarrow \text{RMod}_{\Omega\mathcal{C}}(2\text{Vec})$ with the structure of a lax-monoidal functor.

Since a lax-monoidal functor maps E_1 -algebras to E_1 -algebras, it means that the functor $\text{RMod}_-(\Omega\mathcal{C})$ can be lifted to a functor

$$\begin{aligned} \text{RMod}_-(\Omega\mathcal{C}) : \text{Alg}_{E_2}(\Omega\mathcal{C}) &= \text{Alg}_{E_1}(\text{Alg}_{E_1}(\Omega\mathcal{C})) \rightarrow \text{Alg}_{E_1}(\text{RMod}_{\Omega\mathcal{C}}(2\text{Vec})), \\ A &\mapsto \Sigma A = \text{RMod}_A(\Omega\mathcal{C}). \end{aligned} \quad (5.1.6)$$

Since the separability is defined by 1-morphisms and identities in $\text{Alg}_{E_1}(\Omega\mathcal{C})$, they are mapped to 1-morphisms and identities in $\text{RMod}_{\Omega\mathcal{C}}(2\text{Vec})$. As a consequence, we obtain a restriction of the functor $\text{RMod}_-(\Omega\mathcal{C})$ on the subcategory $\text{Alg}_{E_2}^c(\Omega\mathcal{C})$:

$$\begin{aligned}\Sigma(-) &= \text{RMod}_-(\Omega\mathcal{C}) : \text{Alg}_{E_2}^c(\Omega\mathcal{C}) \rightarrow \text{Alg}_{E_1}^c(\text{RMod}_{\Omega\mathcal{C}}(2\text{Vec})) = \text{Alg}_{E_1}^c(\mathcal{C}). \\ A &\mapsto \Sigma A = \text{RMod}_A(\Omega\mathcal{C}).\end{aligned}\tag{5.1.7}$$

We summarize above result as a mathematical theorem.

Proposition 5.1.1. For a braided fusion 1-category $\mathcal{B}$, there is a well-defined functor

$$\Sigma(-) : \text{Alg}_{E_2}^c(\mathcal{B}) \rightarrow \text{Alg}_{E_1}^c(\Sigma\mathcal{B}) \quad \text{defined by} \quad A \mapsto \Sigma A = \text{RMod}_A(\mathcal{B}).$$

When $\mathcal{B}$ is non-degenerate, this functor maps Lagrangian E_2 -algebras in $\mathcal{B}$ to Lagrangian E_1 -algebras in $\Sigma\mathcal{B}$ bijectively.

Remark 5.1.2. Proposition 5.1.1 provides a powerful way to construct or classify Lagrangian E_2 -algebras via the classification of Lagrangian E_1 -algebras, which can be obtained from Theorem^{ph} 3.3.32. For example, the classification of the Lagrangian algebras in 2+1D finite gauge theories, originally obtained by Davydov in [Dav10b], can be easily recovered in Example 3.3.44. As we show later, it is a powerful tool to study Lagrangian E_2 -algebras in braided fusion higher categories. $\diamond$


Remark 5.1.3. The functor $\Sigma : \text{Alg}_{E_2}^c(\mathcal{B}) \rightarrow \text{Alg}_{E_1}^c(\Sigma\mathcal{B})$ is not surjective in general. This fact is related to the fact that two Witt equivalent non-degenerate braided fusion categories cannot be obtained from each other via anyon condensations in general; however, they can be obtained from each via condensations of 1-codimensional topological defects. $\diamond$

When $\mathcal{C}^3$ is anomalous, the condensation of a 2-codimensional topological defect $A \in \text{Alg}_{E_2}^c(\Omega\mathcal{C})$ is defined by first condensing it on a line to produce a 1-codimensional topological defect ΣA ; then condensing ΣA . In this case, the equivalence (5.1.3) is not true in general because $\Omega\mathcal{C}$ does not have the complete information of $\mathcal{C}$ and some 1-codimensional topological defects in $\mathcal{C}$ are not condensation descendants of $\Omega\mathcal{C}$. However, the equivalence (5.1.4), i.e., $\Omega \text{Mod}_{\Sigma A}^{E_1}(\mathcal{C}) \simeq \text{Mod}_A^{E_2}(\Omega\mathcal{C})$ remains correct because this equivalence is simply a reformulation of the natural physical intuition that all 2-codimensional topological defects in the condensed (also anomalous) phase D^3 can only come from $\Omega\mathcal{C}$ as deconfined 2-codimensional topological defects, i.e. as E_2 - A -modules in $\Omega\mathcal{C}$. Moreover, $\mathcal{C}^3$ and D^3 should share the same gravitational anomaly, i.e., $Z(\mathcal{C})^4 = Z(D)^4$. We summarize and reformulate above results as mathematical results.

Theorem^o 5.1.4. Let $\mathcal{C}$ be a fusion 2-category, $A \in \text{Alg}_{E_2}^c(\Omega\mathcal{C})$ and $\Sigma A = \text{RMod}_A(\Omega\mathcal{C})$. Then we have $\Sigma A \in \text{Alg}_{E_1}^c(\mathcal{C})$ and a natural braided equivalence:

$$\Omega \text{Mod}_{\Sigma A}^{E_1}(\mathcal{C}) \simeq \text{Mod}_A^{E_2}(\Omega\mathcal{C}).\tag{5.1.8}$$

Proof. We have already obtained this result through physical arguments. Now we sketch a mathematical proof. Note that $\Omega \text{Mod}_{\Sigma A}^{E_1}(\mathcal{C}) \simeq \Omega \text{Mod}_{\Sigma A}^{E_1}(\Sigma\Omega\mathcal{C})$. A 1-morphism $\Sigma A \rightarrow \Sigma A$ in $\text{Mod}_{\Sigma A}^{E_1}(\Sigma\Omega\mathcal{C})$ is a functor from ΣA to ΣA intertwining the right $\Omega\mathcal{C}$ -action and the two-side ΣA -actions in x^1 -dimension. Note that a functor from ΣA to ΣA intertwining the right $\Omega\mathcal{C}$ -action is precisely the functor $-\otimes_A^2 m$ for an A - A -bimodule m in $\Omega\mathcal{C}$, where the superscript ‘2’ represents in the x^2 -dimension. Therefore, a 1-morphism $\Sigma A \rightarrow \Sigma A$ in $\text{Mod}_{\Sigma A}^{E_1}(\Sigma\Omega\mathcal{C})$ is precisely a functor $-\otimes_A^2 m$ intertwining the two-side ΣA -actions in x^1 -direction.



m is equipped with horizontal two-side A -actions induced from the vertical right A -action $m \otimes^2 A \rightarrow m$. These three A -actions on m are illustrated as the three arrows from A to m in the top triangles, which mean that the top two triangles in diagram (4.1.6) (for $x = m$) is commutative. It remains to show that the horizontal A -actions are also compatible with the vertical left A -action $A \otimes^2 m \rightarrow m$.

The condition that $-\otimes_A^2 m$ intertwines the horizontal ΣA -actions says that the horizontal A -actions on m can be equivalently defined by the horizontal A -actions on the A -factor in $A \otimes_A^2 m$. Therefore, the horizontal A -actions are compatible with the vertical left A -action $A \otimes^2 (A \otimes_A^2 m) \rightarrow m$ because A itself is an E_2 - A -module (recall the defining condition of an E_2 - A -module in the commutative diagram (4.1.6)). This compatibility is illustrated in the bottom triangles in the above picture. $\square$

Theorem^o 5.1.5. For a non-degenerate braided fusion 1-category $\mathcal{B}$, $A \in \text{Alg}_{E_2}^c(\mathcal{B})$ and $\Sigma A = \text{RMod}_A(\mathcal{B})$, we have a natural monoidal equivalence:

$$\text{Mod}_{\Sigma A}^{E_1}(\Sigma \mathcal{B}) \simeq \Sigma \text{Mod}_A^{E_2}(\mathcal{B}). \quad (5.1.9)$$

Remark 5.1.6. When A is a condensable E_1 -algebra in a fusion 2-category $\mathcal{C}$, $\Sigma A \in \Sigma' \mathcal{C} := \text{LMod}_{\mathcal{C}}(3\text{Vec})$. We have

$$\Omega_{\Sigma A} \Sigma' \mathcal{C} = \text{Fun}_{\mathcal{C}}(\text{RMod}_A(\mathcal{C}), \text{RMod}_A(\mathcal{C})) \simeq \text{Mod}_A^{E_1}(\mathcal{C}). \quad (5.1.10)$$

When A is a condensable E_2 -algebra in braided fusion 2-category $\mathcal{B}$, $\Sigma A \in \text{Alg}_{E_1}^c(\Sigma \mathcal{B})$. We can repeat the argument to obtain $\Sigma^2 A := \text{RMod}_{\Sigma A}(\Sigma \mathcal{B}) \in \Sigma' \Sigma \mathcal{B}$. Then we obtain

$$\Omega_{\Sigma^2 A}^2(\Sigma' \Sigma \mathcal{B}) = \Omega \text{Mod}_{\Sigma A}^{E_1}(\Sigma \mathcal{B}) \simeq \text{Mod}_A^{E_2}(\mathcal{B}). \quad (5.1.11)$$


The canonical equivalence (5.1.11) provides us a convenient way to understand, compute or even define alternatively the category $\text{Mod}_A^{E_2}(\mathcal{B})$ (recall Section 4.1.1). $\diamond$

Remark 5.1.7. We have assumed the topological order $\mathcal{C}^3$ to be anomaly-free and simple. If $\mathcal{C}^3$ is not anomaly-free, then, in order to define a phase transition by condensing a 2-codimensional topological defect $A \in \Omega \mathcal{C}$, we have to first condense A along a line to create a condensed string ΣA , then condense ΣA along the remaining transversal direction. We discuss such a condensation as a special case of more general condensations in Section 5.2. $\diamond$

5.1.2 Examples


Example 5.1.8. Let us consider the 2+1D $\mathbb{Z}_2$ topological order TC^3 , whose 2-category of topological defects is $\mathcal{TC}$. In this case, $\Omega \mathcal{TC} = \mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$ is a non-degenerate braided fusion 1-category with four simple objects $1, e, m, f$, where $1, e$ generate the sub-fusion category $\text{Rep}(\mathbb{Z}_2)$. By [KZ22a], there are six simple strings $1, \vartheta, ss, sr, rs, rr$ in TC^2 with the fusion rules given in (2.1.5).

1. $A_0 = 1$ is the trivial condensable E_2 -algebra in $\Omega \mathcal{TC}$. If we condense A_0 along a line, we obtain a condensed string $\mathbb{1}$.⁵⁸
2. $A_e = 1 \oplus e$ is a condensable E_2 -algebra in $\Omega \mathcal{TC}$. If we condense A_e along a line, we obtain the condensed string $rr = \Sigma A_e \in \mathcal{C}$.



As we have shown in Example 3.3.42, this string rr is a condensable E_1 -algebra that can be further condensed to create the rough boundary of TC^3 .


3. $A_m = 1 \oplus m$ is a condensable E_2 -algebra in $\Omega \mathcal{TC}$. If we condense A_m along a line, we obtain a condensed string ss .



As we have shown in Example 3.3.42, this string ss has a canonical structure of a condensable E_1 -algebra in $\mathcal{C}$ that can be further condensed to create the smooth boundary of TC^3 .


⁵⁸Note that a condensable E_1 -algebra that are Morita equivalent to 1 (e.g., $[x, x] = x \otimes x^*$ for $x \in \Omega \mathcal{TC}$) produces the ‘essentially same’ condensed string $\mathbb{1}$. In general, when A is only a condensable E_1 -algebra, by choosing a unit morphism $\mathbb{1} \rightarrow \Sigma A$ different from the canonical one, the condensed defect ΣA might be further condensed if A is Morita equivalent to a condensable E_2 -algebra.

4. $A_f = 1 \oplus f$ is a condensable E_1 -algebra in $\Omega\mathcal{TC}$. If we condense A_f along a line, we obtain $\Sigma A_f = \vartheta$. This follows from the fact that the condensable E_1 -algebra, as an internal hom, that creates the e - m -duality wall ϑ from the trivial wall is precisely $A_f = 1 \oplus f$ [Bom10, KK12].




In this case, ϑ is not a condensable E_1 -algebra in $\mathcal{TC}$ and can not be further condensed.

5. $A_4 = 1 \oplus e \oplus m \oplus f$ is a condensable E_1 -algebra in $\Omega\mathcal{TC}$ isomorphic to the tensor product algebra $A_e \otimes A_m$. If we condense A_4 along a line, we obtain $\Sigma A_4 = rs$ because the internal hom algebra associated to r is precisely A_4 [JZWK⁺19, CJKYZ20]. Note that ΣA_4 cannot be further condensed.




6. Note that A_4^{op} is also a condensable E_1 -algebra in $\Omega\mathcal{TC}$ isomorphic to the tensor product algebra $A_m \otimes A_e$. Condensing it produce the condensed string $sr = \Sigma A_4^{\text{op}}$, which cannot be further condensed.



Example 5.1.9. We consider 2+1D double Ising topological order.

1. The only Lagrangian E_2 -algebra in $\Omega(\mathcal{I}s \boxtimes \mathcal{I}s^{\text{op}}) = \Omega\mathcal{I}s \boxtimes \Omega\mathcal{I}s^{\text{op}}$ is $A_0 = 1 \boxtimes 1 \oplus \psi \boxtimes \psi \oplus \sigma \boxtimes \sigma$. By condensing it in an open 2-disk, we obtain the trivial phase $\mathbf{1}^3$. This condensation can be achieved in two steps. First, we condense it along a line, we obtain a condensed string ΣA_0 , which can be identified with $\Omega\mathcal{I}s \simeq \text{RMod}_{A_0}(\Omega\mathcal{I}s \boxtimes \Omega\mathcal{I}s^{\text{op}}) \in \Sigma\Omega(\mathcal{I}s \boxtimes \mathcal{I}s^{\text{op}})$, or, coordinate independently, as 1-codimensional defect obtained by stacking of two layers of the unique gapped boundary of the double Ising topological order.



Condensing ΣA_0 produces the trivial phase $\mathbf{1}^3$ and the gapped boundary $M^2 = \mathbf{Is}^2$, i.e.,

$$\text{Mod}_{\Sigma A_0}^{E_1}(\Sigma\Omega(\mathcal{I}s \boxtimes \mathcal{I}s^{\text{op}})) \simeq 2\text{Vec}, \quad (\mathcal{M}, m) = (\text{RMod}_{\Sigma A_0}(\Sigma\Omega(\mathcal{I}s \boxtimes \mathcal{I}s^{\text{op}})), \Sigma A_0) \simeq (\mathcal{I}s, \mathbf{1}).$$

2. The anyon condensation from double Ising to toric code was explained in Example 4.1.13. Now we reconstruct this phase transition in two steps. Recall that $A = 1 \boxtimes 1 \oplus \psi \boxtimes \psi \in \Omega\mathcal{I}s \boxtimes \Omega\mathcal{I}s^{\text{op}} \simeq \mathfrak{Z}_1(\Omega\mathcal{I}s)$ is a condensable E_2 -algebra in $\mathfrak{Z}_1(\Omega\mathcal{I}s)$. We have $\Sigma A = \text{RMod}_A(\mathfrak{Z}_1(\Omega\mathcal{I}s))$. By further condensing ΣA in $\Sigma\mathfrak{Z}_1(\Omega\mathcal{I}s) = \mathcal{I}s \boxtimes \mathcal{I}s^{\text{op}}$, according to (3.3.18), we obtain the $\mathbb{Z}_2$ topological order as the condensed phase, i.e.,

$$\text{Mod}_{\Sigma A}^{E_1}(\Sigma\mathfrak{Z}_1(\Omega\mathcal{I}s)) \simeq \Sigma\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2)) = \mathcal{TC}.$$

Note that $\mathcal{K} = \text{RMod}_A(\mathfrak{Z}_1(\Omega\mathcal{I}s))^{\text{op}} = 1 \oplus \vartheta$ (recall (3.3.28)) is also a condensable E_1 -algebra in $\mathcal{TC}$, which is not the delooping of any condensable E_2 -algebra in $\Omega\mathcal{TC}$. Recall that condensing $\mathcal{K}$ in $\mathcal{TC}^3$ reproduces the 3D double Ising topological order, i.e., $\text{Mod}_{\mathcal{K}}^{E_1}(\mathcal{TC}) \simeq \mathcal{I}s \boxtimes \mathcal{I}s^{\text{op}}$. We explain in Example 5.3.10 that condensing $\mathcal{K}$ in $\mathcal{TC}^3$ can be understood as the gauging of the e - m -duality in $\mathcal{TC}^3$.

Example 5.1.10. All the examples of anyon condensations given in Section 4.1.2 can be redefined by first condensing an anyon A along a line to give a condensed string ΣA then condensing ΣA . We leave them as exercises. We explain only a special case that echo with previous examples (see Example 3.3.44).

- Let G be a finite group and $H \leq G$ a subgroup. Let $A = \text{Fun}(G/H)$. Note that $A \in \text{Alg}_{E_2}^c(\mathfrak{Z}_1(\text{Rep}(G)))$. The central functor $- \otimes A : \mathfrak{Z}_1(\text{Rep}(G)) \rightarrow \text{RMod}_A(\mathfrak{Z}_1(\text{Rep}(G)))$ defines a condensable E_1 -algebra ΣA in $\mathcal{GT}_G^3 = \Sigma \mathfrak{Z}_1(\text{Rep}(G))$; or equivalently, the monoidal functor

$$\text{Rep}(G) \rightarrow \text{Rep}(G) \boxtimes_{\mathfrak{Z}_1(\text{Rep}(G))} \text{RMod}_A(\mathfrak{Z}_1(\text{Rep}(G))) \simeq \text{RMod}_A(\text{Rep}(G)) = \text{Rep}(H)$$

defines a condensable E_1 -algebra $\Sigma A = \text{Rep}(H)^{\text{op}}$ in $\mathcal{GT}_G^3 = \text{BMod}_{\text{Rep}(G)|\text{Rep}(G)}(2\text{Vec})^{\text{op}}$. By condensing ΣA in $\mathcal{GT}_G^3$, we obtain the condensed phase $D^3 = \text{GT}_H^3$ with $\mathcal{D} = \text{BMod}_{\text{Rep}(H)|\text{Rep}(H)}(2\text{Vec})^{\text{op}}$. This coincides with the well-known result that the anyon condensation of the E_2 -algebra A in $\mathfrak{Z}_1(\text{Rep}(G))$ produces the condensed phase $D^3 = \text{GT}_H^3$ (recall Example 4.1.11).

5.2 Condensation of higher codimensional topological defects

5.2.1 General theory

Consider a (potentially anomalous) $n+1$ D simple topological order C^{n+1} . The category of all topological defects $\mathcal{C}$ is a fusion n -category. By definition, for $k \geq 1$, the category $\Omega^{k-1}\mathcal{C}$ of topological defects of codimension k or higher is an E_k -fusion $(n-k+1)$ -category, and $\Sigma\Omega^{k-1}\mathcal{C}$ is an indecomposable separable $(n-k+2)$ -category containing $1_{\mathbb{1}_{\mathcal{C}}}^{k-2}$ and is the full subcategory of $\Omega^{k-2}\mathcal{C}$.

We denote the composition of k -morphisms in BC by $\otimes^k$. For $1 \leq i \leq k$, similar to the discussion in Section 2.3.2, $\otimes^i$ also represents the fusion product of two k -codimensional topological defects along the x^i -direction. For $1 \leq i, j \leq k$, if we want to discuss the fusion among k -codimensional topological defects along the x^i -direction and the x^j -direction, we often use the following picture in the (x^i, x^j) -plane:

where $a, b, c, d \in \Omega^{k-1}\mathcal{C}$ and other directions of these k -codimensional defects a, b, c, d are assumed to lie in the normal direction of above plane. By restricting to the (x^i, x^j) -plane, we see that these k -codimensional topological defects are equipped with the following isomorphism:

$$(a \otimes^i c) \otimes^j (b \otimes^i d) \xrightarrow[\simeq]{\delta_{a,b,c,d}^{(i,j)}} (a \otimes^j b) \otimes^i (c \otimes^j d) \quad (5.2.2)$$

that defines an E_2 -monoidal structure. These fusion products $\otimes^i$ for $1 \leq i \leq k$ define the k -dimensional multiplication of k -codimensional topological defects and, therefore, define the E_k -monoidal structure on $\Omega^{k-1}\mathcal{C}$.

Lemma° 5.2.1. Let $\mathcal{B}$ be an E_k -multi-fusion n -category. The assignment $A \mapsto \Sigma A = \text{RMod}_A(\mathcal{B})$ defines a functor:

$$\Sigma(-) := \text{RMod}_-(\mathcal{B}) : \text{Alg}_{E_k}^c(\mathcal{B}) \rightarrow \text{Alg}_{E_{k-1}}^c(\Sigma\mathcal{B}).$$

Proof. Since $\mathcal{B}$ is E_k -monoidal, it has fusion products $\otimes^i$ in i -th independent direction for $1 \leq i \leq k$. We have shown in Section 5.1.1 that $\Sigma(-) : \text{Alg}_{E_1}^c(\mathcal{B}) \rightarrow \Sigma\mathcal{B}$, defined as follows:


$$\begin{array}{ccc} A & \longrightarrow & \text{RMod}_A(\mathcal{B}) \\ f \downarrow & & \downarrow - \otimes_{A,f} B \\ B & \longrightarrow & \text{RMod}_B(\mathcal{B}) \end{array}$$

is a well-defined functor. We have used the k -th monoidal structure $\otimes^k$ on $\mathcal{B}$ to define the objects in the category $\text{Alg}_{\mathcal{E}_1}^c(\mathcal{B})$. For $1 \leq i < k$, each tensor product $\otimes^i$ endows $\text{Alg}_{\mathcal{E}_1}^c(\mathcal{B})$ with a monoidal structure denoted by $(\text{Alg}_{\mathcal{E}_1}^c(\mathcal{B}), \otimes^i)$, and also endows $\text{RMod}_A(\mathcal{B})$ with a monoidal structure denoted by $(\text{RMod}_A(\mathcal{B}), \otimes^i)$. Restricting to the (x^i, x^k) -plane, we have already shown in Section 5.1.1 that $\Sigma : (\text{Alg}_{\mathcal{E}_1}^c(\mathcal{B}), \otimes^i) \rightarrow (\Sigma\mathcal{B}, \otimes^i)$ is a lax-monoidal functor, which maps algebras in $(\text{Alg}_{\mathcal{E}_1}^c(\mathcal{B}), \otimes^i)$ to algebras in $(\Sigma\mathcal{B}, \otimes^i)$. Since Σ preserves the monoidal structure in all i -th directions for $1 \leq i < k$, it preserves the $\mathcal{E}_{k-1}$ -monoidal structures on $\text{Alg}_{\mathcal{E}_1}^c(\mathcal{B})$ and $\Sigma\mathcal{B}$. $\square$

Now we are ready to state one of the main results of this work, which is a natural consequence of all the previous discussion in this work.

Theorem^{ph} 5.2.2. In a (potentially anomalous) $n+1$ D simple topological order $\mathcal{C}^{n+1}$, a k -codimensional topological defect $A \in \Omega^{k-1}\mathcal{C}$ is fully condensable, if it is equipped with the structure of a condensable $\mathcal{E}_k$ -algebra in $\Omega^{k-1}\mathcal{C}$ (i.e., $A \in \text{Alg}_{\mathcal{E}_k}^c(\Omega^{k-1}\mathcal{C})$). By condensing it, we mean a k -step process that produces a condensed new phase $\mathcal{D}^{n+1}$.


1. In the first step, we condense the defect A along one of the transversal direction x^k . It means that we first proliferate the defect A along the transversal direction x^k , then introduce interactions among these defect, or equivalently, introduce condensation maps defined by the multiplication map $A \otimes A \rightarrow A$ of the $\mathcal{E}_k$ -algebra in the x^k -direction. This produces a $(k-1)$ -codimensional topological defect $\Sigma A \in \Sigma\Omega^{k-1}\mathcal{C} \hookrightarrow \Omega^{k-2}\mathcal{C}$.



If we use the coordinate system $\Sigma\Omega^{k-1}\mathcal{C} = \text{RMod}_{\Omega^{k-1}\mathcal{C}}((n-k+2)\text{Vec})$, the object ΣA can be defined as follows:

$$\Sigma A := \text{RMod}_A(\Omega^{k-1}\mathcal{C}) \in \text{RMod}_{\Omega^{k-1}\mathcal{C}}((n-k+2)\text{Vec}) \hookrightarrow \Omega^{k-2}\mathcal{C}. \quad (5.2.3)$$

2. It turns out that ΣA has a natural structure of a condensable $\mathcal{E}_{k-1}$ -algebra in $\Sigma\Omega^{k-1}\mathcal{C}$ or in $\Omega^{k-2}\mathcal{C}$. In the second step, we condense ΣA along one of the remaining transversal direction x^{k-1} to produce a $(k-2)$ -codimensional topological defect $\Sigma^2 A := \text{RMod}_{\Sigma A}(\Omega^{k-2}\mathcal{C}) \in \Sigma\Omega^{k-2}\mathcal{C} \hookrightarrow \Omega^{k-3}\mathcal{C}$.



3. Repeat above process until $\Sigma^{k-1}A$, which is a condensable $\mathcal{E}_1$ -algebra in $\mathcal{C}$. In the k -th step, we condense $\Sigma^{k-1}A$ in $\mathcal{C}$ along the only remaining transversal direction x^1 . We obtain a new phase $\mathcal{D}^{n+1}$ and a gapped domain wall $M^n = (\mathcal{M}, m)$ such that

$$\begin{array}{c} \mathcal{C} \xrightarrow{L_1} \mathcal{M} \quad \mathcal{M} \xleftarrow{R_1} \mathcal{D} \\ \mathcal{C}^{n+1} \quad \quad \quad \mathcal{D}^{n+1} \\ \quad \quad \quad \uparrow M^n \\ \quad \quad \quad M^n = (\mathcal{M}, m) \end{array} \quad (5.2.4)$$

$$\mathcal{D} \simeq \text{Mod}_{\Sigma^{k-1}A}^{\mathcal{E}_1}(\mathcal{C}) \quad \text{and} \quad \mathcal{M} = \text{RMod}_{\Sigma^{k-1}A}(\mathcal{C}) = \Sigma^k A, \quad m = \Sigma^{k-1} A. \quad (5.2.5)$$

4. For $1 \leq i \leq k$, i -codimensional topological defects in $\Omega^{i-1}\mathcal{C}$ move onto the wall according to the functor


$$L_i := - \otimes \Sigma^{k-i} A : \Omega^{i-1}\mathcal{C} \rightarrow \Omega_m^{i-1}\mathcal{M};$$

and those in D^{n+1} move onto the wall according to the functor

$$R_i := \Sigma^{k-i} A \otimes_{\Sigma^{k-i} A} - : \Omega^{i-1} \mathcal{D} \rightarrow \Omega_m^{i-1} \mathcal{M}.$$

The functors L_1 and R_1 is illustrated in (5.2.4).

5. C^{n+1} and D^{n+1} share the same gravitational anomaly. In other words, they are gapped boundaries of the same anomaly-free $n+2$ D topological order as illustrated below,




where $Z(C)^{n+2}$ and $Z(D)^{n+2}$ denote the bulk of C^{n+1} and D^{n+1} , respectively. Mathematically, it implies the following results.

$$\mathfrak{Z}_1(\mathcal{C}) \simeq \mathfrak{Z}_1(\text{Mod}_{\Sigma^{k-1}A}^{E_1}(\mathcal{C})), \quad (5.2.6)$$

$$\mathfrak{Z}_0(\text{RMod}_{\Sigma^{k-1}A}(\mathcal{C})) \simeq \text{Mod}_{\Sigma^{k-1}A}^{E_1}(\mathcal{C})^{\text{op}} \boxtimes_{\mathfrak{Z}_1(\mathcal{C})} \mathcal{C}, \quad (5.2.7)$$

$$\mathcal{D}^{\text{op}} \simeq \text{Mod}_{\Sigma^{k-1}A}^{E_1}(\mathcal{C})^{\text{op}} \simeq \mathfrak{Z}_1(\mathcal{C}, \mathfrak{Z}_0(\text{RMod}_{\Sigma^{k-1}A}(\mathcal{C}))). \quad (5.2.8)$$

6. If C^{n+1} is anomaly-free, then one can alternatively terminate the process at $\Sigma^{k-2}A$, which is a condensable E_2 -algebra in $\Omega\mathcal{C}$. More precisely, we can condense $\Sigma^{k-2}A \in \text{Alg}_{E_2}^c(\Omega\mathcal{C})$ directly to obtain the same new phase D^{n+1} and the same gapped domain wall such that



$$\Omega\mathcal{D} \simeq \text{Mod}_{\Sigma^{k-2}A}^{E_2}(\Omega\mathcal{C}) \quad \text{and} \quad \Omega_m \mathcal{M} = \text{RMod}_{\Sigma^{k-2}A}(\Omega\mathcal{C}). \quad (5.2.9)$$

The consistency of two approaches immediately implies the following natural equivalences:

$$\Sigma \text{Mod}_{\Sigma^{k-2}A}^{E_2}(\Omega\mathcal{C}) \simeq \text{Mod}_{\Sigma^{k-1}A}^{E_1}(\mathcal{C}) \quad \text{and} \quad \Sigma \text{RMod}_{\Sigma^{k-2}A}(\Omega\mathcal{C}) \simeq \text{RMod}_{\Sigma^{k-1}A}(\mathcal{C}), \quad (5.2.10)$$

where the first equivalence is a monoidal equivalence. $\square$

Remark 5.2.3. When the k -codimensional topological defect $A \in \Omega^{k-1}\mathcal{C}$ is only a condensable E_i -algebra in $\Omega^{k-1}\mathcal{C}$ for $1 \leq i < k$, then condensing A can only produce lower codimensional defects $\Sigma^j A$ in C^{n+1} for $i \leq j < k$ instead of a new phase D^{n+1} . In this sense, A can be called ‘ i -condensable’. When $i = k$, the defect A is called *fully condensable*. $\diamond$

We first provide a physical or working definition of an E_m -module over an E_m -algebra.

Definition 5.2.4. Given $\mathcal{A} \in \text{Alg}_{E_m}((n+1)\text{Vec})$ and $A \in \text{Alg}_{E_m}(\mathcal{A})$. For $m = 1$, an E_1 -module over A in $\mathcal{A}$ is just an A - A -bimodule in $\mathcal{A}$. For $m = 2$, an E_2 -module over A in $\mathcal{A}$ is defined in Section 4.1.1. For $m \geq 3$, an E_m -module over A (or an E_m - A -module) in $\mathcal{A}$ is an object M in $\mathcal{A}$ equipped with two-side A -actions in the x^i -dimension for all $1 \leq i \leq m$ such that M is an E_2 -module over A in each (x^i, x^j) -plane for $1 \leq i < j \leq m$ (recall Remark 4.1.1). $\square$

We denote the category of E_m -module over A by $\text{Mod}_A^{E_m}(\mathcal{A})$. When $m = 0$, we set $\text{Mod}_A^{E_0}(\mathcal{A}) := (\mathcal{A}, A)$. For $m \geq 2$, we would like to provide an alternative way to understand $\text{Mod}_A^{E_m}(\mathcal{A})$. For an E_m -multi-fusion n -category $\mathcal{A}$ and $A, B \in \text{Alg}_{E_m}^c(\mathcal{A})$, there is a natural equivalence of $(n+1)$ -categories (recall Eq. (4.1.4)):

$$\text{BMod}_{A|B}(\mathcal{A}) \simeq \text{LMod}_A(\mathcal{A}) \boxtimes_{\mathcal{A}} \text{RMod}_B(\mathcal{A}).$$

The $\text{BMod}_{A|B}(\mathcal{A})$ inherits a natural structure of an E_{m-1} -multi-fusion n -category from those on $\text{LMod}_A(\mathcal{A})$ and $\text{RMod}_B(\mathcal{A})$. More explicitly, it is equipped with a tensor unit $A \otimes B$ and a tensor product $\otimes_{A|B}^i$ in the x^i -dimension induced from $\otimes_A^i \boxtimes_{\mathcal{A}} \otimes_B^i$ in $\text{LMod}_A(\mathcal{A}) \boxtimes_{\mathcal{A}} \text{RMod}_B(\mathcal{A})$ for all $1 \leq i \leq m-1$. The tensor product $\otimes_A^i$ in $\text{LMod}_A(\mathcal{A})$ is induced from the two-side A -actions on modules $A \otimes^i M \rightarrow M$ and $M \otimes^i A \rightarrow M$, both of which are induced from the left A -action $A \otimes^m M \rightarrow M$ in the x^m -dimension via adiabatically moving A . A special case of $m = 2$ is illustrated in the picture in (4.1.3). The tensor products $\otimes_B^i$ are similar.

When $A = B$, we obtain $A \in \text{Alg}_{E_{m-1}}^c(\text{Mod}_A^{E_1}(\mathcal{A}))$, and A is a direct summand of the tensor unit $A \otimes A$. Note that, at the same time, $\text{Mod}_A^{E_1}(\mathcal{A})$ has another natural structure of E_1 -multi-fusion n -category with the tensor product $\otimes_A^m$ (defined in the x^m -dimension) and the tensor unit A .

If we repeat this process j -times, the category $\text{Mod}_A^{E_1}(\cdots \text{Mod}_A^{E_1}(\mathcal{A}))$ has two different (higher) monoidal structures:

- an E_{m-j} -monoidal structure with the tensor unit $A \otimes A$ and a tensor product in each x^i -direction for $i = 1, \dots, m-j$ such that they are all compatible;
- an E_j -monoidal structure with a tensor unit A and a tensor product $\otimes_A^i$ in the x^i -dimension for $i = m-j+1, \dots, m$ such that they are all compatible.

Therefore, we obtain a generalization of (4.1.7) and an alternative way to define $\text{Mod}_A^{E_m}(\mathcal{A})$ as summarized in the following Proposition.

Proposition 5.2.5. We have an E_m -monoidal equivalence

$$\text{Mod}_A^{E_m}(\mathcal{A}) \simeq \text{Mod}_A^{E_{m-1}}(\text{Mod}_A^{E_1}(\mathcal{A})) \simeq \text{Mod}_A^{E_1}(\cdots \text{Mod}_A^{E_1}(\mathcal{A})).$$

Remark 5.2.6. It is possible to give a more compact mathematical definition of $\text{Mod}_A^{E_m}(\mathcal{A})$ as an iterated looping of the category of E_0 -modules over an E_0 -algebra. We will do that in [KZZ25]. $\diamond$

When C^{n+1} is anomalous, the condensable E_k -algebra A in $\Omega^{k-1}\mathcal{C}$ does not directly determines the phase transition without going through the condensed defect $\Sigma^{k-1}A$. However, it determines some partial data of the condensed phase directly.

1. Since all (deconfined) topological defects of codimension k or higher in the condensed phase D^{n-1} necessarily come from those in $\Omega^{k-1}\mathcal{C}$, and they are necessarily the E_k -modules over A , we must have the following E_k -monoidal equivalence:

$$\text{Mod}_A^{E_k}(\Omega^{k-1}\mathcal{C}) \simeq \Omega^{k-1} \text{Mod}_{\Sigma^{k-1}A}^{E_1}(\mathcal{C}) = \Omega^{k-1} \Sigma^k A. \quad (5.2.11)$$

By [KZZ2b], the right hand side is automatically an E_k -monoidal multi-fusion $(n-k+1)$ -category, so is the left hand side.

2. Topological defects living on the gapped domain wall M^n are confined topological defects. Those k -codimensional topological defect on M^n , i.e., objects in $\Omega_m^{k-1}\mathcal{M}$, are necessarily originated from those in $\Omega^{k-1}\mathcal{C}$. Recall that a deconfined k -codimensional defect needs to support A -actions in $2k$ directions (i.e., two A -actions in each x^i -th dimension for $1 \leq i \leq k$). If we drop the A -action in one single direction among all $2k$ directions, the associated topological defect is not deconfined. On the other hand, the minimal requirement for a topological defect to survive on the wall M^n is that it should support an A -action at least from one single direction (see further discussion in Remark 5.2.7). Therefore, those defects in $\Omega_m^{k-1}\mathcal{M}$ are necessarily right A -modules in $\Omega^{k-1}\mathcal{C}$, i.e.,

$$\text{RMod}_A(\Omega^{k-1}\mathcal{C}) \simeq \Omega^{k-1} \text{RMod}_{\Sigma^{k-1}A}(\mathcal{C}) = \Omega^{k-1} \Sigma^k A. \quad (5.2.12)$$

As a consequence, we also have

$$\Omega_m^i \mathcal{M} = \Omega^i \Sigma^k A = \Omega^i \text{RMod}_{\Sigma^{k-1}A}(\mathcal{C}) \simeq \text{RMod}_{\Sigma^{k-1-i}A}(\Omega^i \mathcal{C}), \quad \text{for } 0 \leq i \leq k-1. \quad (5.2.13)$$

Remark 5.2.7. The condition for an object in $\Omega^{k-1}\mathcal{C}$ to live on the wall M^n is that it supports an A -action from at least a single direction. It can be equipped with compatible A -actions from other directions. When it does not support compatible A -actions in all $2k$ directions in all k -dimensions, say the compatibility of A -action in one of the two directions in the x^k -dimension is dropped, even though one can still define an A -action in this direction using the other A -action in the x^k -dimension via a braiding in (x^i, x^k) -plane, this A -action depends on the paths of the braidings. The incompatibility of different paths causes a strong interference that triggers the confinement. At the same time, one can also see that $(k-1)$ -codimensional defects in $\Omega_m^{k-1}\mathcal{M}$ can have different levels of confinement. More precisely, assume the domain wall M^n is defined by the hyperplane defined by $x^k = x_0^k$ in space. Then all $(k-1)$ -codimensional defects in $\Omega_m^{k-1}\mathcal{M}$ are confined in the x^k -dimension. For $1 \leq i < k$, if a $(k-1)$ -codimensional defect in $\Omega_m^{k-1}\mathcal{M}$ supports a two-side A -action in the x^i -dimension that are compatible with the A -action in one of the x^k -direction, then it is deconfined in the x^i -dimension. $\diamond$

We summarize and reformulate above results as mathematical results.

Theorem 5.2.8. Let $\mathcal{C}$ be a fusion n -category and $A \in \text{Alg}_{E_k}^c(\Omega^{k-1}\mathcal{C})$. For $1 \leq i < k$, we set $\Sigma^0 A := A$ and define inductively $\Sigma^i A = \text{RMod}_{\Sigma^{i-1}A}(\Omega^{k-i}\mathcal{C})$. Then $\Sigma^i A \in \text{Alg}_{E_{k-i}}^c(\Omega^{k-i-1}\mathcal{C})$, and there is a natural E_{k-i} -monoidal equivalence of E_{k-i} -fusion higher categories:

$$\text{Mod}_{\Sigma^i A}^{E_{k-i}}(\Omega^{k-i-1}\mathcal{C}) \simeq \Omega^{k-i-1} \text{Mod}_{\Sigma^{k-1}A}^{E_1}(\mathcal{C}). \quad (5.2.14)$$

Proof. We obtain this result through physical arguments. Now we sketch a mathematical proof of (5.2.14) parallel to that of Theorem 5.1.4. It is enough to prove the following E_k -monoidal equivalence:

$$\text{Mod}_A^{E_k}(\Omega^{k-1}\mathcal{C}) \simeq \Omega \text{Mod}_{\Sigma A}^{E_{k-1}}(\Omega^{k-2}\mathcal{C}). \quad (5.2.15)$$

Note that $\Omega \text{Mod}_{\Sigma A}^{E_{k-1}}(\Omega^{k-2}\mathcal{C}) \simeq \Omega \text{Mod}_{\Sigma A}^{E_{k-1}}(\Sigma \Omega^{k-1}\mathcal{C})$. A 1-morphism $\Sigma A \rightarrow \Sigma A$ in $\text{Mod}_{\Sigma A}^{E_{k-1}}(\Sigma \Omega^{k-1}\mathcal{C})$ is a functor from ΣA to ΣA intertwining the right $\Omega^{k-1}\mathcal{C}$ -action and the $(k-1)$ -dimensional ΣA -action. Since a functor from ΣA to ΣA intertwining the right $\Omega^{k-1}\mathcal{C}$ -action is necessarily equivalent to the functor $-\otimes_A^k m$ for an A - A -bimodule m in $\Omega^{k-1}\mathcal{C}$, a 1-morphism $\Sigma A \rightarrow \Sigma A$ in $\text{Mod}_{\Sigma A}^{E_{k-1}}(\Sigma \Omega^{k-1}\mathcal{C})$ is precisely a functor $-\otimes_A^k m$ intertwining the ΣA -actions in the x^i -dimension for all $1 \leq i < k$, a condition which upgrades m to an E_2 - A -module in $\Omega^{k-1}\mathcal{C}$ in each (x^i, x^k) -plane as shown in the proof of Theorem 5.1.4. Hence, m is an E_k - A -module in $\Omega^{k-1}\mathcal{C}$. $\square$

If $\mathcal{C} = \Sigma^{k-1}\Omega^{k-1}\mathcal{C}$, i.e., topological defects outside $\Omega^{k-1}\mathcal{C}$ are all condensation descendants of those in $\Omega^{k-1}\mathcal{C}$, then we have the following stronger results.

1. It is natural to ask if all deconfined topological defects outside $\Omega^{k-1}\mathcal{D}$ are also condensation descendants of those in $\Omega^{k-1}\mathcal{D}$. It turns out that this is not correct because, for $i < k-1$, $\Omega^i\mathcal{D}$ is not connected in general even though $\Omega^i\mathcal{C}$ is connected. We give an example of this fact. Consider $\mathcal{C}^5 = \mathbf{1}^5$ and $A = \text{Rep}(G) \in 2\text{Vec}$, then $\Sigma A = 2\text{Rep}(G) \in 3\text{Vec} = \Omega\mathcal{C}$ and $\Sigma^2 A = 3\text{Rep}(G)$. In this case, we have $\text{Mod}_{\Sigma A}^{E_2}(3\text{Vec}) \simeq \mathfrak{Z}_1(3\text{Rep}(G))$, which is not indecomposable as a separable 3-category [KTZ20b]. Therefore, $\text{Mod}_{\Sigma A}^{E_2}(3\text{Vec}) \neq \Sigma \text{Mod}_A^{E_3}(2\text{Vec})$. Moreover, for $i = 0$, it was known that every fusion 2-category is Morita equivalent to a connected one (i.e., $\Sigma\mathcal{B}$ for a braided fusion 1-category $\mathcal{B}$) [Déc23a].
2. In this case, all confined topological defects on the gapped domain wall are necessarily condensation descendants of those in $\Omega_m^{k-1}(\mathcal{M})$. This fact will be explained mathematically in [KZZZ25]. Therefore, we obtain the following natural equivalence:

$$\Sigma^{k-1} \text{RMod}_A(\Omega^{k-1}\mathcal{C}) \simeq \text{RMod}_{\Sigma^{k-1}A}(\mathcal{C}). \quad (5.2.16)$$

Note that (5.2.12) is an immediate consequence of (5.2.16), respectively. A special case of (5.2.16) will be useful later. For a subgroup H of a finite group G , we have

$$\text{RMod}_{n\text{Rep}(H)}((n+1)\text{Rep}(G)) \simeq \Sigma^n(\text{RMod}_{\text{Fun}(G/H)}(\text{Rep}(G))) \simeq \Sigma^n(\text{Rep}(H)) \simeq n\text{Rep}(H). \quad (5.2.17)$$

We reformulate (5.2.11), (5.2.12) and (5.2.16) as mathematical results below.

Theorem° 5.2.9. Let $\mathcal{B}$ be an E_k -fusion n -category and $A \in \text{Alg}_{E_k}^c(\mathcal{B})$. We set $\Sigma^0 A = A$ and, for $0 \leq i < k$, we define inductively $\Sigma^i A = \text{RMod}_{\Sigma^{i-1} A}(\Sigma^{i-1} \mathcal{B})$. We have a natural E_{k-i} -monoidal equivalence:

$$\text{Mod}_{\Sigma^i A}^{E_{k-i}}(\Sigma^i \mathcal{B}) \simeq \Omega^{k-i} \text{Mod}_{\Sigma^k A}^{E_0}(\Sigma^k \mathcal{B}), \quad (5.2.18)$$

which is proved in the proof of Theorem 5.2.8. Moreover, we have a natural E_{k-1-i} -monoidal equivalence and an E_{k-1} -monoidal equivalence, respectively, given as follows:

$$\Sigma^i \text{RMod}_A(\mathcal{B}) \simeq \text{RMod}_{\Sigma^i A}(\Sigma^i \mathcal{B}), \quad \text{RMod}_A(\mathcal{B}) \simeq \Omega^i \text{RMod}_{\Sigma^i A}(\Sigma^i \mathcal{B}). \quad (5.2.19)$$

Example 5.2.10. The E_k -monoidal equivalence (5.2.18) provides a convenient way to understand and compute the category $\text{Mod}_A^{E_k}(\mathcal{B})$. It can also serve as a working definition of $\text{Mod}_A^{E_k}(\mathcal{B})$. We give an example. Let $A = n\text{Rep}(G)$, which can be viewed as a condensable E_k -algebra in $(n+1)\text{Vec}$ for all $k \geq 1$. We have the following results.

1. $\text{Mod}_{n\text{Rep}(G)}^{E_1}((n+1)\text{Vec}) \simeq \mathcal{GT}_G^{n+2}$.
2. $\text{Mod}_{n\text{Rep}(G)}^{E_2}((n+1)\text{Vec}) \simeq \Omega \text{Mod}_{(n+1)\text{Rep}(G)}^{E_1}((n+2)\text{Vec}) \simeq \mathfrak{Z}_1((n+1)\text{Rep}(G)) \simeq \Omega \mathcal{GT}_G^{n+2}$.
3. For $k > 2$, we have

$$\text{Mod}_{n\text{Rep}(G)}^{E_k}((n+1)\text{Vec}) \simeq (n+1)\text{Rep}(G) \quad (5.2.20)$$

♡

which follows from

$$\text{Mod}_{n\text{Rep}(G)}^{E_k}((n+1)\text{Vec}) \simeq \Omega^{k-1}(\text{Mod}_{(n+k-1)\text{Rep}(G)}^{E_1}((n+k)\text{Vec})) \simeq \Omega^{k-2} \mathfrak{Z}_1((n+k-1)\text{Rep}(G)).$$

The following result generalizes the equivalence (3.3.5).

Theorem° 5.2.11. Let $\mathcal{B}$ be an E_k -fusion n -category and $A \in \text{Alg}_{E_k}^c(\mathcal{B})$. We have an E_k -monoidal equivalence:

$$\text{Mod}_A^{E_k}(\mathcal{B})^{\text{op}} \simeq \mathfrak{Z}_k(\mathcal{B}, \mathfrak{Z}_{k-1}(\text{RMod}_A(\mathcal{B}))). \quad (5.2.21)$$

When $\mathcal{B} = n\text{Vec}$, we obtain $\text{Mod}_A^{E_k}(n\text{Vec})^{\text{op}} \simeq \mathfrak{Z}_{k-1}(\Sigma A)$ (recall Example 2.3.23).

Proof. It follows from the following equivalences:

$$\begin{aligned} \text{Mod}_A^{E_k}(\mathcal{B})^{\text{op}} &\simeq \Omega^{k-1}(\text{Mod}_{\Sigma^{k-1} A}^{E_1}(\Sigma^{k-1} \mathcal{B})^{\text{op}}) \\ &\simeq \Omega^{k-1} \mathfrak{Z}_1(\Sigma^{k-1} \mathcal{B}, \mathfrak{Z}_0(\text{RMod}_{\Sigma^{k-1} A}(\Sigma^{k-1} \mathcal{B}))) \\ &\simeq \mathfrak{Z}_k(\mathcal{B}, \Omega^{k-1} \mathfrak{Z}_0(\Sigma^{k-1} \text{RMod}_A(\mathcal{B}))) \\ &\simeq \mathfrak{Z}_k(\mathcal{B}, \mathfrak{Z}_{k-1}(\text{RMod}_A(\mathcal{B}))), \end{aligned}$$

where the first ‘ $\simeq$ ’ is due to (5.2.18), the second ‘ $\simeq$ ’ is due to (3.3.5), the third ‘ $\simeq$ ’ is due to (2.3.13) and the 4th ‘ $\simeq$ ’ is due to (2.3.15). □

Remark 5.2.12. The special cases of $n = 2$ and $k = 1, 2, 3$ of Theorem 5.2.11 are also obtained independently in [Xu24a], which appeared on arXiv on the same day as ours. These results generalize similar results for 1-categories obtained much earlier in [Sch01]. ◇

Example 5.2.13. When $\mathcal{A} = (n-1)\text{Rep}(G)$ and $\mathcal{B} = n\text{Vec}$, we have

$$\begin{aligned} \text{Mod}_{(n-1)\text{Rep}(G)}^{E_1}(n\text{Vec})^{\text{op}} &\simeq \mathfrak{Z}_0(n\text{Rep}(G)), \\ \text{Mod}_{(n-1)\text{Rep}(G)}^{E_2}(n\text{Vec})^{\text{op}} &\simeq \mathfrak{Z}_1(n\text{Rep}(G)), \end{aligned}$$

which are also the consequences of (3.3.11) and (4.2.13), respectively. For $k \geq 3$, we obtain the following equivalences:

$$\text{Mod}_{(n-1)\text{Rep}(G)}^{E_k}(n\text{Vec})^{\text{op}} \simeq \mathfrak{Z}_{k-1}(n\text{Rep}(G)) \simeq n\text{Rep}(G), \quad (5.2.22)$$

where the second ‘ $\simeq$ ’ is due to Example 5.2.10 and was rigorously proved in [KTZ20b, Proposition 3.11] for $n = 2$ and $k = 3$. The other cases of this result is new. ♡

Remark 5.2.14. Note that (5.2.22) shows certain stability of higher centers that might hold for more general E_m -fusion n -categories. Actually, for an E_{k+1} -fusion n -category $\mathcal{E}$, we have

$$\mathcal{E} = \Omega \Sigma \mathcal{E} \simeq \mathfrak{Z}_k(\mathcal{E}),$$

if and only if the canonical E_k -monoidal functor $\Sigma \mathcal{E} \rightarrow \mathfrak{Z}_{k-1}(\Sigma \mathcal{E})$ is fully faithful. This condition clearly holds for $\mathcal{E} = n\text{Rep}(G)$ and $k \geq 2$. $\diamond$

5.2.2 General constructions

Similar to Section 3.3.2, in this subsection, we provide some general examples or constructions of the condensations of k -codimensional topological defects in C^{n+1} . We would like to begin with a discussion on how to construct condensable E_k -algebras.

For an $n+1$ D topological order C^{n+1} , some condensable E_k -algebras in $\Omega^{k-1}\mathcal{C}$ can be constructed from the condensable E_{k+1} -algebras in $\Omega^k\mathcal{C}$ by delooping. In general, there are additional condensable E_k -algebra in $\Omega^{k-1}\mathcal{C}$. In particular, those condensable E_k -algebras in the sub-category $\Sigma\Omega^k\mathcal{C}$ of $\Omega^{k-1}\mathcal{C}$ can all be constructed physically. More explicitly, consider D^{n+1} and a gapped (potentially anomalous) domain wall M^n between C^{n+1} and D^{n+1} . For $k+1 \leq n$, we illustrate the $(k+1)$ -th left bulk-to-wall map L_{k+1} as follow.

$$\begin{array}{ccc} C^{n+1} & & D^{n+1} \\ \Omega^k\mathcal{C} \times \Omega_m^k\mathcal{M} \xrightarrow{\odot} \Omega_m^k\mathcal{M} & \xrightarrow{\quad} & \Omega_m^k\mathcal{M} \\ \Omega^k\mathcal{C} \xrightarrow{L_{k+1} = -\odot 1_m^k} \Omega_m^k\mathcal{M} & & \end{array}$$

The category $\Omega_m^k\mathcal{M} := \Omega^{k-1}(\Omega_m\mathcal{M})$ is E_k -monoidal. It is clear that $\Omega_m^k\mathcal{M} \in \text{LMod}_{\Omega^k\mathcal{C}}((n-k+1)\text{Vec})$. Moreover, the left $\Omega^k\mathcal{C}$ -action is compatible with the fusion products $\otimes^i$ in $\Omega_m^k\mathcal{M}$ for $i = 1, \dots, k$ because the $\Omega^k\mathcal{C}$ is acting from the x^{k+1} -dimension that are orthogonal to the $x^1, \dots, x^k$ -dimensions. In other words, $\Omega_m^k\mathcal{M} \in \text{LMod}_{\Omega^k\mathcal{C}}(\text{Alg}_{E_k}^c(n-k+1)\text{Vec})$. Similar to the proof of (3.3.14), by running the arguments in each (x^i, x^k) -plane (for $i < k$), we must have

$$\Omega_m^k\mathcal{M} \in \text{Alg}_{E_k}^c(\text{RMod}_{\Omega^k\mathcal{C}}((n-k+1)\text{Vec})) = \text{LMod}_{\Omega^k\mathcal{C}}(\text{Alg}_{E_k}^c(n-k+1)\text{Vec}).$$

Similar to the discussion in Section 3.3.2, all condensable E_k -algebras in $\Sigma\Omega^k\mathcal{C}$ should arise in this way. We summarize this results as a mathematical theorem.

Theorem^o 5.2.15. For an E_k -fusion n -category $\mathcal{B}$, we have a natural equivalence:

$$\text{Alg}_{E_{k-1}}^c(\text{RMod}_{\mathcal{B}}((n+1)\text{Vec})) \simeq \text{LMod}_{\mathcal{B}}(\text{Alg}_{E_{k-1}}^c((n+1)\text{Vec})).$$

Remark 5.2.16. Note that the action $\odot : \Omega^k\mathcal{C} \times \Omega_m^k\mathcal{M} \rightarrow \Omega_m^k\mathcal{M}$ also induces an internal hom $[1_m^k, 1_m^k]_{\Omega^k\mathcal{C}}$, which is automatically a condensable E_{k+1} -algebra in $\Omega^k\mathcal{C}$, i.e.,

$$[1_m^k, 1_m^k]_{\Omega^k\mathcal{C}} \in \text{Alg}_{E_{k+1}}^c(\Omega^k\mathcal{C}). \quad (5.2.23)$$

We give a brief explanation of (5.2.23) in Appendix A.3. In general, it is not true that condensing $[1_m^k, 1_m^k]_{\Omega^k\mathcal{C}}$ can recover the gapped domain wall M^n . In general, $\text{RMod}_{[1_m^k, 1_m^k]_{\Omega^k\mathcal{C}}}(\Omega^k\mathcal{C})$ is only a sub-fusion category of $\Omega_m^k\mathcal{M}$. However, if $\mathcal{C} = \Sigma^k\Omega^k\mathcal{C}$ and M^n is obtained from the condensation of $A \in \text{Alg}_{E_{k+1}}^c(\Omega^k\mathcal{C})$, i.e., $\mathcal{M} = \text{RMod}_{\Sigma^k A}(\mathcal{C})$, then we have $\Omega_m^k\mathcal{M} \simeq \text{RMod}_A(\Omega^k\mathcal{C})$. As a consequence, we must have $A = [1_m^k, 1_m^k]_{\Omega^k\mathcal{C}}$. $\diamond$

Example 5.2.17. The basic idea behind all explicit constructions of condensable E_k -algebras in an E_k -fusion n -category $\mathcal{B}$ is to find a nice coordinate system for $\mathcal{B}$.

- (1) Consider an E_k -fusion $(n-1)$ -category $\mathcal{A}$. Recall Theorem 5.2.11, the E_k -fusion n -category $\mathfrak{Z}_{k-1}(\Sigma\mathcal{A})^{\text{op}}$ is equipped with a coordinate system $\mathfrak{Z}_{k-1}(\Sigma\mathcal{A})^{\text{op}} = \text{Mod}_{\mathcal{A}}^{E_k}(n\text{Vec})$. In this case, a condensable E_k -algebra in $\text{Mod}_{\mathcal{A}}^{E_k}(n\text{Vec})$ is precisely an E_k -multi-fusion $(n-1)$ -category $\mathcal{P}$ equipped with an E_k -monoidal functor $\phi : \mathcal{A} \rightarrow \mathcal{P}$. Moreover, we have

$$\text{Mod}_{\mathcal{P}}^{E_k}(\mathfrak{Z}_{k-1}(\Sigma\mathcal{A})^{\text{op}}) \simeq \text{Mod}_{\mathcal{P}}^{E_k}(\text{Mod}_{\mathcal{A}}^{E_k}(n\text{Vec})) \simeq \text{Mod}_{\mathcal{P}}^{E_k}(n\text{Vec}) \simeq \mathfrak{Z}_{k-1}(\Sigma\mathcal{P})^{\text{op}}$$

Examples of such $\mathcal{P}$ can be constructed. Let $\mathcal{X} \in \text{LMod}_{\mathcal{A}}(\text{Alg}_{\mathbb{E}_{k-1}}^c(n\text{Vec}))$. The $\mathbb{E}_{k-1}$ -monoidal $\mathcal{A}$ -action $\mathcal{A} \boxtimes \mathcal{X} \rightarrow \mathcal{X}$ induces a canonical $\mathbb{E}_k$ -monoidal functor $\mathcal{A} \rightarrow \mathfrak{Z}_{k-1}(\mathcal{X})$, which defines a condensable $\mathbb{E}_k$ -algebra in $\mathcal{B}$. For example, one can choose $\mathcal{X} = \mathcal{A}$.

- (a) Let $\mathcal{A} = n\text{Rep}(G)$ and $\mathcal{B} = \text{Mod}_{n\text{Rep}(G)}^{\mathbb{E}_k}((n+1)\text{Vec})$ (recall Example 5.2.10). The canonical $\mathbb{E}_k$ -monoidal functors [KZ24]:

$$n\text{Rep}(G) \xrightarrow{f} n\text{Rep}(H), \quad n\text{Rep}(G) \rightarrow \mathfrak{Z}_{k-1}(n\text{Rep}(H)),$$

where $H \leq G$ is a subgroup, define condensable $\mathbb{E}_k$ -algebras in $\mathcal{B}$.

- (2) Assume $\mathcal{B} = \Sigma \mathfrak{Z}_k(\mathcal{A})$ for an $\mathbb{E}_k$ -fusion $(n-1)$ -category $\mathcal{A}$. In this case, we have

$$\Sigma \mathfrak{Z}_k(\mathcal{A}) = \text{RMod}_{\mathfrak{Z}_k(\mathcal{A})}(n\text{Vec}) \simeq \Sigma \Omega \text{Mod}_{\mathcal{A}}^{\mathbb{E}_k}(n\text{Vec}) \simeq \Sigma \Omega \mathfrak{Z}_{k-1}(\Sigma \mathcal{A}). \quad (5.2.24)$$

where we have used the mathematical fact $\mathfrak{Z}_k(\mathcal{A}) \simeq \text{Fun}_{\mathcal{A}}^{\mathbb{E}_k}(\mathcal{A}, \mathcal{A}) \simeq \Omega(\text{Mod}_{\mathcal{A}}^{\mathbb{E}_k}(n\text{Vec}))$ followed directly from the universal property of the $\mathbb{E}_k$ -center $\mathfrak{Z}_k(-)$ (see [Lur17, Fra12]). Therefore, this $\mathcal{B}$ is a full subcategory of $\text{Mod}_{\mathcal{A}}^{\mathbb{E}_k}(n\text{Vec})$. More precisely, a condensable $\mathbb{E}_k$ -algebra in $\mathcal{B} = \Sigma \mathfrak{Z}_k(\mathcal{A})$ is precisely an $\mathbb{E}_k$ -fusion $(n-1)$ -category $\mathcal{P}$ lying in $\Sigma \Omega \text{Mod}_{\mathcal{A}}^{\mathbb{E}_k}(n\text{Vec})$ and equipped with an $\mathbb{E}_k$ -monoidal functor $\phi : \mathcal{A} \rightarrow \mathcal{P}$. $\heartsuit$

Now we provide some general constructions of k -codimensional defect condensations.

- (1) $\mathcal{C}^{n+1} = \mathbf{1}^{n+1}$: In this case, we have $\mathcal{C} = n\text{Vec}$ and $\Omega^{k-1}\mathcal{C} = (n-k+1)\text{Vec}$ for $k \leq n$. A simple condensable $\mathbb{E}_k$ -algebra in $(n-k+1)\text{Vec}$ is precisely an $\mathbb{E}_k$ -fusion $(n-k)$ -category $\mathcal{A}$. Following Theorem^{ph} 5.2.2, we first condense the k -codimensional defect $\mathcal{A}$ along a transversal direction of the defect $\mathcal{A}$ and obtain a $(k-1)$ -codimensional defect $\Sigma \mathcal{A}$, then we repeat the process to obtain a 2-codimensional defect $\Sigma^{k-2}\mathcal{A}$ and a 1-codimensional defect $\Sigma^{k-1}\mathcal{A}$. By condensing $\Sigma^{k-1}\mathcal{A}$ along the remaining transversal direction, we obtain a condensed phase $\mathcal{D}^{n+1}$ and a gapped domain wall $\mathcal{M}^n$ such that $\mathcal{D}^{n+1} = \overline{\mathcal{Z}(\mathcal{M})^{n+1}}$,


$$\mathcal{D} \simeq \text{Mod}_{\Sigma^{k-1}\mathcal{A}}^{\mathbb{E}_1}(n\text{Vec}) \simeq \mathfrak{Z}_0(\Sigma^k \mathcal{A})^{\text{op}}, \quad (\mathcal{M}, m) = (\Sigma^k \mathcal{A}, \Sigma^{k-1} \mathcal{A}).$$

Moreover, we have, for $1 \leq i \leq k-1$,

$$\Omega^i \mathcal{D} \simeq \text{Mod}_{\Sigma^{k-i}\mathcal{A}}^{\mathbb{E}_{i+1}}((n-i)\text{Vec}) \simeq \Omega^i \mathfrak{Z}_0(\Sigma^k \mathcal{A})^{\text{op}} \simeq \mathfrak{Z}_i(\Sigma^{k-i} \mathcal{A})^{\text{op}}, \quad \Omega_m^i \mathcal{M} \simeq \Sigma^{k-i} \mathcal{A}.$$

For $i = k-1$, we recover $\text{Mod}_{\mathcal{A}}^{\mathbb{E}_k}((n-k+1)\text{Vec}) \simeq \mathfrak{Z}_{k-1}(\Sigma \mathcal{A})^{\text{op}}$ obtained previously as a special case in Theorem 5.2.11.

- (2) Now we consider a condensation of k -codimensional defect in $\mathcal{D}^{n+1}$ obtained above. A simple condensable $\mathbb{E}_k$ -algebra B in $\Omega^{k-1}\mathcal{D}$ in the coordinate system $\Omega^{k-1}\mathcal{D} \simeq \text{Mod}_{\mathcal{A}}^{\mathbb{E}_k}((n-k+1)\text{Vec})$ is precisely given by an $\mathbb{E}_k$ -monoidal functor $\mathcal{A} \rightarrow \mathcal{B}$ for an $\mathbb{E}_k$ -fusion $(n-k)$ -category $\mathcal{B}$. Now we condense B viewed as a k -codimensional defect in $\mathcal{D}^{n+1}$ according to the k -step process described in Theorem^{ph} 5.2.2. First, we condensed the k -codimensional defect B along one of its transversal direction. We obtain a $(k-1)$ -codimensional defect ΣB defined by an $\mathbb{E}_{k-1}$ -monoidal functor $\Sigma \mathcal{A} \rightarrow \Sigma \mathcal{B}$. We repeat the process. We obtain a 2-codimensional defect $\Sigma^{k-2}B$ defined by the $\mathbb{E}_2$ -monoidal functor $\Sigma^{k-2}\mathcal{A} \rightarrow \Sigma^{k-2}\mathcal{B}$ and a 1-codimensional defect $\Sigma^{k-1}B$ defined by the $\mathbb{E}_1$ -monoidal functor $\Sigma^{k-1}\mathcal{A} \rightarrow \Sigma^{k-1}\mathcal{B}$. By further condensing such obtained 1-codimensional defect, we obtain a new condensed phase $\mathcal{E}^{n+1}$ and a new gapped domain wall $\mathcal{N}^n$ as illustrated below:



such that

$$\begin{aligned} \mathcal{E} &\simeq \text{Mod}_{\Sigma^{k-1}B}^{\mathbb{E}_1}(\mathcal{D}) \simeq \text{Mod}_{\Sigma^{k-1}\mathcal{B}}^{\mathbb{E}_1}(n\text{Vec}) \simeq \mathfrak{Z}_0(\Sigma^k \mathcal{B})^{\text{op}}, \\ (\mathcal{N}, n) &\simeq (\text{RMod}_{\Sigma^{k-1}B}(\mathcal{D}), \Sigma^{k-1}B) \simeq (\text{BMod}_{\Sigma^{k-1}\mathcal{A}|\Sigma^{k-1}\mathcal{B}}(n\text{Vec}), \Sigma^{k-1}\mathcal{B}). \end{aligned}$$

Moreover, we have, for $1 \leq i \leq k-1$,

$$\Omega^i \mathcal{E} \simeq \text{Mod}_{\Sigma^{k-i-1}\mathcal{B}}^{\text{E}_{i+1}}(\Omega^i \mathcal{D}) \simeq \text{Mod}_{\Sigma^{k-i-1}\mathcal{B}}^{\text{E}_{i+1}}(\text{Mod}_{\Sigma^{k-i-1}\mathcal{A}}^{\text{E}_{i+1}}((n-i)\text{Vec})) \simeq \text{Mod}_{\Sigma^{k-i-1}\mathcal{B}}^{\text{E}_{i+1}}((n-i)\text{Vec}) \simeq \mathfrak{Z}_i(\Sigma^{k-i}\mathcal{B})^{\text{op}}.$$

Moreover, we have the following (monoidal) equivalences:

$$\begin{aligned} \mathcal{M} \boxtimes_{\mathcal{D}} \mathcal{N} &\simeq \text{RMod}_{\Sigma^{k-1}\mathcal{B}}(n\text{Vec}) \simeq \Sigma^k \mathcal{B}, \\ \Omega_m \mathcal{M} \boxtimes_{\Omega \mathcal{D}} \Omega_n \mathcal{N} &\simeq \Sigma^{k-1} \mathcal{A} \boxtimes_{\mathfrak{Z}_1(\Sigma^{k-1}\mathcal{A})} \text{Fun}_{\Sigma^{k-1}\mathcal{A}|\Sigma^{k-1}\mathcal{B}}(\Sigma^{k-1}\mathcal{B}, \Sigma^{k-1}\mathcal{B}) \\ &\simeq \text{Fun}_{(n-1)\text{Vec}|\Sigma^{k-1}\mathcal{B}}(\Sigma^{k-1}\mathcal{B}, \Sigma^{k-1}\mathcal{B}) \simeq \Sigma^{k-1}\mathcal{B} \simeq \Omega_{m \boxtimes_{\mathcal{D}} n}(\mathcal{M} \boxtimes_{\mathcal{D}} \mathcal{N}). \end{aligned}$$

Note that $\text{E}^{n+1} = \mathbf{1}^{n+1}$ if and only if $\Sigma^{k-1}\mathcal{B}$ is non-degenerate. Moreover, by [KZZ1, Corollary* 2.10], we obtain $\text{E}^{n+1} = \mathbf{1}^{n+1}$ if and only if $\Sigma^k \mathcal{B}$ is invertible as a separable n -category. This motivates us to introduce the following notion [KZZZ25].

Definition 5.2.18. An E_k -fusion m -category $\mathcal{B}$ is called non-degenerate if $\Sigma^k \mathcal{B}$ is invertible separable $(m+k)$ -category. $\square$

Lemma^o 5.2.19. If an E_k -fusion m -category $\mathcal{B}$ is non-degenerate, then we have $\text{Mod}_{\mathcal{B}}^{\text{E}_k}((m+1)\text{Vec}) \simeq m\text{Vec}$, which further implies that $\mathfrak{Z}_k(\mathcal{B}) \simeq m\text{Vec}$.

Proof. Note the first condition (i.e., $\mathcal{B}$ being non-degenerate) means $\mathfrak{Z}_0(\Sigma^k \mathcal{B})$ is trivial; and the second condition means $\mathfrak{Z}_{k-1}(\Sigma \mathcal{B}) \simeq \Omega^{k-1} \mathfrak{Z}_0(\Sigma^k \mathcal{B})$ is trivial; and the third condition means $\mathfrak{Z}_k(\mathcal{B}) \simeq \Omega^k \mathfrak{Z}_0(\Sigma^k \mathcal{B})$ is trivial. $\square$

We summarize above results as a theorem that generalizes Theorem^{ph} 3.3.32 and Theorem^{ph} 4.2.25.

Theorem^{ph} 5.2.20. For an E_k -fusion m -category $\mathcal{A}$, a simple condensable E_k -algebra B in $\mathfrak{Z}_{k-1}(\Sigma \mathcal{A})^{\text{op}}$ in the coordinate system (recall Theorem 5.2.11)

$$\mathfrak{Z}_{k-1}(\Sigma \mathcal{A})^{\text{op}} \simeq \text{Mod}_{\mathcal{A}}^{\text{E}_k}((m+1)\text{Vec})$$

is precisely defined by an E_k -monoidal functor $\mathcal{A} \rightarrow \mathcal{B}$ for an E_k -multi-fusion m -category $\mathcal{B}$. The E_k -algebra B is Lagrangian if and only if $\mathcal{B}$ is Lagrangian. Moreover, we have

$$\text{Mod}_B^{\text{E}_k}(\mathfrak{Z}_{k-1}(\Sigma \mathcal{A})^{\text{op}}) \simeq \text{Mod}_{\mathcal{B}}^{\text{E}_k}((m+1)\text{Vec}) \simeq \mathfrak{Z}_{k-1}(\Sigma \mathcal{B})^{\text{op}}.$$

This result is precisely the k -codimensional layer of a physical condensation from the initial phase $\mathcal{C}^{m+k+1}$, which is anomaly-free, simple and non-chiral, such that

$$\mathcal{C} = \text{Mod}_{\Sigma^{k-1}\mathcal{A}}^{\text{E}_1}((m+k)\text{Vec}) \simeq \mathfrak{Z}_0(\Sigma^k \mathcal{A})^{\text{op}}, \quad \Omega^{k-1} \mathcal{C} \simeq \Omega^{k-1} \mathfrak{Z}_0(\Sigma^k \mathcal{A})^{\text{op}} \simeq \mathfrak{Z}_{k-1}(\Sigma \mathcal{A})^{\text{op}}$$

to the condensed phase $\mathcal{D}^{m+k+1}$ such that

$$\mathcal{D} = \text{Mod}_{\Sigma^{k-1}\mathcal{B}}^{\text{E}_1}(\mathcal{C}) \simeq \text{Mod}_{\Sigma^{k-1}\mathcal{B}}^{\text{E}_1}((m+k)\text{Vec}), \quad \Omega^{k-1} \mathcal{D} \simeq \text{Mod}_{\mathcal{B}}^{\text{E}_k}((m+1)\text{Vec}) \simeq \mathfrak{Z}_{k-1}(\Sigma \mathcal{B})^{\text{op}},$$

where $\Sigma^{k-1}\mathcal{B}$ is equipped with an E_1 -monoidal functor $\Sigma^{k-1}\mathcal{A} \rightarrow \Sigma^{k-1}\mathcal{B}$ and should be viewed as a condensable E_1 -algebra in $\mathcal{C}$. Moreover, the gapped domain wall $\mathcal{M}^{m+k}$ produced by this condensation is given by

$$\begin{aligned} (\mathcal{M}, m) &= (\text{BMod}_{\Sigma^{k-1}\mathcal{A}|\Sigma^{k-1}\mathcal{B}}(\Sigma^{k-1}\mathcal{B}, \Sigma^{k-1}\mathcal{B}), \Sigma^{k-1}\mathcal{B}), \\ \Omega_m^{k-1} \mathcal{M} &= \Omega^{k-2} \text{Fun}_{\Sigma^{k-1}\mathcal{A}|\Sigma^{k-1}\mathcal{B}}(\Sigma^{k-1}\mathcal{B}, \Sigma^{k-1}\mathcal{B}) \simeq \text{RMod}_{\mathcal{B}}(\text{Mod}_{\mathcal{A}}^{\text{E}_k}((m+1)\text{Vec})), \end{aligned}$$

which generalizes the $k=2$ case in (4.2.15). $\square$

Example 5.2.21. Recall (5.2.22), for $k \geq 3$, we have following equivalences:

$$\text{Mod}_{n\text{Rep}(G)}^{\text{E}_k}((n+1)\text{Vec}) \simeq \mathfrak{Z}_{k-1}((n+1)\text{Rep}(G))^{\text{op}} \simeq (n+1)\text{Rep}(G)^{\text{op}} = (n+1)\text{Rep}(G), \quad (5.2.25)$$

In this coordinate system of $(n+1)\text{Rep}(G)$, a condensable E_k -algebra B in $(n+1)\text{Rep}(G)$ is precisely given by an E_k -monoidal functor $n\text{Rep}(G) \rightarrow \mathcal{B}$ for an E_k -multi-fusion n -category $\mathcal{B}$. We have

$$\text{Mod}_B^{\text{E}_k}((n+1)\text{Rep}(G)) \simeq \text{Mod}_{\mathcal{B}}^{\text{E}_k}((n+1)\text{Vec}) \simeq \mathfrak{Z}_{k-1}(\Sigma \mathcal{B})^{\text{op}}.$$

For example, let $\mathcal{B} = \text{Rep}(H)$ for a subgroup $H \leq G$. Let B be defined by the forgetful functor $f : \text{Rep}(G) \rightarrow \text{Rep}(H)$. Then for $k \geq 3$, we obtain

$$\text{Mod}_B^{\text{E}_k}((n+1)\text{Rep}(G)) \simeq \text{Mod}_{n\text{Rep}(H)}^{\text{E}_k}((n+1)\text{Vec}) \simeq \mathfrak{Z}_{k-1}((n+1)\text{Rep}(H))^{\text{op}} \simeq (n+1)\text{Rep}(H),$$

where we have used (5.2.22) in the last ‘ $\simeq$ ’. $\heartsuit$

5.2.3 Concrete examples

Example 5.2.22. Consider the 3+1D $\mathbb{Z}_2$ -gauge theory $\mathrm{GT}_{\mathbb{Z}_2}^4$ (recall the notations in Example 4.2.39).

1. Condensing the e -particles:

- (a) We first condense the e -particle along a line, we obtain the $\mathbb{1}_c$ -string. Mathematically, it amounts to condense the condensable E_3 -algebra $A_e^0 = \mathbb{1}_1 \oplus e$ in $\Omega^2 \mathcal{TC} = \mathrm{Rep}(\mathbb{Z}_2)$ along a line and we obtain $\mathbb{1}_c = \Sigma A_e^0 = \mathrm{RMod}_{A_e^0}(\mathrm{Rep}(\mathbb{Z}_2)) \simeq \mathrm{Vec}$ in $2\mathrm{Rep}(\mathbb{Z}_2) = \mathrm{RMod}_{\mathrm{Rep}(\mathbb{Z}_2)}(2\mathrm{Vec})$.
- (b) Then we condense the $\mathbb{1}_c$ -strings along one of the remaining transversal direction. We obtain a condensed defect $\Sigma^2 A_e^0 = \mathrm{rr}$, where rr denotes the 1-codimensional topological defect obtained by stacking two rough boundaries. It is clear that rr is condensable E_1 -algebra in $\mathcal{GT}_{\mathbb{Z}_2}^4$. By further condensing $\Sigma^2 A_e^0$, we obtain the trivial phase $\mathbf{1}^4$ as the condensed phase and the rough boundary as the domain wall.
- (c) Alternatively, we can condense the $\mathbb{1}_c$ -string as a condensable E_2 -algebra in $\Omega \mathcal{GT}_{\mathbb{Z}_2}^4$, we obtain the trivial phase $\mathbf{1}^4$ as the condensed phase and the rough boundary, on which topological defects form the fusion 2-category $2\mathrm{Vec}_{\mathbb{Z}_2}$.

2. Condensing the $A_m = (1 \oplus m)$ -string:

- (a) We first condense A_m along one of the transversal direction. It produces a membrane $\Sigma A_m = \mathrm{ss}$, where ss denotes the 1-codimensional defect obtained by stacking two smooth boundaries. By further condensing ss , we obtain $\mathbf{1}^4$ as the condensed phase and the smooth boundary, on which topological defects form the fusion 2-category $2\mathrm{Rep}(\mathbb{Z}_2)$.
- (b) Alternatively, we can condense A_m directly in the remaining two transversal directions to obtain the smooth boundary.

3. Condensing the $A_m^{\mathrm{tw}} = (1 \oplus m)$ -string:

- (a) We first condense A_m^{tw} along one of the transversal direction. It produces a membrane $\Sigma A_m^{\mathrm{tw}} = \mathrm{tt}$, where tt denotes the 1-codimensional defect obtained by stacking two twist smooth boundaries. By further condensing tt , we obtain $\mathbf{1}^4$ as the condensed phase and the twist smooth boundary, on which topological defects form the fusion 2-category $2\mathrm{Rep}(\mathbb{Z}_2)$.
- (b) Alternatively, we can condense A_m^{tw} directly in the remaining two transversal directions to obtain the twist smooth boundary. $\heartsuit$

Example 5.2.23. Consider the 3+1D G -gauge theory GT_G^4 . As fusion 3-categories, we have the following two different coordinate systems:

$$\mathcal{GT}_G^4 = \mathrm{RMod}_{\mathfrak{Z}_1(2\mathrm{Rep}(G))}(3\mathrm{Vec}), \quad \mathcal{GT}_G^4 = \mathrm{BMod}_{2\mathrm{Rep}(G)|2\mathrm{Rep}(G)}(3\mathrm{Vec})^{\mathrm{op}}.$$

Its particles form a symmetric fusion 1-category $\Omega^2 \mathcal{GT}_G^4 = \mathrm{Rep}(G)$, and its strings form a braided fusion 2-category $\Omega \mathcal{GT}_G^4 = \mathfrak{Z}_1(2\mathrm{Rep}(G))$.

Let $H \leq G$ be a subgroup of G . We have $A = \mathrm{Fun}(G/H) \in \mathrm{Alg}_{E_3}^c(\mathrm{Rep}(G))$. By condensing it along one of the transversal directions, we obtain

$$\Sigma A := \mathrm{RMod}_A(\mathrm{Rep}(G)) \simeq \mathrm{Rep}(H) \in 2\mathrm{Rep}(G) \hookrightarrow \mathfrak{Z}_1(2\mathrm{Rep}(G)).$$

By condensing ΣA along one of the remaining transversal directions, we obtain

$$\Sigma^2 A = \mathrm{RMod}_{\Sigma A}(\mathfrak{Z}_1(2\mathrm{Rep}(G))) \in \mathrm{Alg}_{E_1}^c(\Sigma \mathfrak{Z}_1(2\mathrm{Rep}(G))).$$

Now we translate $\Sigma^2 A$ into the second coordinate system of $\mathcal{GT}_G^4$ as follows:

$$2\mathrm{Rep}(G) \rightarrow 2\mathrm{Rep}(G) \boxtimes_{\mathfrak{Z}_1(2\mathrm{Rep}(G))} \mathrm{RMod}_{\Sigma A}(\mathfrak{Z}_1(2\mathrm{Rep}(G))) \simeq \mathrm{RMod}_{\Sigma A}(2\mathrm{Rep}(G)) \simeq 2\mathrm{Rep}(H), \quad (5.2.26)$$

where the first ' $\simeq$ ' a natural generalization of results in 1-categories (see for example [KZ18a, XY25] and also Remark 5.2.24) and the second ' $\simeq$ ' is due to (5.2.17). By condensing $\Sigma^2 A = 2\mathrm{Rep}(H)^{\mathrm{op}}$ in $\mathcal{GT}_G^4$, we obtain GT_H^4 as the condensed phase because

$$\mathrm{Mod}_{\Sigma^2 A}^{E_1}(\mathcal{GT}_G^4) \simeq \mathrm{BMod}_{2\mathrm{Rep}(H)|2\mathrm{Rep}(H)}(3\mathrm{Vec})^{\mathrm{op}} \simeq \mathcal{GT}_H^4.$$

As a consequence, it also implies that we should have

$$\mathrm{Mod}_{\Sigma A}^{\mathbb{E}_2}(\mathfrak{Z}_1(2\mathrm{Vec}_G)) \simeq \mathfrak{Z}_1(2\mathrm{Vec}_H).$$

This provides a proof of Example 4.2.40. The gapped domain wall $M^3 = (\mathcal{M}, m)$ is given by

$$(\mathcal{M}, m) \simeq (\mathrm{RMod}_{\Sigma^2 A}(\mathcal{GT}_G^4), \Sigma^2 A) \simeq (\mathrm{BMod}_{2\mathrm{Rep}(H)|2\mathrm{Rep}(G)}(3\mathrm{Vec}), 2\mathrm{Rep}(H)).$$

We see that $\Omega_m \mathcal{M} = \mathrm{Fun}_{2\mathrm{Rep}(H)|2\mathrm{Rep}(G)}(2\mathrm{Rep}(H), 2\mathrm{Rep}(H))$. When $A = \mathrm{Fun}(G)$, the condensed phase is trivial and M^2 is a gapped boundary of GT_G^4 such that the boundary defects form the fusion 2-category $\Omega_m \mathcal{M} \simeq 2\mathrm{Vec}_G$. $\heartsuit$

Remark 5.2.24. In Eq. (5.2.26), we have used a natural equivalence without giving a proof. It is actually a special case of a more general result. For $\mathcal{A} \in \mathrm{Alg}_{\mathbb{E}_k}^c(n\mathrm{Vec})$, $\mathcal{M} \in \mathrm{RMod}_{\mathcal{A}}(\mathrm{Alg}_{\mathbb{E}_{k-1}}^c(n\mathrm{Vec}))$ and $A \in \mathrm{Alg}_{\mathbb{E}_k}^c(\mathcal{A})$, we have natural $\mathbb{E}_{k-1}$ -monoidal equivalences:

$$\mathrm{RMod}_A(\mathcal{M}) \simeq \mathrm{RMod}_A(\mathcal{M} \boxtimes_{\mathcal{A}} A) \simeq \mathcal{M} \boxtimes_{\mathcal{A}} \mathrm{RMod}_A(A).$$

When $n = 2$, the proof of the above results, which is quite standard (see for example [KZ18a, XY25]), is based on the colimits theory for 1-categories. It is natural to expect that we need a fully developed colimit theory for separable higher categories for a proof of the $(n > 2)$ -cases. $\diamond$

Example 5.2.25. Consider the $n+2$ D G -gauge theory GT_G^{n+2} . As fusion $(n+1)$ -categories, we have two coordinate systems:

$$\mathcal{GT}_G^{n+2} = \mathrm{RMod}_{\mathfrak{Z}_1(n\mathrm{Rep}(G))}((n+1)\mathrm{Vec}), \quad \mathcal{GT}_G^{n+2} = \mathrm{BMod}_{n\mathrm{Rep}(G)|n\mathrm{Rep}(G)}((n+1)\mathrm{Vec})^{\mathrm{op}}. \quad (5.2.27)$$

Its particles form a symmetric fusion 1-category $\Omega^n \mathcal{GT}_G^{n+2} = \mathrm{Rep}(G)$. In general, we have

$$\Omega^{n-k} \mathcal{GT}_G^{n+2} = (k+1)\mathrm{Rep}(G) \quad \text{for } 0 \leq k < n-1$$

and $\Omega \mathcal{GT}_G^{n+2} = \mathfrak{Z}_1(n\mathrm{Rep}(G))$.

Let $H \leq G$ be a subgroup of G . The commutative algebra $A = \mathrm{Fun}(G/H)$ of all $\mathbb{C}$ -valued functions on G is an $\mathbb{E}_{n+1}$ -algebra in $\Omega^n \mathcal{GT}_G^{n+2} = \mathrm{Rep}(G)$. As a composed particle, $\mathrm{Fun}(G/H)$ can be condensed. More precisely, by condensing the $\mathrm{Fun}(G/H)$ -particles, we mean the following procedures.

1. We first condensing the A -particle along a line, we obtain a string

$$\Sigma A = \mathrm{RMod}_A(\mathrm{Rep}(G)) \simeq \mathrm{Rep}(H) \in \mathrm{Alg}_{\mathbb{E}_n}^c(2\mathrm{Rep}(G)).$$

2. We further condense the ΣA -string along one of the remaining transversal directions, we obtain

$$\Sigma^2 A = \mathrm{RMod}_{\Sigma A}(2\mathrm{Rep}(G)) \simeq \mathrm{RMod}_{\Sigma A}(2\mathrm{Vec}) \simeq 2\mathrm{Rep}(H).$$

3. We repeat the procedure until $\Sigma^{n-1} A = (n-1)\mathrm{Rep}(H) \in n\mathrm{Rep}(G) \hookrightarrow \mathfrak{Z}_1(n\mathrm{Rep}(G))$. By condensing $\Sigma^{n-1} A$ along one of the remaining two transversal directions, we obtain

$$\Sigma^n A = \mathrm{RMod}_{\Sigma^{n-1} A}(\mathfrak{Z}_1(n\mathrm{Rep}(G))) \in \mathrm{Alg}_{\mathbb{E}_1}^c(\Sigma \mathfrak{Z}_1(n\mathrm{Rep}(G))).$$

Now we translate $\Sigma^n A$ into an $\mathbb{E}_1$ -algebra in the second coordinate system of $\mathcal{GT}_G^{n+2}$ in (5.2.27). It is defined by the following monoidal functor (recall Remark 5.2.24):

$$n\mathrm{Rep}(G) \rightarrow n\mathrm{Rep}(G) \boxtimes_{\mathfrak{Z}_1(n\mathrm{Rep}(G))} \mathrm{RMod}_{\Sigma^{n-1} A}(\mathfrak{Z}_1(n\mathrm{Rep}(G))) \simeq \mathrm{RMod}_{\Sigma^{n-1} A}(n\mathrm{Rep}(G)) \simeq n\mathrm{Rep}(H).$$

4. By condensing $\Sigma^n A = n\mathrm{Rep}(H)$, we obtain GT_H^{n+2} as the condensed phase because

$$\mathrm{GT}_H^{n+2} \simeq \mathrm{Mod}_{n\mathrm{Rep}(H)^{\mathrm{op}}}^{\mathbb{E}_1}((n+1)\mathrm{Vec}) \simeq \mathrm{Mod}_{n\mathrm{Rep}(H)^{\mathrm{op}}}^{\mathbb{E}_1}(\mathcal{GT}_G^{n+2}).$$

In this example, we provides the precise mathematical theory behind the folklore that breaking the G -gauge “symmetry” in GT_G^{n+2} to a subgroup H gives the H -gauge theory GT_H^{n+2} . $\heartsuit$




Figure 15: The dashed line is the invertible domain wall of the topological order such that moving an anyon across the domain wall realizes the braided equivalence $\rho_g : \mathcal{C} \rightarrow \mathcal{C}$. The topological defects living on the end of this invertible domain wall are called the g -defects, and they form the category $\mathcal{C}_g$. Equivalently, moving $x \in \mathcal{C}$ around $y \in \mathcal{C}_g$ yields the auto-equivalence ρ_g . One can think of that y is attached with a flux g .

5.3 Gauging a G -symmetry

Recall that $\Omega\mathcal{TC}^3$ has a $\mathbb{Z}_2$ -symmetry defined by the non-trivial braided auto-equivalence of $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$, which is also called the e - m -duality. It was known that the double Ising modular tensor category $\mathfrak{Z}_1(\Omega\mathcal{I}s)$ can be obtained from the toric code $\Omega\mathcal{TC}$ via an equivariantization of this $\mathbb{Z}_2$ -symmetry [DGN010]. In physics, this equivariantization is viewed as a process of ‘gauging’ this e - m -duality in $\Omega\mathcal{TC}$. In this subsection, we provide a mathematical foundation of this alleged ‘gauging’ process by explaining that it is a natural consequence of the condensation of 1-codimensional topological defects or the gauging of a generalized 0-form symmetry. Mathematically, it amounts to establish the a previously unknown relation between G -crossed braided extensions (or minimal non-degenerate extensions) and 1-codimensional defect condensations.

5.3.1 Gauging a G -symmetry in a 2+1D SET order

Let G be a finite group. We use 2+1D G -SET order to mean a symmetry enriched topological order with an onsite G -symmetry (or a generalized 0-form symmetry). There are several different approaches towards the mathematical characterization of a 2+1D G -SET order [BBCW19, LKW16a, KLVZZ20b, KZ22b, LYW24].

The first approach towards 2+1D G -SET orders is based on the idea of equipping a topological order with a G -action. Let $\mathcal{C}$ be a nondegenerate braided fusion category equipped with a 2-group homomorphism (monoidal functor) $\rho : G \rightarrow \text{Aut}^{\text{E}_2}(\mathcal{C})$, where $\text{Aut}^{\text{E}_2}(\mathcal{C})$ is the 2-group of braided auto-equivalences of $\mathcal{C}$. It turns out that this G -action is not enough to characterize the G -SET order. In [BBCW19], Barkeshli, Bonderson, Cheng and Wang showed that a G -SET order is a G -crossed extension of $\mathcal{C}$, i.e., a G -crossed braided fusion category

$$\mathcal{C}_G^\times = \bigoplus_{g \in G} \mathcal{C}_g,$$

where $\mathcal{C}_e = \mathcal{C}$ and the category $\mathcal{C}_g \neq 0$ consists of the g -defects living on the end of an invertible 1+1D domain wall realizing the braided auto-equivalence ρ_g as illustrated in Figure 15. The topological defects living on the invertible 1+1D domain wall form the multi-fusion 1-category $\text{Fun}_{\mathcal{C}}(\mathcal{C}_g, \mathcal{C}_g)^{\text{op}} \simeq \mathcal{C}$ because the wall is invertible. As a consequence, $\mathcal{C}_g$ is an invertible $\mathcal{C}$ - $\mathcal{C}$ -bimodule, i.e., $\mathcal{C}_g \in \text{Pic}(\mathcal{C})$, where $\text{Pic}(\mathcal{C})$ is the 3-group of invertible $\mathcal{C}$ -modules, invertible $\mathcal{C}$ -module functors and invertible $\mathcal{C}$ -module natural transformations. According to [BBCW19], the nondegenerate braided fusion category of the particle-like topological defects in the gauged topological order is the G -equivariantization $(\mathcal{C}_G^\times)^G$ of $\mathcal{C}_G^\times$, which consists of ‘ G -invariant’ symmetry defects.

Remark 5.3.1. According to [ENO10], a G -crossed extension of $\mathcal{C}$ is equivalent to lifting a group homomorphism $g \mapsto \mathcal{C}_g$ to a 3-group homomorphism (i.e., a monoidal 2-functor) $\bar{\rho} : G \rightarrow \text{Pic}(\mathcal{C})$. This lifting can be achieved in two steps.

- To lift it to a 2-group homomorphism $G \rightarrow \Pi_2(\text{Pic}(\mathcal{C}))$ (i.e., a monoidal 1-functor), where $\Pi_2(-)$ denotes the underlying 2-group, we need to endow the 1-functor with a monoidal structure, which is defined by $\mathcal{C}$ -module equivalences

$$\mathcal{C}_g \boxtimes_{\mathcal{C}} \mathcal{C}_h \simeq \mathcal{C}_{gh}, \quad \forall g, h \in G \quad (5.3.1)$$

satisfying the axioms of a monoidal structure. These equivalences are equivalent to the tensor product functor $\otimes: \mathcal{C}_g \times \mathcal{C}_h \rightarrow \mathcal{C}_{gh}$ of $\mathcal{C}_G^\times$, because $\otimes$ is $\mathcal{C}_e$ -balanced. This lifting exists if and only if certain obstruction class $O_3 \in H^3(G; \pi_2(\text{Pic}(\mathcal{C}))) \simeq H^3(G; \pi_1(\mathcal{C}^\times))$ vanishes. When O_3 vanishes, all the liftings form a torsor over $H^2(G; \pi_1(\mathcal{C}^\times))$.

- To lift it to a 3-group homomorphism $G \rightarrow \text{Pic}(\mathcal{C})$, we need to specify some $\mathcal{C}$ -module natural isomorphisms as the monoidal structure, which are precisely the associators and G -crossed braidings of the associated G -crossed extension $\mathcal{C}_G^\times$. This lifting exists if and only if certain obstruction class $O_4 \in H^4(G; \pi_3(\text{Pic}(\mathcal{C}))) \simeq H^3(G; \mathbb{k}^\times)$ vanishes. When O_4 vanishes, all the liftings form a torsor over $H^3(G; \mathbb{k}^\times)$.

Since there is a canonical 3-group homomorphism $\partial: \text{Pic}(\mathcal{C}) \rightarrow \text{Aut}^{\text{E}_2}(\mathcal{C})$ [ENO10, DN13], which restricts to a 2-group equivalence $\Pi_2(\text{Pic}(\mathcal{C})) \rightarrow \text{Aut}^{\text{E}_2}(\mathcal{C})$ when $\mathcal{C}$ is nondegenerate, a G -crossed extension of $\mathcal{C}$ (i.e., a 3-group homomorphism $\bar{\rho}: G \rightarrow \text{Pic}(\mathcal{C})$) automatically defines a G -action on $\mathcal{C}$ by the composed functor

$$G \xrightarrow{\bar{\rho}} \text{Pic}(\mathcal{C}) \xrightarrow{\partial} \text{Aut}^{\text{E}_2}(\mathcal{C}).$$

Hence, the 3-group homomorphism $\bar{\rho}: G \rightarrow \text{Pic}(\mathcal{C})$ lifts the 2-group homomorphism $\rho: G \rightarrow \text{Aut}^{\text{E}_2}(\mathcal{C})$. $\diamond$

The second approach towards 2+1D G -SET orders, due to Lan, Kong and Wen [LKW16a, LKW17], is based on the categorical description of G -invariant particle-like topological defects. Such defects can be fused and braided thus form a braided fusion 1-category $\mathcal{D}$. Symmetry charges $\text{Rep}(G)$ form a full subcategory of $\mathcal{D}$ that is equivalent to $\mathfrak{Z}_2(\mathcal{D})$, i.e. $\text{Rep}(G) \rightarrow \mathfrak{Z}_2(\mathcal{D})$. Gauging the G symmetry amounts to a minimal nondegenerate extension of $\mathcal{D}$, i.e., a nondegenerate braided fusion 1-category $\mathcal{M}$ equipped with a braided embedding $\mathcal{D} \hookrightarrow \mathcal{M}$, which induces an equivalence $\text{Rep}(G) \hookrightarrow \mathfrak{Z}_2(\mathcal{D}, \mathcal{M})$. A minimal nondegenerate extension of $\mathcal{D}$ provides a complete categorical description of a 2+1D (bosonic⁵⁹) G -SET order.

Remark 5.3.2. There are two additional approaches towards SET orders based on boundary-bulk relation [KLWZZ20b] and topological Wick rotations [KZ22b]. We do not need them here. $\diamond$

For a bosonic finite symmetry G , the minimal nondegenerate extension approach is equivalent to the G -crossed extension approach. More precisely, by [DGN010, Theorem 4.44], the following two constructions are mutually inverse of each other:

- given a G -crossed braided fusion category $\mathcal{K}$, its equivariantization $\mathcal{K}^G$ is a braided fusion category containing $\text{Rep}(G)$;
- given a braided fusion category $\mathcal{L}$ containing $\text{Rep}(G)$, its de-equivariantization $\mathcal{L}_G$ is a G -crossed braided fusion category.

Moreover, $\mathcal{K}^G$ is nondegenerate if and only if $\mathcal{K}_e$ is nondegenerate and each $\mathcal{K}_g$ is nonzero by [DGN010, Theorem 4.56]. Therefore, the notion of a G -crossed extension is equivalent to that of a minimal nondegenerate extension. For 2+1D G -SET order, we have $\mathcal{D} \simeq \mathcal{C}^G$ and $\mathcal{M} = (\mathcal{C}^\times)^G$. We summarize the relation between these categories in the following diagram:

$$\begin{array}{ccccc}
\text{Vec} & \xrightarrow{\text{inclusion}} & \mathcal{C} & \xrightarrow{G\text{-crossed extension}} & \mathcal{C}_G^\times \\
\downarrow (-)^G & \uparrow (-)_G & \downarrow (-)^G & \uparrow (-)_G & \downarrow (-)^G \uparrow (-)_G \\
\text{Rep}(G) & \xrightarrow{\text{inclusion}} & \mathcal{D} & \xrightarrow{\text{minimal nondegenerate extension}} & \mathcal{M}
\end{array}$$

The third approach towards $n+1$ D SET orders, preceded by earlier works [DKR11, TW24, KLWZZ20a] on a similar formulation of SPT orders, was developed in [KZ24, Section 7.3] and in [LYW24]⁶⁰. An $n+1$ D SET order can be described by a linear monoidal n -functor

$$\phi: \mathcal{T} \rightarrow \mathcal{B}, \quad (5.3.2)$$

⁵⁹Replacing the bosonic symmetry charges $\text{Rep}(G)$ by fermionic symmetry charges $\text{Rep}(G, z)$, where z is the fermion parity, automatically gives a categorical description of a fermionic (G, z) -SET order [LKW16a, LKW17].

⁶⁰Even just from IR point of view, more general notion of a symmetry was possible and was discussed in some online talks during 2022-2023 by the first author of this work [Kon22] (see also Remark 5.3.16).

where the fusion n -category $\mathcal{T}$ describes (non-invertible) higher form symmetries and the fusion n -category $\mathcal{B}$ is the category of topological defects of a topological phase B^{n+1} . To gauge the symmetry amounts to choose a $\mathcal{T}$ -module $\mathcal{K}$ and the gauged theory is described by the indecomposable multi-fusion n -category $\text{Fun}_{\mathcal{B}}(\mathcal{B} \boxtimes_{\mathcal{T}} \mathcal{K}, \mathcal{B} \boxtimes_{\mathcal{T}} \mathcal{K})$. When $\mathcal{K} \simeq \text{RMod}_A(\mathcal{T})$ for a condensable E_1 -algebra $A \in \mathcal{T}$ (see also [LYW24, Remark 4.5]), the gauged theory is described by $\text{BMod}_{\phi(A)|\phi(A)}(\mathcal{B})$. In other words, this gauging is a special case of 1-codimensional defect condensations and the same as the condensation of $\phi(A) \in \text{Alg}_{E_1}^c(\mathcal{B})$. In the rest of this subsection, we established a previously unknown connection between this gauging and that defined by a G -crossed extension (or that defined by a minimal nondegenerate extension).

In the setting of the 2+1D G -SET order in this subsection. Recall that $\Sigma\mathcal{C}$ is a fusion 2-category describing all topological defects in a 2+1D anomaly-free topological order. Since $\text{Pic}(\mathcal{C}) \simeq (\Sigma\mathcal{C})^\times$, a 3-group homomorphism $\phi : G \rightarrow \text{Pic}(\mathcal{C})$ is equivalent to a linear monoidal 2-functor $\phi : 2\text{Vec}_G \rightarrow \Sigma\mathcal{C}$. Therefore, the G -SET order can be equivalently described by a linear monoidal 2-functor $\phi : 2\text{Vec}_G \rightarrow \Sigma\mathcal{C}$. One can recover the G -crossed extension $\mathcal{C}_G^\times$ of $\mathcal{C}$ canonically as follows. Note that $\text{Vec}_G \in 2\text{Vec}_G$ is a condensable E_1 -algebra with the multiplication induced by the group multiplication of G .

Lemma° 5.3.3. The monoidal structure of ϕ endows its image $\phi(\text{Vec}_G)$ with a structure of a condensable E_1 -algebra in $\Sigma\mathcal{C}$, which is precisely the G -crossed braided fusion category $\mathcal{C}_G^\times$.

Proof. Recall that, in the coordinate system $\Sigma\mathcal{C} = \text{RMod}_{\mathcal{C}}(2\text{Vec})$, the condensable E_1 -algebras in $\Sigma\mathcal{C}$ are multi-fusion $\mathcal{C}$ -modules, i.e., multi-fusion categories $\mathcal{P}$ equipped with a braided functor $\mathcal{C} \rightarrow \mathfrak{Z}_1(\mathcal{P})$ [BJS21, Definition-Proposition 3.2] (see also [D  c24, Lemma 2.2.6]) or Theorem^{ph} 3.3.26. In this coordinate system, the G -crossed extension $\mathcal{C}_G^\times$ of $\mathcal{C}$ is equipped with a canonical braided functor $\mathcal{C} \rightarrow \mathfrak{Z}_1(\mathcal{C}_G^\times)$, which is induced by the G -crossed braiding on $\mathcal{C}_G^\times$. It endows $\mathcal{C}_G^\times$ with a condensable E_1 -algebra structure in $\Sigma\mathcal{C}$.

On the other hand, as an object,

$$\phi(\text{Vec}_G) \simeq \bigoplus_{g \in G} \phi(g) = \bigoplus_{g \in G} \mathcal{C}_g = \mathcal{C}_G^\times.$$

The multiplication of $\phi(\text{Vec}_G)$ is induced by the group multiplication of G and the monoidal structure of ϕ :

$$\mathcal{C}_g \boxtimes_{\mathcal{C}} \mathcal{C}_h = \phi(g) \boxtimes_{\mathcal{C}} \phi(h) \xrightarrow{\phi_{g,h}^2} \phi(gh), \quad g, h \in G,$$

where ϕ^2 is part of the monoidal structure of ϕ . According to the discussion below (5.3.1), this monoidal structure is exactly the tensor product of $\mathcal{C}_G^\times$. Similarly, the higher morphisms of the monoidal structure of ϕ induce the associator and G -crossed braiding of $\mathcal{C}_G^\times$. Therefore, the condensable algebra $\phi(\text{Vec}_G) \in \Sigma\mathcal{C}$ is exactly the G -crossed braided fusion category $\mathcal{C}_G^\times$. $\square$

Proposition° 5.3.4. Condensing the condensable E_1 -algebra $\mathcal{C}_G^\times$ in $\Sigma\mathcal{C}$ produces the condensed phase given by $\Sigma\mathcal{M}$ (where $\mathcal{M} = (\mathcal{C}_G^\times)^G$), i.e., we have a canonical monoidal equivalence

$$\text{BMod}_{\mathcal{C}_G^\times|\mathcal{C}_G^\times}(\Sigma\mathcal{C}) \simeq \Sigma\mathcal{M}. \quad (5.3.3)$$

Proof. Let $A := \text{Fun}(G) \in \text{Rep}(G) \subset \mathcal{M}$. We have $\mathcal{C}_G^\times \simeq \mathcal{M}_G = \text{RMod}_A(\mathcal{M})$. Moreover, we have $\mathcal{C} \simeq \text{Mod}_A^{E_2}(\mathcal{M})$ [Kir01, M  g04] (see also [DGNO10, Proposition 4.55]). Since $\mathcal{M}$ is nondegenerate, by [DMNO13, Corollary 3.30], we obtain a canonical braided equivalence:

$$\mathfrak{Z}_1(\mathcal{C}_G^\times) \simeq \mathcal{C} \boxtimes \overline{\mathcal{M}},$$

which implies that $\mathfrak{Z}_2(\mathcal{C}, \mathfrak{Z}_1(\mathcal{C}_G^\times)) \simeq \overline{\mathcal{M}}$. Using (3.3.5) and (2.3.13), we obtain a canonical equivalence of fusion 2-categories $\text{Mod}_{\mathcal{C}_G^\times}^{E_1}(\Sigma\mathcal{C})^{\text{op}} \simeq \mathfrak{Z}_1(\Sigma\mathcal{C}, \mathfrak{Z}_0(\text{RMod}_{\mathcal{C}_G^\times}(\Sigma\mathcal{C}))) \simeq \Sigma\mathfrak{Z}_2(\mathcal{C}, \mathfrak{Z}_1(\mathcal{C}_G^\times)) \simeq \Sigma\overline{\mathcal{M}}$. $\square$

Remark 5.3.5. Above proposition simply says that gauging the G -symmetry in a 2+1D G -SET order defined by the equivariantization of a G -crossed braided fusion 1-category $\mathcal{C}_G^\times$ is equivalent to gauging the symmetry $2\text{Vec}_G \rightarrow \Sigma\mathcal{C}$ in the setting of (5.3.2) for $\mathcal{T} = 2\text{Vec}_G$, $\mathcal{K} = 2\text{Vec}$ and $A = \text{Vec}_G \in \text{Alg}_{E_1}^c(2\text{Vec}_G)$. $\diamond$

Remark 5.3.6. A generalized 0-from G -symmetry in an n +1D topological order C^{n+1} can be defined by a monoidal functor $n\text{Vec}_G \rightarrow \mathcal{C}$. By [LYW24], gauging it amounts to the gauging process in the case $\mathcal{T} = n\text{Vec}_G$, $\mathcal{K} = n\text{Vec}$ and $A = (n-1)\text{Vec}_G$, where the $\mathcal{T}$ -action on $\mathcal{K}$ is defined by the forgetful functor $f : n\text{Vec}_G \rightarrow n\text{Vec} \simeq \text{Fun}(n\text{Vec}, n\text{Vec})$. $\diamond$

Remark 5.3.7. Recently, we were informed by Nils Carqueville that the relation between G -crossed extensions and orbifold TQFT's was also independently established in [CH25, HPRW25]. $\diamond$

5.3.2 General constructions and examples

Example 5.3.8. A G -SPT (symmetry protected topological) order is a G -SET order with a trivial topological order. Thus 2+1D G -SPT orders are classified by the G -crossed extensions of Vec , which are Vec_G^ω for $\omega \in H^3(G; \mathbb{C}^\times)$ [DGN010, Proposition 4.61]. The physical meaning of Vec_G^ω is that, the only simple g -defect is the flux g for every $g \in G$, and the associators of the fusion of G -fluxes are defined by ω .

The equivariantization gives $\text{Vec}^G = \text{Rep}(G)$ and $(\text{Vec}_G^\omega)^G \simeq \mathfrak{Z}_1(\text{Vec}_G^\omega)$. Note that $(\text{Vec}_G^\omega)^G$ for $\omega \in H^3(G; \mathbb{C}^\times)$ are precisely all the minimal nondegenerate extensions of $\text{Rep}(G)$.

The G -crossed extensions of Vec , i.e., Vec_G^ω , are condensable E_1 -algebras in $2\text{Vec} = \Sigma\text{Vec}$. By condensing Vec_G^ω in the trivial 2+1D topological order, we obtain a non-chiral topological order:

$$\text{Mod}_{\text{Vec}_G^\omega}^{E_1}(2\text{Vec}) \simeq \Sigma \mathfrak{Z}_1(\text{Vec}_G^\omega) \simeq \mathfrak{Z}_0(\Sigma \text{Vec}_G^\omega) \simeq \Sigma(\text{Vec}_G^\omega)^G.$$

Note that the forgetful functor $f : 2\text{Vec}_G \rightarrow 2\text{Vec}$ is a monoidal functor. We can twist its monoidal structure by defining the hexagonator by a non-trivial 3-cocycle $\omega \in Z^3(G, \mathbb{C}^\times)$. We denote the resulting monoidal functor by $f^\omega : 2\text{Vec}_G \rightarrow 2\text{Vec}$. It is easy to check that f^ω maps $\text{Vec}_G \in \text{Alg}_{E_1}^c(2\text{Vec}_G)$ to $\text{Vec}_G^\omega \in \text{Alg}_{E_1}^c(2\text{Vec})$. Therefore, gauging the G -symmetry in the G -crossed extension Vec_G^ω of Vec is equivalent to gauging the 2Vec_G -symmetry defined by the monoidal functor $f^\omega : 2\text{Vec}_G \rightarrow 2\text{Vec}$. $\heartsuit$

Example 5.3.9. Suppose $\mathcal{A}$ is a fusion 1-category and $\mathcal{C} = \mathfrak{Z}_1(\mathcal{A})$. Physically this means that $\mathcal{C}$ is the category of anyons in a 2+1D topological order that has a gapped boundary and the boundary particles form a fusion 1-category $\mathcal{A}$. Note that there is an equivalence of 3-groups

$$\text{BrPic}(\mathcal{A}) = \text{BMod}_{\mathcal{A}|\mathcal{A}}(2\text{Vec})^\times \simeq \text{RMod}_{\mathfrak{Z}_1(\mathcal{A})}(2\text{Vec})^\times \simeq \text{Pic}(\mathfrak{Z}_1(\mathcal{A})),$$

where $\text{BrPic}(\mathcal{A})$ is the 3-group of invertible $\mathcal{A}$ - $\mathcal{A}$ -bimodules [ENO10]. More explicitly, this equivalence sends an invertible $\mathcal{A}$ - $\mathcal{A}$ -bimodule $\mathcal{X}$ to its ‘center’ $\mathfrak{Z}_\mathcal{A}(\mathcal{X}) \simeq \text{Fun}_{\mathcal{A}|\mathcal{A}}(\mathcal{A}, \mathcal{X})$ as defined in [GNN09]. The physical meaning of this equivalence was explained in [KLWZZ20b, Figure 4]. Then a G -crossed extension of $\mathfrak{Z}_1(\mathcal{A})$ is equivalent to a 3-group homomorphism $G \rightarrow \text{BrPic}(\mathcal{A})$, which by [ENO10] is equivalent to a G -graded extension of $\mathcal{A}$:

$$\bigoplus_{g \in G} \mathcal{A}_g, \quad \mathcal{A}_e = \mathcal{A}.$$

The category $\mathcal{A}_g$ consists of the g -defects living on the junction between the invertible domain wall realizing the equivalence ρ_g and the boundary (see Figure 16). The corresponding G -crossed extension of $\mathfrak{Z}_1(\mathcal{A})$ is

$$\mathfrak{Z}_\mathcal{A}(\bigoplus_{g \in G} \mathcal{A}_g) \simeq \text{Fun}_{\mathcal{A}|\mathcal{A}}(\mathcal{A}, \bigoplus_{g \in G} \mathcal{A}_g) \simeq \bigoplus_{g \in G} \mathfrak{Z}_\mathcal{A}(\mathcal{A}_g) \simeq \mathfrak{Z}_1(\mathcal{A}, \bigoplus_{g \in G} \mathcal{A}_g),$$

where $\mathfrak{Z}_1(\mathcal{A}, \bigoplus_{g \in G} \mathcal{A}_g)$ is the E_1 -centralizer (recall Definition 2.3.17). Its G -crossed braided structure was defined in [GNN09, Theorem 3.3]. Moreover, by [GNN09, Theorem 3.5], the equivariantization of the G -crossed braided fusion category $\mathfrak{Z}_1(\mathcal{A}, \bigoplus_{g \in G} \mathcal{A}_g)$ is precisely the Drinfeld center $\mathfrak{Z}_1(\bigoplus_{g \in G} \mathcal{A}_g)$, i.e.,

$$\mathfrak{Z}_1(\mathcal{A}, \bigoplus_{g \in G} \mathcal{A}_g)^G \simeq \mathfrak{Z}_1(\bigoplus_{g \in G} \mathcal{A}_g).$$

Physically, this means that the topological defects on the boundary after gauging the symmetry are all symmetry defects on the boundary. We summarize various relations in the following diagram.

$$\begin{array}{ccccc}
& \mathcal{A} & \xrightarrow{\text{G-graded extension}} & \bigoplus_{g \in G} \mathcal{A}_g & \\
& \downarrow \mathfrak{Z}_1(\mathcal{A}, -) & & \downarrow \mathfrak{Z}_1(\mathcal{A}, -) & \\
\text{Vec} & \xrightarrow{\text{inclusion}} & \mathfrak{Z}_1(\mathcal{A}) & \xrightarrow{\text{G-crossed extension}} & \mathfrak{Z}_1(\mathcal{A}, \bigoplus_{g \in G} \mathcal{A}_g) \\
\downarrow (-)^G \uparrow (-)_G & & \downarrow (-)^G \uparrow (-)_G & & \downarrow (-)^G \uparrow (-)_G \\
\text{Rep}(G) & \xrightarrow{\text{inclusion}} & (\mathfrak{Z}_1(\mathcal{A}))^G & \xrightarrow{\text{minimal nondegenerate extension}} & \mathfrak{Z}_1(\bigoplus_{g \in G} \mathcal{A}_g)
\end{array}$$





Figure 16: The invertible domain wall realizing the braided equivalence $\rho_g : \mathcal{C} \rightarrow \mathcal{C}$ can end on the boundary. The topological defects living on the junction are called the g -defects on the boundary, and they form the category $\mathcal{A}_g$.

The 3-group homomorphism $G \rightarrow \text{BrPic}(\mathcal{A})$ extends to a monoidal 2-functor

$$\phi : 2\text{Vec}_G \rightarrow \text{BMod}_{\mathcal{A}|\mathcal{A}}(2\text{Vec})^{\text{op}} \simeq \Sigma \mathcal{Z}_1(\mathcal{A}).$$

It maps $\text{Vec}_G \in \text{Alg}_{\mathbb{E}_1}^c(2\text{Vec}_G)$ to a condensable $\mathbb{E}_1$ -algebra $\phi(\text{Vec}_G)$ in $\Sigma \mathcal{Z}_1(\mathcal{A})$, which is precisely $\mathcal{Z}_1(\mathcal{A}, \bigoplus_g \mathcal{A}_g)$. Using the coordinate transformation between $\text{BMod}_{\mathcal{A}|\mathcal{A}}(2\text{Vec})^{\text{op}}$ and $\Sigma \mathcal{Z}_1(\mathcal{A})$ (recall Theorem^{ph} 3.3.32),



Two definitions of the same algebra in $\mathcal{C}$:

- (1) the monoidal functor $\mathcal{A} \rightarrow \bigoplus_g \mathcal{A}_g$ defines $(\bigoplus_g \mathcal{A}_g)^{\text{op}} \in \text{Alg}_{\mathbb{E}_1}^c(\text{BMod}_{\mathcal{A}|\mathcal{A}}(2\text{Vec})^{\text{op}})$;
- (2) the central functor $\mathcal{Z}_1(\mathcal{A}) \rightarrow \text{Fun}_{\mathcal{A}|\oplus_g \mathcal{A}_g}(\oplus_g \mathcal{A}_g, \oplus_g \mathcal{A}_g)^{\text{op}}$ defines $\mathcal{Z}_1(\mathcal{A}, \oplus_g \mathcal{A}_g)^{\text{op}} \in \text{Alg}_{\mathbb{E}_1}^c(\text{RMod}_{\mathcal{Z}_1(\mathcal{A})}(2\text{Vec}))$,

where we have used the fact that $\mathcal{Z}_1(\mathcal{A}, \oplus_g \mathcal{A}_g) \simeq \text{Fun}_{\mathcal{A}|\oplus_g \mathcal{A}_g}(\oplus_g \mathcal{A}_g, \oplus_g \mathcal{A}_g)$ as fusion 1-categories. In other words, $\phi(\text{Vec}_G) = (\bigoplus_{g \in G} \mathcal{A}_g)^{\text{op}} \in \text{Alg}_{\mathbb{E}_1}^c(\text{BMod}_{\mathcal{A}|\mathcal{A}}(2\text{Vec})^{\text{op}})$. By condensing $\phi(\text{Vec}_G)$ in $\text{BMod}_{\mathcal{A}|\mathcal{A}}(2\text{Vec})^{\text{op}}$ or $\Sigma \mathcal{Z}_1(\mathcal{A})$, we obtain the condensed topological order determined by $\mathcal{Z}_1(\bigoplus_{g \in G} \mathcal{A}_g)$. $\heartsuit$

Example 5.3.10. This example explains the well-known statement that the double Ising topological order can be obtained from the toric code model by gauging a 0-form $\mathbb{Z}_2$ -symmetry called the e - m -duality (see also [BBCW19, Section X.G.2]) as a special case of Example 5.3.9.

There is a $\mathbb{Z}_2$ -action on the toric code model given by exchanging e and m . This gives a 2-group homomorphism $\mathbb{Z}_2 \rightarrow \text{Aut}^{\mathbb{E}_2}(\mathcal{Z}_1(\text{Rep}(\mathbb{Z}_2)))$. An $\mathbb{Z}_2$ -SET order lifting this symmetry action, by Example 5.3.9, is equivalent to a $\mathbb{Z}_2$ -graded extension of $\text{Rep}(\mathbb{Z}_2)$:

$$\mathcal{A}_0 \oplus \mathcal{A}_1, \quad \mathcal{A}_0 = \text{Rep}(\mathbb{Z}_2).$$

By considering the 2-group homomorphism $\mathbb{Z}_2 \rightarrow \text{Aut}^{\mathbb{E}_2}(\mathcal{Z}_1(\text{Rep}(\mathbb{Z}_2))) \simeq \Pi_2(\text{BrPic}(\text{Rep}(\mathbb{Z}_2)))$, it is not hard to see that $\mathcal{A}_1 \simeq \text{Vec}$. Physically, $\mathcal{A}_1$ consists of the topological defects living on the junction between the e - m -exchange domain wall and the rough boundary of the toric code model. Thus we can also check that $\mathcal{A}_1 \simeq \text{Vec}$ in the concrete lattice model (see for example [KZ22a, Section 4.4.3]).

An $\mathbb{Z}_2$ -graded extension $\text{Rep}(\mathbb{Z}_2) \oplus \text{Vec}$ of $\text{Rep}(\mathbb{Z}_2)$ is the same as an Ising type fusion 1-category, that is, a fusion 1-category with three simple objects $\mathbb{1}, \psi, \sigma$ and the fusion rule

$$\psi \otimes \psi \simeq \mathbb{1}, \quad \psi \otimes \sigma = \sigma = \sigma \otimes \psi, \quad \sigma \otimes \sigma = \mathbb{1} \oplus \psi.$$

By [TY98] (see also [DGNO10, Section B.1] or [ENO10, Example 9.4]), there are two Ising type fusion 1-categories up to equivalence, denoted by $\text{Is}_{\pm}$. They differ in the definition of the associators⁶¹. According to the discussion in Example 5.3.9, the category of particle-like topological defects in the gauged theory is $\mathcal{Z}_1(\text{Is}_{\pm})$. It also means that there are two 3-group homomorphisms $\phi_{\pm} : G \rightarrow \text{BrPic}(\text{Rep}(\mathbb{Z}_2))$. Figure 17 depicts the physical relation between the $\mathbb{Z}_2$ and double Ising topological orders.

⁶¹Note that $\text{Is}_{\pm} = \Omega \mathcal{I}$ s as fusion 1-categories.




Figure 17: The physical relation between the $\mathbb{Z}_2$ and double Ising topological orders.

By Example 5.3.9, we know that the $\mathbb{Z}_2$ -crossed extensions of $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$ are given by

$$\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2), \text{Is}_\pm) \simeq \text{Fun}_{\text{Rep}(\mathbb{Z}_2)|\text{Is}_\pm}(\text{Is}_\pm, \text{Is}_\pm),$$

which can be computed directly. On the other hand, it can also be determined directly by the fact that it is precisely the de-equivariantization of $\mathfrak{Z}_1(\text{Is}_\pm)$. This de-equivariantization can be determined as follows. By [DGN010, Section B.3], there are 8 Ising type braided fusion categories Is_ζ up to equivalence, labeled by complex numbers ζ satisfying $\zeta^8 = -1$, and both of them are nondegenerate. The underlying fusion category of Is_ζ is determined by the sign $\pm$ in the equation $\zeta^2 + \zeta^{-2} = \pm\sqrt{2}$. Therefore, there are braided equivalences $\text{Is}_\zeta \boxtimes \overline{\text{Is}_\zeta} \simeq \mathfrak{Z}_1(\text{Is}_\pm)$ for some choices of ζ . After taking the equivariantization, we obtain two minimal nondegenerate extensions:

$$\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))^{\mathbb{Z}_2} \hookrightarrow \mathfrak{Z}_1(\text{Is}_\pm) = \text{Is}_\zeta \boxtimes \overline{\text{Is}_\zeta}$$

The simple objects of $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))^{\mathbb{Z}_2}$ in $\text{Is}_\zeta \boxtimes \overline{\text{Is}_\zeta}$ are

$$\mathbb{1} \boxtimes \mathbb{1}, \quad \mathbb{1} \boxtimes \psi, \quad \psi \boxtimes \mathbb{1}, \quad \psi \boxtimes \psi, \quad \sigma \boxtimes \sigma.$$

More precisely, we have

$$\langle \mathbb{1} \boxtimes \mathbb{1}, \psi \boxtimes \psi \rangle = \langle \mathbb{1} \rangle^{\mathbb{Z}_2} \simeq \text{Rep}(\mathbb{Z}_2), \quad \langle \mathbb{1} \boxtimes \psi, \psi \boxtimes \mathbb{1} \rangle = \langle f \rangle^{\mathbb{Z}_2}, \quad \langle \sigma \boxtimes \sigma \rangle = \langle e, m \rangle^{\mathbb{Z}_2},$$

where $\langle x_i \rangle$ denotes the full subcategory consisting of the direct sums of the simple objects x_i . Moreover, notice that $\mathfrak{Z}_2(\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))^{\mathbb{Z}_2}) = \langle \mathbb{1} \boxtimes \mathbb{1}, \psi \boxtimes \psi \rangle \simeq \text{Rep}(\mathbb{Z}_2)$. Then the de-equivariantization of $\mathfrak{Z}_1(\text{Is}_\pm)$ is equivalent to take the modules over the condensable E_2 -algebra

$$A := \mathbb{1} \boxtimes \mathbb{1} \oplus \psi \boxtimes \psi = \text{Fun}(\mathbb{Z}_2) \in \text{Rep}(\mathbb{Z}_2) \hookrightarrow \text{Is}_\zeta \boxtimes \overline{\text{Is}_\zeta}.$$

Recall that this algebra defines the well-known anyon condensation from the double Ising topological order to the $\mathbb{Z}_2$ topological order. We obtain two $\mathbb{Z}_2$ -crossed extensions of $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$:

$$\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2), \text{Is}_\pm) \simeq (\mathfrak{Z}_1(\text{Is}_\pm))_G \simeq \text{RMod}_A(\text{Is}_\zeta \boxtimes \overline{\text{Is}_\zeta}) \simeq \langle \mathbb{1}, e, m, f \rangle \oplus \langle \chi_+, \chi_- \rangle, \quad (5.3.4)$$

which was shown in [CJKYZ20, Section 3.2]⁶² to consist of 4 simple local A -modules $\mathbb{1}, e, m, f$ generating the trivial graded component (which is equivalent to $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$), and 2 simple non-local A -modules $\chi_\pm$ generating the nontrivial graded component.

The two 3-group homomorphisms $G \rightarrow \text{BrPic}(\text{Rep}(\mathbb{Z}_2))$ naturally induce two monoidal 2-functors

$$\phi_\pm : 2\text{Vec}_{\mathbb{Z}_2} \rightarrow \text{BMod}_{\text{Rep}(\mathbb{Z}_2)|\text{Rep}(\mathbb{Z}_2)}(2\text{Vec})^{\text{op}} \simeq \Sigma \mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2)).$$

The image of $\phi_\pm(\text{Vec}_{\mathbb{Z}_2})$ are condensable E_1 -algebras in $\mathcal{TC}$. In the coordinate system $\Sigma \mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$, they are precisely the two $\mathbb{Z}_2$ -crossed extensions of $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$, explicitly computed in (5.3.4). In the coordinate system $\text{BMod}_{\text{Rep}(\mathbb{Z}_2)|\text{Rep}(\mathbb{Z}_2)}(2\text{Vec})^{\text{op}}$, they are precisely given by $\text{Is}_\pm$ as illustrated in Figure 17. By condensing $\phi_\pm(\text{Vec}_{\mathbb{Z}_2})$, we obtain the fusion 2-category $\Sigma \mathfrak{Z}_1(\text{Is}_\pm)$ associated to the condensed phases. Therefore, gauging the e - m -duality (i.e., a 0-form symmetry) in the 2+1D $\mathbb{Z}_2$ topological order is equivalent to gauging the $2\text{Vec}_{\mathbb{Z}_2}$ -symmetry in $\mathcal{TC} = \Sigma \mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$. $\heartsuit$

⁶²Indeed, [CJKYZ20, Section 3.2] only consider the case $\zeta = \exp(-\pi i/8)$, but there is no essential difference for other choices of ζ .

5.3.3 Generalizations, examples and remarks

In this subsection, we revisit an example of 1-form symmetry with 't Hooft anomaly and provide an example of gauging a 0-form symmetry in 4D. We remark on further generalizations of the notions of a symmetry and gauging.

Example 5.3.11. Let A be a finite abelian group. Its delooping BA is a 2-group, describing a 1-form symmetry. Let $\mathcal{C}$ be the modular tensor category of anyons in a 2+1D topological order. Then a 1-form symmetry BA on this topological order is a monoidal 2-functor $\phi : BA \rightarrow \Sigma\mathcal{C}$ or $\phi : 2\text{Vec}_{BA} \simeq 2\text{Rep}(\hat{A}) \rightarrow \Sigma\mathcal{C}$, where $\hat{A} := \text{Hom}(A, U(1))$. A monoidal 2-functor $\phi : BA \rightarrow \Sigma\mathcal{C}$ is the same as a braided functor $\phi : A \rightarrow \mathcal{C}$, which maps each $a \in A$ to an abelian anyon (invertible object in $\mathcal{C}$). Moreover, since the braiding on A is trivial, the anyons $\phi(a)$ have trivial double braiding with each other. Therefore, $E := \bigoplus_{a \in A} \phi(a) \in \mathcal{C}$ is a condensable E_2 -algebra.

Gauging this 1-form symmetry is the same as condensing the algebra $\phi(\text{Vec}_{BA}) \in \Sigma\mathcal{C}$. Here Vec_{BA} is the condensation completion (or Karoubi completion) of $\text{Bk}[A]$. In other words, Vec_{BA} can be obtained by condensing $\mathbb{k}[A]$ from the tensor unit $\text{Vec} \in 2\text{Vec}_{BA}$. Thus $\phi(\text{Vec}_{BA}) \in \Sigma\mathcal{C}$ can be obtained by condensing $\phi(\mathbb{k}[A]) = \bigoplus_{a \in A} \phi(a) = E$ from the tensor unit of $\Sigma\mathcal{C}$. In other words, $\phi(\text{Vec}_{BA}) = \Sigma E$, which is the 1+1D domain wall obtained by condensing the condensable E_2 -algebra E along a line. Hence gauging this 1-form symmetry is the same as the 2-codimensional condensation (i.e., anyon condensation) of E .

In literature, a subgroup A of abelian anyons with nontrivial braiding is also called a 1-form symmetry with 't Hooft anomaly (see for example [GKSW15]). In this case, these abelian anyons span a fusion subcategory of $\mathcal{C}$, and its associator and braiding determine a class in the abelian cohomology (i.e., E_2 Eilenberg-MacLane cohomology) $\pi \in H_{E_2}^4(A; \mathbb{C}^\times)$. This fusion subcategory is denoted by Vec_A^π . On the other hand, 2Vec_{BA} can also be twisted by $\pi \in H_{E_2}^4(BA; \mathbb{C}^\times) \simeq H_{E_2}^4(A; \mathbb{C}^\times)$. Then this anomalous 1-form symmetry is a monoidal 2-functor $\phi : 2\text{Vec}_{BA}^\omega \simeq \Sigma\text{Vec}_A^\omega \rightarrow \Sigma\mathcal{C}$.

When π is nontrivial, Vec_{BA} is not a condensable algebra in 2Vec_{BA}^π because there is no fiber 2-functor $2\text{Vec}_{BA}^\pi \rightarrow 2\text{Vec}$. So the BA -symmetry cannot be gauged. We may say that the nontrivial class $\pi \in H_{E_2}^4(A; \mathbb{C}^\times)$ characterizes the 't Hooft anomaly of this 1-form symmetry. $\heartsuit$

Example 5.3.12. The codimension-2 topological defects in the 3+1D toric code model form a E_2 -fusion 2-category $\mathcal{C} := \mathfrak{Z}_1(2\text{Rep}(\mathbb{Z}_2))$ [KTZ20a]. There is an invertible domain wall in the 3+1D toric code model corresponding to the nontrivial 2+1D $\mathbb{Z}_2$ SPT order [KLWZZ20b, Theorem^{ph} 3.34 and Figure 5]. Also, by [KLWZZ20b, Theorem^{ph} 3.37], this invertible domain wall in the coordinate system

$$\Sigma\mathcal{C} \simeq \text{BMod}_{2\text{Rep}(\mathbb{Z}_2)|2\text{Rep}(\mathbb{Z}_2)}(3\text{Vec})$$

is $2\text{Rep}(\mathbb{Z}_2, \omega) = \Sigma\text{Vec}_{\mathbb{Z}_2}^\omega$, where $\omega \in H^3(\mathbb{Z}_2; \mathbb{C}^\times) \simeq \mathbb{Z}_2$ is the nontrivial class.

This invertible domain wall has order 2. It is natural to ask whether this invertible domain wall can be promoted to $\mathbb{Z}_2$ -symmetry. If such a $\mathbb{Z}_2$ -symmetry exists, it is a monoidal 3-functor $\phi : 3\text{Vec}_{\mathbb{Z}_2} \rightarrow \Sigma\mathcal{C}$ which maps the nontrivial simple object to the above invertible domain wall. In the coordinate system $\Sigma\mathcal{C} \simeq \text{BMod}_{2\text{Rep}(\mathbb{Z}_2)|2\text{Rep}(\mathbb{Z}_2)}(3\text{Vec})$, it maps the tensor unit of $3\text{Vec}_{\mathbb{Z}_2}$ to $2\text{Rep}(\mathbb{Z}_2)$ and the nontrivial simple object to $2\text{Rep}(\mathbb{Z}_2, \omega)$. Denote $\mathcal{P}_0 := 2\text{Rep}(\mathbb{Z}_2)$ and $\mathcal{P}_1 := 2\text{Rep}(\mathbb{Z}_2, \omega)$. The monoidal structure of ϕ is equivalent to a $\mathbb{Z}_2$ -graded fusion 2-category structure on $\mathcal{P} = \mathcal{P}_0 \oplus \mathcal{P}_1$.

By the obstruction theory (similar to [ENO10]), such $\mathbb{Z}_2$ -graded fusion 2-categories $\mathcal{P}$ form a torsor over $H^4(\mathbb{Z}_2; \mathbb{C}^\times) = 0$, hence are unique. It is also known that there is a 2-group $\mathcal{G}$ defined by $\pi_1(\mathcal{G}) = \pi_2(\mathcal{G}) = \mathbb{Z}_2$ and the Postnikov class (associator) $\omega \in H^3(\mathbb{Z}_2; \mathbb{Z}_2) \simeq H^3(\mathbb{Z}_2; \mathbb{C}^\times) \simeq \mathbb{Z}_2$ such that $\mathcal{P} = 2\text{Rep}(\mathcal{G})$ (see for example [HZ23, Example 3.36]).

By gauging this $\mathbb{Z}_2$ -symmetry, we obtain a 3+1D topological order with the fusion 3-category of topological defects being

$$\text{BMod}_{\mathcal{P}|\mathcal{P}}(\text{BMod}_{2\text{Rep}(\mathbb{Z}_2)|2\text{Rep}(\mathbb{Z}_2)}(3\text{Vec})) \simeq \text{BMod}_{\mathcal{P}|\mathcal{P}}(3\text{Vec}) \simeq \Sigma\mathfrak{Z}_1(\mathcal{P}).$$

By [DY25, Proposition 4.2.1], $\mathcal{P} = 2\text{Rep}(\mathcal{G})$ is Morita equivalent to $2\text{Vec}_{\mathbb{Z}_2 \times \mathbb{Z}_2}^\pi$, where the 4-cocycle $\pi \in H^4(\mathbb{Z}_2 \times \mathbb{Z}_2; \mathbb{C}^\times)$ is defined by

$$\pi((a_1, b_1), (a_2, b_2), (a_3, b_3), (a_4, b_4)) := \omega(a_2, a_3, a_4)^{b_1}.$$

Then $\mathfrak{Z}_1(\mathcal{P})$ is braided equivalent to $\mathfrak{Z}_1(2\text{Vec}_{\mathbb{Z}_2 \times \mathbb{Z}_2}^\pi)$. Hence the gauged theory is $\text{GT}_{(\mathbb{Z}_2 \times \mathbb{Z}_2, \pi)}^4$, i.e., the 3+1D π -twisted $\mathbb{Z}_2 \times \mathbb{Z}_2$ gauge theory. $\heartsuit$

Remark 5.3.13. It seems that the framework of formulating an SET order as a monoidal functor also works for continuous group G symmetry, but a monoidal n -functor $G \rightarrow \mathcal{C}^\times$ should be understood as a pointed continuous map $BG \rightarrow B\mathcal{C}^\times$, where B denotes the classifying space. For example, consider the 2+1D trivial phase equipped with a compact group G symmetry. The space of invertible topological defects in the 2+1D trivial phase is $\mathcal{C}^\times \simeq B^2\mathcal{C}^\times \simeq B^3\mathbb{Z} = K(\mathbb{Z}, 3)$. The set of homotopy classes of pointed maps $BG \rightarrow B\mathcal{C}^\times \simeq K(\mathbb{Z}, 4)$ is $H^4(BG, \mathbb{Z})$, which exactly classifies the 2+1D Chern-Simons G -gauge theory [DW90]. In other words, the 2+1D SPT orders with a compact group G symmetry should be classified by $H^4(BG, \mathbb{Z})$, and their gauged theories are the Chern-Simons theories. However, the categorical description of the gauging (or condensation) process in this case is not clear. $\diamond$

Remark 5.3.14. Let G be a finite group. Given an $n+1$ D G -SET order described by a monoidal n -functor $\phi : n\text{Vec}_G \rightarrow \mathcal{C}$, its partition functions can be constructed as follows. A spacetime manifold is an $(n+1)$ -dimensional closed manifold M equipped with a principal G -bundle, or equivalently, a continuous map $\gamma : M \rightarrow BG$. If we fix a triangulation on M , then the map γ is equivalent to a flat G -connection, that is, an assignment to each 1-simplex of M a group element such that the holonomy on each 2-simplex vanishes. The monoidal n -functor ϕ is equivalent to a continuous map $B\phi : BG \rightarrow B\mathcal{C}^\times$. The composition of these two maps $(B\phi) \circ \gamma : M \rightarrow B\mathcal{C}^\times$ assigns

- to each 1-simplex e of M the codimension-1 domain wall $\phi(\gamma(e)) \in \mathcal{C}$;
- to each 2-simplex of M the codimension-2 topological defect corresponding to the monoidal structure of ϕ ;
- to each $(n+1)$ -simplex of M a codimension- $(n+1)$ topological defect (a number in $\mathcal{C}^\times$).

This is a $\mathcal{C}$ -colored graph on M , and the monoidal structure of ϕ ensures that this graph is invariant under the Pachner moves. Therefore, this graph can be evaluated to a number using the graph calculus in $\mathcal{C}$ because M is closed. This number is defined to be the partition function $Z(M, \gamma)$. For the gauged theory, the partition function need to sum over all possible G -bundles on M :

$$Z_{\text{gauged}}(M) := \frac{1}{|G|} \sum_{\gamma} Z(M, \gamma).$$

This partition function is the same as a graph in M whose 1-simplices are labeled by the algebra $A := \phi((n-1)\text{Vec}_G) \in \mathcal{C}$, 2-simplices are labeled by the multiplication of A , and higher simplices are labeled by the higher associativity constraints of A .

This construction generalizes the Dijkgraaf-Witten theory [DW90] and the decorated domain wall construction of SPT wave functions [CLV14], and should be a special case of the orbifold construction of defect TQFT [CRS19]. $\diamond$

Remark 5.3.15. Corollary^{ph} 3.3.4 can also be interpreted as a process of gauging the fusion n -category symmetry $\mathcal{C}$. One way to gauge the symmetry is to map $\mathcal{C}$ into a indecomposable multi-fusion n -category $\mathcal{X}$ that is ‘anomaly-free’ in the sense that it is the category of topological defects of an anomaly-free $n+1$ D topological order X^{n+1} . Since this idea is originated from the idea of minimal modular extensions (MME) [LKW16a, LKW16b, KLWZZ20b], as a way of gauging categorical symmetries, we call this gauging an MME-gauging. When $X^{n+1} = Z(M)^{n+1}$, an MME-gauging can be characterized by a monoidal functor $f : \mathcal{C} \rightarrow \mathcal{X} = \mathfrak{Z}_0(\mathcal{M})$, or equivalently, a left $\mathcal{C}$ -module $\mathcal{M}$. By adding a refined data, i.e., a distinguished object $m \in \mathcal{M}$, we provide a wall condition to $\mathcal{C}$ in the sense that the way how defects in C^{n+1} move onto the wall M^n is specified by the wall condition $m \in \mathcal{M}$. As a consequence, the pair $M^n = (\mathcal{M}, m)$ is an E_0 -algebra over the E_1 -algebra $\mathcal{C}$. In this way, we recover the condensed phase D^{n+1} as the centralizer $\mathcal{D} = \mathfrak{Z}_1(f) = \mathfrak{Z}_1(\mathcal{C}, \mathfrak{Z}_0(\mathcal{M}))$ of the monoidal functor $f : \mathcal{C} \rightarrow \mathfrak{Z}_0(\mathcal{M})$ as shown in Theorem^{ph} 3.3.9. For convenience, we call $Z(M)^{n+1}$ the MME-gauged phase and $Z(f)^{n+1} = D^{n+1}$ the condensed phase. The MME-gaugings of a finite onsite symmetry in 1+1D SPT orders were studied in [KLWZZ20b, Section 2.2.1]. More general MME-gauging theory for SPT/SET orders with finite internal symmetries were developed in [KZ24].

By the physical realization of a monoidal functor [KWZ15, KZ18a, KZ24] and by allowing M^n to be gapless, we can recover all possible MME-gaugings, which can be viewed as a generalized condensation (see an example in Theorem^{ph} 6.3.4 and more discussion in Section 6.3.2). We come back to this MME-gauging in Remark 6.3.3, in which we briefly mention an even more general type of MME-gauging such that both the MME-gauged phase and the condensed phase are gappable gapless. $\diamond$

Remark 5.3.16. Further generalizations of the notion of a symmetry are possible.

- (1) According to the mathematical theory of gapped/gapless quantum liquids [KZ18a, KZ20, KZ21, KZ22b, KZ22c], the topological skeleton X_{top}^{n+1} of an n D gapped/gapless quantum liquid X^n is determined by a generalized quiche (B^{n+1}, Q^n) such that the fusion n -category $\mathcal{Q}$ is equipped with a braided monoidal functor $\varphi : \Omega\mathcal{B}^{\text{op}} \rightarrow \mathfrak{Z}_1(\mathcal{Q})$. The pair becomes an ordinary quiche when φ is an equivalence.

The notion of a generalized symmetry can be defined as a morphism between pairs $(\mathcal{A}, \mathcal{P}) \rightarrow (\mathcal{B}, \mathcal{Q})$, or equivalently, a morphism between two ‘quiches’ $(A^{n+1}, P^n) \rightarrow (B^{n+1}, Q^n)$ by a ‘generalized sandwich construction’ as depicted below:

$$\begin{array}{c}
 \begin{array}{c}
 \text{Z(B)}^{n+2} \quad \xrightarrow{D^{n+1}} \quad \begin{array}{c} \text{F}^{n+1} \quad \text{Z(A)}^{n+2} \quad \text{A}^{n+1} \\ \text{G}^n \quad \text{C}^{n+1} \quad \text{P}^n \end{array} \\
 \text{G}^n \boxtimes_{\text{Z(A)}} \text{A}^n = \text{B}^n \\
 \text{G}^n \boxtimes_{\text{C}} \text{P}^n = \text{Q}^n
 \end{array}
 \end{array} \tag{5.3.5}$$

where the generalized ‘quiche’ (F^{n+1}, G^n) defines the morphism, and C^{n+1} (resp. D^{n+1}) is the unique topological order that is Morita equivalent to A^{n+1} (resp. B^{n+1}) via P^n (resp. Q^n).

When $A^{n+1} = B^{n+1} = 1^{n+1}$, by [KZ18a, KZ24], this generalized symmetry recovers the symmetry defined by a monoidal functor $\mathcal{G} : \mathcal{P} \rightarrow \mathcal{Q}$ as proposed in [KZ24, LYW24]. When $n = 1$, this ‘generalized sandwich construction’ provides a physical realization of an enriched monoidal 1-functor, a notion which was introduced in [KYZ24]. For general n , we expect this ‘generalized sandwich construction’ to provide a physical realization of an enriched monoidal higher functor.

- (2) A new type of ‘topological symmetries’, generalizing that of ‘quiche’ [FMT24], was proposed in [Kon22] as an application of topological Wick rotation. This new type of a ‘topological symmetry’ consists of a topological order, a gapped boundary (codimension 1), a gapped corner (a defect of codimension 2 living on the boundary), a higher corner (a defect of codimension > 2 living on the boundary), etc., largely generalizing the notion of a ‘quiche’ (i.e., a topological order + a gapped boundary). Maybe it can be called a “higher quiche”. A symmetry can be defined as a morphism between two generalized ‘topological symmetries’ via a ‘generalized sandwich construction’, i.e., via a topless higher dimensional cubic generalizing the topless square (with three edges labeled by F^{n+1} , C^{n+1} , A^{n+1}) in (5.3.5). This leads to an interesting relation between theories in different dimensions (differ by two or higher), i.e., a generalized topological holography [Kon22].

The theory of gauging these further-generalized symmetries is automatic or tautologically followed from our theory of defect condensations in topological orders because all the ingredients in the ‘generalized sandwich construction’ are all (potentially anomalous) topological orders. In the light of this discussion, it is beneficial to simply define a symmetry by a topological order or a quiche or a higher quiche. In particular, this generalized gauging of symmetry allows us to obtain an SPT/SET/SSB order from a topological order via gauging and vice versa. $\diamond$

6 Generalizations and Applications

In this work, we have developed a general theory of the condensations of topological defects in higher dimensions. In this section, we show that this theory leads to many new interesting questions, generalizations, applications and outlooks. We want to emphasize that this work is not an end of condensation theory but a beginning of a much richer theory in both mathematics and physics. For example, the natural questions in condensations lead us to a general higher Morita theory; the physical intuition of integrating local observables to global ones leads us to a theory generalizing the usual theory of factorization homology; the generalization to gapless liquid-like defects leads us a more complete theory of condensation. In this section, we can only scratch the surfaces of these directions but leave the thorough development to future publications.

6.1 Higher Morita theory

In this subsection, we study some questions that naturally arise from the physics of condensations. The most basic question is when two condensations produce the same condensed phase or the same condensed defect.

6.1.1 Higher Morita equivalences

Given a topological order C^{n+1} and two condensations defined by $A, B \in \text{Alg}_{E_1}^c(\mathcal{C})$, it is natural to ask when they produce the same condensed phase D^{n+1} . We formulate the question in more mathematical terms.

Definition 6.1.1. Let $\mathcal{B} \in \text{Alg}_{E_1}^c((n+1)\text{Vec})$ and $A, B \in \text{Alg}_{E_1}^c(\mathcal{B})$.

- For $\mathcal{M} \in \text{Mod}_{\mathcal{B}}^{E_1}((n+1)\text{Vec})$, A and B are called *Mod $^{E_1}(\mathcal{M})$ -equivalent* if $\text{Mod}_A^{E_1}(\mathcal{M}) \simeq \text{Mod}_B^{E_1}(\mathcal{M})$ in $\text{Alg}_{E_1}^c((n+1)\text{Vec})$. When $\mathcal{M} = \mathcal{B}$, we abbreviate this equivalence to *Mod E_1 -equivalence*.
- For $\mathcal{N} \in \text{LMod}_{\mathcal{B}}((n+1)\text{Vec})$, A and B are called *LMod $(\mathcal{N})$ -equivalent* if $\text{LMod}_A(\mathcal{N}) \simeq \text{LMod}_B(\mathcal{N})$ in $(n+1)\text{Vec}$. When $\mathcal{N} = \mathcal{B}$, we abbreviate this equivalence to *LMod-equivalence*, which is just the usual Morita equivalence. The definition of a *RMod $(\mathcal{P})$ -equivalence* for $\mathcal{P} \in \text{RMod}_{\mathcal{B}}((n+1)\text{Vec})$ is similar.

Example 6.1.2. Recall that Morita equivalent topological orders can be obtained from each other via condensations of 1-codimensional topological defects. Therefore, for fusion n -category $\mathcal{C}$ with a trivial E_1 -center, there is a one-to-one correspondence between the *Mod E_1 -equivalence* classes of simple condensable E_1 -algebras in $\mathcal{C}$ and the isomorphic classes of non-degenerate braided fusion $(n-1)$ -categories within the Witt class of $[\Omega\mathcal{C}]$. $\heartsuit$

Example 6.1.3. Recall Theorem^{ph} 3.3.32, for a fixed $\mathcal{P}$, all the condensable E_1 -algebras $\text{Fun}_{\mathcal{P}^{\text{op}}}(\mathcal{X}, \mathcal{X})$ in $\mathcal{C} = \text{BMod}_{\mathcal{B}|\mathcal{B}}(n\text{Vec})$ for $\mathcal{X} \in \text{BMod}_{\mathcal{B}|\mathcal{P}}(n\text{Vec})$ are all *Mod E_1 -equivalent*. $\heartsuit$

Proposition^o 6.1.4. If A and B are LMod-equivalence, then they are *Mod E_1 -equivalent*.

Proof. When A and B are LMod-equivalent, set $\mathcal{M} = \text{LMod}_A(\mathcal{B}) \simeq \text{LMod}_B(\mathcal{B})$. We have $\text{Mod}_A^{E_1}(\mathcal{B}) \simeq \text{Fun}_{\mathcal{B}}(\mathcal{M}, \mathcal{M})^{\text{op}} \simeq \text{Mod}_B^{E_1}(\mathcal{B})$. $\square$

Remark 6.1.5. Conversely, *Mod E_1 -equivalence* does not imply LMod-equivalence. We give an example. For $\mathcal{B} = \text{Rep}(\mathbb{Z}_2)$ and $A = \mathbb{1}, B = \mathbb{1} \oplus e \in \text{Alg}_{E_1}^c(\text{Rep}(\mathbb{Z}_2))$, we have

$$\text{LMod}_A(\text{Rep}(\mathbb{Z}_2)) \simeq \text{Rep}(\mathbb{Z}_2), \quad \text{LMod}_B(\text{Rep}(\mathbb{Z}_2)) \simeq \text{Vec}.$$

On the other hand, we have $\text{Mod}_A^{E_1}(\text{Rep}(\mathbb{Z}_2)) \simeq \text{Vec}_{\mathbb{Z}_2} \simeq \text{Fun}_{\text{Rep}(\mathbb{Z}_2)}(\text{Vec}, \text{Vec}) \simeq \text{Mod}_B^{E_1}(\text{Rep}(\mathbb{Z}_2))$. $\diamond$

We generalize the equivalences in Definition 6.1.1 to E_k -algebras.

Definition 6.1.6. Let $\mathcal{B} \in \text{Alg}_{E_k}^c((n+1)\text{Vec})$ and $A, B \in \text{Alg}_{E_k}^c(\mathcal{B})$.

1. For $\mathcal{M} \in \text{Mod}_{\mathcal{B}}^{E_k}((n+1)\text{Vec})$, A and B are called *Mod $^{E_k}(\mathcal{M})$ -equivalent* if

$$\text{Mod}_A^{E_k}(\mathcal{M}) \simeq \text{Mod}_B^{E_k}(\mathcal{M}) \in \text{Alg}_{E_k}^c((n+1)\text{Vec}).$$

When $\mathcal{M} = \mathcal{B}$, we abbreviate ‘*Mod $^{E_k}(\mathcal{B})$ -equivalent*’ to ‘*Mod E_k -equivalent*’.

2. For $l < k$ and $\mathcal{N} \in \text{LMod}_{\mathcal{B}}(\text{Alg}_{E_l}^c((n+1)\text{Vec}))$, A and B are called $\text{LMod}(\mathcal{N})$ - E_l -equivalent if

$$\text{LMod}_A(\mathcal{N}) \simeq \text{LMod}_B(\mathcal{N}) \quad \in \text{Alg}_{E_l}^c((n+1)\text{Vec}).$$

When $\mathcal{N} = \mathcal{B} \in \text{LMod}_{\mathcal{B}}(\text{Alg}_{E_{k-1}}^c((n+1)\text{Vec}))$, we abbreviate the $\text{LMod}(\mathcal{B})$ - E_{k-1} -equivalence to $\text{LMod}_{E_{k-1}}$ -equivalence.

3. Recall that $\Sigma\mathcal{B} = \text{RMod}_{\mathcal{B}}((n+1)\text{Vec}) \in \text{Alg}_{E_{k-1}}^c((n+2)\text{Vec})$ and $\Sigma A := \text{RMod}_A(\mathcal{B}) \in \text{Alg}_{E_{k-1}}^c(\Sigma\mathcal{B})$. For $k \geq p > 1$, we define $\Sigma^p A$ iteratively as $\Sigma^p A := \text{RMod}_{\Sigma^{p-1}A}(\Sigma^{p-1}\mathcal{B}) \in \text{Alg}_{E_{k-p}}^c(\Sigma^p\mathcal{B})$. Then A and B are called k -Morita equivalent if $\Sigma^k A \simeq \Sigma^k B$ in $\Sigma^k\mathcal{B}$. $\square$

Example 6.1.7. We give an example of k -Morita equivalence when $\mathcal{B} = n\text{Vec}$. We spell out the k -Morita equivalence in this case more explicitly.

- Two E_k -multi-fusion n -categories $\mathcal{A}$ and $\mathcal{B}$ are called k -Morita equivalent if $\Sigma^k \mathcal{A} \simeq \Sigma^k \mathcal{B}$ as separable $(n+k)$ -categories.

This k -Morita equivalence generalizes the usual Morita equivalence and Witt equivalence.

1. When $k = 1$, 1-Morita equivalence is precisely the usual Morita equivalence.
2. When $k = 2$ and both $\mathcal{A}$ and $\mathcal{B}$ are nondegenerate, the 2-Morita equivalence is precisely the usual Witt equivalence [JF22, KZ22b]. The advantage of the notion of 2-Morita equivalence is that it also works for (not nondegenerate) braided fusion categories.

The k -Morita equivalence is clearly a well-defined equivalence relation. We denote the k -Morita equivalence class associated to $\mathcal{A}$ by $[\mathcal{A}]_k$. $\heartsuit$

Example 6.1.8. Let $\mathcal{B}$ be a non-degenerate braided fusion n -category. Then all Lagrangian E_2 -algebras in $\mathcal{B}$ are Mod^{E_2} -equivalent by definition. It is an interesting question to work out when two condensable E_2 -algebra in $\mathcal{B}$ are Mod^{E_2} -equivalent. When $n = 1$, an interesting answer to this question can be found in [FFRS06]. A work devoted to the study of this question for $n = 1$ appeared recently [XY25]. $\heartsuit$

Remark 6.1.9. When $\mathcal{B}$ is non-degenerate, the functor $\Sigma(-) : \text{Alg}_{E_2}^c(\mathcal{B}) \rightarrow \text{Alg}_{E_1}^c(\Sigma\mathcal{B})$ induces an injection from the set of Mod^{E_2} -equivalence classes of condensable E_2 -algebras in $\Omega\mathcal{B}$ to the set of Mod^{E_1} -equivalence classes of condensable E_1 -algebras in $\mathcal{B}$. $\diamond$

Remark 6.1.10. There is a natural higher category called Morita category, in which many notions and problems of higher Morita theory can be formulated and studied. It consists of E_m -algebras as 0-morphisms, E_{m-1} -algebras over E_m -algebras as 1-morphisms, E_{m-2} -algebras over E_{m-1} -algebras as 2-morphisms, so on and so forth. We will discuss this Morita category elsewhere [KZZZ25]. $\diamond$

We expect the following result to be true. We will provide a proof of this result, together with a more complete theory of k -Morita equivalence, elsewhere [KZZZ25].

Proposition^o 6.1.11. For $\mathcal{A} \in \text{Alg}_{E_k}^c((n+1)\text{Vec})$ and $A \in \text{Alg}_{E_k}^c(\mathcal{A})$, $\mathcal{A}$ and $\text{Mod}_A^{E_k}(\mathcal{A})$ are k -Morita equivalent.

Proposition^o 6.1.12. For $\mathcal{A}, \mathcal{B} \in \text{Alg}_{E_k}^c((n+1)\text{Vec})$, if $\mathcal{A}$ and $\mathcal{B}$ are k -Morita equivalent, then $\mathfrak{Z}_k(\mathcal{A}) \simeq \mathfrak{Z}_k(\mathcal{B})$ in $\text{Alg}_{E_{k+1}}^c((n+1)\text{Vec})$.

Proof. We have E_{k+1} -monoidal equivalences: $\mathfrak{Z}_k(\mathcal{A}) \simeq \Omega^k \mathfrak{Z}_0(\Sigma^k \mathcal{A}) \simeq \Omega^k \mathfrak{Z}_0(\Sigma^k \mathcal{B}) \simeq \mathfrak{Z}_k(\mathcal{B})$. $\square$

Remark 6.1.13. There is a more general notion of the E_k -center of an E_k -algebra in an E_k -monoidal n -category $\mathcal{C}$. It is necessarily living in the category $\mathfrak{Z}_k(\mathcal{C})$. Hence, one can expect a generalization of above proposition in this setting. We will develop this theory in [KZZZ25]. $\diamond$

The following Lemma is a result in [Lur17, Fra12] reformulated in the world of separable n -categories. It is briefly explained in Appendix A.3.

Lemma° 6.1.14. For $\mathcal{A}, \mathcal{B} \in \text{Alg}_{\mathbb{E}_k}^c((n+1)\text{Vec})$ and an $\mathbb{E}_k$ -monoidal functor $F : \mathcal{A} \rightarrow \mathcal{B}$, we have

$$\mathfrak{Z}_k(F) = \mathfrak{Z}_k(\mathcal{A}, \mathcal{B}) \simeq \text{hom}_{\text{Mod}_{\mathcal{A}}^{\mathbb{E}_k}((n+1)\text{Vec})}(\mathcal{A}, \mathcal{B}) \in \text{Alg}_{\mathbb{E}_k}^c((n+1)\text{Vec}).$$

When $F = \text{id}_{\mathcal{A}}$, we obtain $\mathfrak{Z}_k(\mathcal{A}) \simeq \text{hom}_{\text{Mod}_{\mathcal{A}}^{\mathbb{E}_k}((n+1)\text{Vec})}(\mathcal{A}, \mathcal{A}) \in \text{Alg}_{\mathbb{E}_{k+1}}^c((n+1)\text{Vec})$, which is also a consequence of Theorem 5.2.11.

Proposition° 6.1.15. For $\mathcal{A}, \mathcal{B} \in \text{Alg}_{\mathbb{E}_k}^c((n+1)\text{Vec})$, if $\mathcal{A}$ and $\mathcal{B}$ are $\text{Mod}^{\mathbb{E}_k}$ -equivalent, then $\mathfrak{Z}_k(\mathcal{A}) \simeq \mathfrak{Z}_k(\mathcal{B})$.

Proposition° 6.1.16. For $\mathcal{B} \in \text{Alg}_{\mathbb{E}_k}^c((n+1)\text{Vec})$ and $A, B \in \text{Alg}_{\mathbb{E}_k}^c(\mathcal{B})$, if $A, B \in \text{Alg}_{\mathbb{E}_k}^c(\mathcal{B})$ are k -Morita equivalent, then they are $\text{Mod}^{\mathbb{E}_k}$ -equivalent.

Proof. It follows from $\text{Mod}_A^{\mathbb{E}_k}(\mathcal{B}) \simeq \Omega^{k-1} \text{Mod}_{\Sigma^{k-1}A}^{\mathbb{E}_1}(\Sigma^{k-1}\mathcal{B}) \simeq \Omega^{k-1} \text{Mod}_{\Sigma^{k-1}B}^{\mathbb{E}_1}(\Sigma^{k-1}\mathcal{B}) \simeq \text{Mod}_B^{\mathbb{E}_k}(\mathcal{B})$, where the second ‘ $\simeq$ ’ is a consequence of the assumption $\Sigma^k A \simeq \Sigma^k B$ in $\Sigma^k \mathcal{B}$ and Proposition 6.1.4. $\square$

6.1.2 Enveloping algebras

In order to study the $\text{Mod}^{\mathbb{E}_k}$ -equivalence among $\mathbb{E}_k$ -algebras, it is beneficial to introduce an $\text{Mod}^{\mathbb{E}_k}$ -invariant, which is called the enveloping algebra U_A of an $\mathbb{E}_k$ -algebra $A \in \text{Alg}_{\mathbb{E}_k}^c(\mathcal{C})$ for an $\mathbb{E}_k$ -fusion n -category $\mathcal{C}$ such that $\text{Mod}_A^{\mathbb{E}_k}(\mathcal{C}) \simeq \text{LMod}_{U_A}(\mathcal{C})$. This notion allows us to reduce the problem of studying $\text{Mod}^{\mathbb{E}_k}$ -equivalence to that of studying usual Morita equivalence.

Let us start from the $k = 1$ case. Let us first recall an $\mathbb{E}_1$ -algebra A in Vec is just an ordinary $\mathbb{C}$ -algebra. An A - A -bimodule is the same as a left $A \otimes_{\mathbb{C}} A^{\text{op}}$ -module. Therefore, $U_A = A \otimes_{\mathbb{C}} A^{\text{op}}$. However, this construction does not generalize to $A \in \text{Alg}_{\mathbb{E}_1}^c(\mathcal{C})$ naively because $\mathcal{C}$ is only monoidal without braidings.

In order to find the proper definition, we first look at the physical meaning behind the data A and $\mathcal{C}$. When $k = 1$, the fusion n -category $\mathcal{C}$ is the category of topological defects of an $n+1$ D topological order $\mathcal{C}^{n+1}$, which has potentially non-trivial gravitational anomaly $Z(\mathcal{C})^{n+2}$ as illustrated in Figure 18 (a), where $x \in \text{Mod}_A^{\mathbb{E}_1}(\mathcal{C})$ and only 1-spatial-dimension (the horizontal direction) is shown and the remaining $(n-1)$ -spatial-dimensions are orthogonal to the paper. Since 1-codimensional topological defects in $Z(\mathcal{C})^{n+2}$ change the boundary condition, we only consider 2-codimensional topological defects in $Z(\mathcal{C})^{n+2}$. They form a braided fusion n -category given by the Drinfeld center $\mathfrak{Z}_1(\mathcal{C})$ of $\mathcal{C}$ by boundary-bulk relation [KWZ15, KWZ17]. Since the $(n-1)$ -spatial-dimensions (orthogonal to the paper) do not appear in our discussion. Without loss of generality, we can simply assume $n = 1$.

The shaded regions in Figure 18 are $D^1 \times \mathbb{R}_-$, where $\mathbb{R}_- := (-\infty, 0)$ and D^1 is an oriented closed 1-disk and its oriented boundary is a 0-sphere, i.e. $\partial D^1 = \partial D_+^1 \cup \partial D_-^1 = S^0 = \{+, -\}$. Notice that the macroscopic observables in the interior of the shaded region (i.e. $\mathring{D}^1 \times \mathbb{R}_-$) is given by $\mathfrak{Z}_1(\mathcal{C})$. We denote this fact by $\text{Obs}(\mathring{D}^1 \times \mathbb{R}_-) = \mathfrak{Z}_1(\mathcal{C})$. Similarly, we have $\text{Obs}(\partial D_+^1 \times \mathbb{R}_-) = \mathcal{C}$ and $\text{Obs}(\partial D_-^1 \times \mathbb{R}_-) = \mathcal{C}$, which should be viewed as observables living in an open neighborhood of $\partial D_{\pm}^1 \times \mathbb{R}_-$. One can summarize all the observables living in the shaded region $D^1 \times \mathbb{R}_-$ in Figure 18 by integrating them over the whole region [AKZ17]. More explicitly, we have

$$\begin{aligned} \text{Obs}(D^1 \times \mathbb{R}_-) &= \int_{D^1 \times \mathbb{R}_-} (\text{Obs}(\mathring{D}^1 \times \mathbb{R}_-), \text{Obs}(\partial D_+^1 \times \mathbb{R}_-), \text{Obs}(\partial D_-^1 \times \mathbb{R}_-)) \\ &= \int_{D^1 \times \mathbb{R}_-} (\mathfrak{Z}_1(\mathcal{C})|_{\mathring{D}^1 \times \mathbb{R}_-}, \mathcal{C}|_{\partial D_+^1 \times \mathbb{R}_-}, \mathcal{C}|_{\partial D_-^1 \times \mathbb{R}_-}) := \mathcal{C} \boxtimes_{\mathfrak{Z}_1(\mathcal{C})} \mathcal{C}^{\text{op}}, \end{aligned} \quad (6.1.1)$$

where $\boxtimes_{\mathfrak{Z}_1(\mathcal{C})}$ is the relative tensor product over $\mathfrak{Z}_1(\mathcal{C})$. One way to see why $\mathcal{C} \boxtimes_{\mathfrak{Z}_1(\mathcal{C})} \mathcal{C}^{\text{op}}$ is qualified to be called the ‘global observable’ is to show that $\text{Obs}(\mathring{D}^1 \times \mathbb{R}_-), \text{Obs}(\partial D_+^1 \times \mathbb{R}_-), \text{Obs}(\partial D_-^1 \times \mathbb{R}_-)$ naturally map into the global one. Indeed, one can define these three maps explicitly and precisely below.

$$\begin{aligned} \mathcal{C} &\rightarrow \mathcal{C} \boxtimes_{\mathfrak{Z}_1(\mathcal{C})} \mathcal{C}^{\text{op}}, & \mathfrak{Z}_1(\mathcal{C}) &\rightarrow \mathcal{C} \boxtimes_{\mathfrak{Z}_1(\mathcal{C})} \mathfrak{Z}_1(\mathcal{C}) \boxtimes_{\mathfrak{Z}_1(\mathcal{C})} \mathcal{C}^{\text{op}} \simeq \mathcal{C} \boxtimes_{\mathfrak{Z}_1(\mathcal{C})} \mathcal{C}^{\text{op}}, & \mathcal{C}^{\text{op}} &\rightarrow \mathcal{C} \boxtimes_{\mathfrak{Z}_1(\mathcal{C})} \mathcal{C}^{\text{op}} \\ a &\mapsto a \boxtimes_{\mathfrak{Z}_1(\mathcal{C})} \mathbb{1}_{\mathcal{C}}, & z &\mapsto \mathbb{1}_{\mathcal{C}} \boxtimes_{\mathfrak{Z}_1(\mathcal{C})} z \boxtimes_{\mathfrak{Z}_1(\mathcal{C})} \mathbb{1}_{\mathcal{C}}, & b &\mapsto \mathbb{1}_{\mathcal{C}} \boxtimes_{\mathfrak{Z}_1(\mathcal{C})} b \end{aligned}$$

Moreover, all three maps preserve the $\mathbb{E}_1$ -algebraic structure of three observables algebras. In other words, they are all monoidal functors as required by the obvious physical intuitions.

Therefore, it is clear that the two-side A -action on x is the same as the left $A \boxtimes A$ -action on x through the following three logical steps:




Figure 18: Physical intuition of the enveloping algebras of $\mathcal{C}$ and $A \in \text{Alg}_{E_1}^c(\mathcal{C})$

1. $A \boxtimes A$ is an E_1 -algebra in $\mathcal{C} \boxtimes \mathcal{C}^{\text{op}}$.
2. There are two canonical monoidal functors:

$$\begin{aligned} \mathcal{C} \boxtimes \mathcal{C}^{\text{op}} &\xrightarrow{\boxtimes_{Z_1(\mathcal{C})}} \mathcal{C} \boxtimes_{Z_1(\mathcal{C})} \mathcal{C}^{\text{op}} \xrightarrow{\simeq} \text{Fun}(\mathcal{C}, \mathcal{C}) \\ a \boxtimes b &\mapsto a \boxtimes_{Z_1(\mathcal{C})} b \mapsto a \otimes - \otimes b, \end{aligned} \quad (6.1.2)$$

where the relative tensor product $\mathcal{C} \boxtimes_{Z_1(\mathcal{C})} \mathcal{C}^{\text{op}}$ amounts to closing fan in Figure 18 and the equivalence ‘ $\simeq$ ’ is due to the boundary-bulk relation [KWZ15, KWZ17]. Therefore, $A \boxtimes A$ is mapped to an E_1 -algebra in $\text{Fun}(\mathcal{C}, \mathcal{C})$.

3. $\text{Fun}(\mathcal{C}, \mathcal{C})$ acts on $\mathcal{C}$ canonically. Therefore, $A \boxtimes A$ acts on $x \in \mathcal{C}$ canonically. More precisely, by (6.1.2), this action $(A \boxtimes A) \odot x := A \otimes x \otimes A$ is precisely the correct two-side A -action.

Therefore, the correct definition of U_A is $A \boxtimes A \in \mathcal{C} \boxtimes \mathcal{C}^{\text{op}}$.

Remark 6.1.17. Note that the fusion product $\otimes$ in the category $\mathcal{C} \boxtimes_{Z_1(\mathcal{C})} \mathcal{C}^{\text{op}}$ is defined by

$$(a \boxtimes_{Z_1(\mathcal{C})} b) \otimes (a' \boxtimes_{Z_1(\mathcal{C})} b') := (a \otimes_{\mathcal{C}} a') \boxtimes_{Z_1(\mathcal{C})} (b' \otimes_{\mathcal{C}} b').$$

Geometrically, it says that, by integrating over D^1 , the fusion product along this D^1 -direction in $Z_1(\mathcal{C})$ has been integrated out, and no longer makes sense after the integration. However, the fusions along the $\mathbb{R}_-$ -direction remains a well-defined fusion product. By ignoring the fusion products in $\mathcal{C}, Z_1(\mathcal{C}), \mathcal{C}^{\text{op}}$ in the $\mathbb{R}_-$ -direction, it makes sense to integrate over D^1 only. Moreover, in 2Vec, we can simply set:

$$\int_{D^1} (Z_1(\mathcal{C})|_{\hat{D}^1}, \mathcal{C}|_{\partial D_+^1}, \mathcal{C}^{\text{op}}|_{\partial D_-^1}) := \int_{D^1 \times \mathbb{R}_-} (Z_1(\mathcal{C})|_{\hat{D}^1 \times \mathbb{R}_-}, \mathcal{C}|_{\partial D_+^1 \times \mathbb{R}_-}, \mathcal{C}^{\text{op}}|_{\partial D_-^1 \times \mathbb{R}_-}).$$

Since remembering the fusion product in the $\mathbb{R}_-$ -direction in the right hand side causes no confusion, we agree that above identity holds in $\text{Alg}_{E_1}((n+1)\text{Vec})$ as well. We use them interchangeably. $\diamond$

Before we give the official definition of U_A , we first discuss the meaning of $\mathcal{C} \boxtimes \mathcal{C}^{\text{op}}$. Since $\mathcal{C}$ is an E_1 -algebra in $(n+1)\text{Vec}$, mathematically, an $\mathcal{C}$ - $\mathcal{C}$ -module is precisely a left $\mathcal{C} \boxtimes \mathcal{C}^{\text{op}}$ -module. This mathematical fact has a nice physical meaning as shown in Remark 6.1.18. Therefore, $\mathcal{C} \boxtimes \mathcal{C}^{\text{op}}$ is precisely the enveloping algebra of $\mathcal{C}$.

Remark 6.1.18. The physical meaning of a $\mathcal{C}$ - $\mathcal{C}$ -bimodule $\mathcal{M}$ is a the category of all wall conditions on a gapped domain wall M^n between $\mathcal{C}^{n+1}$ and $\overline{\mathcal{C}^{n+1}}$ as illustrated in Figure 18 (b). Moreover, this wall uniquely determines a relative bulk or a relative gravitational anomaly $Z^{(1)}(M)^{n+1}$ such that the category of topological defects on it is given by $\text{Fun}_{C|C}(\mathcal{M}, \mathcal{M})$ (recall Figure 7) [KWZ15, KZ24]. By definition, to say $\mathcal{M}$ is a $\mathcal{C}$ - $\mathcal{C}$ -bimodule is equivalent to say there is a monoidal functor from $f : \mathcal{C} \boxtimes \mathcal{C}^{\text{op}}$ to $\text{Fun}(\mathcal{M}, \mathcal{M})$, which is precisely a morphism from $\mathcal{C}^{n+1} \boxtimes \overline{\mathcal{C}^{n+1}}$ to $Z(M)^{n+1}$. By the definition of a morphism between topological order [KWZ15, KWZ17], this morphism is precisely defined by the domain wall $Z^{(1)}(M)^{n+1}$ between $Z(\mathcal{C})^{n+2}$ and $Z(\mathcal{C})^{n+2}$ (see also [KZ18a, KZ24]). $\diamond$

We summarize above discussion in the following mathematical definition and Lemma.

Definition 6.1.19. For $\mathcal{C} \in \text{Alg}_{E_1}^c(2\text{Vec})$, we define its enveloping algebra $U_{\mathcal{C}}$ in $(n+1)\text{Vec}$ as follows:

$$U_{\mathcal{C}} := \int_{S^0} \mathcal{C} := \mathcal{C} \boxtimes \mathcal{C}^{\text{op}} \in \text{Alg}_{E_1}^c((n+1)\text{Vec}). \quad (6.1.3)$$

□

where S^0 denotes the 0-sphere consisting of two points, i.e. $S^0 = \{+, -\}$. For $A \in \text{Alg}_{E_1}^c(\mathcal{C})$, we define the enveloping algebra U_A of A as follows:

$$U_A := \int_{S^0} A = A \boxtimes A \in \text{Alg}_{E_1}^c(U_{\mathcal{C}}).$$

Remark 6.1.20. One can also view the pair $(\mathcal{C}, A)$ as an E_1 -algebra [Str12] and its enveloping algebra is also defined by a pair $U_{(\mathcal{C}, A)} := (U_{\mathcal{C}}, U_A) = \int_{S^0} (\mathcal{C}, A)$. ◇

Since $\mathcal{C}$ is naturally a left $U_{\mathcal{C}}$ -module, it makes sense to talk about a left U_A -module in $\mathcal{C}$. More precisely, a left U_A -module in $\mathcal{C}$ is an object $x \in \mathcal{C}$ equipped with a unital left U_A -action $\mu_x : U_A \odot x \rightarrow x$ satisfying the usual left unit and associativity properties of a left module. Moreover, the category $\text{LMod}_{U_A}(\mathcal{C})$ of left U_A -modules in $\mathcal{C}$ is well-defined. The following Lemma is tautological.

Lemma 6.1.21. We have $\text{Mod}_{\mathcal{C}}^{E_1}((n+1)\text{Vec}) \simeq \text{LMod}_{U_{\mathcal{C}}}((n+1)\text{Vec})$ and

$$\text{Mod}_A^{E_1}(\mathcal{C}) \simeq \text{LMod}_{U_A}(\mathcal{C}).$$

Since $\text{Alg}_{E_k}^c(-) = \text{Alg}_{E_1}^c(\text{Alg}_{E_{k-1}}^c(-))$ and $\text{Mod}^{E_k}(-) = \text{Mod}^{E_1}(\text{Mod}^{E_{k-1}}(-))$ and the i -th multiplication or i -th action occurs in i -th spatial direction that is orthogonal to the j -th direction for $i \neq j$, the generalization to E_k -algebra is straightforward.

Definition 6.1.22. For $\mathcal{C} \in \text{Alg}_{E_k}^c((n+1)\text{Vec})$ and $A \in \text{Alg}_{E_k}^c(\mathcal{C})$, we define the enveloping algebra $U_{(\mathcal{C}, A)}^{E_k}$ of $(\mathcal{C}, A)$ as follows:

$$U_{(\mathcal{C}, A)}^{E_k} = \int_{S^{k-1}} (\mathcal{C}, A) = (U_{\mathcal{C}}^{E_k}, U_A^{E_k}) = \left(\int_{S^{k-1}} \mathcal{C}, \int_{S^{k-1}} A \right).$$

Proposition 6.1.23. We have $\mathcal{C} \in \text{Mod}_{\mathcal{C}}^{E_k}((n+1)\text{Vec}) \simeq \text{LMod}_{U_{\mathcal{C}}^{E_k}}((n+1)\text{Vec})$ and

$$\text{Mod}_A^{E_k}(\mathcal{C}) \simeq \text{LMod}_{U_A^{E_k}}(\mathcal{C}).$$

As a consequence, two $A, B \in \text{Alg}_{E_k}^c(\mathcal{C})$ are Mod^{E_k} -equivalent if and only if $U_A^{E_k}, U_B^{E_k} \in \text{Alg}_{E_1}^c(U_{\mathcal{C}}^{E_k})$ are $\text{LMod}(\mathcal{C})$ -equivalent.

6.2 Theory of integrals

The study of dimensional reduction process in topological orders is deeply related to the mathematical theory of factorization homology as shown for 2+1D anomaly-free topological orders in [AKZ17]. However, in general anomalous and higher dimensional settings, it is certainly different from the usual theory (the so-called α -version) of factorization homology [Lur17, AF20, AFT17, AFT16] (see Remark 6.2.2). We believe that the theory of integrals emerging from this work is related to the so-called β -version of factorization homology partially developed in [AFR18] (see Remark 6.2.2). We hope to come back to this point in the future. In this subsection, we only briefly discuss some results of integrating physical observables that follow directly from physical intuitions.

Consider two $n+1$ D Morita equivalent topological orders C^{n+1} and D^{n+1} connected by a gapped domain wall M^n . We specify the wall condition by a pair $(\mathcal{M}, m)$, where the separable n -category $\mathcal{M}$ is the category of wall conditions and the distinguished object $m \in \mathcal{M}$ specifies a single wall condition.

Topological defects of codimension k in C^{n+1} can be fused with a defect (of the same dimension) on the wall and become topological defects on the wall. This fusion defines an action

$$\odot : \Omega^{k-1}\mathcal{C} \times \Omega_m^{k-1}\mathcal{M} \rightarrow \Omega_m^{k-1}\mathcal{M} \quad (6.2.1)$$

$$(a, x) \mapsto a \odot x. \quad (6.2.2)$$

Note that the category $\Omega^{k-1}\mathcal{C}$ is E_k -monoidal and the category $\Omega_m^{k-1}\mathcal{M}$ is E_{k-1} -monoidal. The functor $\odot$ is E_{k-1} -monoidal. We can define internal homs associated to this action, i.e. $\forall x, y \in \Omega_m^{k-1}\mathcal{M}$,

$$\text{hom}_{\Omega_m^{k-1}\mathcal{M}}(a \odot x, y) \simeq \text{hom}_{\Omega^{k-1}\mathcal{C}}(a, [x, y]_{\Omega^{k-1}\mathcal{C}}).$$


When $x = y$, the internal hom $[x, x]_{\Omega^{k-1}\mathcal{C}} \in \text{Alg}_{E_1}^{\mathcal{C}}(\Omega^{k-1}(\mathcal{C}))$. Moreover, we have

$$[1_m^{k-1}, 1_m^{k-1}]_{\Omega^{k-1}\mathcal{C}} \in \text{Alg}_{E_k}^{\mathcal{C}}(\Omega^{k-1}(\mathcal{C})).$$

This action functor $\odot$ naturally induces the following bulk-to-wall map (recall $1_m^0 = m$):

$$L_k := - \odot 1_m^{k-1} : \Omega^{k-1}\mathcal{C} \rightarrow \Omega_m^{k-1}\mathcal{M}, \quad (6.2.3)$$

which is again E_{k-1} -monoidal. We illustrate two examples of L_k in the following picture.



bulk to wall maps :

$$\begin{aligned} \Omega^2\mathcal{C} &\xrightarrow{L_3 = -\odot 1_m^2} \Omega_m^2\mathcal{M}, \\ \Omega\mathcal{C} &\xrightarrow{L_2 = -\odot 1_m} \Omega_m\mathcal{M}, \\ \mathcal{C} &\xrightarrow{L_1 = -\odot m} \mathcal{M}. \end{aligned}$$

Note that the right adjoint functor of L_k is given by $L_k^R = [1_m^{k-1}, -]_{\Omega_m^{k-1}\mathcal{M}} : \Omega_m^{k-1}\mathcal{M} \rightarrow \Omega^{k-1}\mathcal{C}$.

In this work, we motivate a theory of integrals defined on space manifolds from a condensed matter point of view (see for example (4.2.1), (4.2.6) and (6.1.1)). More naturally, we should define the integrals on spacetime manifolds. For example, By rolling up M^n and D^{n+1} along the spacetime manifold $N^k \times \mathbb{R}^{n+1-k}$ for $1 \leq k \leq n$, where N^k is a k -dimensional compact manifold, we obtain a k -codimensional topological defect in C^{n+1} as depicted below.

$$\int_{\mathbb{R}^k \times \mathbb{R}^{n+1-k}} (C^{n+1}|_{(\mathbb{R}^k \setminus N^k) \times \mathbb{R}^{n+1-k}}, D^{n+1}|_{N^k \times \mathbb{R}^{n+1-k}}, M^n|_{\partial N^k \times \mathbb{R}^{n+1-k}}) := \int_{\mathbb{R}^{n+1-k}} (C^{n+1}|_{(\mathbb{R}^k \setminus D^k) \times \mathbb{R}^{n+1-k}}, D^{n+1}|_{D^k \times \mathbb{R}^{n+1-k}}, M^n|_{\partial D^k \times \mathbb{R}^{n+1-k}}) \quad (6.2.4)$$

Theorem^{ph} 6.2.1. Let C^{n+1} and D^{n+1} be two Morita equivalent $n+1$ D topological orders connected by an anomaly-free gapped domain wall $M^n = (\mathcal{M}, m)$ and $m \in \mathcal{M}$. When $N^k = D^k$ (i.e., a closed k -disk) for $1 \leq k \leq n$, we have

$$\begin{aligned} &\int_{\mathbb{R}^k \times \mathbb{R}^{n+1-k}} (C^{n+1}|_{(\mathbb{R}^k \setminus D^k) \times \mathbb{R}^{n+1-k}}, D^{n+1}|_{D^k \times \mathbb{R}^{n+1-k}}, M^n|_{\partial D^k \times \mathbb{R}^{n+1-k}}) \\ &:= \int_{\mathbb{R}^k \times \mathbb{R}^{n+1-k}} (\mathcal{C}|_{(\mathbb{R}^k \setminus D^k) \times \mathbb{R}^{n+1-k}}, \mathcal{D}|_{D^k \times \mathbb{R}^{n+1-k}}, (\mathcal{M}, m)|_{\partial D^k \times \mathbb{R}^{n+1-k}}) \\ &= (\mathcal{C}, \mathbb{1}_{\mathcal{C}}, \dots, 1_{\mathbb{1}_{\mathcal{C}}}^{k-2}, [1_m^{k-1}, 1_m^{k-1}]_{\Omega^{k-1}\mathcal{C}}) \end{aligned} \quad (6.2.5)$$

where we used the convention $1_m^0 := m$ and notation (4.2.7). In this case, we can use a short notation $(\Omega^{k-1}\mathcal{C}, [1_m^{k-1}, 1_m^{k-1}]_{\Omega^{k-1}\mathcal{C}})$ for the answer. $\square$

Remark 6.2.2. Our theory of integral is beyond the usual theory of factorization homology [Lur17, AF15], which was also called the α -version of factorization homology (see [AF20] for a review). For example, all $(n-k)$ -dimensional topological defects in M^n form the category $\Omega_m^k \mathcal{M}$, which is an E_k -fusion $(n-k)$ -category; and all $(n-k)$ -dimensional topological defects in D^{n+1} form the category $\Omega^k \mathcal{D}$, which is an E_{k+1} -fusion $(n-k)$ -category. By the α -version of factorization homology [Lur17, AF15], the following factorization homology

$$\int_{\mathbb{R}^k \times \mathbb{R}^{n+1-k}} (\Omega^k \mathcal{C}|_{(\mathbb{R}^k \setminus D^k) \times \mathbb{R}^{n+1-k}}, \Omega^k \mathcal{D}|_{D^k \times \mathbb{R}^{n+1-k}}, \Omega_m^k \mathcal{M}|_{\partial D^k \times \mathbb{R}^{n+1-k}})$$

is well-defined in the usual sense (see also [AKZ17]). Let $P := [1_m^{k-1}, 1_m^{k-1}]_{\Omega^{k-1} \mathcal{C}}$. There is a natural E_1 -monoidal functor

$$\int_{\mathbb{R}^k \times \mathbb{R}^{n+1-k}} (\Omega^k \mathcal{C}|_{(\mathbb{R}^k \setminus D^k) \times \mathbb{R}^{n+1-k}}, \Omega^k \mathcal{D}|_{D^k \times \mathbb{R}^{n+1-k}}, \Omega_m^k \mathcal{M}|_{\partial D^k \times \mathbb{R}^{n+1-k}}) \longrightarrow \Omega_P \mathcal{C},$$

which is, in general, not an equivalence. This non-equivalence is a very common phenomenon, which says that integrating local observables living on each stratum over a stratified manifold N only produces parts of the global observables on N because certain non-local observables become local when we shrink the size of M^n . This phenomenon was called *spatial fusion anomaly*⁶³ appeared for topological defects of codimension 2 or higher in [KZ20, KZ21]. This phenomenon also says that the theory of integral proposed in this work is beyond the α -version of factorization homology [Lur17, AF20],

In the summer of 2025, one of us (Liang Kong) had a private communication with John Francis. It was clarified that results in Theorem^{ph} 6.2.1 are compatible with the predictions from the β -version of factorization homology. It convinces us that the mathematical foundation of the theory of integrals proposed in this work should be precisely the β -version of factorization homology proposed in [AFR18]. According to Francis, the mathematical foundation of the β -version of factorization homology has not yet been fully established. Some recent developments in this program can be found in [AF24]. $\diamond$

Example 6.2.3. Consider an anomaly-free simple 2+1D topological order X^3 , viewed as a special case of (6.2.5) when $n = 3$, $C^4 = D^4 = 1^4$ and $M^3 = (\mathcal{M}, m) = X^3$. In this case, $\Omega_m^2 \mathcal{M} = \Omega \mathcal{X}$ is a non-degenerate braided fusion 1-category.

1. By rolling up a 2+1D topological order X^3 along a cylinder $S^1 \times \mathbb{R}$ in the space manifold $\mathbb{R}^3$, we obtain a 2-codimensional topological defect (i.e. a string-like defect) in 1^4 ;

$$\int_{\mathbb{R}^2 \times \mathbb{R}^2} (1^4|_{(\mathbb{R}^2 \setminus D^2) \times \mathbb{R}^2}, 1^4|_{D^2 \times \mathbb{R}^2}, X^3|_{\partial D^2 \times \mathbb{R}^2}) = (3\text{Vec}, 2\text{Vec}, \Omega \mathcal{X}) = (2\text{Vec}, \Omega \mathcal{X}). \quad (6.2.6)$$

2. By rolling up X^3 onto a 2-sphere S^2 in the space manifold $\mathbb{R}^3$, we obtain a 3-codimensional topological defect (i.e. a particle-like defect) in 1^4 .

$$\int_{\mathbb{R}^3 \times \mathbb{R}^1} (1^4|_{(\mathbb{R}^3 \setminus D^3) \times \mathbb{R}^1}, 1^4|_{D^3 \times \mathbb{R}^1}, X^3|_{\partial D^3 \times \mathbb{R}^1}) = (3\text{Vec}, 2\text{Vec}, \text{Vec}, \Omega^2 \mathcal{X}) \simeq (\text{Vec}, \mathbb{C}).$$

In this case, $\Omega^2 \mathcal{C} = \text{Vec}$, $1_m^2 = \mathbb{1}_{\Omega \mathcal{X}}$ and the internal hom $[1_m^2, 1_m^2]_{\Omega^2 \mathcal{C}}$ becomes the ordinary hom space $\text{hom}_{\Omega \mathcal{X}}(\mathbb{1}_{\Omega \mathcal{X}}, \mathbb{1}_{\Omega \mathcal{X}}) = \Omega^2 \mathcal{X} = \mathbb{C}$. This result is compatible with the α -version of factorization homology [AKZ17]. This is due to a general principle that the fusions of 1-codimensional topological defects are free of the spatial fusion anomaly [KZ21].

3. By rolling up X^3 along a 2-torus $S^1 \times S^1$ in the space manifold $\mathbb{R}^3$, we obtain a 3-codimensional topological defect (i.e. a particle-like defect) in 1^4 . We denote the set of simple objects in $\Omega \mathcal{X}$ by $\text{Irr}(\Omega \mathcal{X})$.

$$\begin{aligned} \int_{\mathbb{R}^3 \times \mathbb{R}^1} (1^4|_{(\mathbb{R}^3 \setminus (D^2 \times S^1)) \times \mathbb{R}^1}, 1^4|_{D^2 \times S^1 \times \mathbb{R}^1}, X^3|_{\partial D^2 \times S^1 \times \mathbb{R}^1}) &= (\text{Vec}, \text{hom}_{\text{Fun}(\Omega \mathcal{X}, \Omega \mathcal{X})}(\text{id}_{\Omega \mathcal{X}}, \text{id}_{\Omega \mathcal{X}})) \\ &\simeq (\text{Vec}, \text{hom}_{\Omega \mathcal{X}}(\mathbb{1}_{\Omega \mathcal{X}}, \oplus_{i \in \text{Irr}(\Omega \mathcal{X})} i \otimes i^R)), \end{aligned} \quad (6.2.7)$$

which was explained in the following two-step reasoning.

⁶³Interestingly, this anomaly vanishes for the fusions of 1-codimensional gapped domain walls among anomaly-free topological orders [AKZ17, KZ20, KZ24, KZ21] (see also Remark 2.4.3).

- (a) We first notice that the same particle can be obtained physically by rolling up the string obtained in (6.2.6) in $\mathbf{1}^3$ along S^1 in the space manifold $\mathbb{R}^2$.
- (b) Applying (6.2.5) again in the case when $n = k = 2$, $C^3 = D^3 = \mathbf{1}^3$ and $M^2 = (2\text{Vec}, \Omega\mathcal{X})$, we obtain the identity (6.2.7).

This result is compatible with the α -version of factorization homology [AKZ17] for the same reason. $\heartsuit$

Example 6.2.4. Consider the 3+1D $\mathbb{Z}_2$ topological order $C^4 = \text{GT}_{\mathbb{Z}_2}^4$ and $D^4 = \mathbf{1}^4$.

1. Let M^3 be the (twist) smooth boundary of $\text{GT}_{\mathbb{Z}_2}^4$ [KTZ20a], i.e. $M^3 = (3\text{Rep}(\mathbb{Z}_2), \mathbb{1}_{3\text{Rep}(\mathbb{Z}_2)})$. We have

$$\begin{aligned} \int_{\mathbb{R}^4} (\text{GT}_{\mathbb{Z}_2}^4|_{(\mathbb{R}^2 \setminus D^2) \times \mathbb{R}^2}, \mathbf{1}^4|_{\mathbb{D}^2 \times \mathbb{R}^2}, M^3|_{\partial D^2 \times \mathbb{R}^2}) &= (\Omega\mathbb{C}, \mathbb{1}_{\mathbb{C}} \oplus m), \\ \int_{D^3 \times \mathbb{R}^1} (\text{GT}_{\mathbb{Z}_2}^4|_{(\mathbb{R}^3 \setminus D^3) \times \mathbb{R}^1}, \mathbf{1}^4|_{\mathbb{D}^3 \times \mathbb{R}^1}, M^3|_{\partial D^3 \times \mathbb{R}^1}) &= (\Omega^2\mathbb{C}, \mathbb{1}_{\mathbb{C}}^2) = (\text{Rep}(\mathbb{Z}_2), \mathbb{C}), \\ \int_{D^2 \times S^1 \times \mathbb{R}^1} (\text{GT}_{\mathbb{Z}_2}^4|_{(\mathbb{R}^3 \setminus D^2 \times S^1) \times \mathbb{R}^1}, \mathbf{1}^4|_{\mathbb{D}^2 \times S^1 \times \mathbb{R}^1}, M^3|_{\partial D^2 \times S^1 \times \mathbb{R}^1}) &= \int_{S^1} \mathbb{1}_{\mathbb{C}} \oplus m = (\text{Rep}(\mathbb{Z}_2), \mathbb{C}^{\oplus 2}), \end{aligned}$$

where $\mathbb{C}$ denotes the trivial $\mathbb{Z}_2$ -representation, and $\mathbb{1}_{\mathbb{C}} \oplus m$ is precisely the Lagrangian algebra in $\Omega\mathcal{GT}_{\mathbb{Z}_2}^4$ that determines the (twist) smooth boundary [KTZ20a, ZLZH⁺23]; the second identity follows from the fact that the left hand side amounts to a line segment of the $(\mathbb{1}_{\mathbb{C}} \oplus m)$ -string; the third identity follows from the fact that an m -loop gives the trivial particle $\mathbb{1}_{\mathbb{C}}^2$ [KTZ20a].

2. Let M^3 be the rough boundary of $\text{GT}_{\mathbb{Z}_2}^4$, $M^3 = (3\text{Rep}(\mathbb{Z}_2), 2\text{Vec})$, where 2Vec is viewed as a right $2\text{Rep}(\mathbb{Z}_2)$ -module, i.e., an object in $3\text{Rep}(\mathbb{Z}_2) = \text{RMod}_{2\text{Rep}(\mathbb{Z}_2)}(2\text{Vec})$. We have

$$\begin{aligned} \int_{D^2 \times \mathbb{R}^2} (\text{GT}_{\mathbb{Z}_2}^4|_{(\mathbb{R}^2 \setminus D^2) \times \mathbb{R}^2}, \mathbf{1}^4|_{\mathbb{D}^2 \times \mathbb{R}^2}, M^3|_{\partial D^2 \times \mathbb{R}^2}) &= (\Omega\mathbb{C}, \mathbb{1}_{\mathbb{C}}), \\ \int_{D^3 \times \mathbb{R}^1} (\text{GT}_{\mathbb{Z}_2}^4|_{(\mathbb{R}^3 \setminus D^3) \times \mathbb{R}^1}, \mathbf{1}^4|_{\mathbb{D}^3 \times \mathbb{R}^1}, M^3|_{\partial D^3 \times \mathbb{R}^1}) &= (\Omega^2\mathbb{C}, \mathbb{1}_{\mathbb{C}}^2 \oplus e), \\ \int_{D^2 \times S^1 \times \mathbb{R}^1} (\text{GT}_{\mathbb{Z}_2}^4|_{(\mathbb{R}^3 \setminus D^2 \times S^1) \times \mathbb{R}^1}, \mathbf{1}^4|_{\mathbb{D}^2 \times S^1 \times \mathbb{R}^1}, M^3|_{\partial D^2 \times S^1 \times \mathbb{R}^1}) &= \int_{S^1} \mathbb{1}_{\mathbb{C}} = (\Omega^2\mathbb{C}, \mathbb{1}_{\mathbb{C}}^2 \oplus e), \end{aligned}$$

where $\mathbb{1}_{\mathbb{C}}$ is the Cheshire String (recall Example 2.2.7) and also the Lagrangian algebra that determines the rough boundary [ZLZH⁺23]; the second identity follows from the fact that the left hand side amounts to a line segment of the $\mathbb{1}_{\mathbb{C}}$ -string; the third identity follows from the fact that an $\mathbb{1}_{\mathbb{C}}$ -loop gives the particle $\mathbb{1}_{\mathbb{C}}^2 \oplus e$ [KTZ20a, Eq. (4.6)]. $\heartsuit$

Note that the pair $(\mathcal{M}, m)$ only specifies a very special wall condition. The most general wall condition M^n can be defined by a stratified n -disk with each p -codimensional stratum (in M^n) labeled by a p -morphism in $\mathcal{M}$ for $0 \leq p \leq n$. In particular, we can label different 0-codimensional strata (i.e. n -cells) by different 0-morphisms in $\mathcal{M}$. We would like to know how to compute the k -codimensional defect in C^{n+1} obtained by rolling up M^n with such a general wall condition along $N^i \times \mathbb{R}^{n+1-i}$ as long as all strata glue properly. At the current stage, the explicit computation in such a general setup is not available. In the remainder of this subsection, we provide some explicit examples in the case $C^{n+1} = D^{n+1}$ and $M^n = C^n$.

Proposition^{ph} 6.2.5. Let X^{n+1-l} be an l -codimensional topological defect in C^{n+1} . For $l \leq k < n+1$, we have a canonical action $\odot : \Omega^{k-l}\mathbb{C} \times \Omega_X^{k-l}(\mathbb{C}) \rightarrow \Omega_X^{k-l}(\mathbb{C})$, which is an E_{k-l} -monoidal functor. By rolling up X along $S^{k-l} \times \mathbb{R}^{n+1-k}$, we obtain a k -codimensional topological defect $\int_{S^{k-l} \times \mathbb{R}^{n+1-k}} X$ given as follows:


$$\int_{S^{k-l} \times \mathbb{R}^{n+1-k}} X = (\Omega^{k-l}\mathbb{C}, [\mathbb{1}_X^{k-l}, \mathbb{1}_X^{k-l}]_{\Omega^{k-l}\mathbb{C}}), \quad (6.2.8)$$

where $[\mathbb{1}_X^{k-l}, \mathbb{1}_X^{k-l}]_{\Omega^{k-l}\mathbb{C}} \in \text{Alg}_{E_{k-l+1}}^{\mathbb{C}}(\Omega^{k-l}\mathbb{C})$ and, when $k = l$, we have $[\mathbb{1}_X^{k-l}, \mathbb{1}_X^{k-l}]_{\Omega^{k-l}\mathbb{C}} = X \otimes X^R$.

Example 6.2.6. In 2+1D $\mathbb{Z}_2$ topological order TC^3 , there are six simple 1-codimensional topological defects $\mathbb{1}, \text{ss}, \text{sr}, \text{rr}, \text{rs}, \vartheta$ and four simple 2-codimensional topological defects $\mathbb{1}_{\perp}, m, e, f$. Then we have

$$\int_{S^1 \times \mathbb{R}^1} X = \begin{cases} (\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2)), \mathbb{1}_{\perp}) & \text{if } X = \mathbb{1}, \vartheta; \\ (\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2)), \mathbb{1}_{\perp} \oplus m) & \text{if } X = \text{ss}, \text{sr}; \\ (\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2)), \mathbb{1}_{\perp} \oplus e) & \text{if } X = \text{rr}, \text{rs}. \end{cases}$$

Remark 6.2.7. If we select a slightly more general wall condition, more precisely, we specify an $(n-1)$ -dimensional topological defect $m_1 \in \Omega_m \mathcal{M}$ on the wall and an $(n-1)$ -dimensional topological defect $d \in \Omega \mathcal{D}$, then, by rolling up the wall along $S^1 \times \mathbb{R}^{n-1}$, we obtain the following formula (recall Theorem^{ph} 4.1.18).



$$\int_{\mathbb{R}^2 \times \mathbb{R}^{n-1}} (\mathcal{C}|_{(\mathbb{R}^2 \setminus \mathbb{D}^2) \times \mathbb{R}^{n-1}}, (\mathcal{D}, d)|_{\mathbb{D}^2 \times \mathbb{R}^{n-1}}, (\mathcal{M}, m, m_1)|_{\partial \mathbb{D}^2 \times \mathbb{R}^{n-1}}) = (\Omega \mathcal{C}, [1_m, m_1 \odot d]_{\Omega \mathcal{C}}).$$

Moreover, $[1_m, - \odot -]$ is a well-defined functor, which maps $\Omega_m \mathcal{M} \times \Omega \mathcal{D} \rightarrow \Omega \mathcal{C}$. In other words, this roll-up process is functorial and $[1_m, - \odot -]$ also tells us how to obtain the higher codimensional topological defect in $\Omega \mathcal{C}$ resulting from rolling up higher codimensional defects in $\Omega_m \mathcal{M}$ and $\Omega \mathcal{D}$. $\diamond$

6.3 Condensations of gapless liquid-like defects

In this subsection, we sketch a generalization of the condensation theory to gapless liquid-like defects. We leave a detailed study to future publications.

6.3.1 General theory

Let C^{n+1} be an $n+1$ D topological order. It can have gapless defects. For example, for C^3 , by stacking a 2D topological defect with a 1+1D anomaly-free gapless phase (e.g., a 2D rational CFT), we obtain a gapless 1-codimensional defect in C^3 . We give another example. Consider two k -codimensional simple topological defects X and Y in C^{n+1} connected by a $(k+1)$ -codimensional topological defect. By proliferating Y within X , we can construct an $(n+1-k)$ D phase transition from X to Y (without altering C^{n+1}). At the critical point of the phase transition, the gap (in a neighborhood of the defect) is closed, thus we obtain a k -codimensional gapless defect in C^{n+1} .

A general gapless defect can be really wild. In general, the fusions among general defects are not necessarily well-defined. Therefore, we limit ourselves to a special class of gapless defects, which we call *liquid-like* defects [KZ22b, KZ22c]. By ‘liquid-like’, we mean that the defect is ‘soft’ enough so that it can be bent freely without altering the defect. Physically, it means that such a defect should be transparent to energy-momentum tensor⁶⁴. Such an n -dimensional liquid-like defect can also be viewed as an anomalous n D quantum phase, which was called an (anomalous) *quantum liquid* [KZ22b]. The mathematical theory of quantum liquids was developed in [KZ22b, KZ22c]. In particular, a quantum liquid⁶⁵ X^{n+1} can be described by a pair $(X_{\text{Iqs}}, X_{\text{top}})$, where

- X_{Iqs} summarizes all the dynamical data (called *local quantum symmetry*), a mathematical theory of which was developed in [KZ22c];
- X_{top} summarizes all the topological (or categorical) data (called *topological skeleton*).

The fusion of the topological skeletons of two quantum liquids is well-defined. It was conjectured that the fusion of local quantum symmetry can also be properly defined so that it is compatible with that of the topological skeletons [KZ22b]. A possible theory to make sense of this conjecture was proposed and developed in [KZ22c] based on a generalization of conformal nets in 1+1D to topological/defect nets in

⁶⁴The ‘liquid-like’ properties are in some sense topological properties. In physics literature, gapless boundaries or domain walls are often called non-topological. However, some gapless defect lines in 1+1D rational CFT’s are also called topological. To avoid confusion, we will not use the term ‘a non-topological defect’ for a gapless liquid-like defect.

⁶⁵Gapped quantum liquids include topological orders and SPT/SET/SSB orders [ZW15, SM16, KLWZZ20b]. Gapless quantum liquids should include fully dualizable CFTs [KZ22b].

arbitrary dimensions. For simplicity, we assume that the fusions among quantum liquids can be properly defined. As a consequence, it means that the category of all gapped and gapless liquid-like defects in a topological order is well-defined. Given an $n+1$ D topological order C^{n+1} , we denote the category of all liquid-like defects by $\hat{\mathcal{C}}$. It is important to notice that a topological defect is always connected to a large family of gappable liquid-like defects (see Example 6.3.1). Mathematically, the softness of the liquid-like defects simply means that the category of such defects is fully dualizable.

Example 6.3.1. A 1+1D rational conformal field theory (RCFT) can be viewed as a 1-codimensional liquid-like defect in 1^3 . In a general 2+1D topological order C^3 , we can stack a 1-codimensional topological defect with a 1+1D RCFT to obtain a 1-codimensional liquid-like defect in C^3 . For 2+1D $\mathbb{Z}_2$ topological order TC^3 , it has two gapped boundaries: the smooth boundary and the rough boundary. In [CJKYZ20], an explicit purely boundary phase transition was constructed via lattice model, and the critical point, viewed as a gappable gapless boundary of TC^3 , was shown to be liquid-like. More liquid-like gapless boundaries of 2+1D $\mathbb{Z}_2$ topological order TC^3 have been constructed in [KZ21]. $\heartsuit$

It is our long term goal to develop a rather complete condensation theory of gapped/gapless quantum liquids with all the information of local quantum symmetries. Two of us made a non-trivial attempt in this direction in [KZ22c]. In particular, we developed a general theory of local quantum symmetries in all dimensions based on the mathematical theory of topological nets and defects nets, which was introduced to generalize that of conformal nets in 1+1D. This theory also includes a condensation theory of topological nets, which, we believe, is a non-trivial step to achieve our goal. Unfortunately, we have not yet applied this theory to study real physical examples of gapless phases or QFTs. This long term goal is far beyond the scope of this work. Instead of pursuing a complete theory, in this subsection, we consider a simplified theory focusing on topological skeletons only. By forgetting the data of local quantum symmetry⁶⁶, we obtain for each topological order (or quantum liquids) C^{n+1} a category $\hat{\mathcal{C}}_{\text{top}}$, which consists of the topological skeletons of all its liquid-like defects. It was shown that this category is monoidal, fully dualizable and condensation complete [KZ22b, KZ22c]. Note that when $C^{n+1} = 1^{n+1}$, it was shown that $\hat{\mathcal{C}}_{\text{top}}$ is equivalent to either the coslice category $\bullet/(n+2)\text{Vec}$ or the slice category $(n+2)\text{Vec}/\bullet$ [KZ22b, KZ22c]⁶⁷. We will leave a mathematically detailed study of condensations in $\hat{\mathcal{C}}_{\text{top}}$ or $\bullet/(n+2)\text{Vec}$ elsewhere. Instead, we give a heuristic discussion of the condensations in $\hat{\mathcal{C}}_{\text{top}}$.

Remark 6.3.2. We want to emphasize that this category $\hat{\mathcal{C}}_{\text{top}}$ is physically defined but not mathematically defined. It is not clear to us how to formulate it mathematically, i.e., define every object and morphisms with mathematical precision. We suspect that this yet-unknown mathematical definition might provide a ‘correct’ mathematical framework to study enriched (monoidal) categories. $\diamond$

We expect that the condensation theory of liquid-like defects in $\hat{\mathcal{C}}_{\text{top}}$ is completely parallel to that of topological defects in a topological order. In particular, by condensing a k -codimensional liquid-like defect in $\hat{\mathcal{C}}_{\text{top}}$, we mean a multiple-step process.

1. We first condense a condensable E_k -algebra A in one of the transversal directions. We obtain a $(k-1)$ -codimensional liquid-like defect ΣA , which is automatically a condensable E_{k-1} -algebra in $\Omega^{k-1}\hat{\mathcal{C}}_{\text{top}}$.
2. We repeat the process $(k-1)$ -times and obtain a 1-codimensional liquid-like defect $\Sigma^{k-1}A$, which is automatically a condensable E_1 -algebra in $\hat{\mathcal{C}}_{\text{top}}$. It can be further condensed to give a new phase D^{n+1} such that $\hat{\mathcal{D}}_{\text{top}} \simeq \text{Mod}_{\Sigma^{k-1}A}^{E_1}(\hat{\mathcal{C}}_{\text{top}})$ and a potentially gapless wall $(\hat{\mathcal{M}}_{\text{top}}, m)$ such that $\hat{\mathcal{M}}_{\text{top}} \simeq \text{RMod}_{\Sigma^{k-1}A}(\hat{\mathcal{C}}_{\text{top}})$ and $m = \Sigma^{k-1}A$.
3. Alternatively, one can condense the condensable E_2 -algebra $\Sigma^{k-2}A$ in $\Omega\hat{\mathcal{C}}_{\text{top}}$ directly to obtain the condensed phase D^{n+1} such that $\Omega\hat{\mathcal{D}}_{\text{top}} \simeq \text{Mod}_{\Sigma^{k-2}A}^{E_2}(\Omega\hat{\mathcal{C}}_{\text{top}})$ and $\Omega_m\hat{\mathcal{M}}_{\text{top}} \simeq \text{RMod}_{\Sigma^{k-2}A}(\Omega\hat{\mathcal{C}}_{\text{top}})$.

Since any topological order can have a gapped or gapless liquid-like boundary, one expect that any topological order can be obtained via a condensation in the trivial phase. This theory largely generalizes the main results in this paper and will be developed in details elsewhere.

⁶⁶It is not completely clear if forgetting the dynamical data (i.e., local quantum symmetry) is physically legitimate. However, the results in [KZ22c] and the fact that the category of topological skeletons alone supports a well-defined condensation theory [KZ22b] suggests that the complete theory should be closely related to the one defined in the category of topological skeletons.

⁶⁷We ignore the unitarity here.




Figure 19: the idea of topological Wick rotation [KZ18b, KZ20, KZ21]

Remark 6.3.3. Recall Remark 5.3.15 on the MME-gauging of a fusion n -category symmetry $\mathcal{A}$, or equivalently, a (potentially anomalous) topological order A^{n+1} , by an $\mathcal{A}$ -module $\mathcal{M}$ or by a monoidal functor $f : \mathcal{A} \rightarrow \mathfrak{Z}_0(\mathcal{M}) = \text{Fun}(\mathcal{M}, \mathcal{M})$, and the MME-gauged phase is given by $\mathfrak{Z}_0(\mathcal{M})$ and the associated condensed phase is given by the centralizer $\mathfrak{Z}_1(f) \simeq \text{Fun}_{\mathcal{A}}(\mathcal{M}, \mathcal{M})^{\text{op}}$. This can be generalized to an MME-gauging process such that the MME-gauged phase and the condensed phase are both gapless. In this generalized setting, one simply replace $\mathcal{A}$ -module by an “enriched $\mathcal{A}$ -module” $\mathcal{M}^\sharp$, which, together with additional dynamical data (i.e., local quantum symmetries), characterizes a gapless quantum liquid M^n , such that the gapless MME-gauged phase is given by $Z(M)^{n+1}$ and the topological skeleton of $Z(M)^{n+1}$ is again given by the E_0 -center of $\mathcal{M}^\sharp$ (see [KYZZ24] for such a notion in the enriched setting). The gauging can be equivalently defined by an enriched monoidal functor $g : \mathcal{A} \rightarrow \mathcal{M}^\sharp$, and the topological skeleton of the associated condensed phase is again given by the centralizer $\mathfrak{Z}_1(g)$. Such a condensed phase is a gappable gapless quantum liquid that is Morita equivalent to A^{n+1} . Some examples of such anomalous gappable gapless quantum liquids in 1+1D were constructed and studied explicitly in [CJKYZ20, KZ21]. This generalized MME-gauging is enough to show how rich this defect condensation theory really is. $\diamond$

The complete mathematical theory of condensation of gapless liquid-like defects are incredibly rich. It relies on some not-yet developed mathematical theory of enriched (higher) categories and needed-to-be-further-developed theory of local quantum symmetries [KZ22c]. In the next subsection, avoiding mathematical details, we give an illustrating example in 2+1D based on the physical intuition of topological Wick rotation [KZ18b, KZ20, KZ21].

6.3.2 An illustrating example in 2+1D

The mathematical theory of gapped/gapless boundaries of 2+1D topological orders was developed in [KZ18b, KZ20, KZ21]. We do not need the complete theory here. We only focus on the topological data of gapless boundaries and the physical intuition provided by the so-called topological Wick rotation as illustrated in Figure 19. More precisely, the left picture depicts in the spatial dimension a gapped domain wall P^2 between two anomaly-free 2+1D simple topological order C^3 and B^3 . The 2-codimensional topological defects in B^3 (resp. C^3) form a non-degenerate braided fusion 1-category $\Omega\mathcal{B}$ (resp. $\Omega\mathcal{C}$). For convenience, we also assume $P^2 = (\mathcal{P}, p)$ is simple. Then the 1-codimensional topological defects in P^2 form a fusion 1-category $\Omega_p\mathcal{P}$. We have a canonical braided equivalence $\Omega\mathcal{B}^{\text{op}} \boxtimes \Omega\mathcal{C} \simeq \mathfrak{Z}_1(\Omega_p\mathcal{P})$. The canonical action of $\Omega\mathcal{B}^{\text{op}}$ on $\Omega_p\mathcal{P}$ defines an enriched fusion category ${}^{\Omega\mathcal{B}^{\text{op}}}\Omega_p\mathcal{P}$ via the so-called canonical construction [MP17]. It turns out that the topological skeleton $X_{\text{top}} = \mathcal{X}$ of a gapless boundary X^2 of a 2+1D topological C^3 is precisely given by an enriched fusion category $\mathcal{X} = {}^{\Omega\mathcal{B}^{\text{op}}}\Omega_p\mathcal{P}$, in which the original finite dimensional hom space $\text{hom}_{\Omega_p\mathcal{P}}(x, y)$ is replaced by an internal hom $[x, y]$ in $\Omega\mathcal{B}^{\text{op}}$. The physical meaning of $\Omega\mathcal{B}$ is the category of modules over a rational vertex operator algebra, and that of the internal hom space $[x, y]$ is an infinite dimensional vector space of defect fields [KZ18b, KZ20, KZ21]. This suggests that the topological skeleton of the gapless boundary X^2 can be obtained by rotating the 2+1D phase B^3 to the time direction as a fictional phase in spacetime as depicted in the second picture in Figure 19. This fictional phase can be viewed as a bookkeeping device for the combination of $\Omega\mathcal{B}^{\text{op}}$ with $\mathcal{P}$ to form an $\Omega\mathcal{B}^{\text{op}}$ -enriched fusion category ${}^{\Omega\mathcal{B}^{\text{op}}}\Omega_p\mathcal{P}$ [KZ18b]. Actually, it suggests something much deeper than a bookkeeping device as shown already in [KZ18b] that it led a powerful formula for computing the fusion of gapless domain walls [KZ18b, Eq. 5.3] and a generalization of holographic duality to higher dimensions [KZ20, KZ21, KZ22b]. From now on, we take it for granted that the second picture in Figure 19 represents a gapless boundary X^2 of C^3 .

Similar to Section 4.1.3, by rolling up the gapless boundary X^2 , we obtain a hole with a gapless boundary (depicted as a cylinder in (6.3.1) based on the geometric intuition of topological Wick rotation) or a 2-

codimensional liquid-like defect in $\Omega\hat{\mathcal{C}}_{\text{top}}$.

$$A = \int_{S^1} \mathcal{X} = \int_{S^1} \Omega\mathcal{B}^{\text{op}}\mathcal{P} = \text{[Diagram: A cylinder } B^3 \text{ with base } P^2 \text{ on a plane } C^3. \text{ The cylinder is labeled } \Omega\mathcal{B}^{\text{op}}\mathcal{P}. \text{]} \in \Omega\hat{\mathcal{C}}. \quad (6.3.1)$$

Note that this 2-codimensional liquid-like defect is not a particle (or anyon) in the usual sense. It should be viewed as a generalized particle (or a virtual⁶⁸ particle). Similar to the discussion in Section 4.1.3, this liquid-like defect is an internal hom $[\mathbb{1}_{\mathcal{X}}, \mathbb{1}_{\mathcal{X}}]_{\Omega\hat{\mathcal{C}}}$. The internal hom is again defined by the natural action $\odot : \Omega\hat{\mathcal{C}}_{\text{top}} \times \mathcal{X} \rightarrow \mathcal{X}$, i.e.

$$\text{hom}_{\mathcal{X}}(a \odot x, y) \simeq \text{hom}_{\Omega\hat{\mathcal{C}}_{\text{top}}}(a, [x, y]_{\Omega\hat{\mathcal{C}}_{\text{top}}}).$$

Therefore, $A = [\mathbb{1}_{\mathcal{X}}, \mathbb{1}_{\mathcal{X}}]_{\Omega\hat{\mathcal{C}}}$ has a canonical structure of a condensable E_2 -algebra in $\hat{\mathcal{C}}_{\text{top}}$. The geometric meaning of the algebraic structure of A was explained in details in [KZ21, Section 6.2, Remark 6.3] as a natural⁶⁹ geometric fusion of two such ‘cylinders’ depicted in (6.3.1). From the geometric construction of this multiplication, it is clear that A is an E_2 -algebra in $\Omega\hat{\mathcal{C}}_{\text{top}}$.

Notice that a defect in $\mathcal{X}$ also defines an object in $\Omega\hat{\mathcal{C}}_{\text{top}}$ as illustrated below:


$$\int_{S^1} (\mathcal{X}, x) = \text{[Diagram: A cylinder with a vertical line through its center labeled } x \in \mathcal{X}. \text{ The cylinder is labeled } \Omega\mathcal{B}^{\text{op}}\mathcal{P}. \text{ The base plane is } C^3. \text{]} \in \Omega\hat{\mathcal{C}}. \quad (6.3.2)$$

The same geometric fusion of two cylinders defined in [KZ21, Section 6.2] defines an A -action on $\int_{S^1} (\mathcal{X}, x)$. As a consequence, $\int_{S^1} (\mathcal{X}, x)$ is only a right A -module (not an E_2 -module) in $\Omega\hat{\mathcal{C}}_{\text{top}}$. Geometrically, it is clear that such a right A -module is an E_2 - A -module if and only if $x \in \mathcal{X}$ can be moved into the interior of the hole (i.e., trivial phase $\mathbf{1}^3$). However, the only simple 2-codimensional liquid-like defect is the trivial particle. We obtain $\text{Mod}_A^{E_2}(\Omega\hat{\mathcal{C}}_{\text{top}}) \simeq \text{Vec}$. We summarize all the results in this subsection in the following theorem.

Theorem^{ph} 6.3.4. The topological skeleton $X_{\text{top}} = \mathcal{X}$ of a gapless boundary X^2 of an anomaly-free 2+1D topological order C^3 can be obtained by condensing a liquid-like defect $A := \int_{S^1} \mathcal{X}$, which is an E_2 -algebra in $\hat{\mathcal{C}}_{\text{top}}$. Moreover, we have

$$\mathcal{X} \simeq \text{RMod}_A(\Omega\hat{\mathcal{C}}_{\text{top}}), \quad \text{Mod}_A^{E_2}(\Omega\hat{\mathcal{C}}_{\text{top}}) \simeq \text{Vec}.$$

One can also condense A along a line in space. As a consequence, we obtain a condensed 1-codimensional liquid-like defect $\Sigma A = X^2 \boxtimes \overline{X^2} \in \hat{\mathcal{C}}_{\text{top}}$, which is a condensable E_1 -algebra in $\hat{\mathcal{C}}_{\text{top}}$ and can be graphically represented by the following picture.



By condensing ΣA in $\hat{\mathcal{C}}_{\text{top}}$, we again obtain $\mathbf{1}^3$ as the condensed phase and a gapless boundary, which is precisely X^2 . $\square$

⁶⁸In physical reality, by shrinking the radius of the hole, the hole will be gapped out and becomes an ordinary particle. However, in our mathematical theory, there is no length scale. The radius of the cylinder does not play any role. One has to gap it out by hand as it was done in [KZ21]. In this sense, without gapping it out, the cylinder in (6.3.1) should be viewed as a virtual particle (more precisely, it lives in the ∞ -category of ‘generalized nerves’). We will provide more details elsewhere.

⁶⁹There is no unnatural choice involved in the geometric definition of the fusion of two cylinders depicted in (6.3.1).

Remark 6.3.5. When $C^3 = 1^3$, its gapless boundary X^2 is just a 2D RCFT (with liquid-like defects). By condensing the condensable E_2 -algebra $A = \int_{S^1} \mathcal{X}$, or equivalently, condensing ΣA , we obtain 1^3 as the condensed phase and a gapless domain wall $M^2 = X^2$. This result says that condensing a 2D RCFT X^2 in 1^3 reproduces X^2 as the domain wall between the initial phase 1^3 and the condensed phase 1^3 . Note that if X^2 is gapped (i.e., $\Omega \mathcal{B}^{\text{op}}$ is trivial), this construction is already beyond usual anyon condensation theory because we have included the condensation of ‘virtual particles’ (in the category of ‘generalized nerves’). For example, if the cylinder-like defect depicted in (6.3.1) and (6.3.2) has a non-trivial interior, then some non-trivial defect $x \in \mathcal{X}$ in (6.3.2) can be moved into the interior and becomes an E_2 - A -module. In this case, the condensed phase D^3 is non-trivial and we have $\Omega \hat{\mathcal{D}}_{\text{top}} = \text{Mod}_A^{E_2}(\Omega \hat{\mathcal{C}}_{\text{top}})$. This special case already largely generalizes the usual anyon condensation theory even when the wall X^2 is gapped. We will provide a more complete story elsewhere. $\diamond$

Remark 6.3.6. Although this illustrating example is about the condensation in a 2+1D topological order with a gapless liquid-like boundary, it tautologically generalizes to higher dimensional topological orders with gapless liquid-like boundaries (or domain walls) because topological Wick rotation also applies to gapless liquid-like boundaries of higher dimensional topological orders [KZ20, KLWZZ20b, KLWZZ20a, KZ22b]. Moreover, such a generalized condensation has already appeared in Remark 5.3.15 as an example of MME-gaugings. We will provide more details elsewhere. $\diamond$

Remark 6.3.7. By re-examining the physical proof of (3.3.14) (especially recall (3.3.16)), we see the proof is insensitive to the assumption that the domain wall M^n is gapped. Therefore, we generalize this result (3.3.14) to the situations that the domain wall M^n are potentially gapless and liquid-like.

The 1-codimensional condensations of liquid-like gapless defects in C^{n+1} are all defined by the gapped or liquid-like gapless domain walls between the initial phase C^{n+1} and the condensed phase D^{n+1} as condensable E_1 -algebras in $\hat{\mathcal{C}}$.	(6.3.3)
---	---------


As explained in (3.3.16), a condensable E_1 -algebra in the initial phase is precisely defined by the E_1 -algebra of observables living on the gapped/gapless domain wall.

Moreover, recall Remark 3.3.25, (6.3.3) also holds when C^{n+1}, M^n, D^{n+1} are gapless quantum liquids. In other words, a domain wall defines a 1-codimensional condensation should be viewed as a general principle for (at least) all quantum liquids. As a special case of this picture, we expect that a modular-invariant closed 1+1D rational CFT can be obtained by ‘condensing’ any one of its 0+1D boundary CFT’s, which is viewed as a condensable E_1 -algebra in the trivial phase 1^2 . Unfortunately, this generalized condensation theory seems to work only when we ignore the local quantum symmetry of quantum liquids. How to incorporate the data of local quantum symmetry into a truly complete condensation theory remains an open problem. A possible solution was developed in [KZ22c] as a part of the general theory of local quantum symmetry. See Remark 6.3.8 for a related discussion. $\diamond$

Remark 6.3.8. We have ignored the dynamical data X_{qts} completely and have provided a ‘generalized condensation’ only for the topological data. The complete story, including the dynamical data, is not yet known but far richer and more interesting. One way to complete the story is to develop the mathematical theory within the categories of topological nets developed in [KZ22c]. There are other possibilities. One possibility is to generalize the finite type of condensations studied in this work to infinite types. For example, by dropping some finiteness or duality conditions, one can consider more general algebraic objects in a monoidal higher category and develop their higher representation theories. After this paper appeared on arXiv, a mathematical attempt in this direction was developed by Devon Stockall in [Sto25], the physical meaning of which is still waiting to be investigated. On the other hand, since the hom spaces in the enriched categories are internal homs in the category of modules over a vertex operator algebra, the physical realization of these hom spaces are infinite dimensional vector space of defect fields. This infinite feature already shows that the condensation theory presented in this subsection is of infinite type and beyond the main results of this work (recall Remark 2.1.13). $\diamond$

Remark 6.3.9. The generalized condensation theory presented in this subsection seems to suggest a geometric way to understand the usual representation theory of vertex operator algebra (VOA) as a geometric condensation theory. More precisely, consider a rational VOA V , i.e., Mod_V is a modular tensor category. Consider a 2+1D topological order C^3 and a braided monoidal equivalence $\phi : \text{Mod}_V \rightarrow \Omega \mathcal{C}$. According to

the mathematical theory of a gapless boundary of a 2+1D topological order, the triple $(V, \phi, {}^{\Omega\mathcal{C}^{\text{op}}}\Omega\mathcal{C})$ provides a complete mathematical characterization of the so-called canonical gapless boundary of $\mathcal{C}^3$. Consider the solid cylinder in spacetime depicted in the following picture.



It is a spacetime picture of the topological order $\mathcal{C}^3$ and its canonical gapless boundary living in an empty 3D spacetime. The meaning of the label $\mathcal{C}^2$ is explained in Example 2.1.10. Then the representation theory of V or the fact that $\mathcal{C} \simeq \text{Mod}_V$ can be physically understood as the condensation of the solid cylinder (as an E_2 -algebra) in 1^3 (see Remark 6.3.10). Recall that a V -module in the usual sense [FHL93] is precisely an E_2 -module over V . The same picture works if we replace a rational VOA by a rational full field algebras [HK07, Kon07]. $\diamond$

Remark 6.3.10. In order to provide a physical meaning or a physical intuition to a mathematical theory, one has to associate a mathematical object to its physical realization precisely according to physical laws. For example, in mathematical literature, it is perfectly fine to say that a VOA defines an algebra over a spherical partial operad in Huang's geometric theory of VOA [Hua95], or, a local observable algebra on a 1+1D Riemann surface as in the theory of chiral algebra by Beilinson and Drinfeld [BD04]. However, from a physical point of view, a VOA with a non-trivial central charge can not live on a 1+1D worldsheet alone because it is physically anomalous in 1+1D even when it is a holomorphic VOA. It can only be physically realized as a 1+1D chiral gapless boundary of a 2+1D topological order. Moreover, the algebraic structure of the cylinder in (6.3.4) is defined by geometrically fusing the cylinder which was defined in [KZ21, Section 6.2]. In other words, we can depict this algebraic structure as a (3D filled) pant with two (filled) holes lying on the plane labeled by 1^3 and the third (filled) hole lying above the plane. Recall that, in [KZ21, Section 6.2], this fusion is defined topologically. In order to understand the algebraic structure of a VOA physically in this context, it is better to equip the surface of the solid cylinder with a complex structure⁷⁰. This yet-conjectural geometric meanings of a VOA is different from Huang's geometric theory of VOA [Hua95] and Beilinson-Drinfeld's chiral algebra [BD04]. Therefore, it becomes a very interesting problem to develop the mathematical foundation of above geometric pictures. Another deeply related problem, proposed by the first author of this paper a few years ago, is to established a functorial quantum field theory (as a functor from cobordism category to a linear category) that combines the usual Reshetikhin-Turaev TQFT's with a gapless boundary. $\diamond$

6.4 Generalized lattice models with categorified local degrees of freedom

We have shown that the condensation theory is essentially about higher algebras and higher representations, the mathematical theory of which is far from being developed. In this subsection, we discuss some physical or philosophical enlightenments from such a general theory of condensations.

Physically, if we reduce the condensation to its simplest form, it is nothing but a way to select a sub-Hilbert space from a Hilbert space as emphasized in [Kon14]. For example, consider the tensor product of two spins spaces V 's, i.e.

$$V \otimes_{\mathbb{C}} V \simeq V_0 \oplus V_0^{\perp}, \quad (6.4.1)$$

which is quantum mechanical system instead of a quantum many-body system. Introducing an interaction between these two spin spaces amounts to introducing a projector to the subspace V_0 , which is more energy favorable than $V_0^{\perp}$. Although condensation is a term applying to quantum many-body systems, we can view an interaction between two spin spaces, or more generally, an interaction among a few spin spaces, as a condensation in a quantum few-body system.

When we have a chain of spin spaces $\otimes_{i \in \mathbb{Z}} V$, by introducing mutually commuting local projectors (MCLP) (or condensation) among adjacent spins, we obtain a 1+1D lattice model (see Remark 6.4.1). We refer to this type of lattice models by *MCLP lattice models*. If a MCLP lattice model realizes a quantum phase $\mathcal{C}^2$,

⁷⁰Such a solid cylinder can be viewed as a special case of manifolds with metric.

it is reasonable to say that this lattice model realize a condensation from the trivial phase $\mathbf{1}^2$ to C^2 . This point of view automatically generalizes to all dimensions. Namely, a MCLP lattice model realization of a quantum phase C^{n+1} is precisely a physical realization of a condensation from the trivial phase $\mathbf{1}^{n+1}$ to C^{n+1} (see Remark 6.4.1). In other words, a MCLP lattice model and a condensation are essentially the same thing.

Remark 6.4.1. Our condensation theory is based on condensation maps, which are mutually commuting projectors. Therefore, it only applies to lattice models with mutually commuting local interactions (or condensation). On the other hand, a MCLP lattice model construction is already more general than what is covered in this work because the quantum phase C^{n+1} can also be a non-liquid phase (see [Cha05, SSWC18, SSC19, VHF16, Wen20] and references therein). A quantum liquid can be viewed as a quantum phase satisfying certain finite properties because fully dualizability condition can be viewed as a generalized finiteness condition. Therefore, when C^{n+1} is a quantum liquid, we can view a condensation from $\mathbf{1}^{n+1}$ to C^{n+1} as a finite-type condensation; when C^{n+1} is a quantum non-liquid, the associated condensation is an infinite-type condensation. We hope to give a precise mathematical definition of a finite-type or infinite-type condensation in the future. If we allow condensations of infinite-type, then the trivial phase $\mathbf{1}^{n+1}$ can already be viewed as a ‘theory of everything’. $\diamond$

This unification of these terminologies or ideas leads us to many natural generalizations of previous ideas. We give some examples.

1. An $n+1$ D MCLP lattice model can be viewed as first introducing an n D MCLP lattice model followed by a layer construction. In other words, higher dimensional MCLP lattice models are layer constructions of lower dimensional MCLP lattice models.




Figure 20: the construction of a universal MCLP lattice model

2. It also suggests that, for a quantum phase C^{n+1} , there is a universal construction of a ‘MCLP lattice model’ that can realize C^{n+1} . The idea is illustrated in Figure 20. Each square depicts an $n+1$ D quantum phase C^{n+1} surrounded by a (potentially gapless) boundary phase X^n . Such a square should be viewed as a generalized local spin space V , and is clearly infinite dimensional. Note that some local degrees of freedom in V are those on the ‘local boundary’ X^n and others are those in the C^{n+1} -phase. We define the interaction between two adjacent X^n -edge in Figure 20 simply by introducing an interactions between the local degrees of freedom in V that belong to the closest X^n -edges belonging to two adjacent squares such that this interaction defines a condensation⁷¹ that erases the local boundary and connects the interior of two adjacent squares (see Remark 6.4.3). This MCLP lattice model realizes a condensation from $\mathbf{1}^{n+1}$.

- When X^n is gapped, this MCLP lattice model realizes a condensation of ‘finite type’ from $\mathbf{1}^{n+1}$. We expect that this condensation is a continuous phase transition⁷².

⁷¹In [KZ21, Section 6.2], two of us showed how to do it categorically as a fusion process when $n = 2$, X^2 is gapless and C^3 is gapped.

⁷²In general, on the phase boundary between two phases on a phase diagram, the transition points are often of the first order instead of the second order. However, as we are approaching a new age of structured new calculus of quantum many-body physics, we suspect that there are much more refined structures underlying a ‘phase transition’ than the usual notions of the first and second order transitions (see Remark 6.4.3) because these usual notions are the byproducts of the usual-calculus-based (or decategorified) physics.

- When X^n is not-gappable gapless and liquid-like⁷³, this MCLP lattice model realizes a condensation from 1^{n+1} that is more violent than the finite type. Therefore, it is reasonable to call such a condensation an ‘infinite type’. We still expect that the phase transition to be continuous. When X^n is not liquid-like, then we expect it to be ‘discontinuous’.
3. Since the notion of condensation is defined not only from 1^{n+1} to C^{n+1} but also from B^{n+1} to C^{n+1} , we should also have a notion of a *lattice model over a background phase* B^{n+1} , which realizes a new phase C^{n+1} and, at the same time, a condensation from B^{n+1} to C^{n+1} . Moreover, the condensation process defined by first proliferating C^{n+1} -phase-islands within the B^{n+1} -phase then introducing proper interactions among C^{n+1} -phase-islands immediately leads us to a universal construction of a MCLP lattice model over the background phase B^{n+1} that realizes the phase C^{n+1} (see Remark 6.4.3). An illustration of this construction is given in Figure 20 with the background phase given by B^{n+1} . An example of such a MCLP lattice model over a non-trivial background phase B^3 is given in [MRV23].




Figure 21: a universal MCLP lattice model with a brackground phase B^{n+1}

- When X^n is gapped, this MCLP lattice model realizes a condensation of ‘finite type’ from B^{n+1} . We expect this condensation to be realized by a continuous phase transition.
- When X^n is not gappable gapless but liquid-like, this MCLP lattice model realizes a condensation from B^{n+1} that is of ‘infinite type’. We still expect that it can be realized by a continuous phase transition. If X^n is not liquid-like, we expect it to be a discontinuous phase transition.

Remark 6.4.2. In a *lattice model over a background phase* B^{n+1} , the tensor product of local spin spaces is a tensor product in $\Omega^{n-1}\mathcal{B}$ instead of the vector space tensor product $\otimes_{\mathbb{C}}$. Therefore, such a model is not local in the usual sense (see [MRV23]). However, we believe that such generalized lattice models are very natural and valuable for theoretical studies. $\diamond$

Remark 6.4.3. We use the interaction between local degrees of freedom in V to effectively erase the local boundaries and connect the interior of two adjacent squares in order to make sure that we are indeed talking about a lattice model in the usual sense. However, it is perhaps more interesting to view the process of erasing the local boundaries and connecting the interior of two adjacent squares as a generalization of the local interactions in a usual lattice model. For example, we can define this process by a condensation, which can be defined categorically without referring to any local Hilbert space or Hamiltonian. As a consequence, we can generalize the notion of a lattice model by replacing local interactions among local degrees of freedom by local condensations or local phase transitions or other categorical or quantum many-body process. That amounts to say that “the space of local degrees of freedom” (i.e., a vector space) can be replaced by “a more structuralized space of local degrees of freedom”, such as a new quantum phases (as we depicted in Figure 20 and Figure 21) or an separable m -category, which is also called an $(m+1)$ -vector space as an object in $(m+1)\text{Vec}$; and “local interactions” can be replaced by “local condensations”. We will develop

⁷³Then we expect that its bulk C^{n+1} is also liquid-like.

such a generalized lattice models elsewhere. We believe that we are at the beginning stage of developing a structuralized new calculus for quantum many-body physics or QFT's. $\diamond$

Conjecture 6.4.4. Above discussion motivates the following two conjectures.

1. For each $n+1$ D anomaly-free topological order C^{n+1} Morita equivalent to B^{n+1} , there is an $n+1$ D MCLP lattice model over the background phase B^{n+1} realizing C^{n+1} such that the local spin space V is finite dimensional.
2. For each $n+1$ D anomaly-free topological order C^{n+1} not Morita equivalent to B^{n+1} , there is an $n+1$ D MCLP lattice model over the background phase B^{n+1} realizing C^{n+1} such that the local spin space V is necessarily infinite dimensional.

Remark 6.4.5. $2+1$ D Levin-Wen models [LW05] or their higher dimensional analogues (see [GJF19] and Remark 5.11 in [KZ22b]) can be viewed as MCLP lattice model realizations of non-chiral topological orders. This observation can be reformulated in various ways and has already inspired new mathematical theories. One example is the condensation theory of topological nets developed in [KZ22c]. We hope to come back to this point in the future. $\diamond$

Remark 6.4.6. The physical configuration depicted in Figure 20 or 21 also shows that, before we introduce the interactions, the system is a non-liquid because the ground state degeneracy scales as the size of the system. This fact also suggests that a complete mathematical theory of condensations needs a mathematical theory of non-liquid phases and that of gapless quantum liquids. We believe that it is reasonable to call this yet-conjectural complete theory of condensations as a ‘theory of everything’ not only because it contains all interesting phases and phase transitions among them but also because it provides an interesting approach towards a theory of quantum gravity. Indeed, a theory of quantum gravity is expected to describe the early universe, which, as it expands and cools down, should produces a topological order or quantum liquid via finitely many (possibly more general) condensations and vice versa. It suggests that the emergence of space and time can be a consequence of (possibly more general) condensations from relatively elementary quantum many-body systems. This approach towards quantum gravity is an untold dream that has been guiding our research for many years. $\diamond$


A Appendix

A.1 Proof of Theorem 4.1.19

The idea of proof is sketched below Theorem 4.1.19. As we mentioned below (4.1.17), it remains to show that the morphism (4.1.17) coincides with the one defined in (4.1.16). The morphism (4.1.17) is defined via the composition of a series of morphisms. We consider a more general condition by replacing the object 1_m by arbitrary object x, y in $\Omega_m \mathcal{M}$. The morphism we need to study is provided by the following composition

$$L_2^R(x) \otimes L_2^R(y) \rightarrow L_2^R(L_2(L_2^R(x) \otimes L_2^R(y))) \simeq L_2^R(L_2 L_2^R(x) \otimes L_2 L_2^R(y)) \rightarrow L_2^R(x \otimes y) \quad (\text{A.1.1})$$


We examine the geometric meaning of each morphism appearing in the above composition. It turns out that the geometric meaning of (A.1.1) is the fusion of two bubbles.




As a result, by taking $x, y = 1_m$, we recover the morphism (4.1.16).

Before we start, we analyze the geometric meaning of the unit and counit of the adjunction $L_2 \dashv [1_m, -]$, which are used throughout the proof. Recall that, the bulk-to-wall map $L_2 : \Omega\mathcal{C} \rightarrow \Omega_m \mathcal{M}$ means we move a point-like excitation a in $\Omega\mathcal{C}$ next to the gapped domain wall $\Omega_m \mathcal{M}$, so that we can view a as an excitation on $\Omega_m \mathcal{M}$, and we denote it by $L_2(a)$.

- Unit of the adjunction $\eta : \text{id}_{\Omega\mathcal{C}} \Rightarrow [1_m, L_2(-)] = L_2^R L_2(-) = - \otimes L_2^R(1_m)$:



- Counit of the adjunction $\varepsilon : L_2 L_2^R(-) L_2([1_m, -]) \Rightarrow \text{id}_{\Omega_m \mathcal{M}}$:



Then we need to analyze the geometric meaning of following morphisms, whose composition gives rise to (A.1.1).

1. The first morphism is a component of the unit η . It is given by

$$\eta_{L_2^R(x) \otimes L_2^R(y)} : L_2^R(x) \otimes L_2^R(y) \rightarrow L_2^R(L_2(L_2^R(x) \otimes L_2^R(y))) \quad (\text{A.1.2})$$

2. The second morphism

$$L_2^R(L_2(L_2^R(x) \otimes L_2^R(y))) \simeq L_2^R(L_2(L_2^R(x)) \otimes L_2(L_2^R(y))) \quad (\text{A.1.3})$$

only exploits the monoidal structure of L_2 , hence is geometrically trivial. We omit this morphism in our geometric analysis.

3. The third morphism is given by

$$L_2^R(\varepsilon_x \otimes \varepsilon_y) : L_2^R(L_2(L_2^R(x)) \otimes L_2(L_2^R(y))) \rightarrow L_2^R(x \otimes y)$$

We study the morphism inside $L_2^R(-)$, that is, the tensor product of two components of ε :

$$\varepsilon_x \otimes \varepsilon_y : L_2(L_2^R(x)) \otimes L_2(L_2^R(y)) \rightarrow x \otimes y. \quad (\text{A.1.4})$$

The analysis of the morphism (A.1.4) occupies the main space of this appendix.

The geometric meaning of the morphism (A.1.2) is:

$$\eta_{L_2^R(x) \otimes L_2^R(y)} : \begin{array}{c} \text{Diagram 1: A blue box labeled } \Omega\mathcal{D} \text{ with a red dot } x \text{ on top, and another blue box labeled } \Omega\mathcal{D} \text{ with a red dot } y \text{ on top, both inside a larger blue box labeled } \Omega\mathcal{C}. \end{array} \longrightarrow \begin{array}{c} \text{Diagram 2: Two blue boxes labeled } \Omega\mathcal{D} \text{ with red dots } x \text{ and } y \text{ on top, and a third blue box labeled } \Omega\mathcal{D} \text{ below them, all inside a larger blue box labeled } \Omega\mathcal{C}. \end{array}$$

The geometric meaning of the morphism (A.1.4) is:

$$\varepsilon_x \otimes \varepsilon_y : \begin{array}{c} \text{Diagram 1: A blue box labeled } \Omega\mathcal{D} \text{ with a red dot } x \text{ on top, and another blue box labeled } \Omega\mathcal{D} \text{ with a red dot } y \text{ on top, both inside a larger blue box labeled } \Omega\mathcal{C}. \end{array} = \int_{z \in \Omega_m \mathcal{M}}^{\Omega\mathcal{D}} \int_{w \in \Omega_m \mathcal{M}}^{\Omega\mathcal{D}} \begin{array}{c} \text{Diagram 2: A blue box labeled } \Omega\mathcal{D} \text{ with a red dot } x \text{ on top, and another blue box labeled } \Omega\mathcal{D} \text{ with a red dot } y \text{ on top, both inside a larger blue box labeled } \Omega\mathcal{C}. \end{array} \longrightarrow \begin{array}{c} \text{Diagram 3: A blue box labeled } \Omega\mathcal{D} \text{ with a red dot } x \text{ on top, and another blue box labeled } \Omega\mathcal{D} \text{ with a red dot } y \text{ on top, both inside a larger blue box labeled } \Omega\mathcal{C}. \end{array} = \begin{array}{c} \text{Diagram 4: A blue box labeled } \Omega\mathcal{D} \text{ with a red dot } x \text{ on top, and another blue box labeled } \Omega\mathcal{D} \text{ with a red dot } y \text{ on top, both inside a larger blue box labeled } \Omega\mathcal{C}. \end{array}$$

To proceed, let us note that there is a canonical isomorphism:

$$\begin{array}{c} \int_{z \in \Omega_m \mathcal{M}}^{\Omega\mathcal{D}} \begin{array}{c} \text{Diagram 1: A blue box labeled } \Omega\mathcal{D} \text{ with a red dot } x \text{ on top, and another blue box labeled } \Omega\mathcal{D} \text{ with a red dot } y \text{ on top, both inside a larger blue box labeled } \Omega\mathcal{C}. \end{array} = \begin{array}{c} \text{Diagram 2: A blue box labeled } \Omega\mathcal{D} \text{ with a red dot } x \text{ on top, and another blue box labeled } \Omega\mathcal{D} \text{ with a red dot } y \text{ on top, both inside a larger blue box labeled } \Omega\mathcal{C}. \end{array} \\ = \begin{array}{c} \text{Diagram 3: A blue box labeled } \Omega\mathcal{D} \text{ with a red dot } x \text{ on top, and another blue box labeled } \Omega\mathcal{D} \text{ with a red dot } y \text{ on top, both inside a larger blue box labeled } \Omega\mathcal{C}. \end{array} = \int_{z \in \Omega_m \mathcal{M}}^{\Omega\mathcal{D}} \begin{array}{c} \text{Diagram 4: A blue box labeled } \Omega\mathcal{D} \text{ with a red dot } x \text{ on top, and another blue box labeled } \Omega\mathcal{D} \text{ with a red dot } y \text{ on top, both inside a larger blue box labeled } \Omega\mathcal{C}. \end{array} \end{array}$$

Algebraically, this means that the object $\int_{z \in \Omega_m \mathcal{M}}^{\Omega\mathcal{D}} z^R \otimes x \otimes z$ is equipped with a half braiding β whose components are $\int_{z \in \Omega_m \mathcal{M}}^{\Omega\mathcal{D}} z^R \otimes x \otimes z \otimes y \simeq y \otimes (\int_{z \in \Omega_m \mathcal{M}}^{\Omega\mathcal{D}} z^R \otimes x \otimes z) \simeq \int_{z \in \Omega_m \mathcal{M}}^{\Omega\mathcal{D}} y \otimes z^R \otimes x \otimes z$. The mathematical details of the construction of the half-braiding are postponed to the next appendix, see Example A.2.15 and Example A.2.16.

Now we would like to show that, the morphism $\varepsilon_x \otimes \varepsilon_y$ is equal to the following morphism:

$$\begin{aligned}
& \text{Diagram 1: } \int_{z \in \Omega_m \mathcal{M}} \int_{w \in \Omega_m \mathcal{M}} \text{Diagram 2} \\
&= \int_{z \in \Omega_m \mathcal{M}} \int_{w \in \Omega_m \mathcal{M}} \text{Diagram 3} \xrightarrow{\beta} \int_{z \in \Omega_m \mathcal{M}} \int_{w \in \Omega_m \mathcal{M}} \text{Diagram 4} \\
&= \int_{z \in \Omega_m \mathcal{M}} \int_{w \in \Omega_m \mathcal{M}} \text{Diagram 5} \rightarrow \text{Diagram 6} \\
&= \text{Diagram 7}
\end{aligned}$$

Mathematically, this amounts to the commutativity of the following diagram


$$\begin{array}{ccc}
\int_{z \in \Omega_m \mathcal{M}} z^R \otimes x \otimes z \otimes \int_{w \in \Omega_m \mathcal{M}} w^R \otimes y \otimes w & \longrightarrow & \int_{w \in \Omega_m \mathcal{M}} w^R \otimes \left(\int_{z \in \Omega_m \mathcal{M}} z^R \otimes x \otimes z \right) \otimes y \otimes w \\
\downarrow \pi_{1_m} \otimes \pi_{1_m} & & \downarrow \pi_{1_m} \\
& & \int_{w \in \Omega_m \mathcal{M}} w^R \otimes x \otimes y \otimes w \\
& & \downarrow \pi_{1_m} \\
x \otimes y & \xlongequal{\quad} & x \otimes y
\end{array}$$

This commutativity can be demonstrated by considering the following diagram:


$$\begin{array}{ccccc}
\int_{z \in \Omega_m \mathcal{M}} z^R \otimes x \otimes z \otimes \int_{w \in \Omega_m \mathcal{M}} w^R \otimes y \otimes w & \longrightarrow & \int_{w \in \Omega_m \mathcal{M}} w^R \otimes \int_{z \in \Omega_m \mathcal{M}} z^R \otimes x \otimes z \otimes y \otimes w & & \\
\downarrow & & \downarrow & \searrow & \\
\int_{z \in \Omega_m \mathcal{M}} z^R \otimes x \otimes z \otimes 1_m \otimes y \otimes 1_m & \longrightarrow & 1_m \otimes \int_{z \in \Omega_m \mathcal{M}} z^R \otimes x \otimes z \otimes y \otimes 1_m & & \int_{w \in \Omega_m \mathcal{M}} w^R \otimes x \otimes y \otimes w \\
\downarrow & & \downarrow & \nearrow & \\
x \otimes y & \longrightarrow & x \otimes y & &
\end{array}$$

The upper square commutes due to the definition of the upper horizontal arrow. The middle horizontal morphism is the half-braiding with the tensor unit 1_m and so is trivial, hence the lower square commutes. The right triangle commutes trivially.


As a result, the morphism $L_2^R(\varepsilon_x \otimes \varepsilon_y)$ can be depicted as (where we use the “tunnels” between bubbles to indicate the way we fuse bubbles):



Summarizing above analysis, we obtain that the composed morphism (A.1.1) can be written as the composition of the following maps:




By the naturality of η we may rewrite the above map as



Note that the composition of the last two maps is just identity, by the zig-zag equation of adjunction:

$$\text{id} : L_2^R(x \otimes y) \rightarrow L_2^R(x \otimes y) \otimes L_2^R(1_m) = L_2^R(L_2 L_2^R(x \otimes y)) \xrightarrow{L_2^R(\varepsilon_{x \otimes y})} L_2^R(x \otimes y)$$

Finally, we see that the morphism $L_2^R(x) \otimes L_2^R(y) \rightarrow L_2^R(x \otimes y)$ is given by fusion of two bubbles:



This completes the proof of Theorem 4.1.19.

A.2 Module Eilenberg-Watts calculus

In this section, we develop a relative version of classical Eilenberg-Watts calculus. It is different from the one developed in [FSS20]. All categories we consider in this appendix are finite 1-categories, i.e. linear

categories that are equivalent to the category of finite dimensional modules over some finite dimensional algebra A .

Convention A.2.1 (Convention for duality). Our convention for left/right duality is different from that in [DR18] [Lur17]. For $\mathcal{C}$ a monoidal category and $x \in \mathcal{C}$ an object, a left dual of x consists of a triple $(x^L, \text{ev}_x, \text{coev}_x)$ where $x^L \in \mathcal{C}$ is an object, $\text{coev}_x : \mathbb{1}_{\mathcal{C}} \rightarrow x \otimes x^L$ is the coevaluation map and $\text{ev}_x : x^L \otimes x \rightarrow \mathbb{1}_{\mathcal{C}}$ is the evaluation map. The (co)evaluation maps are required to satisfy the zig-zag equation. A right dual of x is a left dual in $\mathcal{C}^{\text{rev}}$.

Notation A.2.2. Let $\mathcal{C}$ be a finite tensor category and $\mathcal{M}$ is a left $\mathcal{C}$ -module, whose module structure is denoted by $\odot : \mathcal{C} \times \mathcal{M} \rightarrow \mathcal{M}$. The rigidity of $\mathcal{C}$ induces two right $\mathcal{C}$ -module structures over $\mathcal{M}^{\text{op}}$, defined in the following way:

- The right $\mathcal{C}$ -action is defined by:

$$\odot^L : \mathcal{M}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{M}^{\text{op}}, \quad (m, c) \mapsto c^L \odot m.$$

For simplicity, we denote the right $\mathcal{C}$ -module structure by $\mathcal{M}^{\text{op}L}$.

- The right $\mathcal{C}$ -action is defined by:

$$\odot^R : \mathcal{M}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{M}^{\text{op}}, \quad (m, c) \mapsto c^R \odot m.$$

For simplicity, we denote the right $\mathcal{C}$ -module structure by $\mathcal{M}^{\text{op}R}$.

Notation A.2.3. Let $\mathcal{C}$ be a finite tensor category and $\mathcal{M}, \mathcal{N}$ be finite left $\mathcal{C}$ -modules. We use $\text{Fun}_{\mathcal{C}}^L(\mathcal{M}, \mathcal{N})$ to denote the finite category of left exact $\mathcal{C}$ -module functors from $\mathcal{M}$ to $\mathcal{N}$. Similarly, we use $\text{Fun}_{\mathcal{C}}^R(\mathcal{M}, \mathcal{N})$ to denote the finite category of right exact $\mathcal{C}$ -module functors from $\mathcal{M}$ to $\mathcal{N}$. For two $\mathcal{C}$ -module functors $(F, \eta_F), (G, \eta_G)$ in $\text{Fun}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})$, we write $\text{Nat}_{\mathcal{C}}(F, G)$ for the vector space of $\mathcal{C}$ -module natural transformations from F to G .

Definition A.2.4. Let $\mathcal{C}$ be a finite tensor category and $\mathcal{M}$ a left $\mathcal{C}$ -module. We equip $\mathcal{M}^{\text{op}}$ with the right $\mathcal{C}$ -module structure $\mathcal{M}^{\text{op}L}$. Let $(F, e) : \mathcal{M}^{\text{op}L} \times \mathcal{M} \rightarrow \mathcal{D}$ be a balanced $\mathcal{C}$ -module functor with e being the balancing natural isomorphism. The $\mathcal{C}$ -module end of F is a pair $(\int_{x \in \mathcal{M}}^{\mathcal{C}} F(x, x), \pi)$ with $\int_{x \in \mathcal{M}}^{\mathcal{C}} F(x, x) \in \mathcal{D}$ an object and $\pi : \Delta_{\int_{x \in \mathcal{M}}^{\mathcal{C}} F(x, x)} \xrightarrow{\bullet\bullet} F$ is a dinatural transformation such that the following diagram is commutative:

$$\begin{array}{ccc} \int_{x \in \mathcal{M}}^{\mathcal{C}} F(x, x) & \xrightarrow{\pi_m} & F(m, m) \\ \pi_{c^L \odot m} \downarrow & & \downarrow \text{coev}_c \\ F(c^L \odot m, c^L \odot m) & \xrightarrow{e_{m, c, c^L \odot m}} & F(m, c \odot c^L \odot m) \end{array}$$

and the pair $(\int_{x \in \mathcal{M}}^{\mathcal{C}} F(x, x), \pi)$ is terminal among all such pairs. $\square$

This is a dual notion defined as follows:

Definition A.2.5. Let $\mathcal{C}$ be a finite tensor category and $\mathcal{M}$ a left $\mathcal{C}$ -module. We equip $\mathcal{M}^{\text{op}}$ with the right $\mathcal{C}$ -module structure $\mathcal{M}^{\text{op}R}$. Let $(F, e) : \mathcal{M}^{\text{op}R} \times \mathcal{M} \rightarrow \mathcal{D}$ be a balanced $\mathcal{C}$ -module functor with e being the balancing natural isomorphism. The $\mathcal{C}$ -module coend of F is a pair $(\int_{\mathcal{C}}^{x \in \mathcal{M}} F(x, x), \pi)$ with $\int_{\mathcal{C}}^{x \in \mathcal{M}} F(x, x) \in \mathcal{D}$ an object and $\pi : F \xrightarrow{\bullet\bullet} \Delta_{\int_{\mathcal{C}}^{x \in \mathcal{M}} F(x, x)}$ a dinatural transformation, such that the following diagram is commutative:

$$\begin{array}{ccc} F(m, (c \otimes c^R) \odot m) & \longrightarrow & F(c^R \odot m, c^R \odot m) \\ \downarrow \text{ev}_{c^R} & & \downarrow \\ F(m, m) & \longrightarrow & \int_{\mathcal{C}}^{x \in \mathcal{M}} F(x, x) \end{array}$$

and the pair $(\int_{\mathcal{C}}^{x \in \mathcal{M}} F(x, x), \pi)$ is initial among all such pairs. $\square$

Remark A.2.6. The $\mathcal{C}$ -module end may be written as the following equalizer:

$$\int_{m \in \mathcal{M}}^{\mathcal{C}} F(m, m) \simeq \text{Eq}(\int_{m \in \mathcal{M}} F(m, m) \xrightarrow[f]{g} \int_{m \in \mathcal{M}} \int_{c \in \mathcal{C}} F(m, c \odot c^L \odot m))$$

Componentwise, f and g can be written respectively as (we choose the component labelled by $d \in \mathcal{C}$ and $n \in \mathcal{M}$)

$$f_{d,n} = \int_{m \in \mathcal{M}} F(m, m) \xrightarrow{\pi_n} F(n, n) \rightarrow F(n, d \otimes d^L \odot n)$$

and

$$g_{d,n} = \int_{m \in \mathcal{M}} F(m, m) \xrightarrow{\pi_{d^L \odot n}} F(d^L \odot n, d^L \odot n) \simeq F(n, d^L \otimes d \odot n).$$

where we use the balancing structure of F in the last step. Dually the $\mathcal{C}$ -module coend may be written as a coequalizer of two coends. $\diamond$

The following lemma generalizes the familiar formula for usual ends:

Lemma A.2.7. Let $\mathcal{M}, \mathcal{N}$ be left modules over a finite tensor category $\mathcal{C}$. Let $(F, \eta^F), (G, \eta^G)$ be $\mathcal{C}$ -module functors. Then $\int_{m \in \mathcal{M}}^{\mathcal{C}} \text{hom}_{\mathcal{N}}(F(m), G(m)) \simeq \text{Nat}_{\mathcal{C}}(F, G)$. Note that the functor $\text{hom}_{\mathcal{N}}(F(-), G(-)) : \mathcal{M}^{\text{op}|L} \times \mathcal{M} \rightarrow \text{Vec}$ is equipped with a canonical balancing structure.

Proof. Take $\alpha \in \text{Nat}_{\mathcal{C}}(F, G)$ and $\pi_m(\alpha) = \alpha_m$ be its component at m . Then the commutativity of the following diagram

$$\begin{array}{ccc} \text{Nat}_{\mathcal{C}}(F, G) & \xrightarrow{\pi_m} & \text{hom}_{\mathcal{N}}(F(m), G(m)) \\ \pi_{c^L \odot m} \downarrow & & \downarrow \\ \text{hom}_{\mathcal{N}}(F(c^L \odot m), G(c^L \odot m)) & \xrightarrow{e_{m, c, c^L \odot m}} & \text{hom}_{\mathcal{N}}(F(m), G(c \odot c^L \odot m)) \end{array}$$

is equivalent to be commutativity of the following diagram

$$\begin{array}{ccc} c^L \odot F(m) & \xrightarrow{c^L \odot \alpha_m} & c^L \odot G(m) \\ \downarrow \eta_{c^L, m}^F & & \downarrow \eta_{c^L, m}^G \\ F(c^L \odot m) & \xrightarrow{\alpha_{c^L \odot m}} & G(c^L \odot m) \end{array}$$

The following lemma is a simple corollary of the enriched Yoneda Lemma, which will be useful in our proof of Lemma A.2.9.

Lemma A.2.8. Let $\mathcal{M}, \mathcal{N}$ be left $\mathcal{C}$ -modules and $(K, \eta^K) \in \text{Fun}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})$ is a $\mathcal{C}$ -module functor. There is a canonical isomorphism of vector spaces for each $y \in \mathcal{M}, z \in \mathcal{N}$:

$$\text{Nat}_{\mathcal{C}}([y, -], [z, K(-)]) \simeq \text{hom}_{\mathcal{N}}(z, K(y)).$$

Similarly we have

$$\text{Nat}_{\mathcal{C}}([- , y], [K(-), z]) \simeq \text{hom}_{\mathcal{N}}(K(y), z).$$

Lemma A.2.9 (Yoneda and co-Yoneda Lemma via (co)ends). Let $(G, \eta^G) \in \text{Fun}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})$, then there is a canonical isomorphism of $\mathcal{C}$ -module functors:

$$G(-) \simeq \int_{\mathcal{C}}^{x \in \mathcal{M}} [x, -] \odot G(x), \quad G(-) \simeq \int_{x \in \mathcal{M}}^{\mathcal{C}} [-, x]^R \odot G(x) \quad (\text{A.2.1})$$

Proof.

$$\begin{aligned} \text{hom}(\int_{\mathcal{C}}^{x \in \mathcal{M}} [x, y] \odot G(x), z) &\simeq \int_{x \in \mathcal{M}}^{\mathcal{C}} \text{hom}_{\mathcal{N}}([x, y] \odot G(x), z) \\ &\simeq \int_{x \in \mathcal{M}}^{\mathcal{C}} \text{hom}_{\mathcal{C}}([x, y], [G(x), z]) \\ &\simeq \text{Nat}_{\mathcal{C}}([- , y], [G(-), z]) \\ &\simeq \text{hom}_{\mathcal{N}}(G(y), z). \end{aligned}$$

Example A.2.10. Let us regard $\mathcal{C}$ as a regular left $\mathcal{C}$ -module. In this case the internal hom is easily computed as

$$[x, y] = y \otimes x^L$$

Applying the Yoneda lemma A.2.9 to the identity functor $\text{id} : \mathcal{C} \rightarrow \mathcal{C}$ (which is equipped with a canonical $\mathcal{C}$ -module functor structure), we have

$$y \simeq \int_{x \in \mathcal{C}} [y, x]^R \otimes x \simeq \int_{x \in \mathcal{C}} y \otimes x^R \otimes x, \quad \mathbb{1}_{\mathcal{C}} \simeq \int_{x \in \mathcal{C}} x^R \otimes x. \quad (\text{A.2.2})$$

♡

Theorem A.2.11 (Generalized Eilenberg-Watts calculus). Let $\mathcal{C}$ be a finite tensor category. We use $\text{Fun}_{\mathcal{C}}^R(\mathcal{M}, \mathcal{N})$ to denote the category of right exact $\mathcal{C}$ -module functors from $\mathcal{M}$ to $\mathcal{N}$ and $\text{Fun}_{\mathcal{C}}^L(\mathcal{M}, \mathcal{N})$ the category of left exact ones. Then there are pairs of adjoint equivalences:

$$\begin{aligned} \Phi^l : \mathcal{M}^{\text{op}|R} \boxtimes_{\mathcal{C}} \mathcal{N} &\rightarrow \text{Fun}_{\mathcal{C}}^L(\mathcal{M}, \mathcal{N}), & x \boxtimes_{\mathcal{C}} y &\mapsto [x, -] \odot y \\ \Psi^l : \text{Fun}_{\mathcal{C}}^L(\mathcal{M}, \mathcal{N}) &\rightarrow \mathcal{M}^{\text{op}|R} \boxtimes_{\mathcal{C}} \mathcal{N}, & F &\mapsto \int_{\mathcal{C}}^{m \in \mathcal{M}} m \boxtimes_{\mathcal{C}} F(m). \end{aligned}$$

and

$$\begin{aligned} \Phi^r : \mathcal{M}^{\text{op}|L} \boxtimes_{\mathcal{C}} \mathcal{N} &\rightarrow \text{Fun}_{\mathcal{C}}^R(\mathcal{M}, \mathcal{N}), & x \boxtimes_{\mathcal{C}} y &\mapsto [-, x]^R \odot y \\ \Psi^r : \text{Fun}_{\mathcal{C}}^R(\mathcal{M}, \mathcal{N}) &\rightarrow \mathcal{M}^{\text{op}|L} \boxtimes_{\mathcal{C}} \mathcal{N}, & F &\mapsto \int_{m \in \mathcal{M}}^{\mathcal{C}} m \boxtimes_{\mathcal{C}} F(m). \end{aligned}$$

Example A.2.12. Let $\mathcal{C}$ be a braided fusion category, $\mathcal{M}$ be a multi-fusion category and $L : \mathcal{C} \rightarrow \mathcal{M}$ be a central functor. L equips $\mathcal{M}$ with a structure of left $\mathcal{C}$ -module in a manifest way. According to Theorem A.2.11, there is an equivalence

$$\Psi^r : \text{Fun}_{\mathcal{C}}(\mathcal{M}, \mathcal{M}) \simeq \mathcal{M}^{\text{op}|L} \boxtimes_{\mathcal{C}} \mathcal{M}, \quad F \mapsto \int_{m \in \mathcal{M}}^{\mathcal{C}} m \boxtimes F(m).$$

In particular, Ψ^r sends the identity functor $\text{id}_{\mathcal{M}}$ to $\int_{m \in \mathcal{M}}^{\mathcal{C}} m \boxtimes m$. Taking the equivalence of right $\mathcal{C}$ -modules

$$\begin{aligned} \mathcal{M}^{\text{op}|L} &\simeq \mathcal{M}^{\text{rev}}, \\ m &\mapsto m^R \end{aligned}$$

into consideration, we obtain a composed equivalence:

$$\text{Fun}_{\mathcal{C}}(\mathcal{M}, \mathcal{M}) \simeq \mathcal{M}^{\text{op}|L} \boxtimes_{\mathcal{C}} \mathcal{M} \simeq \mathcal{M}^{\text{rev}} \boxtimes_{\mathcal{C}} \mathcal{M},$$

which sends the identity functor $\text{Id}_{\mathcal{M}}$ to:

$$\text{id}_{\mathcal{M}} \mapsto \int_{m \in \mathcal{M}}^{\mathcal{C}} m \boxtimes_{\mathcal{C}} m \mapsto \int_{m \in \mathcal{M}}^{\mathcal{C}} m^R \boxtimes_{\mathcal{C}} m. \quad (\text{A.2.3})$$

This image of $\text{id}_{\mathcal{M}}$ is used in Section 4.1.3 and Section A.1. ♡

To give a proof of Theorem A.2.11, we first summarize the following results from [KZ18a], Corollary 2.2.6 and Proposition 2.2.7.

Theorem A.2.13. Let $\mathcal{M}$ and $\mathcal{N}$ be left $\mathcal{C}$ -modules. $x, x' \in \mathcal{M}$ and $y, y' \in \mathcal{N}$.

(1) There is a natural isomorphism

$$\text{hom}_{\mathcal{M}^{\text{op}|L} \boxtimes_{\mathcal{C}} \mathcal{N}}(x \boxtimes_{\mathcal{C}} y, x' \boxtimes_{\mathcal{C}} y') \simeq \text{hom}_{\mathcal{C}}(1, [y, y'] \otimes [x', x])$$

(2) The formula $x \boxtimes_{\mathcal{C}} y \mapsto y \boxtimes_{\mathcal{C}} x$ determines an equivalence $\mathcal{M}^{\text{op}|R} \boxtimes_{\mathcal{C}} \mathcal{N} \simeq (\mathcal{N}^{\text{op}|L} \boxtimes_{\mathcal{C}} \mathcal{M})^{\text{op}}$

- (3) The functor $x \boxtimes_{\mathcal{C}} y \mapsto [x, -] \odot y$ determines an equivalence between $\mathcal{M}^{\text{opl}^R} \boxtimes_{\mathcal{C}} \mathcal{N}$ and $\text{Fun}_{\mathcal{C}}^L(\mathcal{M}, \mathcal{N})$; similarly, the functor $x \boxtimes_{\mathcal{C}} y \mapsto [-, x]^R \odot y$ determines an equivalence between $\mathcal{M}^{\text{opl}^L} \boxtimes_{\mathcal{C}} \mathcal{N} \simeq \text{Fun}_{\mathcal{C}}^R(\mathcal{M}, \mathcal{N})$.

Proof of Theorem A.2.11. According to Theorem A.2.13 (3), Φ^l is an equivalence. It suffices to show that Ψ^l is the left adjoint functor of Φ^l . This follows from the following computation:

$$\begin{aligned} \text{Nat}_{\mathcal{C}}(F, [x, -] \odot y) &\simeq \int_{m \in \mathcal{M}}^{\mathcal{C}} \text{hom}_{\mathcal{N}}(F(m), [x, m] \odot y) \\ &\simeq \int_{m \in \mathcal{M}}^{\mathcal{C}} \text{hom}_{\mathcal{C}}(1, [x, m] \otimes [F(m), y]) \simeq \int_{m \in \mathcal{M}}^{\mathcal{C}} \text{hom}_{\mathcal{N}^{\text{opl}^L} \boxtimes_{\mathcal{C}} \mathcal{M}}(y \boxtimes_{\mathcal{C}} x, F(m) \boxtimes_{\mathcal{C}} m) \\ &\simeq \int_{m \in \mathcal{M}}^{\mathcal{C}} \text{hom}_{(\mathcal{N}^{\text{opl}^L} \boxtimes_{\mathcal{C}} \mathcal{M})^{\text{op}}}(F(m) \boxtimes_{\mathcal{C}} m, y \boxtimes_{\mathcal{C}} x) \\ &\simeq \text{hom}_{\mathcal{M}^{\text{opl}^R} \boxtimes_{\mathcal{C}} \mathcal{N}}\left(\int_{\mathcal{C}}^{m \in \mathcal{M}} m \boxtimes_{\mathcal{C}} F(m), x \boxtimes_{\mathcal{C}} y\right). \end{aligned}$$

The first isomorphism follows from Lemma A.2.9; the second isomorphism is routine; the third isomorphism follows from Theorem A.2.13 (1); the fourth isomorphism is trivial and the last isomorphism follows from Theorem A.2.13 (2).

In a similar manner, we can show that Ψ^r is right adjoint to Φ^r . $\square$

Example A.2.14. Let $\mathcal{C}$ and $\mathcal{D}$ be Morita equivalent finite tensor categories and ${}_c\mathcal{M}_{\mathcal{D}}$ be the invertible module. It is well-known that there is a canonical equivalence of monoidal categories:

$$u : \mathcal{D}^{\text{op}} \rightarrow \text{Fun}_{\mathcal{C}}(\mathcal{M}, \mathcal{M}), \quad d \mapsto - \odot d.$$

Now we are able to write down the quasi-inverse (right adjoint) of u :

$$u^R : \text{Fun}_{\mathcal{C}}(\mathcal{M}, \mathcal{M}) \rightarrow \mathcal{D}^{\text{op}}, \quad F \mapsto \int_{m \in \mathcal{M}}^{\mathcal{C}} [m, F(m)]_{\mathcal{D}^{\text{op}}} \quad (\text{A.2.4})$$

which follows from the following easy calculation:

$$\begin{aligned} \text{hom}_{\mathcal{D}}(d, \int_{m \in \mathcal{M}}^{\mathcal{C}} [m, F(m)]_{\mathcal{D}^{\text{op}}}) &\simeq \int_{m \in \mathcal{M}}^{\mathcal{C}} \text{hom}_{\mathcal{D}}(d, [m, F(m)]_{\mathcal{D}^{\text{op}}}) \\ &\simeq \text{Nat}_{\mathcal{C}}(- \odot d, F(-)). \end{aligned} \quad \heartsuit$$

Example A.2.15. Let $\mathcal{C}, \mathcal{D}$ be non-degenerate braided fusion categories, $\mathcal{M}$ be a closed multi-fusion $\mathcal{C}$ - $\mathcal{D}$ -bimodule. Let $F : \mathcal{C} \rightarrow \mathcal{M}$ be the central functor, whose central structure is witnessed by the braided tensor functor $\tilde{F} : \mathcal{C} \rightarrow \mathfrak{Z}_1(\mathcal{M})$. Given $m \in \mathcal{M}$, we define $T(m)$ to be the $\mathcal{C}$ -module end

$$T(m) := \int_{x \in \mathcal{M}}^{\mathcal{C}} x^R \otimes m \otimes x$$

Note that the central structure of F is indispensable for the definition. When m varies, the assignments $m \mapsto T(m)$ can be extended to a functor $T : \mathcal{M} \rightarrow \mathcal{M}$ which actually defines a comonad over $\mathcal{M}$. Let us look at the Eilenberg-Moore category (i.e. the category of comodules) of this comonad. The Eilenberg-Moore category of this comonad consists of pairs (m, δ_m) where $m \in \mathcal{M}$ and δ_m is a half braiding of m subject to the extra property that (m, δ_m) should lie in the double centralizer of the essential image of $\tilde{F} : \mathcal{C} \rightarrow \mathfrak{Z}_1(\mathcal{M})$, where $\tilde{F}$ is the lift of F . As a result, the Eilenberg-Moore category of T is nothing but $\mathcal{D}$. $\heartsuit$

Example A.2.16. The $\mathcal{C}$ -module end $\int_{x \in \mathcal{M}}^{\mathcal{C}} x^R \otimes m \otimes x \in \mathcal{M}$ defined in Example A.2.15 is equipped with a canonical half-braiding γ :

$$\gamma_y : y \otimes \int_{x \in \mathcal{M}}^{\mathcal{C}} x^R \otimes m \otimes x = \int_{x \in \mathcal{M}}^{\mathcal{C}} y \otimes x^R \otimes m \otimes x \simeq \int_{x \in \mathcal{M}}^{\mathcal{C}} x^R \otimes m \otimes x \otimes y.$$

γ_y is induced by the universal property of $\mathcal{C}$ -module end. Let κ_a be the morphism defined by the following composition

$$\kappa_a : \int_{x \in \mathcal{M}}^{\mathcal{C}} x^R \otimes m \otimes x \otimes y \xrightarrow{\pi_{a \otimes y^L}} (a \otimes y^L)^R \otimes m \otimes a \otimes y^L \otimes y \xrightarrow{\text{id} \otimes \text{coev}_y} y \otimes a^R \otimes m \otimes a.$$

for each $a \in \mathcal{M}$. All these κ_a can be organized into a single morphism

$$\int_{x \in \mathcal{M}}^{\mathcal{C}} x^R \otimes m \otimes x \otimes y \rightarrow \int_{a \in \mathcal{M}}^{\mathcal{C}} y \otimes a^R \otimes m \otimes a = y \otimes \int_{a \in \mathcal{M}}^{\mathcal{C}} a^R \otimes m \otimes a.$$

In a similar manner, one can construct a morphism

$$\gamma_y : y \otimes \int_{a \in \mathcal{M}}^{\mathcal{C}} a^R \otimes m \otimes a \rightarrow \int_{x \in \mathcal{M}}^{\mathcal{C}} x^R \otimes m \otimes x \otimes y.$$

The two morphisms are inverse to each other, giving rise to the desired half-braiding. $\heartsuit$

A.3 E_k -center of an E_k -algebra

In this subsection, we give brief explanations (instead of proofs) of the results in Lemma 6.1.14 [Lur17, Fra12] and (5.2.23) that can enhance our understanding.

We explain Lemma 6.1.14 by first recalling Definition 2.3.17. For $\mathcal{A}, \mathcal{B} \in \text{Alg}_{E_k}^{\mathcal{C}}((n+1)\text{Vec})$ and an E_k -monoidal functor $F : \mathcal{A} \rightarrow \mathcal{B}$, the E_k -centralizer of F is the universal E_k -multi-fusion n -category $\mathfrak{Z}_k(F)$ equipped with a unital action $G : \mathfrak{Z}_k(F) \boxtimes \mathcal{A} \rightarrow \mathcal{B}$, i.e. an E_k -monoidal functor exhibiting the outer triangle of the following diagram commutative (up to equivalences) as illustrated in the following commutative diagram.

$$\begin{array}{ccc} & \mathfrak{Z}_k(F) \boxtimes \mathcal{A} & \\ \uparrow \mathbb{1}_{\mathfrak{Z}_k(F)} \boxtimes \text{id}_{\mathcal{A}} & \nearrow \exists! f \boxtimes \text{id}_{\mathcal{A}} & \uparrow G \\ & \mathcal{X} \boxtimes \mathcal{A} & \\ \downarrow \mathbb{1}_{\mathcal{A}} \boxtimes \text{id}_{\mathcal{M}} & \nwarrow H & \downarrow F \\ \mathcal{A} & \xrightarrow{\quad} & \mathcal{B} \end{array} \quad (\text{A.3.1})$$

1. For $\mathcal{A}, \mathcal{O} \in \text{Alg}_{E_k}^{\mathcal{C}}((n+1)\text{Vec})$, an E_k -monoidal functor $u_{\mathcal{O}} : \mathcal{A} \rightarrow \mathcal{O}$ endows $\mathcal{O}$ with a structure of E_k - $\mathcal{A}$ -module. Moreover, $\mathcal{O}$ and $u_{\mathcal{O}}$ are an object and a 1-morphism in $\text{Alg}_{E_k}^{\mathcal{C}}(\text{Mod}_{\mathcal{A}}^{E_k}((n+1)\text{Vec}))$, respectively, and $u_{\mathcal{O}}$ is precisely the unit of $\mathcal{O}$ as an E_k -algebra in $\text{Mod}_{\mathcal{A}}^{E_k}((n+1)\text{Vec})$.
2. An E_k -algebra homomorphism $g : \mathcal{O} \rightarrow \mathcal{B}$ in $\text{Mod}_{\mathcal{A}}^{E_k}((n+1)\text{Vec})$ is equivalent to an E_k -monoidal functor $g : \mathcal{O} \rightarrow \mathcal{B}$ such that the composed functor $(\mathcal{A} \xrightarrow{u_{\mathcal{O}}} \mathcal{O} \xrightarrow{g} \mathcal{B})$ is isomorphic to $u_{\mathcal{B}}$, a condition which means that g preserves the units.
3. The canonical $(n+1)\text{Vec}$ -action on $\text{Mod}_{\mathcal{A}}^{E_k}((n+1)\text{Vec})$ can be lifted to an action on the level of E_k -algebras as illustrated below.

$$\begin{array}{ccc} \text{Alg}_{E_k}^{\mathcal{C}}((n+1)\text{Vec}) \times \text{Alg}_{E_k}^{\mathcal{C}}(\text{Mod}_{\mathcal{A}}^{E_k}((n+1)\text{Vec})) & \xrightarrow{\quad \boxtimes \quad} & \text{Alg}_{E_k}^{\mathcal{C}}(\text{Mod}_{\mathcal{A}}^{E_k}((n+1)\text{Vec})) \\ \downarrow f \times f & & \downarrow f \\ (n+1)\text{Vec} \times \text{Mod}_{\mathcal{A}}^{E_k}((n+1)\text{Vec}) & \xrightarrow{\quad \boxtimes \quad} & \text{Mod}_{\mathcal{A}}^{E_k}((n+1)\text{Vec}) \end{array} \quad (\text{A.3.2})$$

Therefore, we can simplify the universal property of $\mathfrak{Z}_k(F)$ to that of an internal hom:

$$\begin{array}{ccc} & \mathfrak{Z}_k(F) \boxtimes \mathcal{A} & \\ \nearrow \exists! f \boxtimes \text{id}_{\mathcal{A}} & & \searrow G \\ \mathcal{X} \boxtimes \mathcal{A} & \xrightarrow{\quad H \quad} & \mathcal{B}, \end{array}$$

i.e., the pair $(\mathfrak{Z}_k(F), G)$ precisely defines the internal hom $[\mathcal{A}, \mathcal{B}]$ in $\text{Alg}_{E_k}^c(\text{Mod}_{\mathcal{A}}^{E_k}((n+1)\text{Vec}))$. Then Lemma 6.1.14 simply says that the forgetful functors in (A.3.2) creates the internal hom.

Now we explain (5.2.23). Consider $\mathcal{A} \in \text{Alg}_{E_{k+1}}^c((n+1)\text{Vec})$ and $\mathcal{B} \in \text{LMod}_{\mathcal{A}}(\text{Alg}_{E_k}^c((n+1)\text{Vec}))$. By definition, there is an left $\mathcal{A}$ -action on $\mathcal{B}$ given by an E_k -monoidal functor $\mathcal{A} \times \mathcal{B} \xrightarrow{\odot} \mathcal{B}$ such that $(a, b) \mapsto a \odot b$ for $a \in \mathcal{A}, b \in \mathcal{B}$. It is helpful to think of this action $\odot$ to act in the x^{k+1} -direction that is transversal to a (potentially anomalous) topological order associated to $\mathcal{B}$. It turns out that this E_k -monoidal action can be lifted to an action functor $\text{Alg}_{E_k}^c(\mathcal{A}) \times \text{Alg}_{E_k}^c(\mathcal{B}) \xrightarrow{\odot} \text{Alg}_{E_k}^c(\mathcal{B})$. Since $\mathbb{1}_{\mathcal{B}} \in \text{Alg}_{E_k}^c(\mathcal{B})$, we obtain $[\mathbb{1}_{\mathcal{B}}, \mathbb{1}_{\mathcal{B}}] \in \text{Alg}_{E_1}^c(\text{Alg}_{E_k}^c(\mathcal{A})) \simeq \text{Alg}_{E_{k+1}}^c(\mathcal{A})$.

References

- [AF15] David Ayala and John Francis. *Factorization homology of topological manifolds*. Journal of Topology, 8(4):1045–1084, 2015. [arXiv:1206.5522](#).
- [AF20] David Ayala and John Francis. *A factorization homology primer*. In *Handbook of Homotopy Theory*, pages 39–101. Chapman and Hall/CRC, 2020. [arXiv:1903.10961](#).
- [AF24] David Ayala and John Francis. *The tangle hypothesis: Dimension 1*. arXiv preprint, 2024. [arXiv:2410.23965](#).
- [AFR18] David Ayala, John Francis, and Nick Rozenblyum. *Factorization homology I: Higher categories*. Advances in Mathematics, 333:1042–1177, 2018. [arXiv:1504.04007](#).
- [AFT16] David Ayala, John Francis, and Hiro Lee Tanaka. *Factorization homology of stratified spaces*. Selecta Mathematica, 23(1):293–362, 2016. [arXiv:1409.0848](#).
- [AFT17] David Ayala, John Francis, and Hiro Lee Tanaka. *Local structures on stratified spaces*. Advances in Mathematics, 307:903–1028, 2017. [arXiv:1409.0501](#).
- [AKZ17] Yinghua Ai, Liang Kong, and Hao Zheng. *Topological orders and factorization homology*. Advances in Theoretical and Mathematical Physics, 21(8):1845–1894, 2017. [arXiv:1607.08422](#).
- [ALW19] David Aasen, Ethan Lake, and Kevin Walker. *Fermion condensation and super pivotal categories*. Journal of Mathematical Physics, 60(12):121901, 2019. [arXiv:1709.01941](#).
- [ASW84] Daniel Arovas, J. R. Schrieffer, and Frank Wilczek. *Fractional statistics and the quantum Hall effect*. Physical Review Letters, 53(7):722–723, 1984.
- [BBCW19] Maissam Barkeshli, Parsa Bonderson, Meng Cheng, and Zhenghan Wang. *Symmetry fractionalization, defects, and gauging of topological phases*. Physical Review B, 100(11), 2019. [arXiv:1410.4540](#).
- [BBDR25] Mahesh K. N. Balasubramanian, Matthew Buican, Clement Delcamp, and Rajath Radhakrishnan. *Gauging non-invertible symmetries in (2+1)d topological orders*. arXiv preprint, 2025. [arXiv:2507.01142](#).
- [BD04] Alexander Beilinson and Vladimir Drinfeld. *Chiral algebras*. American Mathematical Society, Providence, R.I, 2004.
- [BEK99] Jens Böckenhauer, David E. Evans, and Yasuyuki Kawahigashi. *On α -induction, chiral generators and modular invariants for subfactors*. Communications in Mathematical Physics, 208(2):429–487, 1999. [arXiv:math/9904109](#).
- [BEK00] Jens Böckenhauer, David E. Evans, and Yasuyuki Kawahigashi. *Chiral structure of modular invariants for subfactors*. Communications in Mathematical Physics, 210(3):733–784, 2000. [arXiv:math/9907149](#).
- [BEK01] Jens Böckenhauer, David E. Evans, and Yasuyuki Kawahigashi. *Longo-Rehren subfactors arising from α -induction*. Publications of the Research Institute for Mathematical Sciences, 37(1):1–35, 2001. [arXiv:math/0002154](#).
- [Bén65] Jean Bénabou. *Catégories relatives*. Comptes Rendus Hebdomadaires des Séances de l’Académie des Sciences, Paris, 256:3824–3827, 1965.
- [Bez04] Roman Bezrukavnikov. *On tensor categories attached to cells in affine Weyl groups*. In *Representation Theory of Algebraic Groups and Quantum Groups*. Mathematical Society of Japan, 2004. [arXiv:math/0010089](#).
- [BJLP19] Marcel Bischoff, Corey Jones, Yuan-Ming Lu, and David Penneys. *Spontaneous symmetry breaking from anyon condensation*. Journal of High Energy Physics, 2019(2), 2019. [arXiv:1811.00434](#).

- [BJQ13] Maissam Barkeshli, Chao-Ming Jian, and Xiao-Liang Qi. *Theory of defects in abelian topological states*. Physical Review B, 88(23), 2013. [arXiv:1305.7203](#).
- [BJS21] Adrien Brochier, David Jordan, and Noah Snyder. *On dualizability of braided tensor categories*. Compositio Mathematica, 157(3):435–483, 2021. [arXiv:1804.07538](#).
- [BK98] S. B. Bravyi and A. Yu. Kitaev. *Quantum codes on a lattice with boundary*. arXiv preprint, 1998. [arXiv:quant-ph/9811052](#).
- [BK01] Bojko Bakalov and Alexander Kirillov, Jr. *Lectures on tensor categories and modular functors*. American Mathematical Society, Providence, RI, 2001.
- [BO04] Roman Bezrukavnikov and Viktor Ostrik. *On tensor categories attached to cells in affine Weyl groups II*. In *Representation Theory of Algebraic Groups and Quantum Groups*. Mathematical Society of Japan, 2004. [arXiv:math/0102220](#).
- [Bom10] H. Bombin. *Topological order with a twist: Ising anyons from an Abelian model*. Physical Review Letters, 105(3), 2010. [arXiv:1004.1838](#).
- [BS09] F. A. Bais and J. K. Slingerland. *Condensate-induced transitions between topologically ordered phases*. Physical Review B, 79(4), 2009. [arXiv:0808.0627](#).
- [BSH09] F. A. Bais, J. K. Slingerland, and S. M. Haaker. *Theory of topological edges and domain walls*. Physical Review Letters, 102(22), 2009. [arXiv:0812.4596](#).
- [BSS02] F. A. Bais, B. J. Schroers, and J. K. Slingerland. *Broken quantum symmetry and confinement phases in planar physics*. Physical Review Letters, 89(18), 2002. [arXiv:hep-th/0205117](#).
- [BSS03] Alexander F Bais, Bernd J Schroers, and Joost K Slingerland. *Hopf symmetry breaking and confinement in (2+1)-dimensional gauge theory*. Journal of High Energy Physics, 2003(05):068–068, 2003. [arXiv:hep-th/0205114](#).
- [BSS11] F. J. Burnell, Steven H. Simon, and J. K. Slingerland. *Condensation of achiral simple currents in topological lattice models: Hamiltonian study of topological symmetry breaking*. Physical Review B, 84(12), 2011. [arXiv:1104.1701](#).
- [BT18] Lakshya Bhardwaj and Yuji Tachikawa. *On finite symmetries and their gauging in two dimensions*. Journal of High Energy Physics, 2018(3), 2018. [arXiv:1704.02330](#).
- [Bur18] F.J. Burnell. *Anyon condensation and its applications*. Annual Review of Condensed Matter Physics, 9(1):307–327, 2018. [arXiv:1706.04940](#).
- [Car23] Nils Carqueville. *Orbifolds of topological quantum field theories*. arXiv preprint, 2023. [arXiv:2307.16674](#).
- [CAW24] Arkya Chatterjee, Ömer Mert Aksoy, and Xiao-Gang Wen. *Quantum phases and transitions in spin chains with non-invertible symmetries*. SciPost Physics, 17(4), 2024. [arXiv:2405.05331](#).
- [CCW16] Iris Cong, Meng Cheng, and Zhenghan Wang. *Topological quantum computation with gapped boundaries*. arXiv preprint, 2016. [arXiv:1609.02037](#).
- [CGHP23] Jessica Christian, David Green, Peter Huston, and David Penneys. *A lattice model for condensation in Levin-Wen systems*. Journal of High Energy Physics, 2023(9), 2023. [arXiv:2303.04711](#).
- [CGS23] Clay Cordova and Diego García-Sepúlveda. *Non-invertible anyon condensation and level-rank dualities*. arXiv preprint, 2023. [arXiv:2312.16317](#).
- [CGS24] Clay Cordova and Diego García-Sepúlveda. *Topological cosets via anyon condensation and applications to gapped QCD₂*. arXiv preprint, 2024. [arXiv:2412.01877](#).
- [CGSO25] Clay Cordova, Diego García-Sepúlveda, and Kantaro Ohmori. *Higgsing transitions from topological field theory & non-invertible symmetry in Chern-Simons matter theories*. arXiv preprint, 2025. [arXiv:2504.03614](#).

- [CGW10] Xie Chen, Zheng-Cheng Gu, and Xiao-Gang Wen. *Local unitary transformation, long-range quantum entanglement, wave function renormalization, and topological order*. Physical Review B, 82(15), 2010. [arXiv:1004.3835](#).
- [CH25] Nils Carqueville and Benjamin Haake. *2-group symmetries of 3-dimensional defect tqfts and their gauging*. arXiv preprint, 2025. [arXiv:2506.08178](#).
- [Cha05] Claudio Chamon. *Quantum glassiness in strongly correlated clean systems: An example of topological overprotection*. Physical Review Letters, 94(4), 2005. [arXiv:cond-mat/0404182](#).
- [CJKYZ20] Wei-Qiang Chen, Chao-Ming Jian, Liang Kong, Yi-Zhuang You, and Hao Zheng. *Topological phase transition on the edge of two-dimensional $\mathbb{Z}_2$ topological order*. Physical Review B, 102(4), 2020. [arXiv:1903.12334](#).
- [CJW25] Arkya Chatterjee, Wenjie Ji, and Xiao-Gang Wen. *Emergent generalized symmetry and maximal symmetry topological order*. Physical Review B, 112(11), 2025. [arXiv:2212.14432](#).
- [CLV14] Xie Chen, Yuan-Ming Lu, and Ashvin Vishwanath. *Symmetry-protected topological phases from decorated domain walls*. Nature Communications, 5(1), 2014. [arXiv:1303.4301](#).
- [CM23] Nils Carqueville and Lukas Müller. *Orbifold completion of 3-categories*. arXiv preprint, 2023. [arXiv:2307.06485](#).
- [CMRSS21] Nils Carqueville, Vincentas Mulevicius, Ingo Runkel, Gregor Schaumann, and Daniel Scherl. *Orbifold graph TQFTs*. arXiv preprints, 2021. [arXiv:2101.02482](#).
- [CMRSS24] Nils Carqueville, Vincentas Mulevicius, Ingo Runkel, Gregor Schaumann, and Daniel Scherl. *Reshetikhin–Turaev TQFTs close under generalised orbifolds*. Communications in Mathematical Physics, 405(242), 2024. [arXiv:2109.04754](#).
- [CR16] Nils Carqueville and Ingo Runkel. *Orbifold completion of defect bicategories*. Quantum Topology, 7(2):203–279, 2016. [arXiv:1210.6363](#).
- [Cra98] Sjoerd E. Crans. *Generalized centers of braided and sylleptic monoidal 2-categories*. Advances in Mathematics, 136(2):183–223, 1998.
- [CRS18] Nils Carqueville, Ingo Runkel, and Gregor Schaumann. *Line and surface defects in Reshetikhin–Turaev TQFT*. Quantum Topology, 10(3):399–439, 2018. [arXiv:1710.10214](#).
- [CRS19] Nils Carqueville, Ingo Runkel, and Gregor Schaumann. *Orbifolds of n -dimensional defect TQFTs*. Geometry & Topology, 23(2):781–864, 2019. [arXiv:1705.06085](#).
- [CRS20] Nils Carqueville, Ingo Runkel, and Gregor Schaumann. *Orbifolds of Reshetikhin–Turaev TQFTs*. Theory and Applications of Categories, 35(15):513–561, 2020. [arXiv:1809.01483](#).
- [Cui19] Shawn Cui. *Four dimensional topological quantum field theories from G -crossed braided categories*. Quantum Topology, 10(4):593–676, 2019. [arXiv:1610.07628](#).
- [CW23] Arkya Chatterjee and Xiao-Gang Wen. *Holographic theory for continuous phase transitions: Emergence and symmetry protection of gaplessness*. Physical Review B, 108(7):075105, 2023. [arXiv:2205.06244](#).
- [Dav10a] Alexei Davydov. *Centre of an algebra*. Advances in Mathematics, 225(1):319–348, 2010. [arXiv:0908.1250](#).
- [Dav10b] Alexei Davydov. *Modular invariants for group-theoretical modular data. I*. Journal of Algebra, 323(5):1321–1348, 2010. [arXiv:0908.1044](#).
- [Déc23a] Thibault D. Décoppet. *The Morita theory of fusion 2-categories*. Higher Structures, 7(1):234–292, 2023. [arXiv:2208.08722](#).
- [Déc23b] Thibault D. Décoppet. *Rigid and separable algebras in fusion 2-categories*. Advances in Mathematics, 419:108967, 2023. [arXiv:2205.06453](#).

- [Déc24] Thibault D. Décoppet. *Finite semisimple module 2-categories*. Selecta Mathematica, 31(1), 2024. [arXiv:2107.11037](#).
- [Déc25] Thibault D. Décoppet. *Drinfeld centers and Morita equivalence classes of fusion 2-categories*. Compositio Mathematica, 161(2):305–340, 2025. [arXiv:2211.04917](#).
- [DFMS97] Philippe Di Francesco, Pierre Mathieu, and David Sénéchal. *Conformal Field Theory*. Springer New York, 1997.
- [DGNO10] Vladimir Drinfeld, Shlomo Gelaki, Dmitri Nikshych, and Victor Ostrik. *On braided fusion categories I*. Selecta Mathematica, 16(1):1–119, 2010. [arXiv:0906.0620](#).
- [DHJFN⁺24] Thibault D. Décoppet, Peter Huston, Theo Johnson-Freyd, Dmitri Nikshych, David Penneys, Julia Plavnik, David Reutter, and Matthew Yu. *The classification of fusion 2-categories*. arXiv preprint, 2024. [arXiv:2411.05907](#).
- [DIHVS17] Kasper Duivenvoorden, Mohsin Iqbal, Jutho Haegeman, Frank Verstraete, and Norbert Schuch. *Entanglement phases as holographic duals of anyon condensates*. Physical Review B, 95(23):235119, 2017. [arXiv:1702.08469](#).
- [DKR11] Alexei Davydov, Liang Kong, and Ingo Runkel. *Field theories with defects and the centre functor*. In *Mathematical Foundations of Quantum Field Theory and Perturbative String Theory*, pages 71–128. American Mathematical Society, 2011. [arXiv:1107.0495](#).
- [DKR15] Alexei Davydov, Liang Kong, and Ingo Runkel. *Functoriality of the center of an algebra*. Advances in Mathematics, 285:811–876, 2015. [arXiv:1307.5956](#).
- [DMNO13] Alexei Davydov, Michael Müger, Dmitri Nikshych, and Victor Ostrik. *The Witt group of non-degenerate braided fusion categories*. Journal für die reine und angewandte Mathematik (Crelles Journal), 2013(677), 2013. [arXiv:1009.2117](#).
- [DN13] Alexei Davydov and Dmitri Nikshych. *The Picard crossed module of a braided tensor category*. Algebra & Number Theory, 7(6):1365–1403, 2013. [arXiv:1202.0061](#).
- [DN21] Alexei Davydov and Dmitri Nikshych. *Braided Picard groups and graded extensions of braided tensor categories*. Selecta Mathematica, 27(4), 2021. [arXiv:2006.08022](#).
- [DR18] Christopher L. Douglas and David J. Reutter. *Fusion 2-categories and a state-sum invariant for 4-manifolds*. arXiv preprint, 2018. [arXiv:1812.11933](#).
- [DS17] Alexei Davydov and Darren Simmons. *On Lagrangian algebras in group-theoretical braided fusion categories*. Journal of Algebra, 471:149–175, 2017. [arXiv:1603.04650](#).
- [DS18] Alexei Davydov and Darren A. Simmons. *Third cohomology and fusion categories*. Homology, Homotopy and Applications, 20(1):275–302, 2018. [arXiv:1704.02401](#).
- [Dun88] Gerald Dunn. *Tensor product of operads and iterated loop spaces*. Journal of Pure and Applied Algebra, 50(3):237–258, 1988.
- [DW90] Robbert Dijkgraaf and Edward Witten. *Topological gauge theories and group cohomology*. Communications in Mathematical Physics, 129(2):393–429, 1990.
- [DX24] Thibault D. Décoppet and Hao Xu. *Local modules in braided monoidal 2-categories*. Journal of Mathematical Physics, 65(6), 2024. [arXiv:2307.02843](#).
- [DY25] Thibault D. Décoppet and Matthew Yu. *Fiber 2-functors and Tambara–Yamagami fusion 2-categories*. Communications in Mathematical Physics, 406(3), 2025. [arXiv:2306.08117](#).
- [EGNO15] Pavel Etingof, Shlomo Gelaki, Dmitri Nikshych, and Victor Ostrik. *Tensor categories*. American Math. Soc, Providence, RI, 2015.
- [EN17] Dominic V. Else and Chetan Nayak. *Cheshire charge in (3+1)-dimensional topological phases*. Physical Review B, 96(4), 2017. [arXiv:1702.02148](#).

- [ENO10] Pavel Etingof, Dmitri Nikshych, and Viktor Ostrik. *Fusion categories and homotopy theory*. Quantum Topology, pages 209–273, 2010. [arXiv:0909.3140](#).
- [ENO11] Pavel Etingof, Dmitri Nikshych, and Victor Ostrik. *Weakly group-theoretical and solvable fusion categories*. Advances in Mathematics, 226(1):176–205, 2011. [arXiv:0809.3031](#).
- [ERB14] I. S. Eliëns, J. C. Romers, and F. A. Bais. *Diagrammatics for Bose condensation in anyon theories*. Physical Review B, 90(19), 2014. [arXiv:1310.6001](#).
- [FFK89] G. Felder, J. Fröhlich, and G. Keller. *Braid matrices and structure constants for minimal conformal models*. Communications in Mathematical Physics, 124(4):647–664, 1989.
- [FFRS06] Jürg Fröhlich, Jürgen Fuchs, Ingo Runkel, and Christoph Schweigert. *Correspondences of ribbon categories*. Advances in Mathematics, 199(1):192–329, 2006. [arXiv:math/0309465](#).
- [FFRS07] Jürg Fröhlich, Jürgen Fuchs, Ingo Runkel, and Christoph Schweigert. *Duality and defects in rational conformal field theory*. Nuclear Physics B, 763(3):354–430, 2007. [arXiv:hep-th/0607247](#).
- [FFRS10] Jürg Fröhlich, Jürgen Fuchs, Ingo Runkel, and Christoph Schweigert. *Defect lines, dualities, and generalised orbifolds*. In *XVIth International Congress on Mathematical Physics*, pages 608–613. World Scientific, 2010. [arXiv:0909.5013](#).
- [FG90] J. Fröhlich and F. Gabbiani. *Braid statistics in local quantum theory*. Reviews in Mathematical Physics, 02(03):251–353, 1990.
- [FHJFK⁺24] Giovanni Ferrer, Brett Hungar, Theo Johnson-Freyd, Cameron Krulewski, Lukas Müller, Nivedita, David Penneys, David Reutter, Claudia Scheimbauer, Luuk Stehouwer, and Chetan Vuppulury. *Dagger n -categories*. arXiv preprint, 2024. [arXiv:2403.01651](#).
- [FHL93] Igor B. Frenkel, Yi-Zhi Huang, and James Lepowsky. *On Axiomatic Approaches to Vertex Operator Algebras and Modules*, volume 104 of *Memoirs of the American Mathematical Society*. American Mathematical Society, 1993.
- [FMT24] Daniel S. Freed, Gregory W. Moore, and Constantin Teleman. *Topological symmetry in quantum field theory*. Quantum Topology, 15(3):779–869, 2024. [arXiv:2209.07471](#).
- [Fra12] John Francis. *The tangent complex and Hochschild cohomology of E_n -rings*. Compositio Mathematica, 149(3):430–480, 2012. [arXiv:1104.0181](#).
- [Fre17] Benoit Fresse. *Homotopy of Operads and Grothendieck–Teichmüller Groups: Parts 1 and 2*, volume 217 of *Mathematical Surveys and Monographs*. American Mathematical Society, 2017.
- [Frö76] Jürg Fröhlich. *New super-selection sectors (“soliton-states”) in two dimensional Bose quantum field models*. Communications in Mathematical Physics, 47(3):269–310, 1976.
- [Fro88] Juerg Froehlich. *Statistics and Monodromy in Two- and Three-Dimensional Quantum · Field Theory*, In *Differential Geometrical Methods in Theoretical Physics*, pages 173–186. Springer Netherlands, 1988.
- [FRS89] K. Fredenhagen, K. H. Rehren, and B. Schroer. *Superselection sectors with braid group statistics and exchange algebras*. Communications In Mathematical Physics, 125(2):201–226, 1989.
- [FRS02] Jürgen Fuchs, Ingo Runkel, and Christoph Schweigert. *TFT construction of RCFT correlators I: partition functions*. Nuclear Physics B, 646(3):353–497, 2002. [arXiv:hep-th/0204148](#).
- [FS03] Jürgen Fuchs and Christoph Schweigert. *Category theory for conformal boundary conditions*, In *Vertex Operator Algebras in Mathematics and Physics*, pages 25–71, volume 39 of *Fields Institute Communications*. American Mathematical Society, 2003. [arXiv:math/0106050](#).
- [FSS20] Jürgen Fuchs, Gregor Schaumann, and Christoph Schweigert. *Module Eilenberg-Watts calculus*. arXiv preprint, 2020. [arXiv:2003.12514](#).

- [FSV13] Jürgen Fuchs, Christoph Schweigert, and Alessandro Valentino. *Bicategories for boundary conditions and for surface defects in 3-d TFT*. Communications in Mathematical Physics, 321(2):543–575, 2013. [arXiv:1203.4568](#).
- [GJF19] Davide Gaiotto and Theo Johnson-Freyd. *Condensations in higher categories*. arXiv preprint, 2019. [arXiv:1905.09566](#).
- [GK21] Davide Gaiotto and Justin Kulp. *Orbifold groupoids*. Journal of High Energy Physics, 2021(2), 2021. [arXiv:2008.05960](#).
- [GKSW15] Davide Gaiotto, Anton Kapustin, Nathan Seiberg, and Brian Willett. *Generalized global symmetries*. Journal of High Energy Physics, 2015(2), 2015. [arXiv:1412.5148](#).
- [GMS80] G. A. Goldin, R. Menikoff, and D. H. Sharp. *Particle statistics from induced representations of a local current group*. Journal of Mathematical Physics, 21(4):650–664, 1980.
- [GMS81] G. A. Goldin, R. Menikoff, and D. H. Sharp. *Representations of a local current algebra in nonsimply connected space and the Aharonov–Bohm effect*. Journal of Mathematical Physics, 22(8):1664–1668, 1981.
- [GNN09] Shlomo Gelaki, Deepak Naidu, and Dmitri Nikshych. *Centers of graded fusion categories*. Algebra & Number Theory, 3(8):959–990, 2009. [arXiv:0905.3117](#).
- [GRIPG17] José Garre-Rubio, Sofyan Iblisdir, and David Pérez-García. *Symmetry reduction induced by anyon condensation: A tensor network approach*. Physical Review B, 96(15):155123, 2017. [arXiv:1702.08759](#).
- [GW09] Zheng-Cheng Gu and Xiao-Gang Wen. *Tensor-entanglement-filtering renormalization approach and symmetry-protected topological order*. Physical Review B, 80(15), 2009. [arXiv:0903.1069](#).
- [Haa96] Rudolf Haag. *Local Quantum Physics*. Springer Berlin Heidelberg, 1996.
- [Haa11] Jeongwan Haah. *Local stabilizer codes in three dimensions without string logical operators*. Physical Review A, 83(4), 2011. [arXiv:1101.1962](#).
- [HHHW22] Yuting Hu, Zichang Huang, Ling-Yan Hung, and Yidun Wan. *Anyon condensation: coherent states, symmetry enriched topological phases, Goldstone theorem, and dynamical rearrangement of symmetry*. Journal of High Energy Physics, 2022(3), 2022. [arXiv:2109.06145](#).
- [HJSWZ25] Ling-Yan Hung, Kaixin Ji, Ce Shen, Yidun Wan, and Yu Zhao. *A 2D-CFT factory: Critical lattice models from competing anyon condensation processes in SymTO/SymTFT*. arXiv preprint, 2025. [arXiv:2506.05324](#).
- [HK64] Rudolf Haag and Daniel Kastler. *An algebraic approach to quantum field theory*. Journal of Mathematical Physics, 5(7):848–861, 1964.
- [HK04] Yi-Zhi Huang and Liang Kong. *Open-string vertex algebras, tensor categories and operads*. Communications in Mathematical Physics, 250(3):433–471, 2004. [arXiv:math.QA/0308248](#).
- [HK07] Yi-Zhi Huang and Liang Kong. *Full field algebras*. Communications in Mathematical Physics, 272(2):345–396, 2007. [arXiv:math/0511328](#).
- [HKL15] Yi-Zhi Huang, Alexander Kirillov, and James Lepowsky. *Braided tensor categories and extensions of vertex operator algebras*. Communications in Mathematical Physics, 337(3):1143–1159, 2015. [arXiv:1406.3420](#).
- [HPRW25] Sebastian Heinrich, Julia Plavnik, Ingo Runkel, and Abigail Watkins. *Generalised orbifolds and G-equivariantisation*. arXiv preprint, 2025. [arXiv:2506.08154](#).
- [HPT16] André Henriques, David Penneys, and James Tener. *Categorified trace for module tensor categories over braided tensor categories*. Documenta Mathematica, 21:1089–1149, 2016. [arXiv:1509.02937](#).

- [Hua95] Yi-Zhi Huang. *Two-Dimensional Conformal Geometry and Vertex Operator Algebras*. Birkhäuser Boston, 1995.
- [Hua08] Yi-Zhi Huang. *Rigidity and modularity of vertex tensor categories*. *Communications in Contemporary Mathematics*, 10(supp01):871–911, 2008. [arXiv:math/0502533](#).
- [HW14] Ling-Yan Hung and Yidun Wan. *Symmetry-enriched phases obtained via pseudo anyon condensation*. *International Journal of Modern Physics B*, 28(24):1450172, 2014. [arXiv:1308.4673](#).
- [HW15] Ling-Yan Hung and Yidun Wan. *Generalized ADE classification of topological boundaries and anyon condensation*. *Journal of High Energy Physics*, 2015(7), 2015. [arXiv:1502.02026](#).
- [HZ23] Mo Huang and Zhi-Hao Zhang. *Tannaka-Krein duality for finite 2-groups*. arXiv preprint, 2023. [arXiv:2305.18151](#).
- [HZW05] Alioscia Hamma, Paolo Zanardi, and Xiao-Gang Wen. *String and membrane condensation on three-dimensional lattices*. *Physical Review B*, 72(3), 2005. [arXiv:cond-mat/0411752](#).
- [IO24] Kansei Inamura and Kantaro Ohmori. *Fusion surface models: 2+1d lattice models from fusion 2-categories*. *SciPost Physics*, 16(6), 2024. [arXiv:2305.05774](#).
- [JF20] Theo Johnson-Freyd. *(3+1)D topological orders with only a $\mathbb{Z}_2$ -charged particle*. arXiv preprint, 2020. [arXiv:2011.11165](#).
- [JF22] Theo Johnson-Freyd. *On the classification of topological orders*. *Communications in Mathematical Physics*, 2022. [arXiv:2003.06663](#).
- [JFR23] Theo Johnson-Freyd and David Reutter. *Minimal nondegenerate extensions*. *Journal of the American Mathematical Society*, 2023. [arXiv:2105.15167](#).
- [JPR22] Corey Jones, David Penneys, and David Reutter. *A 3-categorical perspective on G -crossed braided categories*. *Journal of the London Mathematical Society*, 107(1):333–406, 2022. [arXiv:2009.00405](#).
- [JQ14] Chao-Ming Jian and Xiao-Liang Qi. *Layer construction of 3D topological states and string braiding statistics*. *Physical Review X*, 4(4):041043, 2014. [arXiv:1405.6688](#).
- [JR17] Shenghan Jiang and Ying Ran. *Anyon condensation and a generic tensor-network construction for symmetry-protected topological phases*. *Physical Review B*, 95(12):125107, 2017. [arXiv:1611.07652](#).
- [JZWK⁺19] Zhih-Ahn Jia, Yuan-Hang Zhang, Yu-Chun Wu, Liang Kong, Guang-Can Guo, and Guo-Ping Guo. *Efficient machine-learning representations of a surface code with boundaries, defects, domain walls, and twists*. *Physical Review A*, 99(1):012307, 2019. [arXiv:1802.03738](#).
- [Kik23a] Ken Kikuchi. *Classification of connected étale algebras in modular fusion categories up to rank five*. arXiv preprint, 2023. [arXiv:2312.13353](#).
- [Kik23b] Ken Kikuchi. *Classification of connected étale algebras in pre-modular fusion categories up to rank three*. arXiv preprint, 2023. [arXiv:2311.15631](#).
- [Kik24] Ken Kikuchi. *Classification of connected étale algebras in multiplicity-free modular fusion categories up to rank nine*. arXiv preprint, 2024. [arXiv:2404.16125](#).
- [Kir01] Alexander Kirillov, Jr. *Modular categories and orbifold models II*. arXiv preprint, 2001. [arXiv:math/0110221](#).
- [Kit03] Alexei Kitaev. *Fault-tolerant quantum computation by anyons*. *Annals of Physics*, 303(1):2–30, 2003. [arXiv:quant-ph/9707021](#).
- [Kit06] Alexei Kitaev. *Anyons in an exactly solved model and beyond*. *Annals of Physics*, 321(1):2–111, 2006. [arXiv:cond-mat/0506438](#).

- [KK12] Alexei Kitaev and Liang Kong. *Models for gapped boundaries and domain walls*. Communications in Mathematical Physics, 313(2):351–373, 2012. [arXiv:1104.5047](#).
- [KKH24] Ken Kikuchi, Kah-Sen Kam, and Fu-Hsiang Huang. *Classification of connected étale algebras in multiplicity-free modular fusion categories at rank six*. arXiv preprint, 2024. [arXiv:2402.00403](#).
- [KL20] Kyle Kawagoe and Michael Levin. *Microscopic definitions of anyon data*. Physical Review B, 101(11), 2020. [arXiv:1910.11353](#).
- [KLWZZ20a] Liang Kong, Tian Lan, Xiao-Gang Wen, Zhi-Hao Zhang, and Hao Zheng. *Algebraic higher symmetry and categorical symmetry: A holographic and entanglement view of symmetry*. Physical Review Research, 2(4), 2020. [arXiv:2005.14178](#).
- [KLWZZ20b] Liang Kong, Tian Lan, Xiao-Gang Wen, Zhi-Hao Zhang, and Hao Zheng. *Classification of topological phases with finite internal symmetries in all dimensions*. Journal of High Energy Physics, 2020(9), 2020. [arXiv:2003.08898](#).
- [KO02] Alexander Kirillov, Jr. and Viktor Ostrik. *On a q -analogue of the McKay correspondence and the ADE classification of $\hat{\mathfrak{sl}}_2$ conformal field theories*. Advances in Mathematics, 171(2):183–227, 2002. [arXiv:math/0101219](#).
- [Kon07] Liang Kong. *Full field algebras, operads and tensor categories*. Advances in Mathematics, 213(1):271–340, 2007. [arXiv:math/0603065](#).
- [Kon13] Liang Kong. *Some universal properties of Levin-Wen models*. In *XVIIth International Congress on Mathematical Physics*, pages 444–455. World Scientific, 2013. [arXiv:1211.4644](#).
- [Kon14] Liang Kong. *Anyon condensation and tensor categories*. Nuclear Physics B, 886:436–482, 2014. *Erratum and addendum: “Anyon condensation and tensor categories” [Nucl. Phys. B 886 (2014) 436–482]*. Nuclear Physics B, 973:115607, 2021. See a corrected version in, [arXiv:1307.8244](#).
- [Kon22] Liang Kong. *Topological Wick rotation and holographic duality*. Online Lectures, Since 2022. See for example <https://www.youtube.com/watch?v=9tPLFPxjQfc>. The earliest one is in Chinese <https://www.bilibili.com/video/BV1we4y1q7K8>.
- [KR08] Liang Kong and Ingo Runkel. *Morita classes of algebras in modular tensor categories*. Advances in Mathematics, 219(5):1548–1576, 2008. [arXiv:0708.1897](#).
- [KR09] Liang Kong and Ingo Runkel. *Cardy algebras and sewing constraints, I*. Communications in Mathematical Physics, 292(3):871–912, 2009. [arXiv:0807.3356](#).
- [KS11a] Anton Kapustin and Natalia Saulina. *Surface operators in 3d topological field theory and 2d rational conformal field theory*. In *Mathematical Foundations of Quantum Field Theory and Perturbative String Theory*, pages 175–198. American Mathematical Society, 2011. [arXiv:1012.0911](#).
- [KS11b] Anton Kapustin and Natalia Saulina. *Topological boundary conditions in abelian Chern-Simons theory*. Nuclear Physics B, 845(3):393–435, 2011. [arXiv:1008.0654](#).
- [KTZ20a] Liang Kong, Yin Tian, and Zhi-Hao Zhang. *Defects in the 3-dimensional toric code model form a braided fusion 2-category*. Journal of High Energy Physics, 2020(78), 2020. [arXiv:2009.06564](#).
- [KTZ20b] Liang Kong, Yin Tian, and Shan Zhou. *The center of monoidal 2-categories in 3+1D Dijkgraaf-Witten theory*. Advances in Mathematics, 360:106928, 2020. [arXiv:1905.04644](#).
- [KW14] Liang Kong and Xiao-Gang Wen. *Braided fusion categories, gravitational anomalies, and the mathematical framework for topological orders in any dimensions*. arXiv preprint, 2014. [arXiv:1405.5858](#).

- [KWZ15] Liang Kong, Xiao-Gang Wen, and Hao Zheng. *Boundary-bulk relation for topological orders as the functor mapping higher categories to their centers*. arXiv preprint, 2015. [arXiv:1502.01690](#).
- [KWZ17] Liang Kong, Xiao-Gang Wen, and Hao Zheng. *Boundary-bulk relation in topological orders*. Nuclear Physics B, 922:62–76, 2017. [arXiv:1702.00673](#).
- [KWZ22] Liang Kong, Xiao-Gang Wen, and Hao Zheng. *One dimensional gapped quantum phases and enriched fusion categories*. Journal of High Energy Physics, 2022(3), 2022. [arXiv:2108.08835](#).
- [KYZ21] Liang Kong, Wei Yuan, and Hao Zheng. *Pointed Drinfeld center functor*. Communications in Mathematical Physics, 381(3):1409–1443, 2021. [arXiv:1912.13168](#).
- [KYZZ24] Liang Kong, Wei Yuan, Zhi-Hao Zhang, and Hao Zheng. *Enriched monoidal categories I: Centers*. Quantum Topology, 16(2):343–417, 2024. [arXiv:2104.03121](#).
- [KZ17] Liang Kong and Hao Zheng. *Semisimple and separable algebras in multi-fusion categories*. arXiv preprint, 2017. [arXiv:1706.06904](#).
- [KZ18a] Liang Kong and Hao Zheng. *The center functor is fully faithful*. Advances in Mathematics, 339:749–779, 2018. [arXiv:1507.00503](#).
- [KZ18b] Liang Kong and Hao Zheng. *Gapless edges of 2d topological orders and enriched monoidal categories*. Nuclear Physics B, 927:140–165, 2018. [arXiv:1705.01087](#).
- [KZ20] Liang Kong and Hao Zheng. *A mathematical theory of gapless edges of 2d topological orders. Part I*. Journal of High Energy Physics, 2020(2), 2020. [arXiv:1905.04924](#).
- [KZ21] Liang Kong and Hao Zheng. *A mathematical theory of gapless edges of 2d topological orders. Part II*. Nuclear Physics B, 966:115384, 2021. [arXiv:1912.01760](#).
- [KZ22a] Liang Kong and Zhi-Hao Zhang. *An invitation to topological orders and category theory*. arXiv preprint, 2022. [arXiv:2205.05565](#).
- [KZ22b] Liang Kong and Hao Zheng. *Categories of quantum liquids I*. Journal of High Energy Physics, 2022(8), 2022. [arXiv:2011.02859](#).
- [KZ22c] Liang Kong and Hao Zheng. *Categories of quantum liquids III*. arXiv preprint, 2022. [arXiv:2201.05726](#).
- [KZ24] Liang Kong and Hao Zheng. *Categories of quantum liquids II*. Communications in Mathematical Physics, 405(9), 2024. [arXiv:2107.03858](#).
- [KZZZ25] Liang Kong, Zhi-Hao Zhang, Jiaheng Zhao, and Hao Zheng. *Separable algebras in higher fusion categories*. In preparation, 2025.
- [LB24] Chien-Hung Lin and Fiona J. Burnell. *Anyon condensation in string-net models*. Physical Review B, 110(11):115127, 2024. [arXiv:2303.07291](#).
- [Lei02] Tom Leinster. *A survey of definitions of n-category*. Theory and Applications of Categories, 10(1):1–70, 2002. [arXiv:math/0107188](#).
- [Lev13] Michael Levin. *Protected edge modes without symmetry*. Physical Review X, 3(2), 2013. [arXiv:1301.7355](#).
- [LKW16a] Tian Lan, Liang Kong, and Xiao-Gang Wen. *Modular extensions of unitary braided fusion categories and 2+1D topological/SPT orders with symmetries*. Communications in Mathematical Physics, 351(2):709–739, 2016. [arXiv:1602.05936](#).
- [LKW16b] Tian Lan, Liang Kong, and Xiao-Gang Wen. *Theory of (2+1)-dimensional fermionic topological orders and fermionic/bosonic topological orders with symmetries*. Physical Review B, 94(15), 2016. [arXiv:1507.04673](#).

- [LKW17] Tian Lan, Liang Kong, and Xiao-Gang Wen. *Classification of (2+1)-dimensional topological order and symmetry-protected topological order for bosonic and fermionic systems with on-site symmetries*. Physical Review B, 95(23), 2017. [arXiv:1602.05946](#).
- [LKW18] Tian Lan, Liang Kong, and Xiao-Gang Wen. *Classification of 3+1D bosonic topological orders: The case when pointlike excitations are all bosons*. Physical Review X, 8(2), 2018. [arXiv:1704.04221](#).
- [LM77] J. M. Leinaas and J. Myrheim. *On the theory of identical particles*. Il Nuovo Cimento B Series 11, 37(1):1–23, 1977.
- [LSCH21] Jiaqi Lou, Ce Shen, Chaoyi Chen, and Ling-Yan Hung. *A (dummy’s) guide to working with gapped boundaries via (fermion) condensation*. Journal of High Energy Physics, 2021(2), 2021. [arXiv:2007.10562](#).
- [Lur09] Jacob Lurie. *Higher Topos Theory*. Princeton University Press, 2009. [arXiv:math/0608040](#).
- [Lur17] Jacob Lurie. *Higher Algebra*. Book online, 2017. <http://www.math.harvard.edu/~lurie/papers/HA.pdf>.
- [LW05] Michael A. Levin and Xiao-Gang Wen. *String-net condensation: A physical mechanism for topological phases*. Physical Review B, 71(4), 2005. [arXiv:cond-mat/0404617](#).
- [LW19] Tian Lan and Xiao-Gang Wen. *Classification of 3+1D bosonic topological orders (II): The case when some pointlike excitations are fermions*. Physical Review X, 9(2), 2019. [arXiv:1801.08530](#).
- [LY23] Yalei Lu and Holiverse Yang. *The boundary phase transitions of the 2+1D $\mathbb{Z}_N$ topological order via topological Wick rotation*. Journal of High Energy Physics, 2023(10), 2023. [arXiv:2208.01572](#).
- [LYW24] Tian Lan, Gen Yue, and Longye Wang. *Category of SET orders*. Journal of High Energy Physics, 2024(11), 2024. [arXiv:2312.15958](#).
- [MP17] Scott Morrison and David Penneys. *Monoidal categories enriched in braided monoidal categories*. International Mathematics Research Notices, 2019(11):3527–3579, 2017. [arXiv:1701.00567](#).
- [MR91] Gregory Moore and Nicholas Read. *Nonabelions in the fractional quantum hall effect*. Nuclear Physics B, 360(2–3):362–396, 1991.
- [MR23] Vincentas Mulevičius and Ingo Runkel. *Constructing modular categories from orbifold data*. Quantum Topology, 13(3):459–523, 2023. [arXiv:2002.00663](#).
- [MRV23] Vincentas Mulevicius, Ingo Runkel, and Thomas Voß. *Internal Levin-Wen models*. arXiv preprint, 2023. [arXiv:2309.05755](#).
- [MS89a] Gregory Moore and Nathan Seiberg. *Classical and quantum conformal field theory*. Communications in Mathematical Physics, 123(2):177–254, 1989.
- [MS89b] Gregory Moore and Nathan Seiberg. *Naturality in conformal field theory*. Nuclear Physics B, 313(1):16–40, 1989.
- [MS89c] Gregory Moore and Nathan Seiberg. *Taming the conformal zoo*. Physics Letters B, 220(3):422–430, 1989.
- [Müg03a] Michael Müger. *From subfactors to categories and topology I: Frobenius algebras in and Morita equivalence of tensor categories*. Journal of Pure and Applied Algebra, 180(1-2):81–157, 2003. [arXiv:math/0111204](#).
- [Müg03b] Michael Müger. *From subfactors to categories and topology II: The quantum double of tensor categories and subfactors*. Journal of Pure and Applied Algebra, 180(1-2):159–219, 2003. [arXiv:math/0111205](#).

- [Müg03c] Michael Müger. *On the structure of modular categories*. Proceedings of the London Mathematical Society, 87(02):291–308, 2003. [arXiv:math/0201017](#).
- [Müg04] Michael Müger. *Galois extensions of braided tensor categories and braided crossed G -categories*. Journal of Algebra, 277(1):256–281, 2004. [arXiv:math/0209093](#).
- [Mul24] Vincentas Mulevičius. *Condensation inversion and Witt equivalence via generalised orbifolds*. Theory and Applications of Categories, 41(36):1203–1292, 2024. [arXiv:2206.02611](#).
- [NHvKSB16a] Titus Neupert, Huan He, Curt von Keyserlingk, Germán Sierra, and B. Andrei Bernevig. *Boson condensation in topologically ordered quantum liquids*. Physical Review B, 93(11):115103, 2016. [arXiv:1601.01320](#).
- [NHvKSB16b] Titus Neupert, Huan He, Curt von Keyserlingk, Germán Sierra, and B Andrei Bernevig. *No-go theorem for boson condensation in topologically ordered quantum liquids*. New Journal of Physics, 18(12):123009, 2016. [arXiv:1607.06457](#).
- [Ost03a] Victor Ostrik. *Module categories, weak Hopf algebras and modular invariants*. Transformation Groups, 8(2):177–206, 2003. [arXiv:math/0111139](#).
- [Ost03b] Viktor Ostrik. *Module categories over the Drinfeld double of a finite group*. International Mathematics Research Notices, 2003(27):1507, 2003. [arXiv:math/0202130](#).
- [Par95] B. Pareigis. *On braiding and dyslexia*. Journal of Algebra, 171(2):413–425, 1995.
- [Reh90] Karl-Henning Rehren. *Braid group statistics and their superselection rules*. In *The Algebraic Theory of Superselection Sectors*, pages 333–355. World Scientific, 1990.
- [RSS23] Konstantinos Roumpedakis, Sahand Seifnashri, and Shu-Heng Shao. *Higher gauging and non-invertible condensation defects*. Communications in Mathematical Physics, 401(3):3043–3107, 2023. [arXiv:2204.02407](#).
- [RT91] N. Reshetikhin and V. G. Turaev. *Invariants of 3-manifolds via link polynomials and quantum groups*. Inventiones Mathematicae, 103(1):547–597, 1991.
- [Sch01] Peter Schauenburg. *The monoidal center construction and bimodules*. Journal of Pure and Applied Algebra, 158(2-3):325–346, 2001.
- [SM16] Brian Swingle and John McGreevy. *Renormalization group constructions of topological quantum liquids and beyond*. Physical Review B, 93(4), 2016. [arXiv:1407.8203](#).
- [SN23] Sakura Schafer-Nameki. *ICTP lectures on (non-)invertible generalized symmetries*. arXiv preprint, 2023. [arXiv:2305.18296](#).
- [SSC19] Wilbur Shirley, Kevin Slagle, and Xie Chen. *Universal entanglement signatures of foliated fracton phases*. SciPost Physics, 6(1), 2019. [arXiv:1803.10426](#).
- [SSWC18] Wilbur Shirley, Kevin Slagle, Zhenghan Wang, and Xie Chen. *Fracton models on general three-dimensional manifolds*. Physical Review X, 8(3):031051, 2018. [arXiv:1712.05892](#).
- [Sto25] Devon Stockall. *Large condensation in enriched ∞ -categories*. arXiv preprint, 2025. [arXiv:2506.23632](#).
- [Str12] Ross Street. *Monoidal categories in, and linking, geometry and algebra*. Bulletin of the Belgian Mathematical Society - Simon Stevin, 19(5), 2012. [arXiv:1201.2991](#).
- [SW03] Paolo Salvatore and Nathalie Wahl. *Framed discs operads and Batalin-Vilkovisky algebras*. The Quarterly Journal of Mathematics, 54(2):213–231, 2003.
- [Tak25] Shunta Takahashi. *Anyon condensation in Virasoro TQFT: wormhole factorization*. Journal of High Energy Physics, 2025(6), 2025. [arXiv:2412.11486](#).
- [TSG82] D. C. Tsui, H. L. Stormer, and A. C. Gossard. *Two-dimensional magnetotransport in the extreme quantum limit*. Physical Review Letters, 48(22):1559–1562, 1982.

- [Tur00] Vladimir Turaev. *Homotopy field theory in dimension 3 and crossed group-categories*. arXiv preprint, 2000. [arXiv:math/0005291](https://arxiv.org/abs/math/0005291).
- [Tur20] Vladimir G. Turaev. *Quantum Invariants of Knots and 3-Manifolds*. Walter de Gruyter GmbH & Co.KG, 2020.
- [TW24] Ryan Thorngren and Yifan Wang. *Fusion category symmetry. Part I. Anomaly in-flow and gapped phases*. Journal of High Energy Physics, 2024(4), 2024. [arXiv:1912.02817](https://arxiv.org/abs/1912.02817).
- [TY98] Daisuke Tambara and Shigeru Yamagami. *Tensor categories with fusion rules of self-duality for finite abelian groups*. Journal of Algebra, 209(2):692–707, 1998.
- [VHF16] Sagar Vijay, Jeongwan Haah, and Liang Fu. *Fracton topological order, generalized lattice gauge theory, and duality*. Physical Review B, 94(23):235157, 2016. [arXiv:1603.04442](https://arxiv.org/abs/1603.04442).
- [Wen90] Xiao-Gang Wen. *Topological orders in rigid states*. International Journal of Modern Physics B, 04(02):239–271, 1990.
- [Wen91] Xiao-Gang Wen. *Non-Abelian statistics in the fractional quantum Hall states*. Physical Review Letters, 66(6):802–805, 1991.
- [Wen17] Xiao-Gang Wen. *Colloquium: Zoo of quantum-topological phases of matter*. Reviews of Modern Physics, 89(4), 2017. [arXiv:1610.03911](https://arxiv.org/abs/1610.03911).
- [Wen19] Xiao-Gang Wen. *Choreographed entanglement dances: Topological states of quantum matter*. Science, 363(6429):eaal3099, 2019. [arXiv:1906.05983](https://arxiv.org/abs/1906.05983).
- [Wen20] Xiao-Gang Wen. *Systematic construction of gapped nonliquid states*. Physical Review Research, 2(3):033300, 2020. [arXiv:2002.02433](https://arxiv.org/abs/2002.02433).
- [Wen24] Rui Wen. *String Condensation in 3+1D Topological Orders and Applications to Topological Holography*. Talk, 2024. <https://bimsa.net/bimsavideo.html?id=41394.mp4>.
- [Wen25] Rui Wen. *Topological holography for 2+1-D gapped and gapless phases with generalized symmetries*. arXiv preprint, 2025. [arXiv:2503.13685](https://arxiv.org/abs/2503.13685).
- [Wil82] Frank Wilczek. *Magnetic flux, angular momentum, and statistics*. Physical Review Letters, 48(17):1144–1146, 1982.
- [Wil08] Simon Willerton. *The twisted Drinfeld double of a finite group via gerbes and finite groupoids*. Algebraic & Geometric Topology, 8(3):1419–1457, 2008. [arXiv:math/0503266](https://arxiv.org/abs/math/0503266).
- [Wit89] Edward Witten. *Quantum field theory and the Jones polynomial*. Communications in Mathematical Physics, 121(3):351–399, 1989.
- [Wu84] Yong-Shi Wu. *General theory for quantum statistics in two dimensions*. Physical Review Letters, 52(24):2103–2106, 1984.
- [WW17] Yidun Wan and Chenjie Wang. *Fermion condensation and gapped domain walls in topological orders*. Journal of High Energy Physics, 2017(3), 2017. [arXiv:1607.01388](https://arxiv.org/abs/1607.01388).
- [XLWWC23] Wenjie Xi, Tian Lan, Longye Wang, Chenjie Wang, and Wei-Qiang Chen. *On a class of fusion 2-category symmetry: condensation completion of braided fusion category*. arXiv preprint, 2023. [arXiv:2312.15947](https://arxiv.org/abs/2312.15947).
- [Xu24a] Hao Xu. *Higher Witt groups for 2-categories I: Centralizers*. arXiv preprint, 2024. [arXiv:2403.07768](https://arxiv.org/abs/2403.07768).
- [Xu24b] Hao Xu. *On Étale algebras and bosonic fusion 2-categories*. arXiv preprint, 2024. [arXiv:2411.13367](https://arxiv.org/abs/2411.13367).
- [XY25] Rongge Xu and Holiverse Yang. *Classification of gapped domain walls in 2+1D topological orders through 2-Morita equivalence*. Chinese Physics Letters, 42(7):070710, 2025. [arXiv:2403.19779](https://arxiv.org/abs/2403.19779).

- [XZ24] Rongge Xu and Zhi-Hao Zhang. *Categorical descriptions of one-dimensional gapped phases with Abelian onsite symmetries*. Physical Review B, 110(15):155106, 2024. [arXiv:2205.09656](#).
- [XZZ20] Wen-Tao Xu, Qi Zhang, and Guang-Ming Zhang. *Tensor network approach to phase transitions of a non-Abelian topological phase*. Physical Review Letters, 124(13):130603, 2020. [arXiv:1912.07836](#).
- [YWL25] Gen Yue, Longye Wang, and Tian Lan. *Condensation completion and defects in 2+1D topological orders*. Physical Review B, 112(10), 2025. [arXiv:2402.19253](#).
- [Zhe17] Hao Zheng. *Extended TQFT arising from enriched multi-fusion categories*. arXiv preprint, 2017. [arXiv:1704.05956](#).
- [ZHK89] S. C. Zhang, T. H. Hansson, and S. Kivelson. *Effective-field-theory model for the fractional quantum Hall effect*. Physical Review Letters, 62(1):82–85, 1989.
- [ZHWHW23] Yu Zhao, Shan Huang, Hongyu Wang, Yuting Hu, and Yidun Wan. *Characteristic properties of a composite system of topological phases separated by gapped domain walls via an exactly solvable hamiltonian model*. SciPost Physics Core, 6(4), 2023. [arXiv:2209.12750](#).
- [ZLZH⁺23] Jiaheng Zhao, Jia-Qi Lou, Zhi-Hao Zhang, Ling-Yan Hung, Liang Kong, and Yin Tian. *String condensations in 3+1D and Lagrangian algebras*. Advances in Theoretical and Mathematical Physics, 27(2):583–622, 2023. [arXiv:2208.07865](#).
- [ZVW25] Carolyn Zhang, Ashvin Vishwanath, and Xiao-Gang Wen. *Hierarchy construction for non-Abelian fractional quantum Hall states via anyon condensation*. Physical Review B, 112(12), 2025. [arXiv:2406.12068](#).
- [ZW15] Bei Zeng and Xiao-Gang Wen. *Gapped quantum liquids and topological order; stochastic local transformations and emergence of unitarity*. Physical Review B, 91(12), 2015. [arXiv:1406.5090](#).
- [ZW25a] Yu Zhao and Yidun Wan. *Landau-Ginzburg paradigm of topological phases*. arXiv preprint, 2025. [arXiv:2506.05319](#).
- [ZW25b] Yu Zhao and Yidun Wan. *Nonabelian anyon condensation in 2+1d topological orders: A string-net model realization*. Journal of High Energy Physics, 2025(5), 2025. [arXiv:2409.05852](#).
- [ZWG22] Jing-Ren Zhou, Qing-Rui Wang, and Zheng-Cheng Gu. *Towards a complete classification of nonchiral topological phases in two-dimensional fermion systems*. Physical Review B, 106(24):245120, 2022. [arXiv:2112.06124](#).