

[Drug Name]General Name: Shenling Baizhu Tablets Pinyin: Shenling Baizhu Pian [Ingredients]Dangshen (Codonopsis pilosula), Fuling (Poria cocos), Baizhu (Atractylodes macrocephala, stir-fried), Shanyao (Dioscorea opposita), Babian dou (Hyacinth bean, stir-fried), Lianzi (Lotus seed, stir-fried), Yiyiren (Coix seed, stir-fried), Sharen (Amomum villosum), Jiegeng (Platycodon grandiflorus), Zhi Gancao (Fried licorice root), Chenpi (Dried tangerine peel). [Description]This product is a sugar-coated tablet (or film-coated tablet). After removing the coating, it appears brownish-yellow to yellow-brown; it has a fragrant aroma and a slightly sweet taste. [Indications]Strengthen the spleen and replenish qi. Indicated for fatigue, poor appetite, and loose stools. [Dosage and Administration]Take orally: 6–12 tablets each time, twice daily; reduce the dosage for children. [Precautions]1. Contraindicated in patients with diarrhea accompanied by constipation or a sensation of rectal tenesmus. 2. Do not take this medication concurrently with Lilu (Veratrum nigrum), Wulingzhi (Trogopterus dudermskyi), Zaojia (Saposhnikovia divaricata), or their preparations. 3. Avoid tea and radish consumption to prevent interference with drug efficacy. 4. Do not take concurrently with cold medications. 5. Patients with hypertension, heart disease, kidney disease, severe diabetes, or pregnant women should use under medical supervision. 6. Take before meals or with food. 7. Follow the prescribed dosage; children should be administered under medical supervision. 8. If symptoms do not improve after two weeks of medication, seek medical attention. 9. Contraindicated in individuals with known hypersensitivity to this product; use with caution in those with allergic predisposition. 10. Do not use if the product exhibits altered physical properties. 11. Children must use under adult supervision. 12. Keep out of reach of children. 13. Consult a physician or pharmacist before using this product if other medications are being taken. [Drug Interactions]Concomitant use with other drugs may result in interactions; consult a physician or pharmacist for details. [Storage]Keep tightly sealed. [Packaging][Not specified] [Standard Implementation]Volume 20 of the Ministry-issued standards

[Drug Name]General Name: Egui Yangxue Keli Pinyin: Egui Yangxue Keli [Ingredients]Angelica sinensis (Danggui), Donkey-hide gelatin (Ejiao), Prepared Rehmannia root (Shuhuang), white peony root (White Peony Root), Codonopsis pilosula (Dangshen), Astragalus membranaceus (Huangqi), Poria cocos (Fuling), Ligusticum chuanxiong (Chuanxiong), and fried licorice root (Zhi Gancao). [Description]The product appears as brownish-yellow granules with a faint fragrance, sweet taste, slight bitterness, astringency, and pungency. [Functions and Indications]tonifies qi and blood. Indicated for qi and blood deficiency, manifested as sallow complexion, dizziness, fatigue, and scanty or pale menstrual flow. [Specification]Each sachet contains 10 grams. [Dosage and Administration]Take orally: one sachet per dose, three times daily. [Contraindications]Contraindicated in patients with exogenous febrile syndromes. [Precautions] 1. Patients with spleen-stomach dysfunction should reduce the dosage or combine with spleen-stomach regulating herbs, or regulate the spleen-stomach first before using this medicine. 2. Do not take this medicine concurrently with cold medications. 3. Take on an empty stomach or before meals. 4. Follow the recommended dosage; diabetic patients and children should use under physician supervision. 5. If symptoms do not improve after two weeks of medication, seek medical attention. 6. Contraindicated in individuals with known hypersensitivity to this product; use with caution in those with allergic predisposition. 7. Do not use if the product exhibits altered physical properties. 8. Children must use under adult supervision. 9. Keep out of reach of children. 10. Consult a physician or pharmacist before use if taking other medications. [Drug Interactions]Concomitant use with other drugs may result in interactions; consult a physician or pharmacist for details. [Storage]Keep tightly sealed. [Packaging][Not specified] [Standard Compliance]Volume 12 of the National Pharmaceutical Standards

Fuyan Rehabilitation Manual

Drug Name

Generic Name: Fuyan Kangfu

Pinyin: fuyankangfu pian

[Ingredients] This product is a tablet prepared by processing the following medicinal ingredients: Patrinia scabiosifolia, Coix seed, Toosendan fruit, Bupleurum root, Tangerine peel, and Scutellaria baicalensis.

[Properties] This product is a sugar-coated tablet, appearing dark brown after removal of the sugar coating; with a slightly bitter taste.

[Indications]Clear heat and promote diuresis, resolve stasis and alleviate pain. Indicated for leukorrhea in women caused by damp-heat and blood stasis, characterized by yellow, viscous discharge resembling bean dregs with foul odor, accompanied by lower abdominal and lumbosacral pain.

[Specification]The substrate weighs 0.35 g

[Dosage and Administration]Oral administration: 5 tablets each time, three times daily.

[Precautions] 1. Avoid spicy foods and reduce intake of greasy foods. 2. Use with caution in patients with significant spleen and stomach deficiency. 3. Not recommended for deficiency-type leukorrhea, which manifests as clear, odorless discharge accompanied by fatigue, dizziness, pale or sallow complexion, and cold extremities. 4. Seek medical attention if leukorrhea is markedly abnormal or associated with other conditions. 5. Discontinue use and consult a physician if symptoms persist after one week of medication. 6. Elderly individuals, adolescent girls, and pregnant women should use under medical supervision. 7. Contraindicated in patients with known hypersensitivity to this product; use with caution in those with allergic predisposition. 8. Do not use if the product exhibits altered physical properties. 9. Store out of reach of children. 10. If you are taking other medications, consult a physician or pharmacist before using this product.

[Drug Interactions]Drug interactions may occur when used concomitantly with other medications. For details, consult a physician or pharmacist. [Storage]Store in a tightly closed container protected from moisture.