

Ibuprofen Tablet Instructions

Please read the instructions carefully and use them as directed, or purchase and use under the guidance of a pharmacist.

Drug Name

Generic Name: Ibuprofen Tablets

English name: Ibuprofen Tablets

Pinyin: Buluofen Pian

[Ingredients]

The active ingredient of this product is ibuprofen.

The excipients include anhydrous lactose, microcrystalline cellulose, cross-linked carboxymethyl cellulose sodium, colloidal silica, and stearic acid. [Properties] This product is a film-coated tablet, appearing white after removal of the coating.

[Category of Action] This product is an over-the-counter (OTC) antipyretic and analgesic medication.

[Indications] For the relief of mild to moderate pain such as headache, arthralgia, migraine, toothache, myalgia, neuralgia, and dysmenorrhea. Also indicated for fever caused by the common cold or influenza.

Specification: 0.2 grams

[Dosage and Administration] Oral administration. For adults, take 1 tablet per dose. In cases of persistent pain or fever, repeat the dose every 4–6 hours, with a maximum of 4 doses within 24 hours.

【 untoward effect 】

1. A small number of patients may experience nausea, vomiting, gastric burning sensation or mild dyspepsia, gastrointestinal ulcers and bleeding, elevated transaminase levels, headache, dizziness, tinnitus, blurred vision, nervous tension, drowsiness, lower limb edema, or sudden weight gain.
2. Rare rash, allergic nephritis, cystitis, nephrotic syndrome, renal papillary necrosis or renal failure, bronchospasm.

【 taboo 】

1. **Contraindicated in Patients With Hypersensitivity to Other Nonsteroidal Anti-Inflammatory Drugs (NSAIDs).**
2. **Contraindicated in Pregnant and Lactating Women.**
3. **Contraindicated in Asthma Patients With an Allergy to Aspirin.**

【 matters need attention 】

1. This product is a symptomatic treatment agent and should not be used for prolonged or high-dose administration. For analgesic use, the duration should not exceed 5 days; for antipyretic use, it should not exceed 3 days. If symptoms persist, consult a physician or pharmacist.
2. Do not take other medications containing antipyretic and analgesic agents simultaneously (e.g., certain compound cold medicines).
3. Do not consume alcohol or alcoholic beverages while taking this product.
4. The following patients should use with caution: those aged 60 years or older, patients with bronchial asthma, hepatic or renal insufficiency, or coagulation disorders or platelet dysfunction (e.g., hemophilia).
5. Patients with the following conditions should use the medication under physician supervision: history of peptic ulcer, gastrointestinal bleeding, cardiac insufficiency, or hypertension.
6. If an overdose occurs or severe adverse reactions are observed, seek immediate medical attention.
7. Contraindicated in patients with known hypersensitivity to this product. Use with caution in individuals with allergic predisposition.
8. This product must not be used if its physical properties have changed.
9. Keep this product out of reach of children.
10. If you are using other medications, consult a physician or pharmacist before using this product.
11. If gastrointestinal bleeding or ulcers, chest pain, shortness of breath, weakness, or slurred speech occur, discontinue the medication and consult a physician.
12. If a rash or allergic reaction occurs upon first use of this product, discontinue use and consult a physician.

【 drug interaction 】

1. Concomitant use of this product with other antipyretic, analgesic, or anti-inflammatory drugs may increase the risk of gastrointestinal adverse reactions and may lead to ulcer formation.
2. When co-administered with anticoagulants such as heparin or dicoumarol, this product may prolong prothrombin time and increase the risk of bleeding.
3. Concomitant use of this product with digoxin, methotrexate, or oral hypoglycemic agents may increase the plasma concentrations of these drugs; therefore, concomitant administration is not recommended.
4. When co-administered with furosemide (furosemide), the natriuretic and antihypertensive effects of the latter are diminished; similarly, when used concomitantly with antihypertensive agents, the hypotensive efficacy of these drugs is reduced.
5. Drug interactions may occur when used concomitantly with other medications; consult a physician or pharmacist for details. [Pharmacological Effects] This product inhibits prostaglandin synthesis, exhibiting antipyretic, analgesic, and anti-inflammatory effects.

[Storage] Store in a sealed container.

[Packaging] Oral solid dosage form in high-density polyethylene bottles. 100 tablets/bottle.

[Shelf Life] 24 months

【 operative norm 】

[Approval Number] National Drug Approval
Number H37020386 [Revised Date of Instructions]

[Marketing Authorization Holder]

Name of the Marketing Authorization Holder: Shandong
Xinhua Pharmaceutical Co., Ltd.

Registered Address: Chemical Industry Zone, Zibo High-tech
Industrial Development Zone

Postal code: 255000

Tel: 0533-2166666

Fax number: 0533-2184991

Website: www.xhzy.com

【 manufacturing enterprise 】

Company Name: Shandong Xinhua Pharmaceutical Co., Ltd.

Production Address: No.14, East First Road, Zhangdian District,
Zibo City, Shandong Province

Postal code: 255005

Phone number: 0533-2196361

Fax number: 0533-2196365

Website: www.xhzy.com

If any issues arise, please contact the manufacturing company.