

Ahmad Yazdankhah

ahmad.yazdankhah@sjsu.edu
www.cs.sjsu.edu/~yazdankhah

Mathematical Preliminaries

(Part 2)

Lecture 03
Day 03/31

CS 154
Formal Languages and Computability
Fall 2019

Agenda of Day 03

- Waiting List Enrollment ...
- Announcement
- Summary of Lecture 02
- Lecture 03: Teaching ...
 - Covering one slide from the past
 - Mathematical Preliminaries (Part 2)

Announcement

- Our first quiz will be **next Thursday!**
- Some of the questions are **true/false** and **multiple choice**.
- So, please have **Scantron 882 E**.
- **If you forget**, no problem at all! **I'll sell it at:**

~~\$20~~ Each

Now Just \$19.99!

Summary of Lecture 02: We learned ...

Sets

- A **set** is ...
 - ... a collection of objects.
- A **list** is ...
 - ... a collection of **ordered** objects.
- A **set is known** when its **boundary** is clearly defined.
- **Three methods** to represent sets ...
 - Roster method
 - Venn diagram
 - **Set builder**
- **Universal set** of a set is ...
 - ... the set containing all possible elements under consideration.

- The **power set** of the set S is ...
 - ... the set of all subsets of S .
 - It is denoted by 2^S .
 - $|2^S| = 2^{|S|}$
- A set is called **finite** if ...
 - ... its size is a natural number.
- A set is called **infinite** if ...
 - ... we cannot express its size by a natural number.

Any question?

Empty Set Representation by Set Builder

- How to represent empty set by set builder?
- We know: $A - B = \{x : x \in A \text{ AND } x \notin B\}$
- Substitute A for B: $A - A = \{x : x \in A \text{ AND } x \notin A\}$
- $\therefore \phi = \{x : \text{False}\}$
- So, to represent empty set, just put any false statement in the description part of the set builder.
- For example, the following sets represent empty sets:
- $\{x : x \text{ is the 8}^{\text{th}} \text{ day of week}\}$
- $\{x : x \notin U\}$

Mathematical Preliminaries

Recap from Math 42

Cartesian Products

Motivation

- Recall that in sets, order of elements does NOT matter.
- But in practice, we do need ordered collections.
- As we said before, in computer science we use "Lists" for ordered collections.
- The question is how we can mathematically model lists?

Introduction

- Mathematicians defined a new mathematical structure called "n-tuple".
- It is denoted by (a_1, a_2, \dots, a_n) where a_i 's are objects.
 - A special case of n-tuple is 2-tuple aka ordered-pair (a_1, a_2) .
- We use a mathematical operation called Cartesian product to create n-tuples.
- This operation is named after the great French philosopher, mathematician, and physicist René Descartes (1596-1650).



Cartesian Products

Definition

- Let A and B be two sets.
- The Cartesian product of A and B is the set of all ordered-pairs (a, b) , where $a \in A$ and $b \in B$.
- Cartesian product of A and B is denoted by $A \times B$.

⚠ Set Builder Definition

$$A \times B = \{(a, b) : a \in A, b \in B\}$$

- Note that (a, b) is the pattern of the elements.



Cartesian Products: Examples

Example 25

- Let $A = \{0, 1\}$, $B = \{3, 6, 9\}$; $A \times B = ?$

Solution

- $\{0, 1\} \times \{3, 6, 9\} = \{(0, 3), (0, 6), (0, 9), (1, 3), (1, 6), (1, 9)\}$

Example 26

- Let $Q = \{q_0, q_1\}$, $\Sigma = \{a, b\}$; $Q \times (\Sigma \cup \{\lambda\}) = ?$
 - " λ " is pronounced "**lambda**".

Solution



! Cartesian Products: Properties

- What is the result of the following Cartesian product?

$$A = \{1, 2\}, B = \phi; A \times B = ?$$

$$A \times B = \phi$$

– In fact, the result of Cartesian product would be ϕ if one of the sets is ϕ .



- How can you prove it?
- Does Cartesian product have commutative property?
- Is this a true statement: $A \times B = B \times A$
- In general, No! $A \times B \neq B \times A$
- But in the following special cases, they are equal:
 $A \times B = B \times A$ iff $(A = B) \vee (A = \phi) \vee (B = \phi)$

Cartesian Products **Extension**

- We can extend the idea to n sets to produce n -tuple:



$$S_1 \times S_2 \times \dots \times S_n = \{(x_1, x_2, \dots, x_n) : x_1 \in S_1, \dots, x_n \in S_n\}$$

Homework

- Let $Q = \{q_0, q_1\}$, $\Sigma = \{a, b\}$, $\Gamma = \{x, y\}$
- $Q \times (\Sigma \cup \{\lambda\}) \times \Gamma = ?$




Mathematical Preliminaries

Recap from Math 42

Functions

Introduction

- In many situations in real life, there is a **relationship between two sets**.
- For example, we assign a **letter grade to each student of a class**.



- This **relationship** is an example of the concept of **function**.




Functions

Definition

- Let D and R be two **sets**.
- A function g from D to R is a **rule** that **assigns** (or **maps**) to the elements of D a "**unique element**" of R .

- The set D is called the "**domain**" of g .
- The set R is called the "**range**" of g .



- The function g from D to R is **denoted** by: $g : D \rightarrow R$

Functions: Naming Convention


- We usually **name a function** by **lower-case** letters (English or Greek) such as f , g , h , δ (pronounce "**delta**"), etc.

Example 27

$S = \{\text{John, Tom, Marry, Ted, Adam, Eve}\}$, $G = \{A, B, C, D, F\}$

- We can name this function "**h**".

$h : S \rightarrow G$



Functions: Algebraic Notation

Example 27 (cont'd)


- What is the rule of this function in algebraic notation?

$S = \{\text{John, Tom, Marry, Ted, Adam, Eve}\}$, $G = \{A, B, C, D, F\}$

$h : S \rightarrow G$

- We can translate the Venn Diagram into algebraic notation like this:


$$\begin{cases} h(\text{John}) = A \\ h(\text{Tom}) = C \\ h(\text{Marry}) = B \\ h(\text{Adam}) = A \\ h(\text{Eve}) = F \end{cases}$$



Functions: Notes



- $h(\text{Ted}) = ?$
- $h(\text{Ted}) = \text{Undefined}$




- So, it is possible to have some elements in the domain that is **NOT mapped** to any value of the range. (e.g. **Ted** in the domain)
- Also, it is possible to have some elements in the range that is **NOT assigned** by any value of the domain. (e.g. **D** in the range)

Functions: Notes



- Is it possible for Marry to have two grades at the same time?

- Absolutely, NO.
In this universe,
it cannot happen.




- That's why, in the definition of function, we said elements of the domain are **uniquely mapped** to an element of the range.
- In other words, if there is a mapping, it should be unique.

Functions as Sets

Example 27 (cont'd)

$$\begin{cases} h(\text{John}) = A \\ h(\text{Tom}) = C \\ h(\text{Marry}) = B \\ h(\text{Adam}) = A \\ h(\text{Eve}) = F \end{cases}$$




- The above relations can be represented by **a set of ordered-pairs**:
- $H = \{(\text{John}, A), (\text{Tom}, C), (\text{Marry}, B), (\text{Adam}, A), (\text{Eve}, F)\}$
- And the **universal set** for H is $S \times G$ that contains all possible relations between S's members and G's members.
- $U = S \times G = \{(\text{John}, A), (\text{John}, B), \dots, (\text{John}, F), (\text{Tom}, A), \dots, (\text{Tom}, F), \dots, (\text{Ted}, A), \dots, (\text{Ted}, F), \dots, (\text{Eve}, F)\}$

Functions: Example

Example 28: Mixing Cartesian Product and Function

- Let $Q = \{q_0, q_1, q_2\}$, $\Sigma = \{a, b\}$, $\delta : Q \times \Sigma \rightarrow Q$
- What is the domain and range of δ ?
- Domain:** $Q \times \Sigma = \{q_0, q_1, q_2\} \times \{a, b\} = \{(q_0, a), (q_0, b), (q_1, a), (q_1, b), (q_2, a), (q_2, b)\}$
- Range:** $\{q_0, q_1, q_2\}$
- The **rule** of δ is shown in the following Venn diagram.
- Write the rule of δ by using algebraic notation.**

$$\begin{cases} \delta(q_0, a) = q_0 \\ \delta(q_0, b) = q_1 \\ \delta(q_1, b) = q_2 \\ \delta(q_2, b) = q_2 \end{cases}$$



Total Function


Definition



- A function is called **total** if all of its **domain elements** are **defined**.

Example 29

- The following function is **total** because all domain elements are defined.



Partial Function


Definition



- A function is called **partial** if at least one of its domain's elements is **undefined**.

Example 30

- The following function is **partial** because $f(8) = \text{Undefined}$
- Note that **one example is enough** to make a function partial.



Functions

Homework



Functions: Homework #1

- Let $Q = \{q_0, q_1\}$, $\Sigma = \{a\}$, $\Gamma = \{x\}$.
- The function δ is defined as:

$$\delta : Q \times \{\Sigma \cup \{\lambda\}\} \times \Gamma \rightarrow Q$$


- What is the domain and range of δ ?
- Represent the universal set of δ as a set of quadruples (4-tuple).

Functions: Homework #2

- Write the algebraic notation of the following function:

Solution

$$\begin{cases} f(0) = ? \\ f(1) = ? \\ f(2) = ? \\ f(3) = ? \\ \dots \end{cases}$$



- Sometimes, we can find a pattern between the elements of domain and range.
- What is the pattern of the above example?
- $f(x) = ?$

Functions: Homework #3

- Write a simple Java method to implement the function of HW #2.
- Hint: the argument of the method is the domain and whatever it's returning is range.

Solution

Functions: Homework #4

- What is the range and domain of the following Java code?
- `public int g(int x, int z) { return 2*x + z; }`

Solution

Mathematical Preliminaries

Recap from Math 42

Graphs


Graphs Definition

Definition

- ⓘ A graph is a mathematical construct consisting of two sets:
 - A non-empty and finite set of vertices (aka nodes, states)
 $V = \{v_1, v_2, \dots, v_n\}$
 - A finite set of edges $E = \{e_1, e_2, \dots, e_m\}$
 - Each edge connects two vertices.

Example 31

- $V = \{v_1, v_2, v_3\}$
- $E = \{e_{13}, e_{23}\}$




Directed Graphs

- If the **direction** of the edges matters, then we call the graph **directed graph** (aka **digraph**).
- The edges are shown by **ordered-pair** (start-vertex , end-vertex).
 - In **this course**, we only use **directed graphs**.

Example 32

- Draw a **digraph** with the following specifications:

$$V = \{v_1, v_2, v_3\}, E = \{(v_1, v_3), (v_3, v_1), (v_3, v_2), (v_3, v_3)\}$$




Graphs Terminologies



Walk

- A sequence of edges like $(v_i, v_j), (v_j, v_k), \dots, (v_m, v_n)$, is called a **walk** from v_i to v_n .
 - Note that the **end vertex** of e_i is the **start vertex** of e_{i+1} .
 - In other words, in a walk we **cannot jump**!




Example 33

- Each of the following sequences are a **walk** from v_1 to v_3 :
 - Walk 1: (v_1, v_3)
 - Walk 2: $(v_1, v_3), (v_3, v_3)$
 - Walk 3: $(v_1, v_3), (v_3, v_1), (v_1, v_3)$
 - ...



Length of Walks

- It is the total number of edges traversed.




Example 33 (cont'd)

- Walk 1: (v_1, v_3) ; length = 1
- Walk 2: $(v_1, v_3), (v_3, v_3)$; length = 2
- Walk 3: $(v_1, v_3), (v_3, v_1), (v_1, v_3)$; length = 3

Path

- A walk that **no edges are repeated**.




Example 34

- Which one is a **path**?
 - ✓ – Walk 1: (v_1, v_3)
 - ✓ – Walk 2: $(v_1, v_3), (v_3, v_3)$
 - ✗ – Walk 3: $(v_1, v_3), (v_3, v_1), (v_1, v_3)$

Simple Path

- A path that no vertices are repeated.
 - In other words, no vertex should be visited more than once.




Example 35

- Which one is a simple path?
- ✓ – Walk 1: (v_1, v_3)
- ✗ – Walk 2: $(v_1, v_3), (v_3, v_3)$
- ✗ – Walk 3: $(v_1, v_3), (v_3, v_1), (v_1, v_3)$

Loop

- An edge from a vertex to itself.



Example 36


- Which one is a **loop**?

✓ – Walk 1: (v_3, v_3)

- Is there any other loop in this graph?

Cycle

- A walk from a vertex (called **base**) to itself with no repeated edges.
- Recall that: Walk + No repeated edges = **path**
- **Rewording**: A cycle is a **path** from a vertex (called **base**) to itself.




Example 37

- Which one is a **cycle**?
 - ✗ – Walk 1: $(v_1, v_3), (v_3, v_1), (v_1, v_3), (v_3, v_1)$
 - ✓ – Walk 2: $(v_1, v_3), (v_3, v_1)$
 - ✓ – Walk 3: $(v_1, v_3), (v_3, v_3), (v_3, v_1)$



Simple Cycle

- A cycle that **no vertices other than the base** is repeated.
 - Note that the walk starts from the base and ends to the base.
 - During the walk, **the base should not be repeated too.**
 - In other words, in a simple cycle, all vertices (except the base) and the edges are **visited uniquely.**




Example 38

- Which one is a **simple cycle**?
 - ✗ – Walk 1: $(v_1, v_3), (v_3, v_1), (v_1, v_3), (v_3, v_1)$
 - ✓ – Walk 2: $(v_1, v_3), (v_3, v_1)$
 - ✗ – Walk 3: $(v_1, v_3), (v_3, v_3), (v_3, v_1)$


One-Dimensional Projection of a Walk

- One-dimensional projection (or just projection) is another way of representing a walk.



Example 39

- Represent the following walk as a one-dimensional projection.
- Walk from v_1 to v_2 : $(v_1, v_3), (v_3, v_1), (v_1, v_3), (v_3, v_2)$



- The length of this walk (= the number of edges) is clearly shown.

References

1. Linz, Peter, "An Introduction to Formal Languages and Automata, 5th ed.," Jones & Bartlett Learning, LLC, Canada, 2012
2. Kenneth H. Rosen, "Discrete Mathematics and Its Applications, 7th ed.," McGraw Hill, New York, United States, 2012
3. Sipser, Michael, "Introduction to the Theory of Computation, 3rd ed.," CENGAGE Learning, United States, 2013
ISBN-13: 978-1133187790