

Determination of the proton spectral function of ^{12}C from $(e, e'p)$ data. Supplemental Material

Artur M. Ankowski,¹ Omar Benhar,² and Makoto Sakuda³

¹*Institute of Theoretical Physics, University of Wrocław, 50-204 Wrocław, Poland*

²*INFN, Sezione di Roma, I-00185 Roma, Italy*

³*Physics Department, Okayama University, Okayama 700-8530, Japan*

(Dated: November 3, 2024)

The file **pke12_2024.table** provides a table of the proton spectral function of carbon, $P(|\mathbf{k}|, E)$ described in the article, normalised to $Z=6$. The spectral function has been obtained matching the energy region $E < 21.5$ MeV—resulting from a fit to the high-resolution NIKHEF data of Van der Steenhoven *et al.* [1]—to the model of Benhar *et al.* [2], based on a combination of low-resolution Saclay data [3] and accurate nuclear matter calculations [4] at $21.5 < E < 300$ MeV.

The table is structured as follows:

- line 1: number of bins (40) and binwidth (20 MeV) of the momentum grid
- line 2: number of bins and binwidth of the energy grid
 - 340 bins of width 0.025 MeV in the range $13 < E < 21.5$ MeV
 - 2785 bins of width 0.1 MeV in the range $21.5 < E < 300$ MeV
- in the following lines, E [MeV] and $P(|\mathbf{k}|, E)$ [MeV⁻⁴] are given in 40 blocks, one for each value of $|\mathbf{k}|$. Each block consists of 783 lines:
 - the first line of each block gives the value of $|\mathbf{k}|$ in MeV
 - then follow 85 lines containing four pairs of $(E, P(|\mathbf{k}|, E))$ values each, corresponding to $E < 21.5$ MeV, and 697 lines containing the results for $21.5 < E < 300$ MeV.

The total length of the file is 31322 lines.

-
- [1] G. Van der Steenhoven, H. P. Blok, E. Jans, M. De Jong, L. Lapikas, E. N. M. Quint and P. K. A. de Witt Huberts, “Knockout of 1p protons From ^{12}C Induced by the $(e, e'p)$ Reaction”, Nucl. Phys. A **480**, 547-572 (1988). doi:10.1016/0375-9474(88)90463-0.
- [2] O. Benhar, A. Fabrocini, S. Fantoni and I. Sick, “Spectral function of finite nuclei and scattering of GeV electrons”, Nucl. Phys. A **579**, 493-517 (1994). doi:10.1016/0375-9474(94)90920-2.
- [3] J. Mougey, M. Bernheim, A. Bussiere, A. Gillibert, X. H. Phan, M. Priou, D. Royer, I. Sick, and G. J. Wagner, “Quasifree $(e, e'p)$ Scattering on ^{12}C , ^{28}Si , ^{40}Ca and ^{58}Ni ”, Nucl. Phys. A **262**, 461-492 (1976). doi:10.1016/0375-9474(76)90510-8.
- [4] O. Benhar, A. Fabrocini and S. Fantoni, “The Nucleon Spectral Function in Nuclear Matter”, Nucl. Phys. A **505**, 267-299 (1989). doi:10.1016/0375-9474(89)90374-6.