




Figure 1: Effect of the nucleon axial form factor on the MicroBooNE CC1μ1p / CC1μNp argon cross sections with the N/LFG ground state and hA2018 FSI held fixed: F_A^{LQCD} (blue) versus F_A^{Deu} (orange), compared with MicroBooNE data (black points). Panels: (a) δp_T for $\delta\alpha_T < 45^\circ$ and (b) for $135^\circ < \delta\alpha_T < 180^\circ$ (two slices of the 2D δp_T – $\delta\alpha_T$ measurement); (c) δp_T ; (d) $\delta\alpha_T$; (e) p_μ ; (f) p_p . The LQCD form factor raises the predicted cross section and lowers χ^2/ndf in every observable (2D: 35.98/49 vs 44.21/49; p_μ : 16.29/26 vs 20.89/26), favouring F_A^{LQCD} .




Figure 2: As in Fig. 1 but with the spectral-function ground state and INCL FSI held fixed: F_A^{LQCD} (blue) versus F_A^{Deu} (orange). The LQCD form factor again improves every observable (2D: 66.61/49 vs 81.91/49; p_μ : 23.91/26 vs 35.08/26), but with INCL FSI both predictions remain well below the data (p_p : 32.25/15 and 38.83/15), so varying F_A alone does not reconcile the SF+INCL configuration with MicroBooNE.