



中国科学院高能物理研究所
Institute of High Energy Physics, Chinese Academy of Sciences




IHEP SUSY Group Meeting

Chengxin Liao

**Institute of High Energy Physics
Chinese Academy of Sciences**

Jul 16, 2025





- FF already cross-check with wenyi and result can be matched
- BDT training for run2 already finished and apply the model to run2 sample and run3 sample
- Update LCG version and test code(Ongoing)
- Update support-note(Ongoing)

Fake Factor for Run2 and Run3



Selection:

nBaseTau == 1

nBaseLep >= 1, SigLep >= 1

MET trigger, MET >= 200


Same-Signal(Orthogonal with SR)

bVeto


ID: nMediumTau == 1

antiID: nMediumTau < 1


$0 < |\eta| < 1$






$1 < |\eta| < 1.37$



$1.52 < |\eta| < 2.5$



A small bump show in the last bin






HH Pre-selection	LH Pre-selection
≥ 2 medium taus	≥ 1 medium taus
0 base lepton	1 base lepton, 1 signal lepton
$\text{MET} \geq 200$; pass MET trigger	$\text{MET} \geq 200$; pass MET trigger
$1 \leq n_{\text{Jet}}$	$1 \leq n_{\text{Jet}}$
Opposite-sign hadronic-hadronic tau pair	Opposite-sign lepton-hadronic tau pair
bveto	bveto
jet $p_{\text{T}} > 100$ GeV	jet $p_{\text{T}} > 100$ GeV
$\text{M}_{\text{tt_reco}} \leq 40$ GeV $\text{M}_{\text{tt_reco}} \geq 130$ GeV	$\text{M}_{\text{tt_reco}} \leq 40$ GeV $\text{M}_{\text{tt_reco}} \geq 130$ GeV


MC modeling in Pre-Selection(HH)



run2




run3




MC modeling in Pre-Selection(HH)

run2






run3






MC modeling in Pre-Selection(HH)

run2



run3




MC modeling in Pre-Selection(LH)




Institute of High Energy Physics
Chinese Academy of Sciences

run2




run3




MC modeling in Pre-Selection(LH)

run2






run3






MC modeling in Pre-Selection(LH)

run2



run3





Input sample:

bkg: run2 bkg sample passed pre-selection(HH/LH)

sig: 100_70, 120_90, 140_90(only run2)

Hyperparameters:

HH: Ntrees = 300, MaxDepth = 6, MinNodeSize = 1%, Learning rate = 0.05



LH: Ntrees = 200, MaxDepth = 6, MinNodeSize = 1%, Learning rate = 0.05

Weight choose: `abs(physics weight)`



Split strategy: Separate entries by using mod 5, for Fake bkg, if separate follow sequence, all weighted entry will split into first fold

BDT distribution for LH and HH

run2



run3






中国科学院高能物理研究所
Institute of High Energy Physics, Chinese Academy of Sciences




Backup




$$0 < |\eta| < 1$$

FF_1prong_eta0




$$1 < |\eta| < 1.37$$

FF_1prong_eta1




$$1.52 < |\eta| < 2.5$$




My result

Wenyi's result


FF_1prong_eta0



FF_1prong_eta1



FF_1prong_eta2



Same value for first two bins and different in last bin for different rebin strategy

I check FF with same rebin method in case, it turns out we are the same