
DIO2 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000211448 | Nominal screening candidate



Conditional PTN–DIO2 association: b = +0.273 per mediator SD, p = 6.22e-13, BH FDR = 2.7e-10 | Dx attenuation = 37.1%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

IGSF8 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000162729 | Nominal screening candidate



Conditional PTN–IGSF8 association: b = -0.161 per mediator SD, p = 1.61e-10, BH FDR = 2.88e-08 | Dx attenuation = 32.5%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

KCNJ16 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000153822 | Nominal screening candidate



Conditional PTN–KCNJ16 association: b = +0.357 per mediator SD, p = 6.38e-07, BH FDR = 1.92e-05 | Dx attenuation = 24.7%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

PHYHIP [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000168490 | Nominal screening candidate



Conditional PTN–PHYHIP association: b = -0.108 per mediator SD, p = 2.45e-06, BH FDR = 5.43e-05 | Dx attenuation = 21.5%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

CCND3 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000112576 | Nominal screening candidate



Conditional PTN–CCND3 association: b = -0.100 per mediator SD, p = 3.32e-06, BH FDR = 6.9e-05 | Dx attenuation = 21.3%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

ARAP2 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000047365 | Nominal screening candidate



Conditional PTN–ARAP2 association: b = +0.135 per mediator SD, p = 9.67e-06, BH FDR = 0.00016 | Dx attenuation = 23.3%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

TCEAL4 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000133142 | Nominal screening candidate



Conditional PTN–TCEAL4 association: b = +0.090 per mediator SD, p = 1.15e-05, BH FDR = 0.000185 | Dx attenuation = 23.7%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

CAMK1G [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000008118 | Nominal screening candidate



Conditional PTN–CAMK1G association: b = -0.167 per mediator SD, p = 2.11e-05, BH FDR = 0.000301 | Dx attenuation = 23.0%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

EPRS [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000136628 | Nominal screening candidate



Conditional PTN–EPRS association: b = +0.094 per mediator SD, p = 9.33e-05, BH FDR = 0.000958 | Dx attenuation = 20.3%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

HSPD1 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000144381 | Nominal screening candidate



Conditional PTN–HSPD1 association: b = +0.077 per mediator SD, p = 0.000134, BH FDR = 0.00129 | Dx attenuation = 17.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

MAPKAP1 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000119487 | Nominal screening candidate



Conditional PTN–MAPKAP1 association: b = +0.090 per mediator SD, p = 0.000135, BH FDR = 0.0013 | Dx attenuation = 20.9%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

SLC9A6 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000198689 | Nominal screening candidate



Conditional PTN–SLC9A6 association: b = -0.087 per mediator SD, p = 0.000395, BH FDR = 0.00311 | Dx attenuation = 17.7%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

CHURC1 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000258289 | Nominal screening candidate



Conditional PTN–CHURC1 association: b = +0.087 per mediator SD, p = 0.001, BH FDR = 0.00646 | Dx attenuation = 15.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

AC240274.1 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000271254 | Nominal screening candidate



Conditional PTN–AC240274.1 association: b = +0.251 per mediator SD, p = 0.00108, BH FDR = 0.00687 | Dx attenuation = 17.5%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

ASB6 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000148331 | Nominal screening candidate



Conditional PTN–ASB6 association: b = -0.139 per mediator SD, p = 0.0011, BH FDR = 0.00696 | Dx attenuation = 17.6%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

NKAIN4 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000101198 | Nominal screening candidate



Conditional PTN–NKAIN4 association: b = +0.182 per mediator SD, p = 0.00113, BH FDR = 0.00715 | Dx attenuation = 16.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

CEP63 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000182923 | Nominal screening candidate



Conditional PTN–CEP63 association: b = +0.112 per mediator SD, p = 0.00121, BH FDR = 0.00753 | Dx attenuation = 17.1%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

AVL9 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000105778 | Nominal screening candidate



Conditional PTN–AVL9 association: b = -0.085 per mediator SD, p = 0.00126, BH FDR = 0.00779 | Dx attenuation = 17.0%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

CD2BP2 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000169217 | Nominal screening candidate



Conditional PTN–CD2BP2 association: b = -0.087 per mediator SD, p = 0.00166, BH FDR = 0.00967 | Dx attenuation = 16.6%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

RLIM [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000131263 | Nominal screening candidate



Conditional PTN–RLIM association: b = -0.066 per mediator SD, p = 0.0018, BH FDR = 0.0103 | Dx attenuation = 15.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

TNFSF10 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000121858 | Nominal screening candidate



Conditional PTN–TNFSF10 association: b = +0.262 per mediator SD, p = 0.00292, BH FDR = 0.015 | Dx attenuation = 16.3%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

PPP1R21 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000162869 | Nominal screening candidate



Conditional PTN–PPP1R21 association: b = +0.128 per mediator SD, p = 0.00324, BH FDR = 0.0162 | Dx attenuation = 14.6%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

SERF2 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000140264 | Nominal screening candidate



Conditional PTN–SERF2 association: b = +0.073 per mediator SD, p = 0.00573, BH FDR = 0.0253 | Dx attenuation = 15.3%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

ZNHIT1 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000106400 | Nominal screening candidate



Conditional PTN–ZNHIT1 association: b = +0.063 per mediator SD, p = 0.00585, BH FDR = 0.0257 | Dx attenuation = 15.0%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

HIP1 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000127946 | Nominal screening candidate



Conditional PTN–HIP1 association: b = +0.159 per mediator SD, p = 0.00626, BH FDR = 0.0271 | Dx attenuation = 13.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

GUCY1A1 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000164116 | Nominal screening candidate



Conditional PTN–GUCY1A1 association: b = +0.115 per mediator SD, p = 0.00679, BH FDR = 0.0289 | Dx attenuation = 14.0%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

CCDC167 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000198937 | Nominal screening candidate



Conditional PTN–CCDC167 association: b = +0.138 per mediator SD, p = 0.00721, BH FDR = 0.0301 | Dx attenuation = 13.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

TIMP2 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000035862 | Nominal screening candidate



Conditional PTN–TIMP2 association: b = -0.070 per mediator SD, p = 0.00816, BH FDR = 0.0332 | Dx attenuation = 13.2%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

SLC7A8 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000092068 | Nominal screening candidate



Conditional PTN–SLC7A8 association: b = +0.123 per mediator SD, p = 0.0095, BH FDR = 0.0373 | Dx attenuation = 12.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

ZSCAN18 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000121413 | Nominal screening candidate



Conditional PTN–ZSCAN18 association: b = -0.067 per mediator SD, p = 0.0115, BH FDR = 0.0431 | Dx attenuation = 13.4%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

NUTM2A-AS1 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000223482 | Nominal screening candidate



Conditional PTN–NUTM2A-AS1 association: b = +0.127 per mediator SD, p = 0.0167, BH FDR = 0.0576 | Dx attenuation = 12.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

CNNM2 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000148842 | Nominal screening candidate



Conditional PTN–CNNM2 association: b = -0.081 per mediator SD, p = 0.0176, BH FDR = 0.0598 | Dx attenuation = 12.7%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

IGSF9B [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000080854 | Nominal screening candidate



Conditional PTN–IGSF9B association: b = -0.091 per mediator SD, p = 0.0283, BH FDR = 0.0851 | Dx attenuation = 12.2%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

MBP [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000197971 | Nominal screening candidate



Conditional PTN–MBP association: b = -0.072 per mediator SD, p = 0.0313, BH FDR = 0.0922 | Dx attenuation = 11.9%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

RPL35A [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000182899 | Nominal screening candidate



Conditional PTN–RPL35A association: b = +0.067 per mediator SD, p = 0.0314, BH FDR = 0.0923 | Dx attenuation = 11.0%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

SLC25A29 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000197119 | Nominal screening candidate



Conditional PTN–SLC25A29 association: b = -0.075 per mediator SD, p = 0.0356, BH FDR = 0.102 | Dx attenuation = 11.7%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

CABIN1 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000099991 | Nominal screening candidate



Conditional PTN–CABIN1 association: b = -0.061 per mediator SD, p = 0.0367, BH FDR = 0.104 | Dx attenuation = 11.5%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

CYFIP1 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000273749 | Nominal screening candidate



Conditional PTN–CYFIP1 association: b = +0.067 per mediator SD, p = 0.0427, BH FDR = 0.116 | Dx attenuation = 10.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

ZNF48 [neuropil] | PTN [vascular]

Donor means across available matched SpDs · 61 donors · 272 observations

ptn_vasc_neuropil | ENSG00000180035 | Nominal screening candidate



Conditional PTN–ZNF48 association: b = -0.081 per mediator SD, p = 0.0486, BH FDR = 0.128 | Dx attenuation = 10.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

CAMK1G [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000008118 | Nominal screening candidate



Conditional PTN–CAMK1G association: b = -0.200 per mediator SD, p = 1.06e-08, BH FDR = 3.62e-06 | Dx attenuation = 16.2%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

BOD1L1 [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000038219 | Nominal screening candidate



Conditional PTN–BOD1L1 association: b = +0.167 per mediator SD, p = 1.35e-06, BH FDR = 0.000137 | Dx attenuation = 13.9%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

PDP1 [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000164951 | Nominal screening candidate



Conditional PTN–PDP1 association: b = -0.124 per mediator SD, p = 4.14e-06, BH FDR = 0.000327 | Dx attenuation = 13.3%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

CCDC102B [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000150636 | Nominal screening candidate



Conditional PTN–CCDC102B association: b = +0.294 per mediator SD, p = 2.57e-05, BH FDR = 0.00134 | Dx attenuation = 11.5%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

LY6H [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000176956 | Nominal screening candidate



Conditional PTN–LY6H association: b = -0.121 per mediator SD, p = 0.000137, BH FDR = 0.00472 | Dx attenuation = 10.6%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

SMARCA2 [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000080503 | Nominal screening candidate



Conditional PTN–SMARCA2 association: b = +0.089 per mediator SD, p = 0.000368, BH FDR = 0.00958 | Dx attenuation = 10.0%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

CSPP1 [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000104218 | Nominal screening candidate



Conditional PTN–CSPP1 association: b = +0.170 per mediator SD, p = 0.000858, BH FDR = 0.0175 | Dx attenuation = 9.5%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

SPTY2D1OS [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000247595 | Nominal screening candidate



Conditional PTN–SPTY2D1OS association: b = +0.227 per mediator SD, p = 0.00125, BH FDR = 0.0223 | Dx attenuation = 9.1%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

AC016687.2 [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000250723 | Nominal screening candidate



Conditional PTN–AC016687.2 association: b = +0.222 per mediator SD, p = 0.00238, BH FDR = 0.0348 | Dx attenuation = 8.7%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

STMN3 [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000197457 | Nominal screening candidate



Conditional PTN–STMN3 association: b = -0.074 per mediator SD, p = 0.00324, BH FDR = 0.0431 | Dx attenuation = 8.6%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

SRGAP2B [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000196369 | Nominal screening candidate



Conditional PTN–SRGAP2B association: b = +0.189 per mediator SD, p = 0.0034, BH FDR = 0.0444 | Dx attenuation = 8.3%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

DHX36 [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000174953 | Nominal screening candidate



Conditional PTN–DHX36 association: b = +0.086 per mediator SD, p = 0.00429, BH FDR = 0.0514 | Dx attenuation = 8.2%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

RABEP1 [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000029725 | Nominal screening candidate



Conditional PTN–RABEP1 association: b = +0.067 per mediator SD, p = 0.00517, BH FDR = 0.0574 | Dx attenuation = 8.3%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

TGS1 [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000137574 | Nominal screening candidate



Conditional PTN–TGS1 association: b = +0.184 per mediator SD, p = 0.00692, BH FDR = 0.0688 | Dx attenuation = 7.6%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

NAIP [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000249437 | Nominal screening candidate



Conditional PTN–NAIP association: b = +0.237 per mediator SD, p = 0.00725, BH FDR = 0.0707 | Dx attenuation = 7.7%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

KIF2A [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000068796 | Nominal screening candidate



Conditional PTN–KIF2A association: b = +0.149 per mediator SD, p = 0.00976, BH FDR = 0.0857 | Dx attenuation = 7.7%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

NOG [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000183691 | Nominal screening candidate



Conditional PTN–NOG association: b = +0.186 per mediator SD, p = 0.0148, BH FDR = 0.113 | Dx attenuation = 7.2%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

SERBP1 [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000142864 | Nominal screening candidate



Conditional PTN–SERBP1 association: b = +0.057 per mediator SD, p = 0.0159, BH FDR = 0.118 | Dx attenuation = 7.1%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

ZNHIT1 [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000106400 | Nominal screening candidate



Conditional PTN–ZNHIT1 association: b = +0.063 per mediator SD, p = 0.0187, BH FDR = 0.131 | Dx attenuation = 6.9%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

POU3F4 [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000196767 | Nominal screening candidate



Conditional PTN–POU3F4 association: b = +0.174 per mediator SD, p = 0.0242, BH FDR = 0.153 | Dx attenuation = 6.6%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

MATN1 [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000162510 | Nominal screening candidate



Conditional PTN–MATN1 association: b = -0.094 per mediator SD, p = 0.0395, BH FDR = 0.204 | Dx attenuation = 5.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

AC073534.2 [neuronal] | PTN [vascular]

Donor means across available matched SpDs · 58 donors · 220 observations

ptn_vasc_neun | ENSG00000276030 | Nominal screening candidate



Conditional PTN–AC073534.2 association: b = -0.135 per mediator SD, p = 0.0427, BH FDR = 0.214 | Dx attenuation = 6.0%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

AGT [vascular] | FGF1 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf1_neuropil_vasc | ENSG00000135744 | Nominal screening candidate



Conditional FGF1–AGT association: b = +0.430 per mediator SD, p = 2.31e-06, BH FDR = 0.000256 | Dx attenuation = 20.7%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

MT3 [vascular] | FGF1 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf1_neuropil_vasc | ENSG00000087250 | Nominal screening candidate



Conditional FGF1–MT3 association: b = +0.182 per mediator SD, p = 2.91e-06, BH FDR = 0.000283 | Dx attenuation = 19.6%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

FAM107A [vascular] | FGF1 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf1_neuropil_vasc | ENSG00000168309 | Nominal screening candidate



Conditional FGF1–FAM107A association: b = +0.212 per mediator SD, p = 3.09e-06, BH FDR = 0.000283 | Dx attenuation = 23.1%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

CHGB [vascular] | FGF1 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf1_neuropil_vasc | ENSG00000089199 | Nominal screening candidate



Conditional FGF1–CHGB association: b = -0.205 per mediator SD, p = 0.000246, BH FDR = 0.00856 | Dx attenuation = 16.3%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

SLC6A8 [vascular] | FGF1 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf1_neuropil_vasc | ENSG00000130821 | Nominal screening candidate



Conditional FGF1–SLC6A8 association: b = +0.218 per mediator SD, p = 0.000471, BH FDR = 0.0133 | Dx attenuation = 17.1%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

HINT1 [vascular] | FGF1 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf1_neuropil_vasc | ENSG00000169567 | Nominal screening candidate



Conditional FGF1–HINT1 association: b = -0.170 per mediator SD, p = 0.000542, BH FDR = 0.0147 | Dx attenuation = 16.5%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

MT1G [vascular] | FGF1 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf1_neuropil_vasc | ENSG00000125144 | Nominal screening candidate



Conditional FGF1–MT1G association: b = +0.298 per mediator SD, p = 0.000656, BH FDR = 0.0166 | Dx attenuation = 15.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

SLC24A2 [vascular] | FGF1 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf1_neuropil_vasc | ENSG00000155886 | Nominal screening candidate



Conditional FGF1–SLC24A2 association: b = -0.182 per mediator SD, p = 0.00184, BH FDR = 0.0323 | Dx attenuation = 14.6%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

BCL6 [vascular] | FGF1 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf1_neuropil_vasc | ENSG00000113916 | Nominal screening candidate



Conditional FGF1–BCL6 association: b = +0.268 per mediator SD, p = 0.0036, BH FDR = 0.0495 | Dx attenuation = 14.2%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

BCAP29 [vascular] | FGF1 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf1_neuropil_vasc | ENSG00000075790 | Nominal screening candidate



Conditional FGF1–BCAP29 association: b = -0.120 per mediator SD, p = 0.00513, BH FDR = 0.0631 | Dx attenuation = 13.2%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

GAP43 [vascular] | FGF1 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf1_neuropil_vasc | ENSG00000172020 | Nominal screening candidate



Conditional FGF1–GAP43 association: b = -0.084 per mediator SD, p = 0.00571, BH FDR = 0.0668 | Dx attenuation = 13.6%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

DYNLL1 [vascular] | FGF1 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf1_neuropil_vasc | ENSG00000088986 | Nominal screening candidate



Conditional FGF1–DYNLL1 association: b = -0.109 per mediator SD, p = 0.00787, BH FDR = 0.0822 | Dx attenuation = 12.0%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

MTRNR2L8 [vascular] | FGF1 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf1_neuropil_vasc | ENSG00000255823 | Nominal screening candidate



Conditional FGF1–MTRNR2L8 association: b = +0.227 per mediator SD, p = 0.0113, BH FDR = 0.104 | Dx attenuation = 12.7%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

PITPNC1 [vascular] | FGF1 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf1_neuropil_vasc | ENSG00000154217 | Nominal screening candidate



Conditional FGF1–PITPNC1 association: b = +0.177 per mediator SD, p = 0.027, BH FDR = 0.17 | Dx attenuation = 10.3%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

CHIC1 [vascular] | FGF1 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf1_neuropil_vasc | ENSG00000204116 | Nominal screening candidate



Conditional FGF1–CHIC1 association: b = -0.188 per mediator SD, p = 0.0271, BH FDR = 0.171 | Dx attenuation = 10.4%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

BRSK1 [vascular] | FGF1 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf1_neuropil_vasc | ENSG00000160469 | Nominal screening candidate



Conditional FGF1–BRSK1 association: b = +0.139 per mediator SD, p = 0.0279, BH FDR = 0.174 | Dx attenuation = 11.0%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

C2orf80 [vascular] | FGF1 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf1_neuropil_vasc | ENSG00000188674 | Nominal screening candidate



Conditional FGF1–C2orf80 association: b = -0.155 per mediator SD, p = 0.0468, BH FDR = 0.235 | Dx attenuation = 9.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

AGT [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000135744 | Nominal screening candidate



Conditional FGF2–AGT association: b = +0.662 per mediator SD, p = 1.26e-21, BH FDR = 6.35e-19 | Dx attenuation = 37.1%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

MT1M [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000205364 | Nominal screening candidate



Conditional FGF2–MT1M association: b = +0.594 per mediator SD, p = 4.13e-19, BH FDR = 1.2e-16 | Dx attenuation = 31.5%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

MT3 [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000087250 | Nominal screening candidate



Conditional FGF2–MT3 association: b = +0.229 per mediator SD, p = 5.9e-14, BH FDR = 7.15e-12 | Dx attenuation = 28.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

FAM107A [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000168309 | Nominal screening candidate



Conditional FGF2–FAM107A association: b = +0.256 per mediator SD, p = 8.04e-13, BH FDR = 8.74e-11 | Dx attenuation = 32.6%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

HINT1 [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000169567 | Nominal screening candidate



Conditional FGF2–HINT1 association: b = -0.273 per mediator SD, p = 1.06e-12, BH FDR = 1.11e-10 | Dx attenuation = 31.0%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

SLC14A1 [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000141469 | Nominal screening candidate



Conditional FGF2–SLC14A1 association: b = +0.520 per mediator SD, p = 9.99e-09, BH FDR = 4.91e-07 | Dx attenuation = 23.7%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

CHGB [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000089199 | Nominal screening candidate



Conditional FGF2–CHGB association: b = -0.243 per mediator SD, p = 5.65e-08, BH FDR = 2.34e-06 | Dx attenuation = 22.5%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

MT1G [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000125144 | Nominal screening candidate



Conditional FGF2–MT1G association: b = +0.358 per mediator SD, p = 3.21e-07, BH FDR = 1.02e-05 | Dx attenuation = 22.1%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

DYNLL1 [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000088986 | Nominal screening candidate



Conditional FGF2–DYNLL1 association: b = -0.153 per mediator SD, p = 2.73e-06, BH FDR = 6.93e-05 | Dx attenuation = 19.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

ATP5PO [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000241837 | Nominal screening candidate



Conditional FGF2–ATP5PO association: b = -0.161 per mediator SD, p = 4.2e-06, BH FDR = 0.000101 | Dx attenuation = 18.7%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

MTRNR2L8 [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000255823 | Nominal screening candidate



Conditional FGF2–MTRNR2L8 association: b = +0.309 per mediator SD, p = 1.61e-05, BH FDR = 0.000307 | Dx attenuation = 20.1%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

PITPNC1 [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000154217 | Nominal screening candidate



Conditional FGF2–PITPNC1 association: b = +0.274 per mediator SD, p = 1.88e-05, BH FDR = 0.000348 | Dx attenuation = 18.6%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

BCL6 [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000113916 | Nominal screening candidate



Conditional FGF2–BCL6 association: b = +0.314 per mediator SD, p = 2.03e-05, BH FDR = 0.00037 | Dx attenuation = 19.4%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

GAP43 [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000172020 | Nominal screening candidate



Conditional FGF2–GAP43 association: b = -0.099 per mediator SD, p = 4.3e-05, BH FDR = 0.000712 | Dx attenuation = 18.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

C12orf57 [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000111678 | Nominal screening candidate



Conditional FGF2–C12orf57 association: b = -0.157 per mediator SD, p = 5.01e-05, BH FDR = 0.000821 | Dx attenuation = 17.1%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

SOD2 [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000112096 | Nominal screening candidate



Conditional FGF2–SOD2 association: b = +0.115 per mediator SD, p = 9.7e-05, BH FDR = 0.0014 | Dx attenuation = 17.0%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

QDPR [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000151552 | Nominal screening candidate



Conditional FGF2–QDPR association: b = -0.139 per mediator SD, p = 0.000226, BH FDR = 0.00292 | Dx attenuation = 16.9%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

SLIRP [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000119705 | Nominal screening candidate



Conditional FGF2–SLIRP association: b = -0.144 per mediator SD, p = 0.000348, BH FDR = 0.00416 | Dx attenuation = 16.9%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

NDUFA8 [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000119421 | Nominal screening candidate



Conditional FGF2–NDUFA8 association: b = -0.145 per mediator SD, p = 0.00149, BH FDR = 0.0131 | Dx attenuation = 15.0%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

BCAP29 [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000075790 | Nominal screening candidate



Conditional FGF2–BCAP29 association: b = -0.108 per mediator SD, p = 0.00183, BH FDR = 0.0153 | Dx attenuation = 13.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

SLC24A2 [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000155886 | Nominal screening candidate



Conditional FGF2–SLC24A2 association: b = -0.147 per mediator SD, p = 0.00193, BH FDR = 0.016 | Dx attenuation = 13.7%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

SLC6A8 [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000130821 | Nominal screening candidate



Conditional FGF2–SLC6A8 association: b = +0.153 per mediator SD, p = 0.00247, BH FDR = 0.0195 | Dx attenuation = 14.0%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

PSMA7 [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000101182 | Nominal screening candidate



Conditional FGF2–PSMA7 association: b = -0.108 per mediator SD, p = 0.0104, BH FDR = 0.0588 | Dx attenuation = 10.9%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

C2orf80 [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000188674 | Nominal screening candidate



Conditional FGF2–C2orf80 association: b = -0.155 per mediator SD, p = 0.014, BH FDR = 0.0717 | Dx attenuation = 11.4%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

BRSK1 [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000160469 | Nominal screening candidate



Conditional FGF2–BRSK1 association: b = +0.119 per mediator SD, p = 0.0193, BH FDR = 0.0901 | Dx attenuation = 11.0%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

PLEC [vascular] | FGF2 [neuropil]

Donor means across available matched SpDs · 61 donors · 272 observations

fgf2_neuropil_vasc | ENSG00000178209 | Nominal screening candidate



Conditional FGF2–PLEC association: b = +0.118 per mediator SD, p = 0.0198, BH FDR = 0.0915 | Dx attenuation = 10.2%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

FAM107A [vascular] | FGF1 [neuronal]

Donor means across available matched SpDs · 58 donors · 220 observations

fgf1_neun_vasc | ENSG00000168309 | Nominal screening candidate



Conditional FGF1–FAM107A association: b = +0.144 per mediator SD, p = 5.93e-05, BH FDR = 0.0182 | Dx attenuation = 13.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

RGS4 [vascular] | FGF1 [neuronal]

Donor means across available matched SpDs · 58 donors · 220 observations

fgf1_neun_vasc | ENSG00000117152 | Nominal screening candidate



Conditional FGF1–RGS4 association: b = -0.095 per mediator SD, p = 0.000136, BH FDR = 0.0257 | Dx attenuation = 14.1%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

SLC24A2 [vascular] | FGF1 [neuronal]

Donor means across available matched SpDs · 58 donors · 220 observations

fgf1_neun_vasc | ENSG00000155886 | Nominal screening candidate



Conditional FGF1–SLC24A2 association: b = -0.130 per mediator SD, p = 0.000328, BH FDR = 0.0335 | Dx attenuation = 12.9%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

APOLD1 [vascular] | FGF1 [neuronal]

Donor means across available matched SpDs · 58 donors · 220 observations

fgf1_neun_vasc | ENSG00000178878 | Nominal screening candidate



Conditional FGF1–APOLD1 association: b = +0.231 per mediator SD, p = 0.0076, BH FDR = 0.143 | Dx attenuation = 10.0%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

ADCYAP1R1 [vascular] | FGF1 [neuronal]

Donor means across available matched SpDs · 58 donors · 220 observations

fgf1_neun_vasc | ENSG00000078549 | Nominal screening candidate



Conditional FGF1–ADCYAP1R1 association: b = +0.127 per mediator SD, p = 0.0177, BH FDR = 0.209 | Dx attenuation = 8.8%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.


Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.

IFITM2 [vascular] | FGF1 [neuronal]

Donor means across available matched SpDs · 58 donors · 220 observations

fgf1_neun_vasc | ENSG00000185201 | Nominal screening candidate



Conditional FGF1–IFITM2 association: b = +0.186 per mediator SD, p = 0.0352, BH FDR = 0.294 | Dx attenuation = 8.0%

Values remove centered age, sex and SpD effects; middle panel also removes the centered mediator effect.

ΔDx = SCZ − NTC. P-values/FDR are from the original donor-blocked limma fits; boxes supply no new statistical tests.

Each point averages adjusted values across the available SpDs of one donor; this is a descriptive display, not a refit.

Exploratory association: significance loss after adjustment does not establish causal or complete mediation.